TITLE: Postpartum Weight Retention and Mortality Over 50 Years of Follow-up

AUTHORS: Yajnaseni Chakraborti[1], Sunni L. Mumford[1,2], Edwina H. Yeung[3], Katherine L. Grantz[3], Pauline Mendola[4], James L. Mills[3], Ellen C. Caniglia[1], Colleen M. Brensinger[5], Cuilin Zhang[6,7,8,9], Enrique F. Schisterman[1,2], Stefanie N. Hinkle[1,2]

AFFILIATIONS:

[1] Department of Biostatistics, Epidemiology and Informatics, Perelman School of Medicine, University of Pennsylvania

[2] Department of Obstetrics and Gynecology, Perelman School of Medicine, University of Pennsylvania

[3] Epidemiology Branch, Division of Population Health Research, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health

[4] Department of Epidemiology and Environmental Health, School of Public Health and Health Professions, University of Buffalo

[5] Biostatistics Analysis Center, Center for Clinical Epidemiology and Biostatistics, Perelman School of Medicine, University of Pennsylvania

[6] Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA

[7] Global Centre for Asian Women's Health, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Republic of Singapore

[8] Department of Obstetrics and Gynecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Republic of Singapore

[9] Bia-Echo Asia Centre for Reproductive Longevity & Equality, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Republic of Singapore

CORRESPONDING AUTHOR: Stefanie N. Hinkle, Department of Biostatistics, Epidemiology and Informatics and Department of Obstetrics and Gynecology, Perelman School of Medicine, University of Pennsylvania.

Email: Stefanie.Hinkle@Pennmedicine.upenn.edu

ABSTRACT

Objective: This study aimed to evaluate the link between postpartum weight retention (PPWR) and long-term mortality.

Methods: In this secondary analysis of 8165 women with ≥2 pregnancies in the Collaborative Perinatal Project (U.S., 1959-1966) with vital status follow-up through 2016, we used interpregnancy weight change (IPWC) and postpartum weight change (PPWC) as proxies of PPWR. IPWC was defined as the difference between self-reported pre-pregnancy weights from the 1st and 2nd CPP pregnancies, while PPWC was the difference between weight recorded at delivery of the 1st and pre-pregnancy weight of the 2nd CPP pregnancies. All-cause and cause-specific mortality models were adjusted for sociodemographic, behavioral, clinical, and pregnancy-related characteristics from the 1st CPP pregnancy.

Results: Compared to women with IPWC >0 to 1.8 kg (quintile 3), those with IPWC > –1.4 to 0 kg (quintile 2) had a lower risk of all-cause mortality [aHR (95% CI): 0.85 (0.76–0.96)], with the most notable reduction observed in diabetes-related risk of mortality [aHR (95% CI): 0.45 (0.23-0.87)]. Lower quintiles of PPWC (i.e., greater weight loss) were linked to reduced all-cause and cause-specific risk of mortality, though estimates were imprecise.

Conclusions: Modest PPWR was linked with increased mortality risk over 50 years of follow-up.

Introduction

Over the past three decades, the prevalence of overweight and obesity among women of reproductive age has steadily increased(1–3). Currently, more than one in four women between the ages of 20 and 39 in the U.S. are living with overweight (27.5%) or obesity (36.8%)(1, 2, 4). For many women, weight gained during pregnancy may contribute to the development of overweight or obesity(5). In fact, the 5- and 10-year incidence of obesity is twice as high among women who have had a child compared to those who have never given birth(6, 7). A probable mechanism underlying this incremented risk of obesity associated with parity is postpartum weight retention (PPWR), which refers to the weight a person retains after giving birth, in addition to the weight gained during pregnancy.

Only about one in four women returns to their pre-pregnancy weight by 1-year postpartum(8). For instance, a U.S.-based prospective cohort study by Endres et al. (2015), found that approximately 75% of participants weighed more one year after giving birth than they did before pregnancy—at 1-year postpartum, 47.4% retained more than 10 pounds and 24.2% retained more than 20 pounds. Further, among those with a normal pre-pregnancy body mass index (BMI), 33% were living with overweight or obesity at 1-year postpartum(8). Legro et al. (2020) reported similar findings in their study of first-time mothers in Pennsylvania, where 23.7% retained between 1 and 9 pounds, and 23.9% retained 10 or more pounds at 1-year postpartum. Among those with a normal pre-pregnancy BMI, 12% were living with overweight or obesity at 1-year postpartum(9).

Indeed, gestational weight gain (GWG) is a likely contributor to PPWR(5, 10–12), with studies showing associations between excessive GWG and greater overall and abdominal adiposity even 8 to 15 years postpartum(13). However, GWG is not the only factor influencing PPWR; other contributors, such as lifestyle behaviors and breastfeeding, may also affect whether weight is retained postpartum(12, 14–21). Given that multiple factors contribute to PPWR, it is important to examine its impact on health as a distinct exposure, rather than viewing it solely as a consequence of GWG(10). In fact, one study found that post-partum weight trajectories were associated with

excess weight and increased adiposity risk at 3 years postpartum, even after adjusting for GWG(12).

Additionally, the implications of PPWR on cardiometabolic health are well documented with adverse cardiometabolic profiles often emerging within the first year postpartum(22), and elevated risks of pre-diabetes or diabetes(23), poor cardiometabolic profiles(12, 23), as well as hypertensive and cardiovascular disorders(24) continuing to manifest at 3-(12, 23), 5-(23), and 16-years(24) postpartum.

Given the evidence of sustained alterations in weight and cardiometabolic parameters, it is reasonable to hypothesize that PPWR may be related to long-term mortality, potentially independent of pre-pregnancy BMI and gestational weight gain, yet this remains unexplored. To evaluate this hypothesis, we estimated the risk of all-cause mortality over 50 years of follow-up post-pregnancy, in relation to two exposures: 1) interpregnancy weight change, and 2) post-partum weight change. Additionally, given the existing evidence linking PPWR to cardiovascular and metabolic disorders(25, 26), we also examined the cardiovascular, diabetes, and kidney-specific mortality risks associated with these two exposures.

Methods

This study is a secondary analysis of data collected from the Collaborative Perinatal Project (CPP) and the CPP Mortality Linkage Study. The CPP was a multi-center observational prospective cohort study conducted in the U.S. between 1959 and 1966 to assess the effects of pregnancy and perinatal complications on birth and child outcomes(27–32). A total of 48,917 pregnant women were enrolled at their first prenatal visit, and were followed across multiple pregnancies, resulting in 58,760 registered pregnancies. The mortality status of this cohort as of Dec 31, 2016, was later determined through the CPP Mortality Linkage study(33), to explore the long-term implications of gestational exposures. This analysis utilizes data from the first two pregnancies for participants with two or more CPP pregnancies (n = 8165). It is important to note that the 1st pregnancy registered within the CPP was not necessarily an individual's 1st pregnancy

overall. For the purposes of this study, we have used the term "1st pregnancy" to refer specifically to the 1st pregnancy recorded in the CPP.

There were no specific regulations for approving research involving human subjects at the time the CPP began in the late 1950s, yet general informed consent for participation was obtained. IRB approval for the CPP Mortality Linkage Study was obtained by the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (IRB00011862, approved in 2015) and the Emmes Corporation (IRB00000879, approved in 2013)—the two organizations responsible for abstracting identifying information from historic CPP records and facilitating the data linkages to the National Death Index (NDI) and the Social Security Death Master File (SSDMF). Mortality data were linked through December 31, 2016. The University of Pennsylvania IRB entered into a reliance agreement with the NICHD IRB for oversight of this study.

Exposures

Participants self-reported their pre-pregnancy weight at their first prenatal visit for each CPP pregnancy. Participants' weight at each prenatal visit and delivery admission was abstracted from medical records. This analysis conceptualized PPWR using two different weight exposures based on the participants' consecutive 1st and 2nd CPP pregnancies (Figure 1). The primary exposure was interpregnancy weight change (IPWC), which represented the overall weight change between the 1st and 2nd CPP pregnancies. Specifically, IPWC was calculated as the difference between pre-pregnancy weight of the 2nd and 1st pregnancies. We also examined a secondary exposure, postpartum weight change (PPWC), which focused solely on postpartum weight change in the interpregnancy interval (IPI). Specifically, PPWC was calculated as the difference between pre-pregnancy weight of the 2nd CPP pregnancy and weight recorded at admission for delivery of the 1st CPP pregnancy. We preprocessed height and weight measurements by identifying and setting implausible values to missing based on predefined thresholds (detailed specifications in Appendix B).

Outcomes

The primary outcome of interest was time to all-cause mortality, defined as the time from the start of follow up, to the date of death or Dec 31, 2016, whichever occurred first. Follow up started at the date of the last menstrual period (LMP) of the participant's 2nd CPP pregnancy. All-cause mortality was ascertained through the National Death Index (NDI); deaths not identified in the NDI were captured using the Social Security Death Master File (SSDMF). The NDI includes all U.S. death records from 1979 onward, with matches identified through probabilistic linkage. Although the SSDMF includes deaths dating back to 1958, the NDI is considered a more accurate and comprehensive source of mortality and also includes cause of death[33, 34]. Overall, 89% of the cohort's death information was sourced from the NDI. Given the link between PPWR and cardiometabolic health[12, 23], secondary outcomes considered in this study were time to cause-specific mortality related to cardiovascular disease (I00-I78, I80-I99), diabetes (E10-E14), and kidney conditions (N00-N07, N17-N19, N25-N27) where the underlying cause of death was coded using ICD-10 (International Classification of Diseases, 10th Revision) and comparable ICD-9 codes.

Covariates

Covariates were reported by participants during the first prenatal visit of their 1st CPP pregnancy, and included information on age, frequency of smoking, race, marital status, income, education, number of prior pregnancies, hypertensive disorders of pregnancy, diabetes prior to pregnancy, prior cardiovascular diseases, prior respiratory diseases, prior renal disease, prior neurological conditions, prior cancer/tumor status, study site, pre-pregnancy weight, and height. Race was conceptualized as a social determinant of health in this study. Body mass index (BMI) (kg/m²) was calculated using self-reported height and pre-pregnancy weight. Additional 1st CPP pregnancy variables of interest included in this study were pregnancy plurality, gestational age at delivery, pregnancy outcomes, GWG, and the interconception and interpregnancy intervals. Details on how these variables were derived and incorporated at different analytical steps of the study are provided in Appendix A—Supplementary Table 1.

Statistical Analysis

Post data cleaning and processing, the participant characteristics were summarized for each quintile of IPWC. Missing or implausible exposure, covariate and outcome data were imputed via multiple imputation using chained equations. Under the Missing at Random assumption, 10 imputed datasets were generated for estimating pooled hazard ratios and the respective 95% Confidence Interval (CI). Multiple imputation was conducted using the R package *mice*(35). All other data cleaning, processing and analyses were conducted using SAS.

The study sample was a subset of the CPP cohort with at least two pregnancies in the CPP. Thus, to overcome the potential bias due to this inherent selection process and to ensure that the pooled hazard ratio estimates are generalizable to the entire CPP target-population, inverse probability weights (IPW) were used to reweight the outcomes data. The IPW was calculated as the ratio of two probabilities: the marginal probability of having a 2nd CPP pregnancy as the numerator, and the conditional probability of a 2nd CPP pregnancy as the denominator, where the conditional probability model included a wide range of variables as described in Supplementary Table 1.

Analyses with primary exposure IPWC: IPWC was categorized into quintiles, to identify any potential non-linear associations between the exposure and the outcome. The risks of all-cause mortality, associated with quintiles of IPWC were estimated using IPW weighted Cox proportional hazard (PH) model. Three covariate adjustment scenarios were applied: (1) unadjusted, (2) adjusted for confounders, including socio-demographic characteristics, smoking status, reproductive and medical history, and study site; and (3) additionally adjusted for interconception interval (ICI) i.e., the time difference between the LMP of the 1st and 2nd CPP pregnancies, to assess if time between two pregnancies contributed to weight change.

Multiple sensitivity analyses were conducted. First, to elucidate the effect of extreme weight loss or weight gain, we separated the top and bottom 1% of IPWC values following an initial review of the data distribution, thus rendering 7 different categories of IPWC. Second, we conducted a stratified analysis wherein the quintiles of IPWC were reassessed within BMI

categories: <18.5, 18.5-24.9, 25.0-29.9, ≥30.0. However, given the small sample sizes within some of the BMI category-specific quintiles, the risks of all-cause mortality were estimated only for those with normal (18.5-24.9) pre-pregnancy BMI (70.5%, n=5754).

Cause-specific mortality risks due to cardiovascular, diabetes and kidney conditions associated with quintiles of IPWC were estimated using the Cumulative Incidence Function (CIF), which focuses only on the cause of death that is of interest, while treating other causes as censored events, and does not assume that the competing events (i.e., the different cause of deaths) are independent of each other(36). Cause-specific mortality risks associated with the BMI-specific quintiles were not implemented due to small sizes. Details of the model specifications are provided in Supplementary Table 1.

Analyses with secondary exposure PPWC: PPWC was also categorized into quintiles. The risks of all-cause mortality, associated with quintiles of PPWC and BMI-specific quintiles of PPWC were estimated using similar modeling approaches as the IPWC analyses. The adjusted models were additionally accounted for 1st pregnancy GWG and model 3 alternatively adjusted for the IPI i.e., the time difference between 1st pregnancy delivery date and LMP of the 2nd pregnancy.

We completed multiple sensitivity analysis on PPWC. Given that the majority of the study participants lost weight and only a small fraction had weight gain during IPI, a third scenario distinguishing weight gain as a separate category from the 5th quintile of PPWC was also evaluated to tease out the risk associated with postpartum weight gain. This third scenario was also implemented to models that adjusted for birth weight and gestational age at delivery, and in the analysis with BMI-specific quintiles of PPWC. In addition, we repeated the above all-cause mortality analysis with PPWC quintiles, and with the weight gain group separated out from Q5, to include additional covariate adjustment and accounts for two potential confounding mechanisms. First, we adjusted for birth weight as a covariate given that our derivation of PPWC incorporates weight of the baby and therefore may account partially for the weight change patterns observed during IPI in this study. Secondly, there might have been an impact of shorter gestation among women delivering preterm on PPWC measures as well as on mortality—women delivering preterm

were likely to have gained less gestational weight, and preterm delivery is known to be associated with an increased mortality by cardiovascular disease(37). Therefore, we also ran a separate model including gestational age at delivery as a covariate.

Results

The distributions of most participant characteristics were similar across quintiles (Table 1). Some notable exceptions include the increase in the percentage of non-smokers as IPWC increased, with the highest percentage of non-smokers in the highest quintile of IPWC. The interval between conceptions for most quintiles of IPWC (1st, 3rd and 4th quintiles) averaged around two years (~24 months) but was slightly shorter for the 2nd quintile (~22 months) and longer (~27 months) for the 5th quintile of IPWC. The median follow-up for this study sample was 54 years [inter-quartile range (IQR): 47-56] and 41.7% (n=3405) died during the study period (Table 2).

Risk of mortality associated with IPWC

The majority of women (~51%, n=4141) had a higher weight at the beginning of their 2nd pregnancy than their 1st pregnancy and the median IPWC was 0.91 kg [IQR:-0.45, 3.63] (Supplementary Figure 1). Results from the adjusted model (Figure 2) demonstrated that compared to those with IPWC > 0 kg to 1.8 kg (Q3), all-cause mortality was lower among those in Q2 with IPWC > -1.4 to 0 kg [aHR (95% CI): 0.85 (0.76 – 0.96)]. Adjusting for ICI did not yield any meaningful changes in the estimates. And although the results were imprecise, compared to those in Q3, all-cause mortality was higher among those with IPWC > 1.8 kg (i.e., Q4 and Q5 of IPWC).

Distinguishing the top and bottom 1% of IPWC (Supplementary Figure 2) indicated an increased risk of all-cause mortality for those with very high IPWC > 17.2 kg [aHR (95% CI): 1.57 (1.10 – 2.23)] and very low IPWC ≤ -10.4 kg [aHR (95% CI): 1.15 (0.83 – 1.59)], compared to Q3 of IPWC, albeit the results lacked precision. Adjusting for ICI made only a marginal difference in the estimates.

Among those with normal pre-pregnancy BMI at their 1st CPP prenatal visit, IPWC > -0.9 and 0 kg was associated with a reduced risk of all-cause mortality [aHR (95% CI): 0.85 (0.73 – 0.99)] compared to IPWC > 0 to 1.8 kg (Supplementary Table 2). Consistent with the unstratified

analysis, women in Q4 and Q5 of IPWC exhibited a higher risk of all-cause mortality, with comparable trends in Q1; however, all of these estimates were imprecise.

Similar patterns of cause-specific mortality risk among women in Q2 and Q5 of IPWC were observed, although the estimates were imprecise due to limited sample sizes (Supplementary Table 3), with the notable exception of the risk of mortality due to diabetes [aHR (95% CI): 0.45 (0.23 – 0.87)] which was lower for those with IPWC > -1.4 to 0 kg (Q2). ICI adjusted results were similar.

Risk of mortality associated with PPWC

Approximately 94% of participants experienced weight loss during the postpartum period (median PPWC: -8.16 kg [IQR:-4.9, -11.34]) (Supplementary Figure 1). Supplementary Table 4 describes the participant characteristics according to PPWC. Those in quintile 1 with PPWC <-11.8 kg had the highest gestational weight gain with an average of 13.8 kg. Comparatively those in quintile 5 with PPWC > 4.5 kg had the lowest gestational weight gain with an average of 6.3 kg. The IPI increased slightly from 1.2 years in the 1st PPWC quintile to 1.4 years in the 5th quintile. We found no significant associations between any quintiles of PPWC and all-cause mortality (Figure 3). Nevertheless, reduced risks of all-cause mortality were suggested among those in Q1, Q2, and Q4 of PPWC. Adjusting for IPI did not yield any significant changes in the estimates.

Further separation of the 5th quintile into weight loss and gain (Supplementary Figure 3) showed an elevated but imprecise risk of all-cause mortality among those with weight gain [aHR (95% CI): 1.17 (0.95 – 1.43)]. The IPI-adjusted analysis results were similar. Among those with normal pre-pregnancy BMI, weight loss of less than 5 kg during the IPI (i.e., Q5 of PPWC) was linked to a higher risk of all-cause mortality [aHR, 95% CI: 1.21, 1.02 – 1.43] (Supplementary Table 6). Additionally, those with weight retention or gain during the IPI (i.e. PPWC ≥ 0 kg) had an elevated risk of all-cause mortality [aHR, 95% CI: 1.35, 1.01 – 1.79]. Adjustment for birth weight, and gestational age at delivery (Supplementary Table 7) yielded similar results to those observed in Figure 3 and Supplementary Figure 3.

The cause-specific analysis also revealed comparable patterns of risk across PPWC quintiles, although estimates lacked precision due to small sample sizes (Supplementary Table 8). In particular, the risk of mortality from kidney conditions [aHR (95% CI): 0.37 (0.08 – 1.75)] trended lower for those with PPWC ≥ -11.8 to < -9.1 kg, whereas among those with PPWC ≥ -4.5 kg, the risk trended higher for mortality from kidney conditions [aHR (95% CI): 1.75 (0.68 – 4.47)]. The IPI-adjusted results were consistent across both the sensitivity and cause-specific analyses.

Discussion

In a large, U.S. study with more than 50 years of follow-up after pregnancy, maternal postpartum weight retention independently contributed to long-term mortality risk, beyond the risks attributable to pre-pregnancy BMI and GWG. Specifically, we found that women who returned to their pre-pregnancy weight or maintain a weight slightly below that level, had a lower risk of all-cause mortality and experienced 55% lower diabetes-related mortality compared to those with moderate postpartum weight retention between pregnancies. Furthermore, women who entered a subsequent pregnancy at a higher weight than their pre-pregnancy baseline showed an increased risk of mortality, with this risk becoming markedly greater when weight retained or gained between pregnancies exceeded 17.2 kg. Our analysis of postpartum weight change, independent of gestational weight gain, further indicated that maintaining or modestly reducing weight in the postpartum period may be beneficial, whereas weight retention or gain during this time may increase the risk of mortality.

Our findings are novel and contribute to the growing body of literature on PPWR which highlights that women with high levels of PPWR face an increased risk of chronic conditions(12, 22–24). For instance, Kew et al. (2014) reported increased markers of adverse cardiometabolic health as early as 12 months postpartum, evidenced by higher mean adjusted diastolic blood pressure, insulin resistance, LDL cholesterol, and apolipoprotein B, along with declining adiponectin levels with higher postpartum weight gain trajectories(22). Studies with longer follow-up periods have reached similar conclusions regarding the long-term health risks associated with

postpartum weight retention. For example, Soria-Contreras et al. (2020) found that increasing postpartum weight trajectories were associated with higher levels of central adiposity, inflammation, and LDL cholesterol levels in later postpartum years(12). Similarly, Kramer et al. (2024) reported that PPWR was associated not only with a higher risk of pre-diabetes/diabetes at 5-years postpartum, but also predicted worse characterization of cardiovascular risk indices such as triglycerides, high-/low-density lipoprotein, apolipoprotein-B, Matsuda index, insulin resistance, fasting glucose, and C-reactive protein(23). A study using data from the Danish National Birth Cohort (DNBC; 1997–2002) found elevated risks of hypertension and cardiovascular disease 16 years postpartum among women in the weight retention or gain groups(24). Due to the small sample size, which may have contributed to the lack of precision in our cause-specific results, we could not draw definitive conclusions about the increased risks of mortality from cardiovascular, and kidney conditions associated with IPWC or PPWC within this cohort. However, our finding of a 55% reduction in diabetes-related mortality from postpartum weight loss is notable. These results suggest that weight retention or gain during the postpartum period may increase the risk of mortality, while weight loss may reduce this risk. Thus, our study not only complements previous studies that have linked PPWR to morbidity but also represents a novel contribution—being the first to map the association between PPWR and long-term mortality, and to suggest a potential survival benefit from healthy postpartum weight loss.

Our findings also underscore the need to implement postpartum weight management strategies across the full range of pre-pregnancy BMI categories. In 2021, the US Preventive Services Task Force found strong evidence(38–40) that behavioral interventions reduce the likelihood of excessive weight gain during pregnancy, recommending that healthcare providers offer behavioral counseling to pregnant women to promote healthy weight gain in line with NAM guidelines. However, such guidance for maintaining a healthy weight during the postpartum period remains limited. The few trials that have explored this area have focused on postpartum weight management for women living with overweight or obesity given their elevated risk of long-term health complications, as is evident in this systematic review(41) of 18 randomized controlled trials

(2559 participants). However, BMI-specific analyses from this study suggest that women with low risk at baseline i.e., a normal pre-pregnancy BMI may also face an increased risk of earlier mortality if weight retention or gain occurs during the postpartum period. Therefore, postpartum weight management interventions should also consider normal BMI groups and not just women with BMI > 24.9 kg/m^2.

This study focused on pregnant women from the 1950s and 1960s, utilizing a historic cohort to examine the long-term health implications of PPWR, with follow-up extending over 50 years—an important and unique strength of this study given it is unfeasible to study long-term mortality utilizing contemporary cohorts. Yet, some characteristics, such as prevalence of smoking and underlying causes of mortality in CPP, which differ from current obstetric populations may limit generalizability. However, CPP's rich covariate data and updated ICD-coded mortality enable future application of transportability methods(42) – a valuable direction for future research to extend these findings to modern populations. Another strength of this study is its analytical approach. Although the analytical sample was restricted to women with at least two CPP-registered pregnancies, the use of inverse probability weighting (IPW) accounted for potential selection bias introduced by this restriction, allowing the results to remain generalizable to the overall CPP population. The study's racial diversity, with nearly 50% of participants identifying as Black, is also a key strength, as much of the existing research on pregnancy exposures and long-term outcomes has predominantly focused on White populations. Finally, while the PPWC analyses were adjusted for ICI, the strength of the association between PPWC and long-term mortality may vary across different durations of ICI. Identifying such heterogeneity could help determine optimal time frames for postpartum weight management before a subsequent pregnancy. Future studies should explore potential effect modification by ICI, which was beyond the scope of this study due to sample size constraints.

Conclusion

In conclusion, weight retention or weight gain in the postpartum period may elevate the risk of earlier mortality. This is true irrespective of pre-pregnancy BMI status, thus underscoring

the benefit of postpartum weight loss and the need for effective strategies to prevent postpartum weight retention or new weight gain for all women in the post-partum period. Future research with larger sample sizes and greater statistical power is needed to confirm these findings.

Acknowledgments

References

1. Anon. (2025). Overweight & Obesity Statistics - NIDDK. [WWW document]. URL https://www.niddk.nih.gov/health-information/health-statistics/overweight-obesity

2. Anon. (2025). Products - Health E Stats - Prevalence of Overweight, Obesity, and Extreme Obesity Among Adults Aged 20 and Over: United States, 1960–1962 Through 2017–2018. [WWW document]. URL https://www.cdc.gov/nchs/data/hestat/obesity-adult-17-18/obesity-adult.htm

3. Kent L, McGirr M, Eastwood K-A. Global trends in prevalence of maternal overweight and obesity: A systematic review and meta-analysis of routinely collected data retrospective cohorts. *Int J Popul Data Sci* 2024;9.

4. Statistics NC for H. *National Health Interview Survey: questionnaires, datasets, and related documentation*. 2010.

5. Davis E, Olson C. Obesity in Pregnancy. *Prim Care Clin Off Pract* 2009;36:341–356.

6. Davis EM, Zyzanski SJ, Olson CM, Stange KC, Horwitz RI. Racial, Ethnic, and Socioeconomic Differences in the Incidence of Obesity Related to Childbirth. *Am J Public Health* 2009;99:294–299.

7. Williamson DF, Madans J, Pamuk E, Flegal KM, Kendrick JS, Serdula MK. A prospective study of childbearing and 10-year weight gain in US white women 25 to 45 years of age. *Int J Obes Relat Metab Disord* 1994;18:561–569.

8. Endres LK, Straub H, McKinney C, *et al.* Postpartum Weight Retention Risk Factors and Relationship to Obesity at 1 Year. *Obstet Gynecol* 2015;125:144.

9. Legro NR, Lehman EB, Kjerulff KH. Mode of first delivery and postpartum weight retention at 1 year. *Obes Res Clin Pract* 2020;14:241–248.

10. Nehring I, Schmoll S, Beyerlein A, Hauner H, von Kries R. Gestational weight gain and long-term postpartum weight retention: a meta-analysis1234. *Am J Clin Nutr* 2011;94:1225–1231.

11. Ketterl TG, Dundas NJ, Roncaioli SA, Littman AJ, Phipps AI. Association of Pre-pregnancy BMI and Postpartum Weight Retention Before Second Pregnancy, Washington State, 2003-2013. *Matern Child Health J* 2018;22:1339–1344.

12. Soria-Contreras DC, Rifas-Shiman SL, Aris IM, *et al.* Weight Trajectories After Delivery are Associated with Adiposity and Cardiometabolic Markers at 3 Years Postpartum Among Women in Project Viva. *J Nutr* 2020;150:1889–1898.

13. McClure CK, Catov JM, Ness R, Bodnar LM. Associations between gestational weight gain and BMI, abdominal adiposity, and traditional measures of cardiometabolic risk in mothers 8 y postpartum123. *Am J Clin Nutr* 2013;98:1218–1225.

14. Tahir MJ, Haapala JL, Foster LP, *et al.* Association of Full Breastfeeding Duration with Postpartum Weight Retention in a Cohort of Predominantly Breastfeeding Women. *Nutrients* 2019;11:938.

15. Baker JL, Gamborg M, Heitmann BL, Lissner L, Sørensen TI, Rasmussen KM. Breastfeeding reduces postpartum weight retention34. *Am J Clin Nutr* 2008;88:1543–1551.

16. Foster SF, Vazquez C, Cubbin C, Nichols AR, Rickman RR, Widen EM. Breastfeeding, socioeconomic status, and long-term postpartum weight retention. *Int Breastfeed J* 2023;18:1.

17. Jiang M, Gao H, Vinyes-Pares G, *et al.* Association between breastfeeding duration and postpartum weight retention of lactating mothers: A meta-analysis of cohort studies. *Clin Nutr* 2018;37:1224–1231.

18. Wagner KA, Whitcomb BW, Marcus B, *et al.* The impact of a lifestyle intervention on postpartum weight retention among US Hispanic women with overweight and obesity. *Prev Med Rep* 2024;38:102633.

19. Makama M, Skouteris H, Moran LJ, Lim S. Reducing Postpartum Weight Retention: A Review of the Implementation Challenges of Postpartum Lifestyle Interventions. *J Clin Med* 2021;10:1891.

20. Liu J, Wilcox S, Hutto B, Turner-McGrievy G, Wingard E. Effects of a lifestyle intervention on postpartum weight retention among women with elevated weight. *Obes Silver Spring Md* 2022;30:1370–1379.

21. Brown SD, Kiernan M, Hedderson MM, *et al.* Greater intrinsic motivation for engaging in healthy lifestyle behaviors is associated with reduced postpartum weight. *Obesity* 2025;33:927–935.

22. Kew S, Ye C, Hanley AJ, *et al.* Cardiometabolic implications of postpartum weight changes in the first year after delivery. *Diabetes Care* 2014;37:1998–2006.

23. Kramer CK, Ye C, Hanley AJ, *et al.* Postpartum weight retention and the early evolution of cardiovascular risk over the first 5 years after pregnancy. *Cardiovasc Diabetol* 2024;23:101.

24. Kirkegaard H, Bliddal M, Støvring H, *et al.* Maternal weight change from prepregnancy to 18 months postpartum and subsequent risk of hypertension and cardiovascular disease in Danish women: A cohort study. *PLoS Med* 2021;18:e1003486.

25. Hillemeier MM, Weisman CS, Chuang C, Downs DS, McCall-Hosenfeld J, Camacho F. Transition to Overweight or Obesity Among Women of Reproductive Age. *J Womens Health* 2011;20:703–710.

26. Schon SB, Cabre HE, Redman LM. The impact of obesity on reproductive health and metabolism in reproductive-age females. *Fertil Steril* 2024;122:194–203.

27. Grandi SM, Hinkle SN, Mumford SL, *et al.* Long-Term Mortality in Women With Pregnancy Loss and Modification by Race/Ethnicity. *Am J Epidemiol* 2022;191:787–799.

28. Yeung EH, Saha A, Zhu C, *et al.* Placental characteristics and risks of maternal mortality 50 years after delivery. *Placenta* 2022;117:194–199.

29. Rahman ML, Feng Y-CA, Fiehn O, *et al.* Plasma lipidomics profile in pregnancy and gestational diabetes risk: a prospective study in a multiracial/ethnic cohort. *BMJ Open Diabetes Res Care* 2021;9:e001551.

30.	Hinkle SN, Schisterman EF, Liu D, *et al.* Pregnancy Complications and Long-Term Mortality in a Diverse Cohort. *Circulation* 2023;147:1014–1025.

31.	Grandi SM, Hinkle SN, Mumford SL, *et al.* Infant sex at birth and long-term maternal mortality. *Paediatr Perinat Epidemiol* 2023;37:229–238.

32.	Mitro SD, Sundaram R, Qiao Y, *et al.* History of multifetal gestation and long-term maternal mortality. *Paediatr Perinat Epidemiol* 2024;38:219–226.

33.	Pollack AZ, Hinkle SN, Liu D, *et al.* Vital Status Ascertainment for a Historic Diverse Cohort of U.S. Women. *Epidemiol Camb Mass* 2020;31:310–316.

34.	Lash TL, Silliman RA. A Comparison of the National Death Index and Social Security Administration Databases to Ascertain Vital Status. *Epidemiology* 2001;12:259.

35.	Buuren S van, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained Equations in R. *J Stat Softw* 2011;45:1–67.

36.	Kalbfleisch JD, Prentice RL. *The statistical analysis of failure time data*. 2nd ed. Hoboken, N.J: John Wiley & Sons; 2002.

37.	Wu P, Gulati M, Kwok CS, *et al.* Preterm Delivery and Future Risk of Maternal Cardiovascular Disease: A Systematic Review and Meta-Analysis. *J Am Heart Assoc* 2018;7:e007809.

38.	Kent-Marvick J, Cloyes KG, Meek P, Simonsen S. Racial and ethnic disparities in postpartum weight retention: A narrative review mapping the literature to the National Institute on Minority Health and Health Disparities Research Framework. *Womens Health Lond Engl* 2023;19:17455057231166822.

39.	Li HZ, Boyle JA, Harrison CL. Preventing postpartum weight retention following antenatal lifestyle intervention: One year postpartum follow up of the Healthy Lifestyles in Pregnancy (HeLP-her) randomised controlled trial. *Aust N Z J Obstet Gynaecol* 2022;62:319–322.

40.	Toro-Ramos T, Heaner M, Yang Q, *et al.* Postpartum Weight Retention: A Retrospective Data Analysis Measuring Weight Loss and Program Engagement with a Mobile Health Program. *J Womens Health 2002* 2021;30:1645–1652.

41.	Dalrymple KV, Flynn AC, Relph SA, O'Keeffe M, Poston L. Lifestyle Interventions in Overweight and Obese Pregnant or Postpartum Women for Postpartum Weight Management: A Systematic Review of the Literature. *Nutrients* 2018;10:1704.

42.	Vuong Q, Metcalfe RK, Ling A, Ackerman B, Inoue K, Park JJ. Systematic review of applied transportability and generalizability analyses: A landscape analysis. *Ann Epidemiol* 2025;104:61–70.

Table 1: Summary of participant characteristics across quintiles of interpregnancy weight change (IPWC)

Participant Characteristics at 1st CPP Pregnancy	IPWC quintile					
	Q1: ≤ -1.4 kg (n = 1501)	Q2: >-1.4 to 0 kg (n = 1780)	Q3: > 0 to 1.8 kg (n = 1269)	Q4: >1.8 to 4.5 kg (n = 1621)	Q5: > 4.5 kg (n = 1251)	Missing (n = 743)
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Age (years)						
Mean (SD)	22.6 (5.3)	22.8 (5.3)	23.3 (5.5)	23.1 (5.5)	22.4 (5.1)	23.8 (5.7)
Missing	-	-	-	-	-	-
Race and Ethnicity						
Black	690 (46.0)	811 (45.6)	549 (43.3)	738 (45.5)	623 (49.8)	321 (43.2)
Asian/Other	18 (1.2)	18 (1.0)	12 (0.9)	8 (0.5)	5 (0.4)	8 (1.1)
Puerto Rican	43 (2.9)	63 (3.5)	23 (1.8)	65 (4.0)	41 (3.3)	57 (7.7)
White	750 (50.0)	888 (49.9)	685 (54.0)	810 (50.0)	582 (46.5)	357 (48.0)
Missing	-	-	-	-	-	-
Number of Prior Pregnancies						
0	615 (41.0)	650 (36.5)	485 (38.2)	612 (37.8)	520 (41.6)	244 (32.8)
1	280 (18.7)	382 (21.5)	252 (19.9)	337 (20.8)	222 (17.7)	126 (17.0)
2	184 (12.3)	266 (14.9)	187 (14.7)	211 (13.0)	157 (12.5)	87 (11.7)
3	155 (10.3)	153 (8.6)	118 (9.3)	157 (9.7)	114 (9.1)	96 (12.9)
4	261 (17.4)	315 (17.7)	216 (17.0)	297 (18.3)	229 (18.3)	154 (20.7)
Missing	*6 (0.4)*	*14 (0.8)*	*11 (0.9)*	*7 (0.4)*	*9 (0.7)*	*36 (4.8)*
Marital Status						
Married/ Common Law	1156 (77.0)	1406 (79.0)	1044 (82.3)	1295 (79.9)	992 (79.3)	597 (80.3)
Single	264 (17.6)	271 (15.2)	166 (13.1)	236 (14.6)	181 (14.5)	104 (14.0)
Widowed/Divorced/Separated	81 (5.4)	103 (5.8)	59 (4.6)	90 (5.6)	78 (6.2)	42 (5.7)
Missing	-	-	-	-	-	-
Smoking Status						
Nonsmoker	697 (46.4)	859 (48.3)	654 (51.5)	883 (54.5)	736 (58.8)	354 (47.6)
<1 pack per day	548 (36.5)	655 (36.8)	431 (34.0)	512 (31.6)	371 (29.7)	244 (32.8)
>= 1 pack per day	241 (16.1)	247 (13.9)	168 (13.2)	210 (13.0)	133 (10.6)	103 (13.9)
Missing	*15 (1.0)*	*19 (1.1)*	*16 (1.3)*	*16 (1.0)*	*11 (0.9)*	*42 (5.7)*
Annual Income						
<= $1999	234 (15.6)	256 (14.4)	139 (11.0)	241 (14.9)	193 (15.4)	93 (12.5)
$2000 - $3999	613 (40.8)	675 (37.9)	457 (36.0)	614 (37.9)	472 (37.7)	272 (36.6)

Participant Characteristics at 1st CPP Pregnancy	IPWC quintile					
	Q1: ≤ -1.4 kg (n = 1501)	Q2: >-1.4 to 0 kg (n = 1780)	Q3: > 0 to 1.8 kg (n = 1269)	Q4: >1.8 to 4.5 kg (n = 1621)	Q5: > 4.5 kg (n = 1251)	Missing (n = 743)
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
$4000 - $5999	284 (18.9)	389 (21.9)	279 (22.0)	337 (20.8)	236 (18.9)	137 (18.4)
$6000 - $7999	113 (7.5)	144 (8.1)	154 (12.1)	151 (9.3)	127 (10.2)	65 (8.7)
$8000 - $9999	47 (3.1)	54 (3.0)	50 (3.9)	70 (4.3)	40 (3.2)	17 (2.3)
>= $10000	20 (1.3)	37 (2.1)	26 (2.0)	29 (1.8)	23 (1.8)	13 (1.7)
Missing	*234 (15.6)*	*256 (14.4)*	*139 (11.0)*	*241 (14.9)*	*193 (15.4)*	*93 (12.5)*
Education						
< High School	251 (16.7)	268 (15.1)	144 (11.3)	235 (14.5)	223 (17.8)	154 (20.7)
Some High School	622 (41.4)	765 (43.0)	480 (37.8)	664 (41.0)	509 (40.7)	259 (34.9)
High School Grad	421 (28.0)	504 (28.3)	435 (34.3)	522 (32.2)	391 (31.3)	178 (24.0)
>= Some College	148 (9.9)	181 (10.2)	155 (12.2)	143 (8.8)	89 (7.1)	84 (11.3)
Missing	*59 (3.9)*	*62 (3.5)*	*55 (4.3)*	*57 (3.5)*	*39 (3.1)*	*68 (9.2)*
Hypertensive disorders of pregnancy						
Normotensive	1395 (92.9)	1692 (95.1)	1215 (95.7)	1505 (92.8)	1145 (91.5)	682 (91.8)
Chronic hypertension	33 (2.2)	37 (2.1)	21 (1.7)	38 (2.3)	50 (4.0)	25 (3.4)
Gestational hypertension	33 (2.2)	27 (1.5)	22 (1.7)	39 (2.4)	32 (2.6)	12 (1.6)
Preeclampsia/Eclampsia	17 (1.1)	11 (0.6)	6 (0.5)	20 (1.2)	11 (0.9)	9 (1.2)
Superimposed Preeclampsia/Eclampsia	23 (1.5)	13 (0.7)	4 (0.3)	19 (1.2)	13 (1.0)	7 (0.9)
Missing	-	-	*1 (0.1)*	-	-	*8 (1.1)*
Prior Diabetes Ⱡ						
No	1472 (98.1)	1755 (98.6)	1245 (98.1)	1599 (98.6)	1229 (98.2)	700 (94.2)
Yes	29 (1.9)	25 (1.4)	24 (1.9)	22 (1.4)	22 (1.8)	36 (4.8)
Missing	-	-	-	-	-	*7 (0.9)*
Prior Cardiovascular Conditions ‡						
No	1294 (86.2)	1561 (87.7)	1121 (88.3)	1420 (87.6)	1087 (86.9)	653 (87.9)
Yes	207 (13.8)	218 (12.2)	147 (11.6)	201 (12.4)	164 (13.1)	77 (10.4)
Missing	-	*1 (0.1)*	*1 (0.1)*	-	-	*13 (1.7)*
Prior Respiratory Conditions §						
No	1390 (92.6)	1636 (91.9)	1183 (93.2)	1503 (92.7)	1171 (93.6)	677 (91.1)
Yes	111 (7.4)	143 (8.0)	85 (6.7)	118 (7.3)	80 (6.4)	53 (7.1)
Missing	-	*1 (0.1)*	*1 (0.1)*	-	-	*13 (1.7)*

Participant Characteristics at 1st CPP Pregnancy	IPWC quintile					
	Q1: ≤ -1.4 kg (n = 1501)	Q2: >-1.4 to 0 kg (n = 1780)	Q3: > 0 to 1.8 kg (n = 1269)	Q4: >1.8 to 4.5 kg (n = 1621)	Q5: > 4.5 kg (n = 1251)	Missing (n = 743)
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Prior Renal Conditions ¶						
No	1438 (95.8)	1714 (96.3)	1213 (95.6)	1553 (95.8)	1206 (96.4)	689 (92.7)
Yes	63 (4.2)	65 (3.7)	55 (4.3)	68 (4.2)	45 (3.6)	41 (5.5)
Missing	-	*1 (0.1)*	*1 (0.1)*	-	-	*13 (1.7)*
Prior Neurological Conditions ||						
No	1385 (92.3)	1646 (92.5)	1165 (91.8)	1508 (93.0)	1151 (92.0)	663 (89.2)
Yes	116 (7.7)	133 (7.5)	103 (8.1)	113 (7.0)	100 (8.0)	67 (9.0)
Missing	-	*1 (0.1)*	*1 (0.1)*	-	-	*13 (1.7)*
Prior Cancer or Tumors **						
No	1447 (96.4)	1732 (97.3)	1216 (95.8)	1576 (97.2)	1213 (97.0)	708 (95.3)
Yes	54 (3.6)	47 (2.6)	52 (4.1)	45 (2.8)	38 (3.0)	22 (3.0)
Missing	-	*1 (0.1)*	*1 (0.1)*	-	-	*13 (1.7)*
Pregnancy Plurality						
Singleton	1487 (99.1)	1766 (99.2)	1254 (98.8)	1605 (99.0)	1245 (99.5)	703 (94.6)
Multiples	12 (0.8)	13 (0.7)	13 (1.0)	14 (0.9)	6 (0.5)	4 (0.5)
Missing	*2 (0.1)*	*1 (0.1)*	*2 (0.2)*	*2 (0.1)*	*0 (0.0)*	*36 (4.8)*
Study Site						
Boston	408 (27.2)	474 (26.6)	444 (35.0)	536 (33.1)	363 (29.0)	248 (33.4)
Buffalo	61 (4.1)	99 (5.6)	64 (5.0)	64 (3.9)	31 (2.5)	22 (3.0)
New Orleans	29 (1.9)	31 (1.7)	16 (1.3)	31 (1.9)	30 (2.4)	4 (0.5)
New York/Columbia	11 (0.7)	13 (0.7)	11 (0.9)	20 (1.2)	10 (0.8)	5 (0.7)
Baltimore	111 (7.4)	118 (6.6)	97 (7.6)	110 (6.8)	94 (7.5)	61 (8.2)
Virginia	98 (6.5)	123 (6.9)	71 (5.6)	81 (5.0)	70 (5.6)	32 (4.3)
Minnesota	70 (4.7)	115 (6.5)	69 (5.4)	65 (4.0)	43 (3.4)	25 (3.4)
New York/Medical College	50 (3.3)	67 (3.8)	19 (1.5)	52 (3.2)	42 (3.4)	46 (6.2)
Oregon	141 (9.4)	134 (7.5)	51 (4.0)	104 (6.4)	90 (7.2)	27 (3.6)
Pennsylvania	352 (23.5)	411 (23.1)	284 (22.4)	368 (22.7)	324 (25.9)	228 (30.7)
Providence	133 (8.9)	143 (8.0)	103 (8.1)	132 (8.1)	106 (8.5)	42 (5.7)
Tennessee	37 (2.5)	52 (2.9)	40 (3.2)	58 (3.6)	48 (3.8)	3 (0.4)
Missing	-	-	-	-	-	-

Participant Characteristics at 1st CPP Pregnancy	IPWC quintile					
	Q1: ≤ -1.4 kg (n = 1501)	Q2: >-1.4 to 0 kg (n = 1780)	Q3: > 0 to 1.8 kg (n = 1269)	Q4: >1.8 to 4.5 kg (n = 1621)	Q5: > 4.5 kg (n = 1251)	Missing (n = 743)
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Height (cm)						
Mean (SD)	161.5 (6.2)	160.8 (6.3)	160.7 (6.2)	160.8 (6.3)	161.3 (6.4)	160.1 (7.1)
Missing	*20 (1.3)*	*17 (1.0)*	*8 (0.6)*	*19 (1.2)*	*8 (0.6)*	*68 (9.2)*
Pre-pregnancy Weight (kg)						
Mean (SD)	62.1 (12.1)	56.8 (9.9)	55.8 (9.2)	57.1 (9.9)	60.5 (11.8)	59.2 (11.7)
Missing	-	-	-	-	-	*131 (17.6)*
Pre-pregnancy BMI (kg/m2)						
Mean (SD)	23.8 (4.4)	22.0 (3.6)	21.6 (3.4)	22.1 (3.5)	23.2 (4.2)	23.1 (4.3)
Missing	20 (1.3)	17 (1.0)	8 (0.6)	19 (1.2)	8 (0.6)	169 (22.7)
Interconception Interval (days) *						
Mean (SD)	710.6 (352.3)	662.3 (330.6)	693.6 (356.3)	703.0 (372.5)	793.3 (426.3)	725.4 (398.1)
Missing	*5 (0.3)*	*2 (0.1)*	-	*1 (0.1)*	-	*655 (88.2)*

Ɨ Type unknown. ‡Cardiovascular conditions included hypertension, rheumatic fever, and any other cardiovascular diseases. §Respiratory conditions included tuberculosis, asthma, other chronic pulmonary diseases, and other conditions requiring thoracic surgery. ¶Renal conditions included pyelitis, glomerulonephritis, and other conditions requiring kidney, urinary, or bladder surgery. ||Neurological conditions included neuromuscular diseases, convulsive disorders psychosis, alcohol or drug addiction, or other neurological diseases. **Cancer and tumors included any history of cancer, and gastrointestinal, kidney, urinary, bladder, or gynecological tumors.

* Inter-conception Interval calculated as the time difference between the last menstrual period of the 1st and 2nd CPP pregnancies.

Table 2: Distribution of outcome of interest across quintiles of interpregnancy weight change

Outcomes of Interest		IPWC quintile					
		Q1: ≤ -1.4 kg (n = 1501)	Q2: >-1.4 to 0 kg (n = 1780)	Q3: > 0 to 1.8 kg (n = 1269)	Q4: >1.8 to 4.5 kg (n = 1621)	Q5: > 4.5 kg (n = 1251)	Missing (n = 743)
		n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Vital Status	Alive	865 (57.6)	1067 (59.9)	752 (59.3)	941 (58.1)	712 (56.9)	423 (56.9)
	Dead	636 (42.4)	713 (40.1)	517 (40.7)	680 (41.9)	539 (43.1)	320 (43.1)
	Missing	-	-	-	-	-	-
Cause of Death	Cardiovascular	162 (25.5)	196 (27.5)	119 (23.0)	184 (27.1)	149 (27.6)	89 (27.8)
	Diabetes	28 (4.4)	19 (2.7)	22 (4.3)	22 (3.2)	27 (5.0)	18 (5.6)
	Kidney	8 (1.3)	8 (1.1)	10 (1.9)	14 (2.1)	14 (2.6)	5 (1.6)
	Missing	*61 (4.1)*	*68 (3.8)*	*46 (3.6)*	*64 (3.9)*	*56 (4.5)*	*30 (4.0)*

Figure 1: Exposures and timeline of measurement

Figure 2: Risk of all cause mortality associated with quintiles of interpregnancy weight change (IPWC)

Reference – Q3: IPWC > 0 to 1.8 kg

Adjusted for CPP 1st Pregnancy variables: age, race and ethnicity, number of prior pregnancies, marital status, smoking status, annual income, education, prior diabetes, prior cardiovascular conditions, prior respiratory conditions, prior renal conditions, prior neurological conditions, prior cancer or tumors, pregnancy plurality, study site, and pre-pregnancy BMI

Figure 3: Risk of all cause mortality associated with quintiles of Postpartum Weight Change (PPWC)

Quintiles of PPWC	HR	95% CI
	1.11	(0.97, 1.27)
Q1: < -11.8 kg	0.93	(0.79, 1.09)
	0.92	(0.79, 1.08)
	0.92	(0.80, 1.06)
Q2: ≥ -11.8 to < -9.1 kg	0.88	(0.75, 1.03)
	0.88	(0.75, 1.04)
	1.00	(0.88, 1.14)
Q4: ≥ -7.3 to < -4.5 kg	0.95	(0.82, 1.11)
	0.95	(0.81, 1.11)
	1.17	(1.03, 1.33)
Q5: ≥ -4.5 kg	1.01	(0.87, 1.17)
	0.99	(0.85, 1.14)

Estimate Type
- Crude
▲ Adjusted
■ IPI-Adjusted

Reference – Q3: PPWC ≥ -9.1 to < -7.3 kg

Adjusted for CPP 1st Pregnancy variables: age, race and ethnicity, number of prior pregnancies, marital status, smoking status, annual income, education, hypertensive disorders of pregnancy, prior diabetes, prior cardiovascular conditions, prior respiratory conditions, prior renal conditions, prior neurological conditions, prior cancer or tumors, pregnancy plurality, study site, pre-pregnancy BMI, and gestational weight gain