Community Consultation and Public Disclosure for the Randomized Trial of Sedative

Choice for Intubation

Ariel A. Lewis MPH[1]; Tiffany L. Israel MSSW[2]; Kevin P. Seitz MD, MSc[3]; Brian E. Driver MD[4] ; Kevin W. Gibbs MD[5]; Adit A Ginde MD, MPH[6]; Stacy A. Trent MD, MPH[6,7]; Derek W. Russell MD[8,9]; Matthew E. Prekker MD, MPH[4,10,]; Aaron E. Robinson MD, MPH[4] ; Jessica A. Palakshappa MD, MS[5]; John P. Gaillard MD[11,12]; L. Jane Stewart MD, JD, MPH[13,14]; Logan L. Beach MD[15]; Bradley D. Lloyd, RRT-ACCS[1]; Stephanie C. DeMasi MD[2]; Margaret A. Hays, MSN[3]; Cori Withers[6]; Amy E. Sullivan[6]; Carolynn Lyle PA-C, MPH[7]; Micah R. Whitson MD[8,15]; Barbara Gould, MSW[17];Todd W. Rice MD MSc[1,3]; Wesley H. Self MD MPH[1,2]; Jin H. Han MD, MSc[2,18] ; Matthew W. Semler MD, MSc[1,3]; Jonathan D. Casey MD, MSc[1,3] for the RSI investigators and the Pragmatic Critical Care Research Group

Affiliations:

1Vanderbilt Institute for Clinical and Translational Research, Vanderbilt University Medical Center, Nashville, TN
2 Department of Emergency Medicine, Vanderbilt University Medical Center, Nashville, TN
3Division of Allergy, Pulmonary, and Critical Care Medicine, Vanderbilt University Medical Center, Nashville, TN
4Department of Emergency Medicine, Hennepin County Medical Center, Minneapolis, MN
Department of Medicine, Section of Pulmonary, Critical Care, Allergy and Immunologic Disease, Wake Forest School of Medicine, Winston-Salem, North Carolina
Department of Emergency Medicine, University of Colorado School of Medicine, Aurora, Colorado
Department of Emergency Medicine, Denver Health Medical Center, Denver, Colorado
Department of Medicine, Division of Pulmonary, Allergy and Critical Care Medicine, University of Alabama at Birmingham, Birmingham, AL
Pulmonary Section, Birmingham Veteran's Affairs Medical Center, Birmingham, AL
Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine, Hennepin County Medical Center, Minneapolis, MN
Department of Anesthesiology-Critical Care, Wake Forest School of Medicine, Winston-Salem, NC
Department of Emergency Medicine, Wake Forest School of Medicine, Winston-Salem, NC
Department of Anesthesiology, University of Colorado School of Medicine, Aurora, Colorado
Department of Anesthesiology, Denver Health Medical Center, Denver, Colorado
Department of Emergency Medicine, University of Alabama at Birmingham, Birmingham AL
Department of Medicine, Division of Pulmonary Sciences and Critical Care Medicine, University of Colorado Anschutz Medical Center, Aurora, CO
Denver, Colorado
Geriatric Research, Education, and Clinical Center, Tennessee Valley Healthcare System, Nashville, TN

Drs. Semler and Casey contributed equally to this manuscript as senior authors.
Collaborators for the RSI Investigators are listed in Appendix 1.

Community Consultation and Public Disclosure for the Randomized Trial of Sedative Choice for Intubation

47 **Corresponding Author:**

48 Jonathan D. Casey, MD, MSc
49

Email:

jonathan.d.casey@vumc.org

Address:

Vanderbilt Institute for Clinical and Translational Research, 2525 West End Ave., Suite 600,

Room 672, Nashville, TN, 37203

Telephone: (615) 875-4681

Manuscript Word Count (body only): 3894/5000

Date: September 26, 2025

CONTRIBUTORS

All study authors approved the final version of this manuscript.

All study authors critically reviewed the manuscript for important intellectual content.

Study concept and design: MWS, JDC, AAL

Acquisition of data: All authors

Analysis and interpretation of data: AAL, TLI, SCD, MWS, JDC

Drafting of the manuscript: AAL, TLI, MWS, JDC

Revision of the manuscript: All authors

Acquisition of funding: MWS, JDC

FUNDING

Research reported in this article was funded through a Patient-Centered Outcomes Research

Institute (PCORI) Award (BPS-2022C3-30021). The views and statements presented in this

article are solely the responsibility of the authors and do not necessarily represent the views of

PCORI. Funding for the trial was also provided by the National Heart, Blood, and Lung Institute

(NHLBI) (K23HL153584). The views and statements presented in this article are solely the

responsibility of the authors and do not necessarily represent the views of the NHLBI.

COMPETING INTERESTS AND FINANCIAL DISCLOSURES

The authors have no financial conflicts of interest relevant to the current work.

B.E.D. reported having received compensation from UpToDate unrelated to the current work.

A.A.G. reported having received compensation from Biomeme, SeaStar, and National Institutes

of Health, unrelated to the current work.

S.A.T. reported having received compensation from the American College of Emergency

Physicians unrelated to the current work.

D. W. R. reported having received compensation from ABSS Solutions, Quantum Leap

Healthcare Collaborative, Critical Care Clinical Trial Network, Achieve Life Sciences, and Direct

Biologics unrelated to the current work.

J.A.P. reported having received compensation from American College of Chest Physicians

unrelated to the current work.

J.P.G. reported having received compensation from American College of Chest Physicians

unrelated to the current work.

T.W.R. reported having received compensation from Cumberland Pharmaceuticals, Cytovale,

Nestle, and Sanofi unrelated to the current work.

M.W.S. reported having received compensation from Baxter Healthcare Corporation,

DynaMedex, and Reprieve Cardiovascular, Inc unrelated to the current work.

J.D.C. reported compensation from Reprieve Cardiovascular, Inc unrelated to the current work.

ABSTRACT

Background

Randomized trials evaluating emergency treatments may be conducted with Exception

from Informed Consent (EFIC) when prospective informed consent is infeasible. In EFIC trials,

a period of community consultation and public disclosure precedes initiation of enrollment. The

Randomized Trial of Sedative Choice for Intubation (RSI) is a 2,364-patient randomized trial

being conducted with EFIC in 14 emergency departments and intensive care units across the

United States. This manuscript reports the approach to community consultation and public

disclosure in the RSI trial.

Methods

Community consultation and public disclosure were conducted locally in each of the 5

regions of enrolling sites. The coordinating center provided sites with templates and access to

an engagement coordinator to assist with developing and executing site plans.

Results

Community consultation and public disclosure occurred at the coordinating center from

February 2021-January 2022 and in the regions of the five additional enrolling sites from

September 2023-June 2024. Community consultation included in-person surveys with 789

patients or family members in emergency department or intensive care unit waiting rooms and

invitation of more than 200 local groups to town halls or community engagement studios. Public

disclosure included (i) social media advertisements viewed more than 1.2 million times, (ii) a

trial website with more than 16,000 unique visitors, (iii) informational flyers in hospitals and

public settings, and (iv) featured information in traditional media. Completing local community

consultation and public disclosure required an average of 139 hours of research personnel time

and approximately $18,822 at each site, in addition to coordinating center effort and costs.

Conclusions

Pre-trial community consultation activities in the RSI trial engaged over 1,000 patients, families,

and community members and public disclosure reached over 1.2 million community members.

While the total cost and duration of activities at sites were substantially lower than reported in

prior EFIC trials, these costs remained significant.

INTRODUCTION

In response to the ethical dilemma posed by emergency care research in which

prospective written informed consent is infeasible and the research is greater than minimal risk,

the U.S. Food and Drug Administration (FDA) created the Exception from Informed Consent

(EFIC) process in 1996. To qualify for EFIC, a study must focus on a life-threatening medical

condition for which available treatments are unproven or unsatisfactory and evaluate an

intervention that must be delivered within a time window that is too short for prospective, written

informed consent to occur. Studies conducted with EFIC are required to include additional

patient protections and mechanisms of addressing ethical principles that would normally be

addressed by prospective informed consent. Two important components of studies conducted

with EFIC are community consultation and public disclosure.

Community consultation occurs during study design and is a process of engaging

representatives of the communities from which patients will be recruited for the study.

Community consultation includes bidirectional communication between community members

and the research team through mechanisms like town hall meetings. It is intended to show

respect for persons while also providing an opportunity for communities affected by the research

to give meaningful input to investigators on the design of the study and to the institutional review

board (IRB) on the acceptability of the study. Community consultation, therefore, is deep

engagement with a narrow portion of the public. Public disclosure, by comparison, is intended to

disseminate information about the research to as large a population as possible through

channels like websites or advertisements. The primary goal of public disclosure is transparency,

and communication is generally one-way from researchers to the public.(1) In addition to the

pre-trial public disclosure activities, EFIC trials are also required to disseminate trial results

through post-trial public disclosure.

Federal regulations provide limited detail regarding how to conduct community

consultation and public disclosure for a trial conducted with EFIC.(2) In the 20 years following

the release of EFIC regulations, only 41 randomized trials were registered with the FDA with a

plan to use EFIC (approximately 2 trials per year).(3) Few of these trials have published

detailed information on the design and execution of their community consultation and public

disclosure. To add to the available literature on approaches to community consultation and

public disclosure in trials conducted with EFIC, this manuscript describes the community

consultation and public disclosure activities for the Randomized Trial of Sedative Choice for

Intubation (RSI) (NCT05277896), a randomized trial comparing ketamine versus etomidate for

emergency tracheal intubation that is being conducted with EFIC.(4) This is intended to provide

an example of a relatively cost-efficient, federated approach to community consultation and

public disclosure that attempts to match the scope, nature, and content of community

engagement to the risks and burdens of the research.

METHODS

The RSI trial

The RSI trial is an ongoing multicenter, pragmatic, randomized trial comparing the use of

ketamine versus etomidate for induction of anesthesia during emergency tracheal intubation of

critically ill adults in an emergency department or intensive care unit. The RSI trial is being

conducted by the Pragmatic Critical Care Research Group (PCCRG), a network of 25 health

systems across the U.S. that conducts comparative effectiveness trials focused on acute

care.(5) The RSI trial is designed to enroll 2,364 patients from 14 emergency departments or

intensive care units at six hospitals across five regions of the U.S. (Table 1). Enrollment began

in April of 2022 and is expected to be completed in August of 2025. Enrollment of the first 464

patients at Vanderbilt University Medical Center was funded by the NIH (K23HL153584) and

enrollment of the final 1,900 patients at 6 sites in 5 regions across the U.S. is funded by the

Patient-Centered Outcomes Research Institute (PCORI) (BPS-2022C3-30021).

Development of the Plan for Community Consultation and Public Disclosure

Development of the plan for community consultation and public disclosure for the RSI

trial began in 2020. The initial draft of the plan was informed by two community engagement

studios, one with patients and community members and one with clinicians. Community

engagement studios are an engagement approach developed by the Meharry-Vanderbilt

Community-Engaged Research Core in which a unique panel of 8-12 patients, community

members, or clinicians (selected for their firsthand knowledge of a particular condition) serve as

paid experts during a 2-hour face-to-face session with researchers.(6) In the initial community

engagement studios for the RSI trial, patients recommended (i) preferentially using social media

rather than traditional media formats (e.g., television and radio), (ii) collaborating with local

clinics including federally qualified health centers and faith-based clinics, and (iii) leveraging

local and national patient and caregiver support networks.

In consultation with the single IRB overseeing the trial, personnel at the coordinating

center developed a community consultation and public disclosure template that provided sites

with a menu of (i) required and (ii) optional activities (Table 2). This approach was intended to

provide structure to the community consultation and public disclosure processes while allowing

sites to tailor their activities to the unique context of their local communities.(7) Required

community consultation activities included face-to-face engagement with patients and

community members using structured surveys to solicit input on the acceptability of the

proposed RSI trial, which were modeled on the surveys used in prior EFIC trials.(8–11)

Required public disclosure activities included developing and maintaining a website for the trial

that provided lay language summaries of the trial design, how patients would be enrolled without

the ability to provide prospective informed consent, and how to obtain an "opt-out" medical

bracelet alert from trial personnel.

Development of Site-Specific Plans

Upon funding of the multi-center portion of the trial in 2023, development of site-specific

plans for community consultation and public disclosure at each participating site began.

Because sites had varying levels of experience with patient and community engagement, the

coordinating center staff included an expert in stakeholder engagement who provided

assistance to sites in 1) developing site-specific community consultation and public disclosure

plans, 2) developing recruitment materials for community consultation activities, 3) identifying

community members and groups to invite to town hall meetings and community engagement

studios, 4) coaching site investigators in presentation and communication strategies that should

be used when presenting potentially sensitive topics during meetings with community members,

and 5) collating community feedback for reporting to local and central institutional review

boards.

Approved plans set a minimum bar for planned activities while allowing flexibility for sites

to conduct additional activities in response to feedback received from the community during the

execution of the planned activities. The coordinating center provided sites with a structured

template to track and summarize their completed community consultation and public disclosure

activities.

RESULTS

Vanderbilt University Medical Center, which serves as both the coordinating center and

an enrolling site for the RSI trial, completed its community consultation and public disclosure

activities over a 12-month period from February 2021 to January 2022. Community consultation

and public disclosure activities at the five remaining sites occurred between September 2023

and June 2024. The activities are summarized below and full details are available on FDA public

docket 95S-0158.(12)

Community Consultation Results

Community consultation activities resulted in rich quantitative and qualitative data from a

diverse group of community members across the 6 trial sites. A total of 789 English-speaking or

Spanish-speaking patients or family members in emergency department or intensive care unit

waiting rooms completed face-to-face surveys with trial personnel. Demographics of

respondents are shown in Table 3. Approximately 96% of survey respondents agreed with the

importance of the RSI trial and 92% agreed with or were indifferent to the conduct of the study

without prospective informed consent. Detailed results of the surveys will be reported

separately.

More than 200 local groups were invited to attend community engagement studios or

town halls across the 6 trial sites in 5 regions. Invited groups included faith-based, government,

first responder, and disease-specific patient and family support groups (such as support groups

for patients with lung disease, hypertension, or stroke). A total of 9 community engagement

studios were held across 5 regions, with 82 attendees. These sessions were transcribed by site

personnel. Feedback on the trial was generally positive overall and was used to revise patient-

facing documents, revise messaging about the trial to participants and families, and inform the

way study coordinators were trained to approach and interact with participants and their

families. At a community engagement studio dedicated to soliciting input specifically from Black

and African American community members, participants highlighted that, given the history of

scientific research misconduct involving Black and African American communities, members of

the research team should deliberately and actively work to build trust and maintain transparency

with members of these communities.

Nine town hall meetings were conducted across 5 regions and included both in-person

and online engagement, with a total of 156 attendees. These sessions were transcribed by site

personnel. At one site, the patient investigator for the site (a survivor of emergency tracheal

intubation who serves on the Steering Committee for the RSI trial) attended the town hall to help

community members understand the experience of critical illness and the perspective of a

survivor. Attendees of the town halls expressed broad support for the trial at all sites. Attendees

also emphasized the importance of timely notification of enrollment in the trial and expressed a

desire that more research teams would disseminate information about ongoing research

activities within their communities. Selected quotes from the community engagement studios

and town halls are available in Table 4. In summary, a majority of consulted community

members across the five regions found the RSI trial to be addressing an important research

question and acceptable to conduct with EFIC.

Public Disclosure Results

All six sites used paid social media advertisements on Facebook (Meta) or Instagram to

target information about the RSI trial to the geographic catchment area of the participating

hospital. All sites posted advertisements in English; two sites also posted advertisements in

Spanish. Three sites posted information about the study on social media pages on Linkedin,

Twitter (X), Facebook (Meta), or NextDoor. Sites varied in terms of the institutional requirements

for marketing or press office involvement in finalizing the social media advertisements from the

template provided. Social media posts and advertisements included details of the study,

including the fact that patients would be enrolled without informed consent. Posts and

advertisements also included a link to the overall study website and a method of requesting opt-

out bracelets. The cost of the social media advertising ranged from $750-$2250 across the

sites. The cumulative cost of social media advertising across all trial sites was $8,291. Metrics

and costs for social media advertisements are shown in Table 5.

Flyers presenting information about the RSI trial were posted in English and Spanish by

five sites in emergency department, critical care, and community clinic settings. Two sites also

posted flyers in public spaces, such as local libraries and recreation centers, and mailed flyers

to local student and community groups.

Press releases were issued by each of the six hospitals, resulting in two local newspaper

articles at one site and one local news television segment at another. One site ran an

advertisement about the study in two local print newspapers with an additional advertisement in

the online version of one of the newspapers. The study website was available throughout the

public disclosure process. During the 6-month period of the most intense public disclosure

activities, from January to June 2024, the RSI trial webpage had 16,635 unique visitors and

35,557 page visits.

Effort and Cost to Conduct Community Consultation and Public Disclosure

The time spent by research teams and principal investigators for community consultation

varied considerably across the six sites. Based on self-reported metrics, research staff at sites

spent an average of 30 hours (range, 15-60 hours) conducting in-person surveys, an average of

5 hours (range, 0-15 hours) coordinating and hosting community engagement studios, and an

average of 22 hours (range 0-110 hours) preparing for and conducting town halls. Site principal

investigators spent an average of 2 hours conducting in-person surveys (range 0-10 hours), an

average of 25 hours (range, 3-120 hours) coordinating and hosting community engagement

studios and an average of 25 hours (range, 6-55 hours) coordinating and hosting town halls.

Cumulatively at all sites, we estimate that approximately 341 hours of research staff time and

310 hours of site principal investigator time were dedicated to completing community

consultation. Considering base salary, funder salary caps, fringe benefits, and the 40% indirect

rate set by the funder, the average hourly rate for a research coordinator was approximately $80

per hour and the average hourly rate for a physician site investigator was approximately $180

per hour. Using these estimates, the total cost of completing community consultation was

$83,080 (approximately $13,847 per site).

The time spent by research teams for public disclosure activities was more consistent

across sites than the time spent on community consultation. Based on self-reported metrics,

research staff spent an average of 8 hours (range, 0-30 hours) finalizing social media posts and

advertisements, an average of 8 hours (range, 2-20 hours) posting flyers, and an average of 2

hours (range,0-10 hours) finalizing press releases. Site principal investigators spent an average

of 7 hours (range, 3-10 hours) finalizing social media posts and advertisements, an average of 4

hours (range 2-10 hours) finalizing press releases, and a cumulative total of 4 hours preparing

for and conducting a TV segment. Cumulatively, the time spent on public disclosure across the

sites for the trial was estimated to include 112 hours of study coordinator time and 70 hours of

site principal investigator time. The estimated total personnel cost of completing public

disclosure was approximately $21,560. Combining the costs of social media advertising and the

time spent by research coordinators and physician site investigators, the estimated total cost of

performing public disclosure at all sites was $29,851 (approximately $4,975 per site).

In summary, the estimated total costs of completing community consultation and public

disclosure prior to beginning enrollment in the RSI trial was approximately $112,931

(approximately $18,822 per site). These estimates do not include the time spent by coordinating

center staff to develop the resources needed for public disclosure or the time spent by

coordinating center staff assisting sites with development and execution of their plans, each of

which were significant.

Beyond Community Consultation and Public Disclosure

While the regulations for EFIC require pre-trial community consultation and public

disclosure and post-trial public disclosure of trial results, they do not require or recommend

patient or community involvement during the conduct of the trial. The RSI trial, however, opted

to engage patients, families, and community members at all stages of the research, using a

previously published multilevel approach to patient and community engagement (Figure 1).(13)

First, trial leadership includes 6 patient investigators, one patient from a patient

advocacy group focused on survivors of acute respiratory distress syndrome and one patient

from each geographic area in which the trial is enrolling. Each patient investigator participates

as full voting members of the Steering Committee and provides input into all major trial

decisions. Second, the PCORI-funded Science, Technology and Research partnership (STAR)

Clinical Research Network (CRN) provides oversight of the trial through their Stakeholder

Advisory Council. The council consists of patients, community members, and clinicians who

review research studies to evaluate their person-centeredness and provide advice regarding

additional opportunities to improve trial conduct.(14) The STAR-CRN Stakeholder Advisory

Council has provided input and feedback on the RSI trial twice yearly. Third, when the trial

encounters an issue that would benefit from broader input from patients or community members

with lived experience in emergency tracheal intubation (e.g., development of consent forms,

amount of financial compensation for participants), trial leadership use the ResearchMatch

Expert Advice Tool(15) to survey a geographically representative sample of community

members with the relevant prior experience who have volunteered to be available for research

studies. Fourth, when the RSI trial needs deep input into a specific study decision (e.g., design

of the study's newsletter for participants), trial leadership convenes a community engagement

studio. Studios for the RSI trial have occurred thrice yearly and have addressed topics such as

best practices for notification of enrollment, best practice for approaching for consent, and

identification of subgroups of interest for subgroup analyses. All of the engagement activities are

overseen by the trial engagement coordinator, an expert with more than 20 years of experience

leading community engagement in research who provides critical input into early drafts of

patient-facing documents, identifies patients and community members for inclusion on the study

steering committee and community engagement groups, and leads training and engagement of

patient investigators in trial conduct.

DISCUSSION

361   Community consultation and public disclosure have been envisioned as methods of

362   addressing the ethical principles of consent in studies in which prospective informed consent is

363   infeasible. Although the regulations outlining the requirements for community consultation and

364   public disclosure in trials conducted under EFIC were published nearly 30 years ago,

365   uncertainty remains regarding the optimal methods and approaches for accomplishing these

366   activities.(1,3,8,10,16,17)

367   Few studies have reported the full details of their community consultation and public

disclosure activities or the associated financial cost, personnel time cost, and duration to

complete. Available publications of such metrics for community consultation and public

disclosure in a EFIC trials have estimated a cost of $50,000 to $60,000 per site and up to 3

years to complete.(18,19) By contrast, the estimated total fixed and personnel cost to complete

community consultation and public disclosure in the RSI trial was approximately $112,931

overall (approximately $18,822 per site), excluding time spent by coordinating center staff to

develop the resources and templates or assist sites with development and execution of their

plans. Compared to prior EFIC trials, personnel costs were similar, but the fixed costs of public

disclosure appear to have been significantly lower. This likely results from the decision to move

toward the use of social media for public disclosure as opposed to more costly, traditional forms

of media (e.g., newspaper and radio advertisements). This is consistent with a growing body of

literature on the effectiveness of a social media campaign as a method for public disclosure in

research.(9,18,20–25)

Some experts have expressed concerns that the cost and complexity of community

consultation and public disclosure have slowed the conduct of randomized trials in emergency

care(26). FDA guidance does suggest, however, that "IRBs may consider more or less

community consultation necessary, depending on the unique circumstances of a particular study

and the community," giving the example that "less extensive community consultation may be

appropriate for . . . . a study comparing an available treatment . . . with another product about

which much is known."(27) The relatively cost-efficient completion of community consultation

and public disclosure in the RSI trial represents an attempt by trial leadership to follow this

guidance, matching the extent of community consultation to the risks and burdens of

participating in a trial comparing standard-of-care treatments. Ketamine and etomidate are the

two most commonly used sedative medications during emergency tracheal intubation in many

settings in the United States, both have been received by millions of patients each year for

decades, the risk profiles of both medications are well known, and both medications are

approved by the FDA for the studied indication at the studied doses. While the extent of

community consultation and public disclosure in RSI were significant, reaching more than 1.2

million adults, they may not have been as intensive as prior trials evaluating novel,

experimental, unapproved drugs or devices with potentially greater risks.

We also found that the costs of community consultation and public disclosure were

driven largely by personnel costs rather than by fixed costs. Completion of community

consultation and public disclosure required an average of 139 hours of research coordinator and

physician site investigator time per site; a total of 833 hours across all sites. The time required

varied significantly across sites, ranging from 63 hours to 261 hours. Several factors may

explain this variability, including: site experience with community consultation and public

disclosure processes, availability of non-study team local support (e.g. press office, local

community engagement teams), variable reliance on the central engagement coordinator based

at the coordinating center, and community-specific differences in acceptability of the studied

interventions or the conduct of trials without prospective informed consent. Because the hours

devoted to each activity were collected retrospectively, recall bias may also have contributed to

the variability reported across sites.

Previous trials have taken different approaches to coordinating the completion of

community consultation and public disclosure activities at participating sites, including: (i) a

central conduct model(10) (site activities conducted centrally by the coordinating center), (ii) a

federated model(19) (in which plans are designed centrally with efforts to oversee and align

approaches while allowing local autonomy in planning and conduct), and (iii) a fully local

model(28) (in which all sites generate plans without significant guidance or oversight from the

coordinating center). The RSI trial used a federated model in which sites worked with their local

community and IRB to develop a plan for community consultation and public disclosure using

standardized templates developed by the coordinating center, which included required activities

and a menu of optional activities and associated minimum standards. We found this approach to

be successful in identifying and addressing community-specific concerns while facilitating

efficient completion. All community consultation and public disclosure activities during the

PCORI-funded multi-center stage of the trial were completed in nine months. The success of

this model was greatly facilitated by having an experienced, central engagement coordinator to

develop templates, identify community members and work with sites with less experience with

EFIC trials.(1,9)

LIMITATIONS

This article has several limitations. First, the pre-existing patient and community

engagement infrastructure available at some participating sites in the RSI trial likely improved

efficiency and decreased cost compared to sites that do not have pre-existing infrastructure,

such as many non-academic hospital systems. Second, community consultation and public

disclosure activities vary greatly in cost and effort, and decisions regarding which activities are

chosen for a given trial may be related to the trial's specific funding mechanism and budget.

Third, the extent of the patient and community engagement during trial conduct (not a

requirement of EFIC) is based, in part, on recommendations regarding engagement provided by

the funder, PCORI, and the resources required to conduct this level of engagement may not be

available in every EFIC trial. Fourth, the time and cost of community consultation and public

disclosure activities were estimated retrospectively, after completion of all activities, and recall

bias could have resulted in underestimation of the total effort and cost. Fifth, estimates do not

account for substantial effort from (i) non-research personnel at sites (e.g., media office

personnel); (ii) coordinating center personnel (e.g., trial primary investigator, project managers)

who led the design of templates, regulatory approval, oversite of site execution of activities, and

submission of completed activities to the IRB and FDA; and (iii) the engagement coordinator

who led the recruitment and preparation for local community engagement studios and town halls

at sites that did not manage these locally.

CONCLUSIONS

In preparation for conducting the RSI trial, study personnel completed community

consultation that included bidirectional interactions with over 1,000 patients, families, and

community members and public disclosure that reached over 1.2 million people living in the

communities in which the trial is being conducted. The use of digital channels of communication

for community consultation and public disclosure and an approach that matched the intensity of

engagement activities to the risk of the research resulted in a total cost and duration of activities

that were lower than reported in prior EFIC trials. However, these costs remained significant and

may inform which treatments are feasible to examine in trials with EFIC.

Community Consultation and Public Disclosure for the Randomized Trial of Sedative Choice for Intubation

REFERENCES

1. Tisherman SA. Defining "Community" and "Consultation" for Emergency Research that
Requires an Exception from Informed Consent. AMA J Ethics. 2018 May 1;20(5):467–74.

2. O'Malley GF, Giraldo P, Deitch K, Aguilera EA, Cadar S, Lares C, et al. A Novel Emergency
Department-based Community Notification Method for Clinical Research Without Consent.
Acad Emerg Med Off J Soc Acad Emerg Med. 2017 Jun;24(6):721–31.

3. Feldman WB, Hey SP, Kesselheim AS. A Systematic Review Of The Food And Drug
Administration's 'Exception From Informed Consent' Pathway. Health Aff (Millwood). 2018
Oct;37(10):1605–14.

4. DeMasi SC, Imhoff B, Lewis AA, Seitz KP, Driver BE, Gibbs KW, et al. Protocol and
Statistical Analysis Plan for a Multicenter Randomized Trial of Ketamine vs Etomidate for
Emergency Tracheal Intubation. CHEST Crit Care [Internet]. 2025 Sep;3(3). Available from:
https://www.chestcc.org/article/S2949-7884(25)00050-4/fulltext

5. About – Pragmatic Critical Care Research Group PCCRG [Internet]. [cited 2025 Jan 10].
Available from: https://www.pragmaticcriticalcare.org/about/

6. Joosten YA, Israel TL, Williams NA, Boone LR, Schlundt DG, Mouton CP, et al. Community
Engagement Studios: A Structured Approach to Obtaining Meaningful Input From
Stakeholders to Inform Research. Acad Med J Assoc Am Med Coll. 2015 Dec;90(12):1646–
50.

7. Carlson JN, Zive D, Griffiths D, Brown KN, Schmicker RH, Herren H, et al. Variations in the
application of exception from informed consent in a multicenter clinical trial. Resuscitation.
2019 Feb;135:1–5.

8. Dickert NW, Metz K, Fetters MD, Haggins AN, Harney DK, Speight CD, et al. Meeting
unique requirements: Community consultation and public disclosure for research in
emergency setting using exception from informed consent. Acad Emerg Med.
2021;28(10):1183–94.

9. Hsu CH, Fowler J, Cranford JA, Thomas MP, Neumar RW. Integration of social media with
targeted emails and in-person outreach for exception from informed consent community
consultation. Acad Emerg Med Off J Soc Acad Emerg Med. 2022 Feb;29(2):217–27.

10. Stephens SW, Carroll-Ledbetter C, Duckert S, Coffman T, Nelson M, Brown KN, et al.
Interactive Media-Based Approach for an Exception From Informed Consent Trial Involving
Patients With Trauma. JAMA Surg. 2024 Sep 1;159(9):1051–8.

11. Ward CE, Adelgais KM, Holsti M, Jacobsen KK, Simon HK, Morris CR, et al. Public support
for and concerns regarding pediatric dose optimization for seizures in emergency medical
services: An exception from informed consent (EFIC) trial. Acad Emerg Med Off J Soc Acad
Emerg Med. 2024 Jul;31(7):656–66.

12. Regulations.gov [Internet]. 2024 [cited 2025 Jan 10]. Public Disclosure from Vanderbilt
University Medical Center. Available from: https://www.regulations.gov/document/FDA-
1995-S-0036-0230

13. Boyer AP, Fair AM, Joosten YA, Dolor RJ, Williams NA, Sherden L, et al. A Multilevel
Approach to Stakeholder Engagement in the Formulation of a Clinical Data Research
Network. Med Care. 2018 Oct;56 Suppl 10 Suppl 1(10 Suppl 1):S22–6.

14. Wilkins CH, Villalta-Gil V, Houston MM, Joosten Y, Richmond A, Vaughn YC, et al.
Development and validation of the Person-Centeredness of Research Scale. J Comp Eff
Res. 2018 Dec;7(12):1153–9.

15. Tischbein M, Cook SK, Shyr C, Benhoff K, Baig A, Quarles K, et al. Expert advice from
ResearchMatch volunteers: Recruitment Innovation Center use cases and innovation. J Clin
Transl Sci. 2024;8(1):e223.

16. Gugel T, Adams K, Baranoski M, Yanez ND, Kampp M, Johnson T, et al. Design and
implementation of community consultation for research conducted under exception from
informed consent regulations for the PreVent and the PreVent 2 trials: Changes over time
and during the COVID-19 pandemic. Clin Trials Lond Engl. 2024 Dec;21(6):671–80.

17. Chisolm-Straker M, Nassisi D, Daya MR, Cook JNB, Wilets IF, Clesca C, et al. Exception
from Informed Consent: How IRB Reviewers Assess Community Consultation and Public
Disclosure. AJOB Empir Bioeth. 2021;12(1):24–32.

18. Matchett G, Ryan TJ, Sunna MC, Lee SC, Pepe PE. Measuring the cost and effect of
current community consultation and public disclosure techniques in emergency care
research. Resuscitation. 2018 Jul 1;128:37–42.

19. Silbergleit R, Biros MH, Harney D, Dickert N, Baren J. Implementation of the Exception from
Informed Consent Regulations in a Large Multicenter Emergency Clinical Trials Network;
the RAMPART Experience. Acad Emerg Med. 2012 Apr;19(4):448–54.

20. Stephens SW, Carroll-Ledbetter C, Duckert S, Coffman T, Nelson M, Brown KN, et al.
Interactive Media-Based Approach for an Exception From Informed Consent Trial Involving
Patients With Trauma. JAMA Surg. 2024 Sep 1;159(9):1051–8.

21. Stephens SW, Carroll-Ledbetter CR, Duckert S, Coffman TW, Nelson MA, Rodgers J, et al.
Analysis of time to regulatory approval in an exception from informed consent trial in trauma
patients. J Trauma Acute Care Surg. 2025 Mar;98(3):492.

22. Eubank L, Lee KS, Seder DB, Strout T, Darrow M, MacDonald C, et al. Approaches to
community consultation in exception from informed consent: Analysis of scope, efficiency,
and cost at two centers. Resuscitation. 2018 Sep;130:81–7.

23. Harvin JA, Podbielski JM, Vincent LE, Liang MK, Kao LS, Wade CE, et al. Impact of Social
Media on Community Consultation in Exception from Informed Consent Clinical Trials. J
Surg Res. 2019 Feb;234:65–71.

24. Farley P, Stephens SW, Crowley B, Collins SP, Wong MD, Panas AB, et al. Exception from
informed consent trials: social-media-based community consultation campaigns are
representative of target communities. Trauma Surg Acute Care Open. 2021 Nov
29;6(1):e000830.

25. Inam ME, Sanzgiri A, Lekka E, Sheth SA, Barreto AD, Savitz SI, et al. Exception from
informed consent in the era of social media: The SEGA stroke trial experience. Brain Circ.
2021;7(4):253–8.

26. Silbergleit R, Biros MH, Harney D, Dickert N, Baren J, NETT Investigators. Implementation
of the exception from informed consent regulations in a large multicenter emergency clinical
trials network: the RAMPART experience. Acad Emerg Med Off J Soc Acad Emerg Med.
2012 Apr;19(4):448–54.

27. Office of the Commissioner. Exception from Informed Consent Requirements for Emergency
Research [Internet]. FDA; 2013 [cited 2025 Jun 26]. Available from:
https://www.fda.gov/regulatory-information/search-fda-guidance-documents/exception-
informed-consent-requirements-emergency-research

28. Sloan EP, Koenigsberg M, Gens D, Cipolle M, Runge J, Mallory MN, et al. Diaspirin cross-
linked hemoglobin (DCLHb) in the treatment of severe traumatic hemorrhagic shock: a
randomized controlled efficacy trial. JAMA. 1999 Nov 17;282(19):1857–64.

TABLES

Table 1. Hospitals that are enrolling patients in the RSI trial.

Hospital	Location	Description
Vanderbilt University Medical Center	Nashville, TN	Tertiary care hospital that serves patients with complex medical illnesses and patients from rural communities.
Hennepin County Medical Center	Minneapolis, Minnesota	Public teaching hospital and Minnesota's largest safety-net hospital.
University of Alabama – Birmingham (UAB)	Birmingham, Alabama	Public teaching hospital and Alabama's largest hospital; serves rural communities state-wide.
University of Colorado -UCHealth University of Colorado Hospital	Aurora, Colorado	Large tertiary care hospital that is part of the largest health system in Colorado.
-Denver Health Medical Center Campus	Denver, Colorado	Urban safety net hospital in downtown Denver.
Atrium Health – Wake Forest Baptist	Winston-Salem, North Carolina	Large tertiary care hospital that is part of the largest health system in North Carolina.

584   **Table 2. Community Consultation and Public Disclosure Templates.**

Community Consultation Template*	
A (Required)	**B (Optional)**
Web-based survey (provided by coordinating center)	Town hall presentation and discussion by an investigator. Conducted by inviting community groups or visiting a meeting of an existing group
Community Engagement Studios	In-person individual interviews or meetings with community representatives
	In-person solicited survey e.g. waiting room (template provided by coordinating center)
Public Disclosure Plan Template[†]	
A (Required)	**B (Optional)**
Study website including details on obtaining opt-out medical alert bracelet	Paid online advertisements (banner ads purchased from Google, Facebook, Youtube, etc.)
Press release	Social media/internet postings (Youtube, Facebook, Twitter, etc.)
	Postings on website or social media groups for community members at increased risk of critical illness (e.g., patients following transplant, patients with lung disease, narcotics anonymous, nursing homes)
	Mailings (including email circulars/bursts and direct paper mailings)
	Outdoor advertising (placards, bus ads, billboards, etc.)
	Television and radio ads (broadcast advertising)
	Newspaper advertisement (and similar print advertising)
	Newsletters (articles or informational ads, print or electronic)
	Brochures, flyers, handouts, bulletin boards
	Radio or TV PSA (public service announcements)

[*] Each site was required to complete at least 4 total community consultation activities including
both activities in column A and at least 2 activities from column B
[†] Each site was required to complete at least 4 total public disclosure activities, including both
activities in column A, and at least 2 types from column B.

Table 3. Characteristics of the patients or family members who responded to in-person
surveys in ED or ICU waiting rooms.
591

Characteristic	Survey Cohort (N=789)
Age, median (IQR) – yr	44 (28, 60) (N=734)
Female sex – no. (%)	427/781 (54.7)
Race and ethnicity – no. (%)	
Non-Hispanic Black	199 (25.2)
Non-Hispanic White	364 (46.1)
Hispanic	147 (18.6)
Other	54 (6.8)
Prefer not to answer or missing	25 (3.2)
Highest Level of Education – no. (%)	
Less than 9th Grade	14/781 (1.8)
9-12th grade, no diploma	71/781 (9.1)
High school graduate	179/781 (22.9)
Technical or Vocational Degree	77/781 (9.9)
Some college credit, no degree	164/781 (21.0)
Bachelor's Degree	152/781 (19.5)
Graduate or Professional Degree	94/781 (12.0)
Prefer not to answer	30/781 (3.8)
Place of Residence– no. (%)	
Urban	293/780 (37.6)
Suburban	283/780 (36.3)
Rural	123/780 (15.8)
Homeless	15/780 (1.9)
Other	27/780 (3.5)
Prefer not to answer	39/780 (5.0)
Household Income– no. (%)	
< $10,000	81/782 (10.4)
$10,000-$25,000	99/782 (12.7)
$26,000-$50,000	131/782 (16.8)
$51,000-$75,000	131/782 (16.8)
> $75,000	184/782 (23.5)
Prefer not to answer	156/782 (20.0)

594 **Table 4. Selected Quotes from the Community Consultation Process**
595

Quotes from community engagement studios and town halls

Importance of the study
- "I wish this was something I had when I was in the ICU, or we had the results when I was in the ICU. I hope you can do more studies in the future about the side effects to see if they are long term."
- "I personally am encouraged because I want to know the results."
- "I'm surprised this study has not been done before because we use both meds every single day."
- "I'm shocked there is more than one medication the doctors can use."
- "I would love to know if one medication would be better for this than the other."
- "My father woke up early in the OR while still intubated and still experiences extreme stress by this, I support the need to find out which medication is better."

Acceptability of exception from informed consent
- "I don't think it would bother me to be told that I was given a particular medication. When I'm in the hospital for a procedure, I don't necessarily give consent for the use of a particular medication because I don't understand them anyway."
- "It boils down to a matter of saving my life. The medications are already tested, so the doctor's decision to use either medication would be okay with me. Hoping and pretty sure that the doctors are focused on saving a person's life—if they are concerned about taking care of me, I am not concerned with which medication is used."
- "As a mom who had a kid in the NICU, being at [RSI trial site] specifically, there was an awareness that it's a research hospital and all the things that may or may not happen with our data."
- "As long as the trial doesn't interfere with the rest of the patient care it seems OK."
- "Just telling people that you had to do it [because it's an emergency] and knowing this information could help people in a similar situation in the future."
- "People are afraid of research because they don't understand it, but we know it's important for us to support this kind of thing."

Patient notification
- "The sooner we start [notifying] the better, but it has to be with a lot of transparency… Be truthful and transparent".

Sharing of research results
- "I've participated in research before, and it was really important to me when I got the results. Communicating with the people who have been part of the study is really important. As an African American, I would like to know how the medications affect African Americans, so communication should specifically point out the effect on people of my race. This is important and is often overlooked."
- "A lot of stigmas have been put on people of lower income, etc. because research has been done on populations without consent in the past. In emergency situations, the doctor has to do what they do, but people might want to know what they did…"
- "You have to be transparent with patients so patients feel like this is ethical."
- "The trust line is broken with minority communities so there has to be a lot of

transparency."
- "It's important for researchers to share what you are doing with us so we can be more aware."

598   **Table 5. Details of Social Media Campaign used for Public Disclosure**
599

Trial site	Social media used for advertisement (length posted)	Metrics	Cost
Site 1	Facebook (2 months)	• Viewed by 176,100 people • 8,000 people clicked the link	$2,000
Site 2	Facebook (6 weeks)	• Viewed by 121,752 people • 584 people interacted with the ad • 532 people clicked the link	$1000
Site 3	Facebook and Instagram (8 days)	English • Viewed by 194,389 people • 144 people interacted with the ad • 136 people clicked the link Spanish • Viewed by 106,505 people • 79 people interacted with the ad • 73 people clicked the link	$800
Site 4	Facebook and Instagram (1 month)	English and Spanish • Viewed by 113,256 people • 898 people clicked the link	$1491
Site 5	Facebook and Instagram (1 month)	• Viewed by 45,769 people • 955 people interacted with the ad • 875 people clicked the link	$750
Site 6	Facebook (2 weeks)	• Viewed by 454,158 people	$2250
Total	---	**1,211,929 views**	**$8,291**

608 **Figure 1. Multilevel approach to stakeholder engagement**

609 Reprinted from DeMasi SC, Imhoff B, Lewis AA, Seitz KP, Driver BE, Gibbs KW, et al. Protocol
610 and Statistical Analysis Plan for a Multicenter Randomized Trial of Ketamine vs Etomidate for
611 Emergency Tracheal Intubation. CHEST Crit Care [Internet]. 2025 Sep;3(3). Available from:
612 https://www.chestcc.org/article/S2949-7884(25)00050-4/fulltext, with permission from Elsevier.

Level of Stakeholder Engagement	Frequency	Research Activity	
Co-Investigators 6 patients & 5 clinicians	Monthly Steering Committee Meetings	Identify research question Refine research question	Year 1 Pre-Award
Stakeholder Advisory Council 15 patients, caregivers, community members, & health system leaders	Semi-annual Advisory Council Meetings	Select outcomes Finalize trial protocol Draft patient materials	Years 2-4
Community Engagement Studios 82 patients, caregivers, community members, & clinicians	Meet with stakeholders every 3-4 months	Advise trial implementation Advise retention Inform analysis plan	
Surveys and Town Halls 945 patients, caregivers, community members, & clinicians	Directed, intermittent in-person and online surveys & town halls	Interpret results Disseminate findings Implement findings	Year 5