Title: Contribution of moderation foods to total energy intake by sociodemographic

characteristics in U.S. adults: National Health and Nutrition Examination Survey, 2017 – 2020

Leah M. Lipsky, Amara J. Channell Doig, Tonja R. Nansel

Social and Behavioral Sciences Branch, Division of Population Health Research, *Eunice*

Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, MD.

Correspondence: Leah Lipsky, 6710B Rockledge Dr MSC 7004, Bethesda MD 20817, 301-

435-6951, lipskylm@mail.nih.gov

Funding: This work was funded in part by the Intramural Research Program of the *Eunice*

Kennedy Shriver National Institute of Child Health and Human Development. The funder had no

involvement or restrictions regarding publication.

Abbreviations:

HEI – Healthy Eating Index

IPR – Income-Poverty Eatio

NHANES – National Health and Nutrition Examination Survey

NCHS – National Center for Health Statistics

Abstract

Background: U.S. adults exceed recommended intakes of multiple nutrients of concern.

Objective: This study uses a recently-developed food-level classification method for identifying

foods exceeding thresholds for nutrients of concern to investigate sociodemographic differences

in their contribution to total energy intake among U.S. adults.

Design: Cross-sectional.

Participants/setting: Participants 20 – 80 years of age in the 2017–2020 National Health and

Nutrition Examination Survey (n=7706).

Main outcome measures: Data from 24-hour dietary recalls were used to identify "moderation

foods" meeting at least one threshold for high content of nutrients of concern: added sugar >20%

energy, sodium >460mg per serving, refined grains >50% of total grains or >10:1 ratio of

carbohydrate to fiber content, saturated fat >20% energy, total fat >9% by weight (applied only

to vegetables, sweets, and snacks), and all alcoholic beverages. The percentage of energy intake

from moderation foods in total and from foods meeting each of the moderation food thresholds

were calculated and compared across age, education, income-to-poverty ratio (IPR), and

race/ethnicity.

Statistical analyses performed: Linear regression models with multiplicative interaction terms

tested whether race/ethnicity or education modified relationships of IPR with moderation food

intake.

Results: Moderation foods contributed 76% (95%CI:75%-77%) of energy intake overall. Intake

was higher in younger adults (mean=79%) versus older (72%-75%) adults, males (77%) versus

females (74%), and adults in the lowest (79%) vs. highest (74%) income quartile. Moderation

food intake was lowest in Non-Hispanic Asian adults (66%) and among adults with the highest

education (mean=71%). Foods meeting thresholds for sodium, refined grains, and fat contributed

the most to overall energy intake. Foods meeting thresholds for added sugar, sodium, and refined

grains contributed a smaller share of energy intake in non-Hispanic Asians versus other

racial/ethnic groups and in older versus younger adults. Income was inversely associated with

moderation food intake only among high school graduates (second highest education group; β = -

1.1 95%CI = -2.1 to -0.2) and modestly inversely associated with moderation food intake in all

race/ethnic groups except Mexican Americans/Other Hispanic adults.

Conclusions: High intake of moderation foods across all subgroups of U.S. adults, with only

modest between-group differences in the contributions of different categories of moderation

foods, suggest a need for population-wide changes in food selection to align with dietary

guidelines. In particular, interventions and policies to assist in identifying foods high in sodium,

refined grains, and fat may help consumers improve food choices.

Introduction

The 2020-2025 Dietary Guidelines for Americans (DGA) recommend limiting intake of added sugar, saturated fat, sodium, refined grains, and alcoholic beverages ("moderation components") due to their associations with multiple adverse health outcomes and their tendency to displace intake of foods that provide essential nutrients[1,2]. However, most Americans exceed recommended maximum intakes of refined grains (>90%), added sugar (70%), saturated fat (70%), and sodium (90%), contributing to the high public health burden of diet-related chronic diseases[1].

While sociodemographic differences in overall diet quality have been reported in multiple studies, these differences are generally modest in magnitude. For example, 2018 – 2018 NHANES data indicate relatively small mean differences (statistical comparisons not provided) in Healthy Eating Index-2015 scores (an indicator of conformance to the 2015-2020 Dietary Guidelines for Americans) between females (mean HEI=60) and males (mean HEI=56), non-Hispanic Asian (mean HEI=65) and Hispanic (mean HEI=58) adults versus non-Hispanic white (mean HEI=57) and non-Hispanic black (mean HEI=54) adults, and in adults with household income greater than 350% of the federal poverty level (mean HEI=60) versus those with incomes less than 350% of the federal poverty level (mean HEI=56)[3]. Education is also positively associated with better diet quality[4-8], and some evidence suggests that associations of income with diet quality differ by education and race/ethnicity[8,9]. In the U.S., educational attainment is positively correlated with income[10], and both income and education differ by race/ethnicity[11,12]. However, few previous studies on sociodemographic correlates of diet quality have examined the potential for confounding or interactions between these variables.

Additionally, research on sociodemographic differences in intake of moderation dietary

components has focused on individual nutrients or food groups (such as sugar-sweetened

beverages or snack foods)[13-16], ultra-processed foods (based on processing rather than nutritional

components)[4,17,18], or "junk"[19] or "fast"[20] foods (using multiple definitions and typically applied

only to select food groups). Given the lack of a guidelines-based definition that can be applied to

all foods in the Food and Nutrient Database for Dietary Studies (FNDDS), we previously

developed a method to identify moderation foods based on evidence-based thresholds for added

sugar, sodium, fat, refined grains, and alcohol. Although some moderation foods may also

provide beneficial nutrients, initial validation analyses indicated that the method effectively

captures foods that are disproportionately high in nutrients of concern (by distinguishing between

foods with low versus high nutrient density) and that greater intake of moderation foods is

associated with lower overall adherence to dietary guidelines in U.S. adults (r = -0.72, p<0.0001

with Healthy Eating Index-2020 scores)[21].

Examining the distribution of moderation food intake across population subgroups using a

systematic, guidelines-based indictor will help clarify how moderation food contributes to

socioeconomic differences in overall diet quality and identify subgroups who may benefit most

from targeted interventions. This study aimed to compare moderation food intake across

sociodemographic characteristics among U.S. adults and investigate whether education or

race/ethnicity modifies associations of income with moderation food intake.

Methods

Study design and population

This study used publicly available data from the pre-pandemic cycle (2017 – 2020) of the
National Health and Nutrition Examination Survey (NHANES), administered by the National
Center for Health Statistics (NCHS). NHANES is a population-based, cross-sectional survey that
uses a complex, stratified, multistage probability sample design to generate a representative
sample of the civilian, non-institutionalized U.S. population. Details on the study design,
protocol, and data collection have been documented elsewhere[22]. The protocols are approved by
the NCHS Research Ethics Review Board, and informed consent was obtained from participants
≥ 18 years of age.

The analytic sample for this study included adults at least 20 years old with at least one 24-hour
dietary recall determined to be reliable by NCHS[23], and data on at least one of the
sociodemographic variables. Of the n = 9332 total adult participants, n = 7706 were included in
the final analytic data set. This secondary analysis of deidentified publicly available data was
exempt from institutional review board review.

Data collection

Participants self-reported sociodemographic characteristics during an in-home interview
administered by trained personnel. Dietary intake data come from the What We Eat In America
dietary interview component of NHANES; trained interviewers conducted two 24-hour dietary
recalls using the USDA's Automated Multiple-Pass Method[24], with the first recall administered
in-person at the mobile exam center and the second by telephone approximately 3 – 10 days after
the initial recall.

Moderation food intake

Foods reported in the 24-hour diet recalls were classified as moderation foods if they met any of

the previously established criteria, described in detail elsewhere[21]. Briefly, component thresholds

included: added sugar >20% of energy; sodium >460 mg per serving; refined grains >50% of

total grains (ounce equivalents) and >10:1 carbohydrate: fiber ratio (g); saturated fat >20% of

energy (excluding vegetables); total fat >9% of energy (applied only to vegetables, sweets, and

snacks), and all alcoholic beverages[21]. Foods that did not exceed any threshold were classified as

non-moderation. Total moderation food intake (% total energy intake, kcal) was calculated from

the sum of energy intake from moderation food relative to total energy intake across both diet

recalls. The percent of energy intake (% kcal) from foods meeting each of the individual

moderation food component thresholds was also calculated.

Sociodemographic characteristics

NHANES categorizes age as 20 – 39 y, 40 – 59 y, and ≥ 60 y[25]. Race/ethnicity is categorized as

Mexican American/Hispanic, non-Hispanic white, non-Hispanic black, non-Hispanic Asian, and

multi/other. Education categories include less than high school, high school degree/general

equivalency diploma, some college, college graduate or above. Income poverty ratio (IPR)[26] is a

continuous measure that reflects household income relative to the federal poverty threshold,

accounting for household size and composition. NHANES truncates the maximum IPR at 5 due

to disclosure concerns; IPR quartiles were used for analysis.

Statistical analysis

All analyses used survey methods to account for the complex sampling design of NHANES

using StataSE version 18. Frequencies and percentages were calculated for sociodemographic

characteristics, and means and 95% confidence intervals of moderation food intake (% kcal) and

intake (% kcal) from foods meeting each of the moderation food thresholds, were calculated for

each sociodemographic subgroup. Differences in mean moderation food intake by

sociodemographic characteristics were evaluated using linear regressions and post-estimation

pairwise comparisons; Sidak adjustment was used to account for multiple comparisons. Separate

models with multiplicative interaction terms were used to investigate whether education or

race/ethnicity modified associations of IPR (examined as a continuous variable) with moderation

food intake.

Results

The sample was approximately evenly split according to age group and sex (**Table 1**). Sixty-

three percent of the sample was non-Hispanic white. Ten percent of the weighted sample earned

less than a high school degree, and the remainder was approximately evenly split between

education groups. IPR was disproportionately concentrated at the highest truncated value (5),

with the mean $\pm$ SD of the upper quartile = 4.9$\pm$0.1.

Mean moderation food intake (% kcal) in the full sample was 76% (95% CI = 75% - 77%).

Intakes from foods meeting thresholds for sodium, refined grains, and fat were 12 to 31

percentage points higher than that from foods meeting thresholds for sugar and alcohol

(**Supplemental Table 1**).

Moderation food intake by sociodemographic characteristics are presented in **Figures 1 – 5** and

Supplemental Table 1. Mean moderation food intake varied from 79% (95%CI: 78% - 80%) in

adults 20 – 39 years of age to 72% (95%CI: 71% - 73%) in adults $\geq$ 60 years of age (p < 0.001

for all pairwise comparisons) (**Figure 1a**). Mean moderation food intake was 4.1% kcal (SE =

0.7% kcal) lower in 40 – 59-year-olds versus 20 – 39-year-olds (p<0.001), 7.2% kcal (SE = 0.6%

kcal) lower in those ≥ 60 years-old versus 20 – 39-year-olds (p<0.001), and 3.1% kcal (SE =

0.7% kcal) lower in those ≥ 60 years-old versus 40 – 59 years old (p = 0.01) (Figure 1a). **Figure**

1b and Supplemental Table 1 indicate that these differences were primarily attributable to

differences in intakes of foods meeting thresholds for sodium (3 – 9 percentage points higher in

middle aged and younger adults versus older adults) and refined grains (2 to 6 percentage points

higher in middle aged and young adults versus older adults).

Mean moderation food intake was 3.2% kcal higher (SE=0.6% kcal) (p < 0.001) in males (77%,

95%CI: 76% - 78%) than females (74%, 95%CI: 73% - 75%) (**Figure 2**). These differences were

primarily attributable to differences in intakes of foods meeting thresholds for sodium (four

percentage points higher in males than females) and refined grains (two percentage points higher

in males than females) (**Figure 2b** and Supplemental Table 1).

When comparing by race/ethnicity (**Figure 3**), moderation food intake was lowest in non-

Hispanic Asian adults (66%, 95%CI: 64% - 69%) and highest in multi/other (80%, 95%CI: 77%

- 82%) and non-Hispanic black adults (79%, 95%CI: 77% - 80%). Moderation food intake was

73% (95%CI: 72% - 75%) in Mexican American/Hispanic adults and 76% (95%CI: 75% - 78%)

in non-Hispanic white adults. While differences in mean moderation food intake of multi/other

as compared to non-Hispanic white (p=0.08) and black (p=0.998) adults were not statistically

different, all other pairwise comparisons were statistically different (p<0.02). The relatively low

moderation food intake in non-Hispanic Asians was due to lower intake of foods exceeding

thresholds for refined grains, added sugar, sodium, and fat (**Figure 3b** and Supplemental Table

1). Mexican American/Hispanic adults also had relatively low intakes of refined grains, added

sugar, and fat, but consumed a relatively higher amount of foods meeting the sodium threshold.

Moderation food intake was lower in adults with at least a college degree (71%, 95%CI: 70% -

73%) than in adults with less education (p < 0.001), but mean intake did not significantly differ

across the three lower education groups (75% - 79%) (**Figure 4a**). Mean moderation food intake

in adults with the highest education was 4.2% kcal (SE=0.9% kcal) lower than in those with less

than a high school degree (p=0.008), 7.5% kcal (SE=1.1% kcal) lower than those with a high

school degree only (p<0.001), and 6.1% kcal (SE=0.8% kcal) lower than those with some

college education (p<0.001). Mean moderation food intake in adults with a high school degree

was 3.3% (SE=0.9%) higher than in those with less than a high school degree (p=0.008). **Figure**

4b and Supplemental Table 1 indicate that the lower intake of moderation foods in the highest

education group was due primarily to lower intake of foods meeting the refined grains threshold

(31% versus 35% - 38% kcal). Group mean intakes from foods meeting other component

thresholds were within four percentage points.

Mean moderation food intake was inversely associated with IPR quartile (**Figure 5a**), with a 4.6-

point difference between the 1st (79%, 95%CI: 77% - 80%) and 4th quartile (74%, 95%CI: 72% -

76%) (p=0.005. However, only the pairwise differences between the 1st and 3rd quartiles (76%,

95%CI: 74% - 78%) and 1st and 4th quartiles were statistically significant (p = 0.02 and 0.005,

respectively). These differences were primarily due to the relatively lower intake in the highest

IPR quartile of foods meeting the added sugar threshold (17% versus 20% for the 2nd and 3rd

quartile and 22% kcal for the lowest quartile) and the refined grains threshold (31% versus 35% -

38% in the lower quartiles) (**Figure 5b** and Supplemental Table 1).

There was no statistically significant interaction overall between IPR and race/ethnicity in

relation to moderation food intake (F(4, 22) = 1.56, Prob > F = 0.22). However, the coefficient

estimate for non-Hispanic Black adults was significantly more negative than that for Mexican

American/Hispanic adults (**Table 2**). **Figure 6** shows that the estimated marginal slopes were

negative for all groups, but the relationship was not statistically significant among Mexican

Americans/Hispanic adults or those categorized as multi/other.

There was no statistically significant interaction between IPR and education in relation to

moderation food intake overall ($F(3, 23) = 2.6$, Prob $> F = 0.08$) (**Table 2**). **Figure 7** shows a null

association of income with moderation food intake in all education groups, except for an inverse

association in high school graduates.

Discussion

Mean moderation food intake ranged from 70% - 80% of energy intake across most segments of

the U.S. adult population in 2017 – 2020. While some statistically significant differences were

attributable to sex, age, income, race/ethnicity, and education, all subgroups obtained the

majority of energy intake from moderation food, and the only subgroup with mean moderation

food intake less than 70% kcal was non-Hispanic Asian adults (66%). Associations of income

with moderation food intake did not vary by race or education, and moderation food intake

differed more by race/ethnicity and education than by income. In the full sample, foods meeting

thresholds for sodium, refined grains, and fat contributed more to total energy intake than foods

meeting thresholds for added sugar or alcohol. Sociodemographic differences in overall

moderation food intake were primarily due to differences in intakes of foods meeting thresholds

for refined grains, sodium, and added sugar, with fewer differences in intakes of foods meeting

thresholds for fat or alcohol.

These findings are consistent with previous studies that reported small but statistically significant

socioeconomic disparities in adult diet quality[4,18,27-32]. In those studies, diet quality scores across

all subgroups were far from optimal, with mean group differences ranging from <1% to 9%.

Reports have often emphasized statistically significant differences while giving less attention to

their small magnitudes, potentially contributing to the misconception that low diet quality affects

only specific population subgroups. In contrast, our findings indicate that suboptimal dietary

patterns are widespread, regardless of sex, income, race/ethnicity, or education, despite some

statistically significant differences. While the DGA recommend that 85% of daily energy intake

come from nutrient-dense foods, leaving only 15% for added sugars, fats, or alcohol [1], our

findings suggest that dietary behaviors across all sociodemographic subgroups of U.S. adults are

essentially reversed, with moderation food accounting for at least two-thirds of energy intake and

non-moderation foods contributing only 20% to 30%. The magnitudes of subgroup differences

were smallest by income-poverty ratio (1 – 4 points) and sex (3 points), and largest by age (8

points between the youngest and oldest adults) and race/ethnicity (7 – 13 points between non-

Hispanic Asians versus other groups). The magnitude of the differences in intakes from foods

meeting each moderation food component threshold were also generally less than 5 percentage

points.

The consistently high moderation food intake observed across different sociodemographic

subgroups suggests that sociodemographic factors may not be primary drivers of low diet quality

in U.S. adults. This aligns with findings from the Consumer Expenditure Survey, which

demonstrate that, across all income groups, the largest share of food-at-home spending (35%)

goes to miscellaneous foods (e.g. frozen meals, snacks, canned and packaged soups, potato

chips, condiments, seasonings), while 20% - 25% goes to meat, poultry, fish, and eggs; and 19%

to fruits and vegetables[33]. These purchasing patterns suggest that households may prioritize

spending on foods for convenience, social, cultural, emotional, or other purposes[34-36], rather than

adherence to dietary guidelines. Additionally, while food accessibility has been identified as a

barrier to healthy eating for underserved groups, national data indicate that 6.1% of the U.S.

population lives in areas with limited supermarket access, with 1.7% of households being both

far from a food store and without vehicle access[37].

The DGA's focus on nutrients rather than foods to limit may have inadvertently weakened its

effectiveness in discouraging intake of nutrients of concern in the U.S population. Since the

1970s, guidelines related to decreasing intakes have provided nutrient-based rather than food-

based recommendations - a practice influenced by industry pressures[38] and enabled by the closed

nature of DGA meetings prior to the 2010 – 2015 cycle. Nutrient-based guidelines are more

difficult for consumers to understand and implement than food-based recommendations[39], partly

since such guidelines require viewing and correctly interpreting nutrition labels that present

nutrients per serving, which consumers must then accurately extrapolate to their total daily

intake. Consequently, consumers may be unaware when intake of nutrients of concern exceeds

recommended amounts. Although the shift to public DGA Committee meetings beginning with

the 2010 – 2015 cycle has offered the opportunity to incorporate food-based recommendations

for moderation components, the nutrient-based approach established during previous closed

meetings has persisted. Another potential reason for the lack of current food-based guidelines to

limit nutrients of concern may involve assertions, largely propagated by the food industry, that

communicating health risks of specific foods inflicts psychological harm (e.g., "food

shaming")[40,41]. In contrast, empirical evidence shows that interpretive front-of-package labels

improve consumer food choices without increasing stigma[42-44]. It will remain important to assess

and monitor adverse psychological effects of any labels on at-risk groups, such as individuals

with eating disorders, given limited evidence.

By contrast, the Chilean government adopted thresholds to classify packaged foods high in kcal,

sugar, sodium, or saturated fat, and applied clear front-of-package labels along with restrictions

on marketing to children and school sales. These policies significantly reduced consumer

purchases and industry supply of high-sugar and high-sodium products, while also decreasing

unhealthy marketing to children, limiting availability in schools, and improving consumers'

ability to identify and avoid unhealthy foods[45-50]. Collectively, this evidence demonstrates how

policies that influence food environments can affect dietary behaviors, even though individual

food choices are motivated by complex social, cultural, and emotional factors. Furthermore, it

illustrates how regulatory approaches that use systematic food classification based on nutrient

thresholds can drive meaningful changes in the food environment and serve as a powerful tool to

improve diet quality and reduce chronic disease risk at the population level.

Several elements of this study strengthen the internal and external validity of the findings. The

study used data from a large, nationally representative sample of the U.S. population that

included administration of two dietary recalls using the Automated Multi-Pass Method. One

limitation is that the use of self-reported dietary intake is susceptible to reporting bias that may

weaken internal validity. However, we combined four years of NHANES data and used analytic

methods to obtain reliable and generalizable estimates of overall intake of moderation food for

multiple sociodemographic subgroups. We also examined potential confounding and effect

modification between correlated characteristics (income, education, and race/ethnicity), which

provides more valid estimates than bivariate analyses. As this was an observational study, it was

descriptive rather than hypothesizing causal relationships.

Conclusion

Moderation food intake is high in U.S. adults across multiple sociodemographic characteristics,

indicating an urgent, population-wide need to replace moderation food with core foods that are

aligned with nutritional requirements. Foods high in sodium, refined grains, and fat were the

largest contributors to overall energy intake. The moderation food classification method may

facilitate moving towards more explicit, food-based guidance and policy changes that modify the

food environment to improve diet quality and chronic disease outcomes at the population level.

Acknowledgments

References

1.   U.S. Department of Agriculture and U.S. Department of Health and Human Services.

Dietary Guidelines for Americans, 2020-2025. 2020. December. Available at

DietaryGuidelines.gov

2.   Collaborators GBDD. Health effects of dietary risks in 195 countries, 1990-2017: a

systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. May 11

2019;393(10184):1958-1972. doi:10.1016/S0140-6736(19)30041-8

3.   U.S. Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy

and Promotion. Average Healthy Eating Index-2020 Scores for the U.S. Population - Total Ages

2 and Older and by Age Groups, WWEIA, NHANES 2017-2018. https://fns-

prod.azureedge.us/cnpp/hei-scores-americans

4.   Baraldi LG, Martinez Steele E, Canella DS, Monteiro CA. Consumption of ultra-

processed foods and associated sociodemographic factors in the USA between 2007 and 2012:

evidence from a nationally representative cross-sectional study. *BMJ Open*. Mar 9

2018;8(3):e020574. doi:10.1136/bmjopen-2017-020574

5.   Beydoun MA, Wang Y. Do nutrition knowledge and beliefs modify the association of

socio-economic factors and diet quality among US adults? *Prev Med*. Feb 2008;46(2):145-53.

doi:10.1016/j.ypmed.2007.06.016

6.   Hiza HA, Casavale KO, Guenther PM, Davis CA. Diet quality of Americans differs by

age, sex, race/ethnicity, income, and education level. *J Acad Nutr Diet*. Feb 2013;113(2):297-

306. doi:10.1016/j.jand.2012.08.011

7.   Lee E, Fitzgerald N, Downs S, et al. Misperception of diet quality among US adults:

implications for cardiometabolic health promotion. *Eur J Clin Nutr*. Aug 2025;79(8):762-769.

doi:10.1038/s41430-025-01605-1

8.   McCullough ML, Chantaprasopsuk S, Islami F, et al. Association of Socioeconomic and

Geographic Factors With Diet Quality in US Adults. *JAMA Netw Open*. Jun 1

2022;5(6):e2216406. doi:10.1001/jamanetworkopen.2022.16406

9.   Rippin HL, Hutchinson J, Greenwood DC, et al. Inequalities in education and national

income are associated with poorer diet: Pooled analysis of individual participant data across 12

European countries. *PLoS One*. 2020;15(5):e0232447. doi:10.1371/journal.pone.0232447

10.   Guzman G, Kollar M. *Income in the United States: 2022*. 2023. *Current Population*

Reports. P60-279. September.

https://www.census.gov/content/dam/Census/library/publications/2023/demo/p60-279.pdf

11.   Chetty R, Friedman JN, Saez E, Turner N, Yagan D. Income Segregation and

Intergenerational Mobility across Colleges in the United States. *Quarterly Journal of Economics*.

Aug 2020;135(3):1567-1633. doi:10.1093/qje/qjaa005

12.   Kim JH, Soler MC, Zhao Z, Swirsky E. *Race and ethnicity in higher education: 2024*

status report. 2024.

13.   DiFrancesco L, Fulgoni VL, 3rd, Gaine PC, Scott MO, Ricciuto L. Trends in added

sugars intake and sources among U.S. adults using the National Health and Nutrition

Examination Survey (NHANES) 2001-2018. *Front Nutr*. 2022;9:897952.

doi:10.3389/fnut.2022.897952

14.   Dunford EK, Popkin BM. Disparities in Snacking Trends in US Adults over a 35 Year

Period from 1977 to 2012. *Nutrients*. Jul 27 2017;9(8):809. doi:10.3390/nu9080809

15.   Lee SH, Zhao L, Park S, et al. High Added Sugars Intake among US Adults:

Characteristics, Eating Occasions, and Top Sources, 2015-2018. *Nutrients*. Jan 4 2023;15(2):265.

doi:10.3390/nu15020265

16.   Lundeen EA, Park S, Pan L, Blanck HM. Daily Intake of Sugar-Sweetened Beverages

Among US Adults in 9 States, by State and Sociodemographic and Behavioral Characteristics,

2016. *Prev Chronic Dis*. Dec 13 2018;15:E154. doi:10.5888/pcd15.180335

17.   Juul F, Parekh N, Martinez-Steele E, Monteiro CA, Chang VW. Ultra-processed food

consumption among US adults from 2001 to 2018. *Am J Clin Nutr*. Jan 11 2022;115(1):211-221.

doi:10.1093/ajcn/nqab305

18.   Pozveh SH, Aktary ML, Polsky JY, Moubarac JC, Vanderlee L, Olstad DL.

Socioeconomic Inequalities in Intakes of Ultraprocessed and Minimally Processed Foods in

Nationally Representative Samples of Adults in Canada: An Analysis of Trends between 2004

and 2015. *J Nutr*. Oct 2024;154(10):3088-3104. doi:10.1016/j.tjnut.2024.07.029

19.   Dunford EK, Popkin B, Ng SW. Junk Food Intake Among Adults in the United States. *J*

Nutr. Feb 8 2022;152(2):492-500. doi:10.1093/jn/nxab205

20.   Fryar CD, Hughes JP, Herrick KA, Ahluwalia N. Fast Food Consumption Among Adults

in the United States, 2013-2016. *NCHS Data Brief*. Oct 2018;(322):1-8.

21.   Channell Doig AJ, Lipsky LM, Choe A, Nansel TR. Development and evaluation of a

nutrient-based method of classifying moderation foods. *medRxiv*. 2025:2025.03.20.25324317.

doi:10.1101/2025.03.20.25324317

22.   Akinbam L, Chen T-C, Davy O, et al. *National Health and Nutrition Examination Survey,*

2017–March 2020 prepandemic file: Sample design, estimation, and analytic guidelines. Vol.

190. 2022. *Vital Health Stat 2*.

23.   Group FSR. Key points using What We Eat In America, NHANES 2017-2018.

Agricultural Research Service, USDA. Accessed 07/01/2025, 2025.

24.   Blanton CA, Moshfegh AJ, Baer DJ, Kretsch MJ. The USDA Automated Multiple-Pass

Method accurately estimates group total energy and nutrient intake. *J Nutr*. Oct

2006;136(10):2594-9. doi:10.1093/jn/136.10.2594

25.   Centers for Disease Control and Prevention, National Center for Health Statistics. 2017-

March 2020 Pre-Pandemic Demographics Data - Continuous NHANES. Accessed 07/02/2025,

2025.

https://wwwn.cdc.gov/nchs/nhanes/search/datapage.aspx?Component=Demographics&Cycle=20

17-2020

26.   2 National Health and Nutrition Examination Survey: Analytic guidelines, 1999-2010

(National Center for Health Statistics) (2013).

27.   Choi YJ, Crimmins EM, Kim JK, Ailshire JA. Food and nutrient intake and diet quality

among older Americans. *Public Health Nutr*. May 2021;24(7):1638-1647.

doi:10.1017/S1368980021000586

28.   Guenther PM, Juan W, Lino M, Carlson A, Hiza HH, Krebs-Smith SM. *Diet quality of*

low-income and higher income Americans in 2003-04 as measured by the Healthy Eating Index-

2005. Vol. 42. 2008. *Nutrition Insight*. December.

29.   Parker HW, Tovar A, McCurdy K, Vadiveloo M. Socio-economic and racial prenatal diet

quality disparities in a national US sample. *Public Health Nutr*. Apr 2020;23(5):894-903.

doi:10.1017/S1368980019003240

30.   Raffensperger S, Kuczmarski MF, Hotchkiss L, Cotugna N, Evans MK, Zonderman AB.

Effect of race and predictors of socioeconomic status on diet quality in the HANDLS Study

sample. *J Natl Med Assoc*. Oct 2010;102(10):923-30. doi:10.1016/s0027-9684(15)30711-2

31.   Rehm CD, Penalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults,

1999-2012. *JAMA*. Jun 21 2016;315(23):2542-53. doi:10.1001/jama.2016.7491

32.   Wang DD, Leung CW, Li Y, et al. Trends in dietary quality among adults in the United

States, 1999 through 2010. *JAMA Intern Med*. Oct 2014;174(10):1587-95.

doi:10.1001/jamainternmed.2014.3422

33.   Carlson A, Frazao E. Food costs, diet quality and energy balance in the United States.

Physiol Behav. Jul 2014;134:20-31. doi:10.1016/j.physbeh.2014.03.001

34.   Foods US. The Diner Dispatch: 2024 American Dining Habits. Accessed 04/16/2025,

2025. https://www.usfoods.com/our-services/business-trends/american-dining-out-habits-

2024.html

35.   Raskind IG, Woodruff RC, Ballard D, et al. Decision-making processes shaping the home

food environments of young adult women with and without children. *Appetite*. Jun 1

2017;113:124-133. doi:10.1016/j.appet.2017.02.027

36.   Goorin J, Venezky E. How Dining Out Differs Across Generations. Eater. Accessed

04/16/2025, 2025. https://www.eater.com/press-room/24388888/future-generations-restaurant-

dining-survey-research

37.   Rhone A, Williams R, Dicken C. *Low-Income and Low-Foodstore-Access Census Tracts,*

2015–19. 2022. *Economic Information Bulletin*.

38.   Nestle M. Food lobbies, the food pyramid, and U.S. nutrition policy. *Int J Health Serv*.

1993;23(3):483-96. doi:10.2190/32F2-2PFB-MEG7-8HPU

39.   Tapsell LC, Neale EP, Satija A, Hu FB. Foods, Nutrients, and Dietary Patterns:

Interconnections and Implications for Dietary Guidelines. *Adv Nutr*. May 2016;7(3):445-54.

doi:10.3945/an.115.011718

40.   Musicus AA, Moran AJ. Front-of-Package Labeling and Stigma. *JAMA Netw Open*. Jun 2

2025;8(6):e2516835. doi:10.1001/jamanetworkopen.2025.16835

41.   Mialon M, Mialon J. Analysis of corporate political activity strategies of the food

industry: evidence from France. *Public Health Nutr*. Dec 2018;21(18):3407-3421.

doi:10.1017/S1368980018001763

42.   D'Angelo Campos A, Grummon AH, Ng SW, Puhl RM, Golden SD, Hall MG. Front-of-

Package Food Labels and Perceived Weight Stigmatization: A Randomized Clinical Trial. *JAMA*

Netw Open. Jun 2 2025;8(6):e2516821. doi:10.1001/jamanetworkopen.2025.16821

43.   Grummon AH, Musicus AA, Moran AJ, Salvia MG, Rimm EB. Consumer Reactions to

Positive and Negative Front-of-Package Food Labels. *Am J Prev Med*. Jan 2023;64(1):86-95.

doi:10.1016/j.amepre.2022.08.014

44.   Grummon AH, Gibson LA, Musicus AA, Stephens-Shields AJ, Hua SV, Roberto CA.

Effects of 4 Interpretive Front-of-Package Labeling Systems on Hypothetical Beverage and

Snack Selections: A Randomized Clinical Trial. *JAMA Netw Open*. Sep 5 2023;6(9):e2333515.

doi:10.1001/jamanetworkopen.2023.33515

45.   Rebolledo N, Ferrer-Rosende P, Reyes M, Smith Taillie L, Corvalan C. Changes in the

critical nutrient content of packaged foods and beverages after the full implementation of the

Chilean Food Labelling and Advertising Law: a repeated cross-sectional study. *BMC Med*. Jan

27 2025;23(1):46. doi:10.1186/s12916-025-03878-6

46.   Reyes M, Smith Taillie L, Popkin B, Kanter R, Vandevijvere S, Corvalan C. Changes in

the amount of nutrient of packaged foods and beverages after the initial implementation of the

Chilean Law of Food Labelling and Advertising: A nonexperimental prospective study. *PLoS*

Med. Jul 2020;17(7):e1003220. doi:10.1371/journal.pmed.1003220

47.   Taillie LS, Bercholz M, Popkin B, Reyes M, Colchero MA, Corvalan C. Changes in food

purchases after the Chilean policies on food labelling, marketing, and sales in schools: a before

and after study. *Lancet Planet Health*. Aug 2021;5(8):e526-e533. doi:10.1016/S2542-

5196(21)00172-8

48.   Dillman Carpentier FR, Correa T, Reyes M, Taillie LS. Evaluating the impact of Chile's

marketing regulation of unhealthy foods and beverages: pre-school and adolescent children's

changes in exposure to food advertising on television. *Public Health Nutr*. Mar 2020;23(4):747-

755. doi:10.1017/S1368980019003355

49.   Massri C, Sutherland S, Kallestal C, Pena S. Impact of the Food-Labeling and

Advertising Law Banning Competitive Food and Beverages in Chilean Public Schools, 2014-

2016. *Am J Public Health*. Sep 2019;109(9):1249-1254. doi:10.2105/AJPH.2019.305159

50.   Taillie LS, Reyes M, Colchero MA, Popkin B, Corvalan C. An evaluation of Chile's Law

of Food Labeling and Advertising on sugar-sweetened beverage purchases from 2015 to 2017: A

before-and-after study. *PLoS Med*. Feb 2020;17(2):e1003015.

doi:10.1371/journal.pmed.1003015

Table 1. Sample characteristics

Characteristic	Adults (≥ 20 y) n = 6640	
	N	%
Age, years		
20 – 39	1998	36
40 – 59	2212	34
60+	2429	30
Sex		
Male	3169	48
Female	3470	52
Race/ethnicity		
Mexican American/Hispanic	1392	16
Non-Hispanic White	2398	63
Non-Hispanic Black	1866	11
Non-Hispanic Asian	666	6
Multi/other	317	4
Education		
Some high school	1079	9
High school graduate	1568	27
Some college	2260	30
College graduate or more	1725	34
Income poverty ratio quartile		

1 (mean $\pm$ SD = 0.7 $\pm$ 0.4)	1475	17
2 (mean $\pm$ SD = 1.8 $\pm$ 0.3)	1462	19
3 (mean $\pm$ SD = 3.3 $\pm$ 0.5)	1457	27
4 (mean $\pm$ SD = 4.9 $\pm$ 0.1)	1464	37

Table 2. Coefficient estimates from regression models estimating the interactions of income poverty ratio (IPR) with race/ethnicity and education on moderation food intake (% kcal) in U.S. adults.

Model	Independent variable	β	95% CI	*P*-value
Race/ethnicity	IPR	-0.3	-1.4 – 1.0	0.70
	Race/ethnicity			
	Mexican American/Hispanic		REF	
	Non-Hispanic White	7.2	2.9 – 11.4	0.002
	Non-Hispanic Black	9.2	5.6 – 12.8	<0.001
	Non-Hispanic Asian	-1.3	-5.0 – 2.5	0.50
	Other/multi	11.9	5.4 – 18.4	0.001
	Interaction of IPR with race/ethnicity			
	Mexican American/Hispanic		REF	
	Non-Hispanic White	-1.2	-2.6 – 0.3	0.11
	Non-Hispanic Black	-1.5	-2.9 - -0.2	0.03
	Non-Hispanic Asian	-1.5	-2.5 - -0.4	0.01
	Other/multi	-1.8	-4.1 – 0.5	0.13
	Intercept	73.9	70.5 – 77.2	<0.001
Education	IPR	0.1	-1.1 – 1.3	0.84
	Education			
	Less than high school	6.2	1.3 – 11.2	0.02
	High school graduate	11.7	7.0 – 16.5	<0.001
	Some college	7.9	2.9 – 12.8	0.003

College graduate or more		REF	
Interaction of IPR with education			
Less than high school degree	-0.8	-2.7 – 1.1	0.42
High school graduate	-1.9	-3.3 - -0.5	0.01
Some college	-0.6	-1.9 – 0.8	0.41
College graduate or more		REF	
Intercept	71.1	66.4 – 75.8	<0.001

IPR – income-poverty ratio