Title: The contribution of moderation foods to total energy intake by sociodemographic

characteristics in U.S. adults: National Health and Nutrition Examination Survey, 2017 –

2020

Leah M. Lipsky, Amara Channel-Doig, Tonja R. Nansel

Social and Behavioral Sciences Branch, Division of Population Health Research, *Eunice*

Kennedy Shriver National Institute of Child Health and Human Development, Bethesda,

MD.

Correspondence: Leah Lipsky, 6710B Rockledge Dr MSC 7004, Bethesda MD 20817,

301-435-6951, lipskylm@mail.nih.gov

Funding: This work was funded in part by the Intramural Research Program of the

Eunice Kennedy Shriver National Institute of Child Health and Human Development.

The funder had no involvement or restrictions regarding publication.

Abbreviations:

HEI – Healthy Eating Index

IPR – Income-Poverty Eatio

NHANES – National Health and Nutrition Examination Survey

NCHS – National Center for Health Statistics

Abstract

Background: Intake of foods high in nutrients of concern (moderation foods) contribute a large proportion of total energy intake in the U.S, but little is known about how moderation food intake varies across sociodemographic subgroups.

Objective: This observational study assessed sociodemographic differences in the contribution of moderation food to energy intake in U.S. adults.

Methods: Data from 24-hour dietary recalls of participants 20 – 80 years of age in the 2017–2020 National Health and Nutrition Examination Survey (n=7706) were used to calculate the percent of energy intake from foods meeting any of the following criteria: added sugar >20% energy, sodium >460mg per serving, refined grains >50% of total grains or >10:1 ratio of carbohydrate to fiber content, saturated fat >20% energy, total fat >9% by weight (applied only to vegetables, sweets, and snacks), and all alcoholic beverages. Moderation food intake was compared by age, sex, education, income poverty ratio (IPR), and race/ethnicity, and linear regressions with multiplicative interaction terms investigated whether race/ethnicity or education modified relationships of IPR with moderation food intake.

Results: Mean moderation food contributed 75% (95%CI:75%-77%) of energy intake in U.S. adults, and was higher in younger (mean=79%) vs. older (mean=72%-75%) adults, males (mean=77%) vs. females (mean=74%), and adults in the lowest (mean=79%) vs. highest (mean=74%) income quartile. Moderation food intake was lowest in Non-Hispanic Asian adults (mean=66%) versus other race/ethnicity groups (m=73% - 80%) and in the highest educated (mean=71%) vs. less-educated adults (m=75%-79%). The relationship of income with moderation food intake did not differ by race/ethnicity

overall, but the relationship was significantly more negative for non-Hispanic Black

adults than in Mexican American/Hispanic adults. Similarly, while the relationship of

income with moderation food intake did not differ overall by education, marginal slopes

estimates indicated an inverse association among high school graduates (second

highest education group; β = -1.1 95%CI = -2.1 to -0.2) and null associations for all

other adults.

Conclusions: High moderation food intake across all U.S. adult subgroups suggests a

need for population-wide changes in food selection to meet dietary guidelines.

Introduction

The 2020-2025 Dietary Guidelines for Americans (DGA) recommend limiting intake of added sugar, saturated fat, sodium, refined grains, and alcoholic beverages ("moderation components") due to their associations with multiple adverse health outcomes and their tendency to displace intake of foods that provide essential nutrients (1, 2). However, most Americans exceed recommended maximum intakes of refined grains (>90%), added sugar (70%), saturated fat (70%), and sodium (90%) contributing to widespread increased risk for chronic diseases (1).

Sociodemographic differences in overall diet quality have been reported in multiple studies. Analyses of recent NHANES data show modestly higher overall diet quality (as indicated by Healthy Eating Index-2015 scores) in females (mean HEI=60) versus males (mean HEI=56), non-Hispanic Asian (mean HEI=65) and Hispanic (mean HEI=58) adults versus non-Hispanic white (mean HEI=57) and non-Hispanic black (mean HEI=54) adults, and in adults with household income greater than 350% of the federal poverty level (mean HEI=60) versus those with incomes less than 350% of the federal poverty level (mean HEI=56) (3). Education is also positively associated with better overall diet quality (4-8), and some evidence suggests that associations of income with diet quality differ by education and race/ethnicity (8, 9). In the U.S., educational attainment is positively correlated with income (10), and both income and education differ by race/ethnicity (11, 12). However, few previous studies on sociodemographic correlates of diet quality have examined the potential for confounding or interactions between these variables.

Research on sociodemographic differences in intake of moderation dietary components

has focused on individual nutrients or food groups (such as sugar-sweetened

beverages or snack foods) (13-16), ultra-processed foods (based on processing rather

than nutritional components) (4, 17, 18), or "junk" (19) or "fast" (20) foods defined in

multiple ways and applied only to select food groups. Given the lack of a guidelines-

based definition for foods high in nutrients of concern that can be applied to all foods in

the Food and Nutrient Database for Dietary Studies (FNDDS), we previously developed

a method to identify moderation foods based on thresholds for added sugar, sodium, fat,

refined grains, and alcohol, which effectively distinguished between foods with high

versus low nutrient density (21). Examining the distribution of moderation food intake

across population subgroups using a systematic, guidelines-based indictor will help

clarify how moderation food contributes to socioeconomic differences in overall diet

quality and identify subgroups who may benefit most from targeted interventions. This

study aimed to compare moderation food intake across sociodemographic

characteristics among U.S. adults and investigate whether education or race/ethnicity

modifies associations of income with moderation food intake.

Methods

<u>Study design and population</u>

This study used publicly available data from the pre-pandemic cycle (2017 – 2020) of

the National Health and Nutrition Examination Survey (NHANES), administered by the

National Center for Health Statistics (NCHS). NHANES is a population-based, cross-

sectional survey that uses a complex, stratified, multistage probability sample design to

generate a representative sample of the civilian, non-institutionalized U.S. population.

Details on the study design, protocol, and data collection have been documented

elsewhere (22). The protocol was approved by the NCHS Research Ethics Review

Board, and informed consent was obtained from participants ≥ 18 years of age.

The analytic sample for this study included adults at least 20 years old with at least one

24-hour dietary recall determined to be reliable by NCHS (23), and data on at least one

of the sociodemographic variables. Of the n = 9,232 total adult participants, n = 688 had

no dietary recall data and n = 837 had no valid recalls. The final analytic data set

included n = 7707 participants. This secondary analysis of deidentified publicly available

data was exempt from institutional review board review.

Data collection

Participants self-reported sociodemographic characteristics during an in-home interview

administered by trained personnel. Dietary intake data come from the What We Eat In

America dietary interview component of NHANES; trained interviewers conducted two

24-hour dietary recalls using the USDA's Automated Multiple-Pass Method (24), with the

first recall administered in-person at the mobile exam center and the second by

telephone approximately 3 – 10 days after the initial recall.

Moderation food intake

Foods reported in the 24-hour diet recalls were classified as moderation foods if they

met any of the previously established criteria: added sugar >20% of energy; sodium

>460mg per serving; refined grains >50% of total grains and >10:1 carbohydrate: fiber

ratio by weight; saturated fat >20% of energy (excluding vegetables); total fat >9% by

weight (applied only to vegetables, sweets, and snacks), and all alcoholic beverages

(25). Foods that did not meet any criteria were classified as non-moderation. Total

moderation food intake (% kcal) was calculated from the sum of energy intake from

moderation foods relative to total energy intake across both diet recalls.

<u>Sociodemographic characteristics</u>

NHANES categorizes age as 20 – 39 y, 40 – 59 y, and ≥ 60 y (26). Race/ethnicity is

categorized as Mexican American/Hispanic, non-Hispanic white, non-Hispanic black,

non-Hispanic Asian, and multi/other. Education is categorized as less than high school,

high school degree/general equivalency diploma, some college, college graduate or

above. Income poverty ratio (IPR) (27) is a continuous measure that reflects household

income relative to the federal poverty threshold, accounting for household size and

composition. NHANES truncates the maximum IPR at 5 due to disclosure concerns;

IPR quartiles were used for analysis.

<u>Statistical analysis</u>

All analyses used survey methods to account for the complex sampling design of

NHANES using StataSE version 18. Frequencies and percentages were calculated for

sociodemographic characteristics, and means and 95% confidence intervals of

moderation food intake (% kcal) were calculated for each sociodemographic subgroup.

Differences in mean moderation food intake by sociodemographic characteristics were

evaluated using linear regressions and post-estimation pairwise comparisons; Sidak

adjustment was used to account for multiple comparisons. We used separate models

with multiplicative interaction terms to investigate whether education or race/ethnicity

modified associations of IPR (examined as a continuous variable) with moderation food

intake.

Results

The sample was approximately evenly split according to age group and sex (**Table 1**).

Sixty-three percent of the sample was non-Hispanic white. Ten percent of the weighted

sample earned less than a high school degree, and the rest of the sample was

approximately evenly split between education groups. IPR was disproportionately

concentrated at the highest truncated value (5), with the mean ± SD of the upper

quartile = 4.9±0.1.

Mean moderation food intake for all adults was 75% (95% CI = 75% - 77%). **Figures 1**

– 5 show mean (95% CI) moderation food intake (% kcal) by sociodemographic

characteristics. Pairwise comparisons of mean moderation food intake by

sociodemographic characteristics are shown in **Table 2**. Group mean differences were

generally < 5 percentage points. The largest mean differences (>10 points) were

observed between race/ethnic groups, particularly when comparing non-Hispanic Asian

adults versus non-Hispanic white, non-Hispanic black, and other/multi adults. Pairwise

comparisons between other race/ethnic groups, income-poverty ratio quartiles, and

between sexes were <5 points. Comparisons between age and education groups were

< 10 points.

Mean moderation food intake decreased with age, ranging from 79% (95%CI: 78% -

80%) in adults 20 – 39 years of age to 72% (95%CI: 71% - 73%) in adults ≥ 60 years of

age ($p < 0.001$ for all pairwise comparisons), and was higher in males than females

(Figure 2, Table 2).

Mean moderation food intake was lower in non-Hispanic Asian adults (mean=66%,

95%CI: 64% - 69%) than adults in other racial/ethnic groups and highest in multi/other

(mean=80%, 95%CI: 77% - 82%) and non-Hispanic black adults (mean=79%, 95%CI:

77% - 80%) (Figure 3, Table 2). Moderation food intake was lower in Mexican

American/Hispanic adults (mean=73%, 95%CI: 72% - 75%) than in non-Hispanic white

adults (mean=76%, 95%CI: 75% - 78%). Mean moderation food intake in adults with

multi/other race/ethnicity was not statistically different than in non-Hispanic white

(p=0.08) or black (p=0.998) adults. All other pairwise differences by race/ethnicity were

statistically significant (p≤0.02).

Mean moderation food intake in adults with at least a college degree was lower than in

those with less education (Figure 4, Table 2). Mean moderation food intake in adults

with some college education was not different from in those with a high school degree

or less (Figure 4, Table 2).

Mean moderation food intake was inversely associated with IPR quartile (Figure 5,

Table 2). However, only the pairwise differences between the 1st and 3rd (76%, 95%CI:

74% - 78%) quartiles and 1st and 4th quartiles were statistically significant.

Income was inversely associated with moderation food intake in all race/ethnic groups

except Mexican American/Hispanic and other/multi adults (Figure 6). Results from the

adjusted Wald test ($F(4, 22) = 1.56$, Prob > F = 0.22) indicated no statistically significant

interaction overall between IPR and race/ethnicity in relation to moderation food intake.

However, the coefficient estimate for non-Hispanic Black adults was significantly more

negative than that for Mexican American/Hispanic adults (**Table 3**).

Income was inversely associated with moderation food intake in high school graduates

only (Figure 7). There was no statistically significant overall interaction between IPR and

education in relation to moderation food intake ($F_{(3, 23)} = 2.6$, Prob > F = 0.08) (Table

3). The association was modestly more negative in high school graduates, but the

estimate was not statistically different from the referent group.

Discussion

Mean moderation food intake ranged from 70% - 80% of energy intake across most

segments of the U.S. adult population in 2017 – 2020. While some statistically

significant differences were attributable to sex, age, income, race/ethnicity, and

education, all subgroups obtained the majority of energy intake from moderation food,

and the only subgroup with mean moderation food intake less than 70% kcal was non-

Hispanic Asian adults (66%). Moderation food intake differed more by race/ethnicity and

education than other socioemographic characteristics, and associations of income with

moderation food intake did not vary by race or education.

Our findings are consistent with previous studies that reported small but statistically

significant socioeconomic disparities in adult diet quality (4, 18, 28-33). In those studies,

diet quality scores across all subgroups were far from optimal, with mean group

differences ranging from <1% to 9%. Reports have often emphasized statistically

significant differences while giving less attention to their small magnitudes, potentially

contributing to the misconception that low diet quality affects only specific population

subgroups. In contrast, these findings indicate that suboptimal dietary patterns are

widespread, regardless of sex, income, race/ethnicity, or education, despite some

statistically significant differences. While the DGA recommend that 85% of daily energy

intake come from nutrient-dense foods, leaving only 15% for added sugars, fats, or

alcohol (1), these findings suggest that dietary behaviors across all sociodemographic

subgroups of U.S. adults are essentially reversed, with moderation food accounting for

at least two-thirds of energy intake and non-moderation foods contributing only 20% to

30%. The magnitudes of subgroup differences were smallest by income-poverty ratio (1

– 4 points) and sex (3 points), and largest by age (8 points between the youngest and

oldest adults) and race/ethnicity (7 – 13 points between non-Hispanic Asians versus

other groups).

The consistently high moderation food intake observed across different

sociodemographic subgroups suggests that sociodemographic factors may not be

primary drivers of low diet quality in U.S. adults. This aligns with findings from the

Consumer Expenditure Survey, which demonstrate that, across all income groups, the

largest share of food-at-home spending (35%) goes to miscellaneous foods (e.g. frozen

meals, snacks, canned and packaged soups, potato chips, condiments, seasonings),

while only 20% - 25% goes to meat, poultry, fish, and eggs; and only 19% to fruits and

vegetables (34). These purchasing patterns suggest that households may prioritize

spending on foods for convenience, social, cultural, emotional, or other purposes (35-

37), rather than adherence to dietary guidelines. Additionally, while food accessibility is

a barrier to healthy eating, national data indicate that 6.1% of the U.S. population lives

in areas with limited supermarket access, with only 1.7% of households being both far

from a foodstore and without vehicle access (38).

The DGA focus on nutrients rather than foods to limit may have inadvertently weakened

its effectiveness in discouraging intake of moderation foods in the U.S population. Since

the 1970s, guidelines related to decreasing intakes have provided nutrient-based rather

than food-based recommendations, a practice influenced by industry pressures (39) and

enabled by the closed nature of DGA meetings prior to the 2010 – 2015 cycle. Nutrient-

based guidelines are more difficult for consumers to understand and implement than

food-based recommendations (40), partly since nutrient-based guidelines require

viewing and correctly interpreting nutrition labels that present nutrients per serving,

which consumers must then accurately extrapolate to their total daily intake.

Consequently, consumers may be unaware when intake of nutrients of concern exceeds

recommended amounts. Although the shift to public DGA Committee meetings

beginning with the 2010 – 2015 cycle has offered the opportunity to incorporate food-

based recommendations for moderation components, the nutrient-based approach

established during previous closed meetings has persisted.

By contrast, the Chilean government adopted thresholds to classify packaged foods

high in kcal, sugar, sodium, or saturated fat, and applied clear front-of-package labels

along with restrictions on school food and marketing to children. These policies

significantly reduced consumer purchases and industry supply of high-sugar and high-

sodium products, while also decreasing unhealthy marketing to children, limiting

availability in schools, and improving consumers' ability to identify and avoid unhealthy

foods (41-46). Collectively, this evidence demonstrates how, even though individual

food choices are motivated by complex social, cultural, and emotional factors, policies

that influence food environments can affect dietary behaviors. Furthermore, it illustrates

how regulatory approaches that use systematic food classification based on nutrient

thresholds can drive meaningful changes in the food environment and serve as a

powerful tool to improve diet quality and reduce chronic disease risk at the population

level.

Several elements of this study strengthen the internal and external validity of the

findings. The study used data from a large, nationally representative sample of the U.S.

population that included administration of two dietary recalls using the Automated Multi-

Pass Method. One limitation is that the use of self-reported dietary intake is susceptible

to reporting bias that may weaken internal validity. However, we combined four years of

NHANES data and used analytic methods to obtain reliable and generalizable estimates

of overall intake of moderation food for multiple sociodemographic subgroups. We also

examined potential confounding and effect modification between correlated

characteristics (income, education, and race/ethnicity), which provides more valid

estimates than bivariate analyses. As this was an observational study, it was descriptive

rather than hypothesizing causal relationships.

Conclusion

Moderation food intake is high in U.S. adults across multiple sociodemographic

characteristics, indicating an urgent, population-wide need to replace moderation food

with core foods that are aligned with nutritional requirements. The moderation food

classification method may facilitate moving towards more explicit, food-based guidance

and policy changes that modify the food environment to improve diet quality and chronic

disease outcomes at the population level.

Acknowledgments

LL- designed research, analyzed data, wrote paper, had primary responsibility for final

content; ACD – designed research, wrote paper; TRN – designed research, wrote

paper. All authors have read and approved the manuscript.

Declaration of generative AI and AI-assisted technologies in the writing process:

During the preparation of this work, LL used ChatGPT [https://chat.openai.com/] and

Perplexity [https://perplexity.ai] to assist in improving the readability and clarity of this

manuscript. After using these tools, LL reviewed and edited the content and takes full

responsibility for the content of the publication.

References

1.   U.S. Department of Agriculture and U.S. Department of Health and Human Services. Dietary Guidelines for Americans, 2020-2025. 9th Edition ed. Washington, DC, 2020.

2.   G. B. D. Diet Collaborators. Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet. 2019;393(10184):1958-72. doi: 10.1016/S0140-6736(19)30041-8.

3.   U.S. Department of Agriculture, Food and Nutrition Service, Center for Nutrition Policy and Promotion. *Average Healthy Eating Index-2020 Scores for the U.S. Population - Total Ages 2 and Older and by Age Groups, WWEIA, NHANES 2017-2018* [Internet]. 2023; Available from: https://fns-prod.azureedge.us/cnpp/hei-scores-americans.

4.   Baraldi LG, Martinez Steele E, Canella DS, Monteiro CA. Consumption of ultra-processed foods and associated sociodemographic factors in the USA between 2007 and 2012: evidence from a nationally representative cross-sectional study. BMJ Open. 2018;8(3):e020574. doi: 10.1136/bmjopen-2017-020574.

5.   Beydoun MA, Wang Y. Do nutrition knowledge and beliefs modify the association of socio-economic factors and diet quality among US adults? Prev Med. 2008;46(2):145-53. doi: 10.1016/j.ypmed.2007.06.016.

6.   Hiza HA, Casavale KO, Guenther PM, Davis CA. Diet quality of Americans differs by age, sex, race/ethnicity, income, and education level. J Acad Nutr Diet. 2013;113(2):297-306. doi: 10.1016/j.jand.2012.08.011.

7.   Lee E, Fitzgerald N, Downs S, Kuhrt N, Visaria A, Kalbag A, Setoguchi S. Misperception of diet quality among US adults: implications for cardiometabolic health promotion. Eur J Clin Nutr. 2025. doi: 10.1038/s41430-025-01605-1.

8.   McCullough ML, Chantaprasopsuk S, Islami F, Rees-Punia E, Um CY, Wang Y, Leach CR, Sullivan KR, Patel AV. Association of Socioeconomic and Geographic Factors With Diet Quality in US Adults. JAMA Netw Open. 2022;5(6):e2216406. doi: 10.1001/jamanetworkopen.2022.16406.

9.   Rippin HL, Hutchinson J, Greenwood DC, Jewell J, Breda JJ, Martin A, et al.

Inequalities in education and national income are associated with poorer diet:

Pooled analysis of individual participant data across 12 European countries.

PLoS One. 2020;15(5):e0232447. doi: 10.1371/journal.pone.0232447.

10.   Guzman G, Kollar M. Income in the United States: 2022. In: Bureau USC, ed.

Current Population Reports. Washington, D.C.: U.S. Government Publishing

Office, 2023.

11.   Chetty R, Friedman JN, Saez E, Turner N, Yagan D. Income Segregation and

Intergenerational Mobility Across Colleges in the United States*. The Quarterly

Journal of Economics. 2020;135(3):1567-633. doi: 10.1093/qje/qjaa005.

12.   Kim JH, Soler MC, Zhao Z, Swirsky E. Race and ethnicity in higher education:

2024 status report. In: Education ACo, ed., 2024.

13.   DiFrancesco L, Fulgoni VL, 3rd, Gaine PC, Scott MO, Ricciuto L. Trends in

added sugars intake and sources among U.S. adults using the National Health

and Nutrition Examination Survey (NHANES) 2001-2018. Frontiers in nutrition.

2022;9:897952. doi: 10.3389/fnut.2022.897952.

14.   Dunford E, Popkin B. Disparities in Snacking Trends in US Adults over a 35 Year

Period from 1977 to 2012. Nutrients. 2017;9(8):809. doi: 10.3390/nu9080809.

15.   Lee SH, Zhao L, Park S, Moore LV, Hamner HC, Galuska DA, Blanck HM. High

Added Sugars Intake among US Adults: Characteristics, Eating Occasions, and

Top Sources, 2015–2018. Nutrients. 2023;15(2):265. doi: 10.3390/nu15020265.

16.   Lundeen EA, Park S, Pan L, Blanck HM. Daily Intake of Sugar-Sweetened

Beverages Among US Adults in 9 States, by State and Sociodemographic and

Behavioral Characteristics, 2016. Prev Chronic Dis. 2018;15:E154. doi:

10.5888/pcd15.180335.

17.   Juul F, Parekh N, Martinez-Steele E, Monteiro CA, Chang VW. Ultra-processed

food consumption among US adults from 2001 to 2018. Am J Clin Nutr.

2022;115(1):211-21. doi: 10.1093/ajcn/nqab305.

18.   Pozveh SH, Aktary ML, Polsky JY, Moubarac JC, Vanderlee L, Olstad DL.

Socioeconomic Inequalities in Intakes of Ultraprocessed and Minimally

Processed Foods in Nationally Representative Samples of Adults in Canada: An

Analysis of Trends between 2004 and 2015. J Nutr. 2024;154(10):3088-104. doi:
10.1016/j.tjnut.2024.07.029.

19.   Dunford EK, Popkin B, Ng SW. Junk Food Intake Among Adults in the United
States. J Nutr. 2022;152(2):492-500. doi: 10.1093/jn/nxab205.

20.   Fryar CD, Hughes JP, Herrick KA, Ahluwalia N. Fast Food Consumption Among
Adults in the United States, 2013-2016. NCHS Data Brief. 2018(322):1-8.

21.   Channell Doig AJ, Lipsky LM, Choe A, Nansel TR. Development and evaluation
of a nutrient-based method of classifying moderation foods. medRxiv.
2025:2025.03.20.25324317. doi: 10.1101/2025.03.20.25324317.

22.   Akinbam L, Chen T-C, Davy O, Ogden C, Fink S, Clark J, Riddles M, Mohadjer L.
National Health and Nutrition Examination Survey, 2017–March 2020
prepandemic file: Sample design, estimation, and analytic guidelines. In: National
Center for Health Statistics, ed. Vital Health Stat 2, 2022.

23.   Group FSR. *Key points using What We Eat In America, NHANES 2017-2018*
[Internet]. Beltsville Human Nutrition Research Center, Agricultural Research
Service, USDA; 2021.

24.   Blanton CA, Moshfegh AJ, Baer DJ, Kretsch MJ. The USDA Automated Multiple-
Pass Method accurately estimates group total energy and nutrient intake. J Nutr.
2006;136(10):2594-9. doi: 10.1093/jn/136.10.2594.

25.   Channell Doig AJ, Lipsky LM, Choe A, Nansel TR. Development and evaluation
of a nutrient-based method of classifying moderation foods. Cold Spring Harbor
Laboratory, 2025.

26.   Centers for Disease Control and Prevention, National Center for Health
Statistics. *2017-March 2020 Pre-Pandemic Demographics Data - Continuous*
NHANES [Internet]. 2021; Available from:
https://wwwn.cdc.gov/nchs/nhanes/search/datapage.aspx?Component=Demogra
phics&Cycle=2017-2020.

27.   Johnson CL, Paulose-Ram R, Ogden CL, Carroll MD, Kruszon-Moran D,
Dohrmann SM, Curtin LR. National Health and Nutrition Examination Survey:
Analytic guidelines, 1999-2010. National Center for Health Statistics, 2013.

28. Choi YJ, Crimmins EM, Kim JK, Ailshire JA. Food and nutrient intake and diet
quality among older Americans. Public Health Nutr. 2021;24(7):1638-47. doi:
10.1017/S1368980021000586.
29. Guenther PM, Juan W, Lino M, Carlson A, Hiza HH, Krebs-Smith SM. Diet quality
of low-income and higher income Americans in 2003-04 as measured by the
Healthy Eating Index-2005. Nutrition Insight, 2008.
30. Parker HW, Tovar A, McCurdy K, Vadiveloo M. Socio-economic and racial
prenatal diet quality disparities in a national US sample. Public Health Nutr.
2020;23(5):894-903. doi: 10.1017/s1368980019003240.
31. Raffensperger S, Kuczmarski MF, Hotchkiss L, Cotugna N, Evans MK,
Zonderman AB. Effect of race and predictors of socioeconomic status on diet
quality in the HANDLS Study sample. J Natl Med Assoc. 2010;102(10):923-30.
doi: 10.1016/s0027-9684(15)30711-2.
32. Rehm CD, Penalvo JL, Afshin A, Mozaffarian D. Dietary Intake Among US Adults,
1999-2012. JAMA. 2016;315(23):2542-53. doi: 10.1001/jama.2016.7491.
33. Wang DD, Leung CW, Li Y, Ding EL, Chiuve SE, Hu FB, Willett WC. Trends in
dietary quality among adults in the United States, 1999 through 2010. JAMA
internal medicine. 2014;174(10):1587-95. doi:
10.1001/jamainternmed.2014.3422.
34. Carlson A, Frazao E. Food costs, diet quality and energy balance in the United
States. Physiol Behav. 2014;134:20-31. doi: 10.1016/j.physbeh.2014.03.001.
35. Foods US. *The Diner Dispatch: 2024 American Dining Habits* [Internet]. 2024;
Available from: https://www.usfoods.com/our-services/business-trends/american-
dining-out-habits-2024.html.
36. Raskind IG, Woodruff RC, Ballard D, Cherry ST, Daniel S, Haardorfer R, Kegler
MC. Decision-making processes shaping the home food environments of young
adult women with and without children. Appetite. 2017;113:124-33. doi:
10.1016/j.appet.2017.02.027.
37. Goorin J, Venezky E. *How Dining Out Differs Across Generations* [Internet].
Eater; 2025; Available from: https://www.eater.com/press-room/24388888/future-
generations-restaurant-dining-survey-research.

38.   Rhone A, Williams R, Dicken C. Low-Income and Low-Foodstore-Access Census
Tracts, 2015–19. Economic Information Bulletin, 2022.
39.   Nestle M. Food lobbies, the food pyramid, and U.S. nutrition policy. Int J Health
Serv. 1993;23(3):483-96. doi: 10.2190/32F2-2PFB-MEG7-8HPU.
40.   Tapsell LC, Neale EP, Satija A, Hu FB. Foods, Nutrients, and Dietary Patterns:
Interconnections and Implications for Dietary Guidelines. Adv Nutr.
2016;7(3):445-54. doi: 10.3945/an.115.011718.
41.   Rebolledo N, Ferrer-Rosende P, Reyes M, Smith Taillie L, Corvalan C. Changes
in the critical nutrient content of packaged foods and beverages after the full
implementation of the Chilean Food Labelling and Advertising Law: a repeated
cross-sectional study. BMC Med. 2025;23(1):46. doi: 10.1186/s12916-025-
03878-6.
42.   Reyes M, Smith Taillie L, Popkin B, Kanter R, Vandevijvere S, Corvalan C.
Changes in the amount of nutrient of packaged foods and beverages after the
initial implementation of the Chilean Law of Food Labelling and Advertising: A
nonexperimental prospective study. PLoS Med. 2020;17(7):e1003220. doi:
10.1371/journal.pmed.1003220.
43.   Taillie LS, Bercholz M, Popkin B, Reyes M, Colchero MA, Corvalan C. Changes
in food purchases after the Chilean policies on food labelling, marketing, and
sales in schools: a before and after study. Lancet Planet Health. 2021;5(8):e526-
e33. doi: 10.1016/S2542-5196(21)00172-8.
44.   Dillman Carpentier FR, Correa T, Reyes M, Taillie LS. Evaluating the impact of
Chile's marketing regulation of unhealthy foods and beverages: pre-school and
adolescent children's changes in exposure to food advertising on television.
Public Health Nutr. 2020;23(4):747-55. doi: 10.1017/S1368980019003355.
45.   Massri C, Sutherland S, Kallestal C, Pena S. Impact of the Food-Labeling and
Advertising Law Banning Competitive Food and Beverages in Chilean Public
Schools, 2014-2016. Am J Public Health. 2019;109(9):1249-54. doi:
10.2105/AJPH.2019.305159.
46.   Taillie LS, Reyes M, Colchero MA, Popkin B, Corvalan C. An evaluation of Chile's
Law of Food Labeling and Advertising on sugar-sweetened beverage purchases

from 2015 to 2017: A before-and-after study. PLoS Med. 2020;17(2):e1003015.

doi: 10.1371/journal.pmed.1003015.

Table 1. Sample characteristics (n = 7707)

Characteristic	Adults (≥ 20 y)	
	N	**%**
Age, years		
20 – 39	2358	36
40 – 59	2548	34
60+	2801	30
Gender		
Male	3745	48
Female	3962	52
Race/ethnicity		
Mexican American/Hispanic	1669	16
Non-Hispanic White	2758	63
Non-Hispanic Black	2071	11
Non-Hispanic Asian	841	6
Multi/other	368	4
Education		
Some high school	1358	10
High school graduate	1863	28
Some college	2567	31
College graduate or more	1910	32
Income poverty ratio quartile		
1 (mean ± SD = 0.7 ± 0.4)	1689	17

2 (mean ± SD = 1.8 ± 0.3)	1685	19
3 (mean ± SD = 3.3 ± 0.5)	1690	27
4 (mean ± SD = 4.9 ± 0.1)	1672	37

Table 2. Sidak-adjusted pairwise comparisons of mean moderation food intake (% kcal) in U.S. adults by sociodemograpic characteristics (National Health and Nutrition Examination Survey, 2017 – 2020)

Characteristic	Comparison	Mean difference (% kcal)	95% CI	*p* - value
Age, y	40 – 59 vs 20 – 39	-4.1	-5.9, -2.3	<0.001
	>=60 vs 20 – 39	-7.2	-8.8, -5.6	<0.001
	>=60 vs 40 – 59	-3.1	-4.8, -1.5	<0.001
Sex	Female vs Male	-3.2	-4.4, -1.9	<0.001
Race/ethnicity	NH white vs Mex/Am Hisp	3.2	0.4, 6.0	0.02
	NH black vs Mex/Am Hisp	5.6	2.9, 8.4	<0.001
	NH Asian vs Mex/Am Hisp	-6.9	-10.5, -3.2	<0.001
	other/multi vs Mex/Am Hisp	6.4	2.6, 10.3	<0.001
	NH black vs NH white	2.5	0.4, 4.6	0.01
	NH Asian vs NH white	-10.0	-13.2, -6.8	<0.001
	other/multi vs NH white	3.3	-2.5, 6.8	0.08
	NH Asian vs NH black	-12.5	-16.9, -8.1	<0.001
	other/multi vs NH black	0.8	-2.6, 4.2	0.998
	other/multi vs NH Asian	13.3	8.7, 17.9	<0.001
Education	HS grad vs. < HS	3.3	0.7, 5.9	0.008
	Some college vs < HS	2.0	-0.6, 4.6	0.22
	College graduate vs < HS	-4.2	-6.9, -1.5	0.001
	Some college vs HS grad	-1.3	-3.4, 0.8	0.45
	College graduate vs HS grad	-7.5	-10.7, -4.2	<0.001
	College graduate vs some college	-6.2	-8.5, -3.8	<0.001
Income-poverty ratio quartiles	2 vs 1	-1.8	-4.7, 1.2	0.47
	3 vs 1	-2.7	-5.9, -0.4	0.02
	4 vs 1	-4.6	-8.1, -1.2	0.005
	3 vs 2	-1.0	-3.7, 1.8	0.91
	4 vs 2	-2.9	-6.6, 0.9	0.21
	4 vs 3	-1.9	-5.6, 1.8	0.63

Abbreviations: NH - non-Hispanic, Mex/Am Hisp – Mexican American or other Hispanic, HS – high school

Table 3. Coefficient estimates from regression models estimating interactions of income poverty ratio (IPR) with race/ethnicity and education on moderation food intake (% kcal) in U.S. adults.

Model	Independent variable	β	95% CI	*P*-value
Race/ethnicity	IPR	-0.3	-1.3, 0.7	0.57
	Race/ethnicity			
	Mexican American/Hispanic		REF	
	Non-Hispanic White	6.1	2.1, 10.1	0.005
	Non-Hispanic Black	8.6	5.4, 11.8	<0.001
	Non-Hispanic Asian	-3.6	-7.6, 0.4	0.08
	Other/multi	9.3	3.5, 15.1	0.003
	Interaction of IPR with race/ethnicity			
	Mexican American/Hispanic		REF	
	Non-Hispanic White	-0.9	-2.2, 0.5	0.21
	Non-Hispanic Black	-1.4	-2.6, -0.3	0.02
	Non-Hispanic Asian	-0.7	-1.9, 0.4	0.21
	Other/multi	-1.2	-3.5, 1.1	0.28
	Intercept	74.1	71.4, 76.8	<0.001
Education	IPR	0.1	-1.0, 1.2	0.79
	Education			
	Less than high school	5.2	0.8, 9.7	0.02
	High school graduate	10.3	5.7, 14.9	<0.001
	Some college	7.4	3.0, 11.8	0.002

College graduate or more		REF	
Interaction of IPR with education			
Less than high school degree	-0.3	-1.8, 1.1	0.63
High school graduate	-1.3	-2.6, 0.02	0.05
Some college	-0.4	-1.5, 0.7	0.43
College graduate or more		REF	
Intercept	71.1	66.5, 75.6	<0.001

Figure 1. Moderation food intake (% kcal) in U.S. adults by age (means and 95%CIs).

Different subscripts denote statistically different pairwise comparisons (using Sidak adjustment for multiple comparisons).

Figure 2. Moderation food intake (% kcal) in U.S. adults by sex (means and 95%CIs).

Figure 3. Moderation food intake (% kcal) in U.S. adults by race/ethnicity (means and 95%CIs).

Different subscripts denote statistically different pairwise comparisons (using Sidak adjustment for multiple comparisons).

Figure 4. Moderation food intake (% kcal) in U.S. adults by education (means and 95%CIs).

Different subscripts denote statistically different pairwise comparisons (using Sidak adjustment for multiple comparisons).

Figure 5. Moderation food intake (% kcal) in U.S. adults by income poverty ratio quartiles (means and 95%CIs).

Different subscripts denote statistically different pairwise comparisons (using Sidak adjustment for multiple comparisons).

Figure 6. Relationship of income with moderation food intake (% kcal) in U.S. adults by education (means and 95%CIs).

Coefficient estimates for the marginal associations of income poverty ratio with moderation food intake (%kcal) by education are shown.

Figure 7. Relationship of income to poverty ratio with moderation food intake (% kcal) in U.S. adults by race/ethnicity (means and 95%CIs).

Coefficient estimates for the marginal associations of income poverty ratio with moderation food intake (% kcal) by race/ethnicity are shown.

Moderation Food intake by Age in U.S. Adults

Moderation Food intake by Sex in U.S. Adults

Moderation Food intake by Income Poverty Ratio quartiles in U.S. Adults

Relations of Income-Poverty Ratio with Moderation Food Intake by Race/Ethnicity

Relationship of Income-Poverty Ratio with Moderation Food Intake by Education