A Qualitative Descriptive Study of Motivation States for Physical Activity Among Middle-Aged Adults with Type 1 Diabetes Mellitus

Xinyi (Vera) Wang, RN, BSN[1]; Garrett I. Ash, PhD[2]*; Estefania Hernandez, MA, MPH[3]; Matthew Stults-Kolehmainen, PhD[4]; Stephanie A. Griggs, PhD, RN, FAAN[5]

[1] Case Western Reserve University, Cleveland, OH, USA

[2] Yale University School of Medicine, West Haven, CT, USA

[3] Case Western Reserve University, Cleveland, OH, USA

[4] Yale-New Haven Hospital, New Haven, CT, USA

[5] Emory University Nell Hodgson Woodruff School of Nursing, Atlanta, GA, USA

*Corresponding author: Garrett I. Ash, PhD, Yale University School of Medicine, 950 Campbell Ave, West Haven, CT 06516, USA. Email: garrett.ash@yale.edu

Author Emails: xxw700@case.edu; Garrett.ash@yale.edu; Estefania.hernandez@case.edu; matthew.stults@ynhh.org; stephanie.griggs2@emory.edu

ORCID: 0009-0008-9610-0570 (Wang); 0000-0002-8655-7525 (Ash); 0000-0002-8925-911X (Hernandez); 0000-0001-5361-9614 (Stults-Kolehmainen); 0000-0001-9901-3853 (Griggs)

Declarations: The authors have no conflicts of interest to disclose.

Garrett Ash and Stephanie Griggs report financial support was provided by the National Institutes of Health. Garrett Ash reports a relationship with American Heart Association Inc that includes: funding grants. Garrett Ash reports a relationship with The Patterson Foundation that includes: funding grants. Garrett Ash reports a relationship with Calm.com that includes: professional services for a nominal fee. Stephanie Griggs reports a relationship with the National Institutes of Health that includes: funding grants. Other authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethics approval and consent to participate: Participants signed a Yale IRB-approved informed consent (#2000035846).

Consent for publication: Yes.

Competing Interests: The authors have no competing interests to disclose.

Funding: GIA and the study were funded by the National Institute of Diabetes, Digestive, and Kidney Diseases of the National Institutes of Health (NIDDK/NIH) under a mentored research scientist development award (K01DK129441). SG is funded by grant NIDDK/NIH, R01DK136604. The sponsor NIDDK/NIH had no role in the design of the study nor collection, analysis, and interpretation of data nor in writing the manuscript.

Availability of data and materials: The data will be available immediately following publication, with no end dates. The data will be shared with researchers who provide a methodologically sound proposal to achieve the aims of the approved proposal. Data will be truncated to meet the proposal aims without compromising participant confidentiality. The proposals should be directed to Dr. Garrett Ash at Yale University (garrett.ash@yale.edu). To gain access, data requesters must sign a data access agreement.

Authors' contributions (CRediT Statement): GIA, conceptualization, data curation, funding acquisition, investigation, project administration, resources, writing – reviewing and editing. EH, data curation, formal analysis, validation, visualization, writing – original draft. XW, formal

analysis, validation, visualization, writing – original draft. MSK, validation, writing – review and editing. SG, conceptualization, data curation, formal analysis, investigation, methodology, project administration, supervision, validation, visualization, writing – original draft.

Acknowledgments: The authors acknowledge that this abstract was presented at the American Diabetes Association (ADA) Scientific Sessions, [2025].

Word count: 6822.

Figure Count: 1.

Table Count: 2

Supplemental Table Count: 3

Glossary: HbA1c, hemoglobin A1c; T1DM, type 1 diabetes mellitus; WANT, Wants and Aversions for Neuromuscular Tasks model.

A Qualitative Descriptive Study of Motivation States for Physical Activity Among Middle-Aged Adults with Type 1 Diabetes Mellitus

Abstract

Background: Adults with type 1 diabetes mellitus (T1DM) have barriers to physical activity (PA), including physiological, psychological, emotional, and/or contextual. To complement prior quantitative work, we employed a qualitative study to explore momentary motivational states for movement and rest and to understand how internal and external factors shape these experiences among middle-aged adults with T1DM who had recently completed an exercise-support intervention.

Methods: We interviewed 23 middle-aged adults with T1DM (mean age 48, SD 11 years, 78% female, 91% non-Hispanic white, 35% overweight, and 26% obese), who recently completed an exercise intervention, using semi-structured interviews guided by the Wants and Aversions for Neuromuscular Tasks (WANT) model. This model addresses desires and aversions (i.e., fears) for movement and rest. We utilized directed content analysis to identify common themes, with both deductive and inductive coding.

Results: Participants exhibited simultaneous desires to move and rest, opting for activities like yoga or outdoor walks for active recovery. Competing motivations also emerged, such as the internal drive to exercise conflicting with physical fatigue or glycemic levels necessitating rest.

Conclusions: Our findings align with previous studies using the WANT model, underscoring the dynamic nature of motivation influenced by physiological parameters, aversive states, and recent activities. These insights gained can inform future interventions targeting psychological aspects of physical activity for this unique population.

Abstract Word Count: 216 (limit 250).

Keywords: adults; diabetes, type 1; physical activity; motivation

1. Introduction

Diabetes management, particularly in Type 1 Diabetes Mellitus (T1DM), has increasingly recognized the pivotal role of physical activity in enhancing health outcomes. Movement and regular physical activity are associated with improved achievement of glycemic targets, cardiovascular health, and overall quality of life in individuals with T1DM (1). However, despite these benefits, many adults with T1DM struggle to engage in regular movement, necessitating a deeper understanding of the psychological, behavioral, and contextual factors that influence their motivations (2, 3).

In previous research, many environmental, social, and psychological factors have been significantly associated with physical activity behaviors in individuals with T1DM (2). For instance, access to safe exercise spaces and social support are critical determinants of physical activity behaviors (3). Psychological factors include a variety of cognitive, behavioral (e.g., readiness to change) and motivation-related variables, all of which interact in complex ways to facilitate or hinder physical activity (4).

There are many barriers that hinder physical activity among individuals with T1DM, such as employment, family responsibilities, and time constraints, which can complicate the ability to prioritize physical activity (5). Other barriers range from physiological factors, such as fear of hypoglycemia and fluctuating glucose levels, to psychological barriers, including anxiety and decreased motivation (6, 7). Fear of hypoglycemia, in particular, can severely deter individuals from engaging in activities that may cause their blood sugar levels to drop (8). Additionally,

negative emotional states, such as depression and anxiety, are prevalent among individuals with

T1DM and can further diminish motivation for physical activity (9, 10).

Cognitive theories of PA behavior, such as the Theory of Planned Behavior, Social Cognitive

Theory, and other traditional expectancy–value models, have shown limited ability to account for

variability in future PA behavior (11), and dual-process (e.g., Automatic-Reflective Theory

(ART), Affective Heath Behavior Framework, Integrative Framework, model from Berry and

Conroy, 2017 (12), and others) and organismic theories such as Self-Determination Theory have

generally taken their place. These theories better accommodate the ideas of automaticity, urges,

desires, aversions, and the corollaries of approach, withdrawal, etc. Self-Determination Theory is

often highlighted because it provides a clear framework for understanding the balance between

intrinsic and extrinsic motivation, as well as situational and long-term motivation, which is central

to sustaining physical activity behaviors in adults with T1DM (13). Furthermore, compared with

other commonly used models, such as Social Cognitive Theory or the Theory of Planned Behavior,

Self-Determination Theory more directly addresses the psychological needs (autonomy,

competence, relatedness) that are especially relevant to maintaining long-term engagement in

physical activity while living with a demanding condition like T1DM. This framework has also

been previously applied to adolescents with T1DM in prior qualitative work by Mak and

colleagues, demonstrating its relevance for diabetes-specific movement experiences (14).

However, motivation in the context of T1DM remains less well understood. To date, prior

qualitative studies have examined attitudes, barriers, and emotional responses related to physical

activity among adults with T1DM (5, 15, 16). For instance, Vlcek et al. identified "emotional

responses to planning for exercise," describing how adults with T1DM experienced shifting

feelings, such as "groaning… a reluctance knowing this is going to take work," when anticipating

physical activity or managing potential glycemic consequences. Their analysis also noted that

"pre- and post-exercise organization and emotional responses" shape readiness for activity and

that exercise is "embedded in a broader socio-cultural context," with preparation sometimes

"negatively affect[ing] other areas of the person's life." (16) These findings highlight that emotions

and inclinations toward activity vary across situations. However, prior work has not systematically

conceptualized these fluctuations as momentary motivational states nor examined them through a

theoretical model, such as the WANT framework, that is specifically designed to capture dynamic

desires and aversions for movement and rest. While Self-Determination Theory provides a strong

foundation for understanding psychological needs relevant to sustained physical activity, it offers

limited guidance for explaining rapid, moment-to-moment motivational shifts that often occur in

response to fluctuating glucose levels, physical symptoms, or safety concerns. These dynamic

motivational processes are central to understanding movement and rest in T1DM and provide the

rationale for incorporating additional frameworks, such as the WANT model, that account for

transient motivational states.

The framework used for this study was the WANT Model (Wants and Aversions for

Neuromuscular Tasks), initially developed by Stults-Kolehmainen and colleagues (17). The model

includes 10 tenets (18) that describe dynamic and transitory motivational states (e.g., desires,

urges, cravings, aversions) relevant to both physical activity and sedentary behaviors, the latter

referring to waking low-energy activities performed in sitting or reclining postures. It provides a

nuanced understanding of how individuals perceive, experience, and navigate their motivational

impulses and emotions in the context of shifting physical and psychological stresses throughout

the day (19). The want model has four quadrants, each named after the extreme motivational points

within each quadrant.

1. High dread to move/high desire to rest: describes motivational states often associated with

injury, illness, and similar conditions.

2. High dread to move/high dread to rest: describes motivational states often associated with

surprise, shock, and uncertain situations,

3. High desire to move/high desire to rest: describes motivational states often associated with most

daily experiences and motivational conflicts

4. High desire to move, high dread of rest: describes motivational states often associated with

stressful/fight or flight experiences.

For adults with T1DM, it is useful to emphasize the ongoing shifts and interplays between wants

and aversions for both movement and sedentary behaviors emanating from managing this

condition. Although the WANT model was originally developed and validated in general adult

samples, recent pilot work in adults with T1DM shows that transient motivation states are also

highly relevant for this population. In an ecological momentary assessment study using WANT-

based items along with continuous glucose monitoring (CGM) and activity data, (20) motivation

to move or rest fluctuated across the day and varied with glycemic patterns, fear of hypoglycemia,

aversion to hyperglycemia, and recent physical activity. These momentary states were also related

to subsequent movement behavior. Together, these findings indicate that the motivational

processes described by the WANT model are particularly relevant in the T1DM context,

supporting its use as the guiding framework for this study.

Despite the growing body of literature on T1DM management and physical activity, significant

gaps remain regarding the nuanced motivations of adults with T1DM. Much of the existing

research is predominantly based on quantitative methodologies, which may overlook the

variability and complexity of individual experiences. Qualitative research can provide a deeper

understanding of the lived experiences of these individuals, highlighting the motivations and

barriers they encounter in their daily lives (21). This study aims to address these gaps by employing

a qualitative descriptive approach to explore the motivational states that influence physical activity

engagement through an understanding of movement and rest among adults with T1DM. Building

on the WANT framework, the purpose of this qualitative descriptive study was to elucidate internal

and external factors that shape motivation states affecting physical activity among middle-aged

adults with T1DM. We aimed to 1) understand perspectives and personal values related to physical

activity and 2) explore experiences of motivation states related to movement and rest.

Understanding influences on motivation states can provide insight into how to create supportive

environments that encourage physical activity among adults living with T1DM.

2. Materials and Methods

2.1. Participants

This study is a descriptive qualitative study conducted with participants drawn from a parent

clinical trial testing an exercise-support intervention for adults with T1DM. While recruitment

occurred through the parent trial, the interviews and the qualitative analysis presented here were

designed specifically to address the research questions guiding this study. A purposive sample of

24 adults was recruited via secure patient messaging from a regional medical group,

ClinicalTrials.gov listings, and snowball sampling. Eligible participants met the following criteria:

(1) Aged 30-65. This age range allowed us to capture motivational dynamics among middle-aged

adults, who often balance substantial work, income, caregiving, and chronic disease

responsibilities that shape "have to" and "want to" motivations. Prior work has shown that younger

adults (e.g., college students) also experience motivational states, but these emerge from life

pressures that are quite different from those in midlife. To avoid mixing distinct age-related

patterns, we focused on middle-aged adults, whose life-stage demands align with the motivational

processes examined in this study. (2) ≥1yr T1DM or other insulin deficiency diabetes (hereafter

T1DM). (3) Lack of consistent engagement in structured aerobic or resistance exercise, which in

the parent trial was defined as participating in fewer than one structured exercise session per week.

This criterion reflected the parent study's focus on adults with low baseline activity, (4)

smartphone user, (5) English literate. and (6) had a healthcare provider who confirmed medical

readiness for home-based exercise. Each participant provided the interview 1 week after

completing the trial. Participants signed an IRB-approved informed consent (IRB #2000035846).

2.2. Study Design and Data Collection

Methods reporting follows COREQ criteria (Table S1). A qualitative descriptive approach was

chosen to explore the motivations for movement and sedentarism/rest, providing an in-depth

understanding while staying closely aligned with the data (22). We invited all 24 trial participants,

of whom 23 (96%) accepted. Semi-structured interviews were conducted between June and

October 2024 using a flexible interview guide via audio-recorded Zoom, allowing for in-depth

insights into T1DM management and the motivational states for movement and rest. Interview

questions, guided by the WANT model, aimed to assess desires and aversions (i.e., fears) for

movement and rest (Table S2).

Participants were interviewed for an average of 28 minutes (range 14-73), and most (78%) kept

their cameras on. Interviews were transcribed verbatim and reviewed for accuracy by the research

team through comparison with the audio recordings, with any transcription errors corrected during

this process. Participants were encouraged to reflect on past experiences where initial effort

contrasted with eventual enjoyment or frustration related to healthy behaviors. Additionally, they

discussed how fluctuating glucose levels influenced their motivation and fears regarding physical

activity and rest. The faculty researchers included one exercise physiologist and one scientist in

nursing experienced in T1DM clinical care and several advanced qualitative methods, both with

postdoctoral training in T1DM behavioral science.

2.3. Data Analysis

Interviews were transcribed verbatim, cleaned, and checked for accuracy. A directed content

analysis approach was used along with deductive (to align with the four quadrants of the WANT

model) and inductive coding (23). The directed content analysis approach involved the close

examination of data to identify common themes and patterns of meaning and ideas. Data analysis

was conducted using ATLAS.ti (24). Interviewers were trained in qualitative interviewing. Two

trained qualitative interviewers with backgrounds in anthropology, public health, and nursing

conducted the interviews. They had no prior knowledge of the participants' experience in the study

nor any prior personal interaction with them.

The first step involved familiarization with the data, where transcripts were read multiple times to

gain a deeper understanding of the content. A codebook was developed early in the process and

revised as interviews were being conducted. The codebook ensured consistency in the coding

process as it included definitions and interview questions as they relate to the theme. For example,

the code 'Barriers to Movement – Diabetes Specific' was paired with the question "Do you find

managing your diabetes gets in the way of your goals related to movement/rest?" alongside others.

Two members of the research team independently coded the transcripts using directed qualitative

content analysis. Coding discrepancies were discussed and resolved through team consensus. The

team refined the coding guidelines to ensure that codes were applied to comparable meaning units.

After initial coding, conceptually similar codes were grouped into higher-order categories. These

categories were then compared across transcripts and interpreted in relation to the WANT model

and relevant prior literature on motivation and physical activity. Through this constant comparative

process, the categories were refined and elevated into broader themes that captured patterned

meaning across participants' experiences. Review of memos (collected before and after interview)

and reflexive journal notes were included in this process to address personal bias, assumptions,

and reasons for involvement in the research topic. We used concept mapping to illustrate the

themes and facilitate theme interpretation.

3. Results

3.1. Participant Characteristics

The final sample comprised 23 participants (Table 1), with a majority identifying as female (78%)

and non-Hispanic White (91%). Substantial fractions were above recommended clinical thresholds

for average glycemic values (57% above 7.0% or 53 mmol/mol HbA1c), blood pressure (43%

above 130/85 mmHg), and/or body mass index (35% overweight, 26% obese). Regarding

socioeconomic status, a majority (61%) reported a household income of $100,000 or more, while

74% earned over $60,000 per year. Most had private health insurance (87%) and automated insulin

delivery (AID) systems (78%). Participants varied in their self-reported occupations and their

impact on activity and rest (Table 2).

3.2. Key Motivation States Themes by Theme Domain

The Wants and Aversions in Neuromuscular Tasks (WANT) Model was used to describe

participant movement-related behaviors and aversions (Figure 1). More specifically, the model

helped to describe how participants approached or avoided (e.g., wants versus aversions)

movement and/or sedentarism/rest, and the nature in which participants could vary in both states

at the same moment. For example, someone could be in a state of high craving to both rest and

move simultaneously when people chose to actively recover after a long day through yoga,

meditation, or an outdoor walk. Another example of competing motivations is when someone has

the strong internal drive to exercise every day; however, their muscles may be fatigued, or their

blood sugars signal a need for rest and recovery. The following sections represent each of the four

quadrants of the WANT model shown in Figure 1 and describe feelings reported by participants

associated with these states, precursors to being in that state, and the behavioral outcomes of being

in that state.

3.2.1. Quadrant 1: High dread to move, high crave to rest

Participants in the current study described feelings of discouragement and hesitance to move, often

experiencing a push-pull dynamic between external factors prompting movement and a strong

desire to rest (i.e., motivational conflict). For instance, using electronic devices or other passive

activities amplifies the urge to rest, thereby reducing motivation to initiate movement. At the same

time, participants also described moments when the need for restoration led them to select light

movement, such as stretching or gentle chores, as a way to feel better without committing to more

demanding physical activity. Engaging in household chores served as a meaningful break,

allowing participants to restore their energy. Some individuals felt uncomfortable exercising in

front of others, leading to an aversion that hindered their motivation to continue movement.

Additionally, workouts that exceeded their usual routines often tested their willpower, making

them prone to giving up, highlighting the internal struggle in this quadrant.

Individuals expressed that when feeling unwell, their bodies typically did not function at their

regular capacity, causing them to hesitate in their decision to exercise as they weigh potential

benefits against drawbacks. In such instances, the body's instinct is often to rest for a quicker

recovery. However, some participants maintained that movement while feeling sick could aid in

detoxification through sweating and accelerating recovery.

For participants who were parents, balancing exercise with the demands of caring for toddlers or

children under time constraints, physical activity could feel burdensome. After expending energy

playing with their children, their need for rest significantly increased.

Participants also mentioned that out-of-range glucose levels often hindered their ability to move.

They mentally prioritized stabilizing their glucose levels to avoid further deterioration,

understanding that both hypo- and hyperglycemia can disrupt normal bodily functions and

necessitate rest. Conversely, some individuals indicated that they intentionally remain active as a

non-medical strategy to bring their glucose levels back to a safe range, regardless of their

predominant motivational state. As one participant noted, "There are times when I was not

intending to do a workout, but my blood sugars are really high and stay elevated for a period, and

I know that a workout will help break that cycle of high blood sugar. So, it prompts me to exercise

even when I may not have wanted to or chosen to" (ID 14, Male, age 50's, using AID for insulin

delivery).

When participants were preparing for bed, malfunctioning diabetes management devices were

reported to contribute to elevated glucose levels, forcing them to stay awake and troubleshoot

despite feeling drowsy. For instance, one participant shared, "Sometimes you can have fluctuations

in your blood sugar that make you stay up longer at night to manage it, or you might need to change

your equipment because something isn't right. That can impact your performance or sleep the next

day… You're up because your Dexcom wasn't working, or you had a site failure, or your pump

wasn't functioning, or maybe you just ate too much…" (ID 07, Female, age 60's, AID).

Psychologically, some participants reported burnout from managing their condition as they grapple

with fluctuating glucose levels without understanding the causes. This frustration was described

to drain their enthusiasm for physical activity or any movement and heightened their need for rest.

Additionally, unfavorable weather, such as rainy days and extreme temperatures, significantly

discouraged outdoor activities and diminished overall activity levels. Sleep deprivation led to

fatigue, which in turn decreased motivation for physical activity, causing individuals to prioritize

recuperation over movement.

For some, excessive time spent socializing could deplete one's social energy, resulting in mental

fatigue that diminished the desire for physical activity. As one participant expressed, "I'm an

introvert, and such social situations make me more likely to want to rest" (ID 03, Female, age 40's,

sensor augmented pump [SAP]). In these moments, individuals often turn to relaxing activities

like reading, listening to music, lying in bed, taking a bath or shower, stretching, napping, or

watching TV as means to rest and recharge. However, in some cases of needing rest, individuals

utilized strength and core workouts to relax. The simultaneous desire to rest and increased

motivation for movement can lead to disrupted sleep patterns, delayed bedtimes, and unintended

exercise.

3.2.2. Quadrant 2: High dread to move, high dread to rest

A vast majority of participants who described this state reported often feeling a strong sense of

obligation to complete tasks rather than take a break. However, once they finished, they

experienced a sense of pride and accomplishment for having chosen productivity over relaxation.

Some individuals, who did not have a prior exercise routine, felt compelled to stick to the study's

exercise plan, and, although they may have resisted at first, they found gratification and enjoyment

in the process. This was particularly noted during intervention exercises, where participants

expressed that their initial reluctance transformed into enthusiasm. As one participant shared, "One

day I didn't really want to go in the pool because it looked really dark and cloudy and wasn't that

warm, but I was on the phone with [STUDY MEMBER], and I kind of forced myself. And then

once I was in, I was able to stay in for a while" (ID 12, Female, age 50's, AID).

Similarly, some participants viewed taking a shower as a chore, motivated by the pressure to

maintain personal hygiene. Yet, once they started, they often emerged feeling refreshed and

rejuvenated. The same theme persisted with yard work; despite initial resistance to maintaining

their household and landscape, individuals pushed themselves to engage in these activities. After

completing the work, they were rewarded with the sight of a well-kept yard. Positive weather

conditions played a significant role, often prompting participants to prioritize outdoor chores over

resting indoors. One participant remarked, "I'm going to have to do yard work, and I just don't

want to because I'm tired but it's a beautiful day out so it urges me to not want to rest because it's

so nice outside, and I want to go do something" (ID 11, Male, age 30's, AID).

In cases where individuals struggled to motivate themselves, they often began with aerobic

activities. Gradually, as they engaged more fully, they began to take pride in their progress and the

improvement in their physical condition. Furthermore, while balanced meals are crucial for

managing T1DM, participants did not always cook out of desire but rather necessity. They opted

to spend time preparing healthy meals to regulate their glucose levels, rather than skipping meals

or choosing easier alternatives. Ironically, many found enjoyment in the cooking process once they

started, revealing how initial aversions can shift to positive experiences.

3.2.3. Quadrant 3: High crave to move, high crave to rest

Individuals often described themselves in a state of energized tranquility when their bodies

signaled a need for recovery, prompting a desire to balance this need with dynamic activities. The

craving for rest could lead to an instinctive counteraction, where individuals engaged in movement,

thereby fostering a harmonious flow between action and relaxation. For instance, during long car

or air travels, discomfort could arise from narrow spaces, prolonged sitting, and insufficient

opportunities for quality rest, culminating in fatigue. The confined environment placed a strain on

muscles and increased feelings of restlessness. These factors combined to create stiffness and

fatigue, amplifying the urge to move to alleviate tension and restore comfort.

As one participant articulated, "When we were driving for three days straight, after about seven

hours in the car, you're like, 'I need to stop driving,' but it's an hour and a half until we reach our

destination, and I'm like, 'I need to move.' We are stopping right now because I have to move. I

have to get out and walk. I have to do something. I gotta jog. I gotta do some jumping jacks. I need

to do something because I'm going to go crazy in the last 90 minutes" (ID 06, Female, age 50's,

AID).

Similarly, after engaging in strenuous exercise, individuals often resorted to active relaxation

techniques, such as stretching or walking, to meet their body's recovery needs. One participant

noted, "After exercise, I usually rest and recover by walking" (ID 02, Female, age 40's, AID). This

interplay between the desire to move and the need for rest illustrates the complex dynamic

encapsulated by the Wants and Aversions in Neuromuscular Tasks (WANT) Model, highlighting

how individuals navigate competing desires simultaneously.

3.2.4. Quadrant 4: High crave to move, high dread to rest

Several participants articulated the experienced heightened sense of mental engagement when

participating in enjoyable physical activities, leading to a reluctance to rest. The satisfaction

derived from completing these activities fostered a sense of contentment. For those who described

themselves as naturally competitive, the effort needed to achieve movement goals was a significant

source of pleasure. When setting such goals, motivation stemmed from the desire to reach new

heights. People who took pleasure in their work frequently expressed enjoyment during their tasks,

making them less inclined to seek rest. Engaging in personal hobbies was also reported to induce

a flow state, wherein individuals become fully absorbed in the activity without perceiving a need

for rest, as the enjoyment and focus they derive sustained their energy levels.

Quality time spent being active with family was described as creating strong emotional

connections, reinforcing family bonds, and cultivating a sense of contentment. This positive

interaction spurred individuals to continue moving, buoyed by the energy of shared experiences

rather than seeking out moments of rest. Team leaders in sports reported feeling a commitment to

their peers, motivating them to attend practices and participate actively rather than opting for rest.

Furthermore, a sense of responsibility to keep teammates engaged and serve as a role model was

described to enhance their motivation to stay active.

In the context of managing blood sugar levels, participants oftentimes gravitated toward moderate-

intensity exercise as a practical means to lower glucose levels, opting for movement over rest to

address their needs. As one participant expressed, "If I could somehow have the freedom, which I

can't, my diabetes doesn't let me, to exhaust myself with physical activity on a daily basis, I

would" (ID 09, Female, age 60's, HCL). Another noted, "I'm conscious of [my sugars] all the time.

If they're on the lower side, I probably wouldn't start with vigorous exercise or workout or

anything" (ID 08, Female, age 30's, using injections for insulin delivery).

To achieve feelings of contentment and enjoyment, individuals typically chose activities that

promote movement and interaction rather than rest. Such activities encompassed outdoor

excursions, group engagement, dancing, playing tennis, interacting with grandchildren, or even

pursuing hobbies like reading or crocheting. However, before embarking on physical activities,

participants' glucose levels could influence the intensity of their exercise, which could at times

cause delays or hinder their movement goals. They often required periods of rest to recheck and

adjust their glucose levels to ensure they are within a safe range before resuming physical activity.

"Every once in a while, I'd run into that fact of where I'd have a spike in my blood sugar to be

high, and when I want to do a workout, which I know is not the best. And then it's like, I got to

wait for it to catch up" (ID 11, Male, age 30's, AID). Similarly, another participant shared,

"Sometimes I might feel an urge or need to do a physical activity later than I would have wanted,

because my blood sugars are high" (ID 14, Male, age 30's, AID).

Given the significant impact of movement on glucose levels, participants reported the need to

consistently monitor their levels throughout activities, which often disrupted their overall

experience. These necessary adjustments lessened enjoyment and reduced feelings of contentment

as activities came to an end. "When I was doing the evening walks, there were several days where

I had dinner a little bit later than I planned, and I still had so much insulin on board that I didn't

think it was a good idea to go out and walk away from home because of the insulin on board" (ID

10, Female, age 40's, AID). Another participant recounted, "We were swimming and I couldn't

[continue] because I was starting to go low" (ID 18, Female, age 50's, AID).

4. Discussion

The current study aimed to explore motivation states for physical activity and sedentary behaviors among middle-aged adults living with Type 1 Diabetes (T1DM) using the WANT Model framework (18). Understanding the dynamic interplay of psychological factors influencing movement and rest among adults with T1DM is crucial for developing effective interventions. This discussion highlights the implications of our findings, the strengths and limitations of our study design, and recommendations for future research.

Our findings illustrate how the interaction between desires (e.g., urges, cravings) and aversions (e.g., dread, fear) informs participants' choices about engaging in movement versus resting behaviors. Additionally, our results align with a mixed methods study of 17 undergraduate students revealing three similarities: (1) there was a dynamic nature of motivation noted where a fluctuating nature of motivation states was found, frequently with rapid changes in desires to move and rest, (2) the influence of external factors (e.g., pressure of maintaining personal hygiene) was significant, and (3) strong urges manifested from the deprivation of both movement and rest (17). These patterns are consistent with the idea that challenging or uncomfortable conditions can make people more likely to choose lower-energy activities. However, the nuances introduced in the T1DM context highlight the unique challenges and conflicts individuals face when managing this complex condition, such as having a low blood sugar level, preventing exercise from occurring simultaneously with an urge to exercise. Further, the current study focused on middle-aged adults who were grappling with a complex chronic condition, and these demographic differences may have led to varied motivational experiences due to differences in life stage, responsibilities, and health management. This suggests that future interventions or research would benefit from

considering these complex influences on motivation alongside physiological states to better

address the needs of diverse populations.

The most hypothesized diseases to be impacted would be movement-related disorders, and a

scoping search found language consistent with the WANT model in 95 studies of such conditions

(25) (Table S3). But available evidence has also suggested more complex pathways involved in

chronic psychiatric disease, such as in anorexia nervosa, where the drive to be active is propelled

by body-conscious ideations (26), or those involved with state anxiety and other mental disorders

(27, 28). Meanwhile, studies of healthy populations have generated evidence that momentary

placement within the WANT model is associated with stimuli similar to what we observed in the

present study including energy availability (a correlate of glucose levels in T1DM), aversive

valence states (akin to anticipation of unpleasant glucose trends), time of day, and recent physical

activity or sedentary time (29). Elevated glucose may therefore influence motivation by signaling

increased energy availability, or, as participants suggested, by prompting movement to maintain

or restore glucose to target ranges with less insulin.

The findings of this study contribute to the growing body of literature emphasizing the profound

connection between physiological states, such as glucose levels, psychological outcomes,

particularly motivation, and behavior. Existing research has demonstrated that individuals with

T1DM often experience fluctuations in mood and motivation due to shifts in their blood glucose

levels. For instance, studies have illustrated how both hyperglycemia and hypoglycemia can lead

not only to physical symptoms but also to cognitive impairments and aversions that affect decision-

making processes (30, 31).

Furthermore, aligning with findings from a study by McCarthy et al. (32), which pointed out the

behavioral impacts of diabetes management on physical activity, the current study reinforces the

argument that glucose fluctuations not only influence physical health but also the psychosocial

dimensions of living with diabetes. This suggests a critical gap in current interventions, which

often focus primarily on the physiological aspects of diabetes management, neglecting the

psychological and motivational layers that are equally crucial for effective self-management. A

number of studies have addressed the fear of hypoglycemia by educational or cognitive behavioral

approaches (33, 34), but studies like the current one reveal the complexity of physical activity

behavior and determinants beyond fear of hypoglycemia (35, 36).

Overall, the emphasis on tailored interventions that address these motivational nuances aligns with

recommendations from researchers advocating for holistic, patient-centered approaches in diabetes

care (37). Such interventions could enhance adherence to recommended physical activity

guidelines, ultimately improving both psychological well-being and physical health outcomes for

individuals with T1DM.

When interpreting the results of this study, several limitations must be acknowledged. We enrolled

individuals at the end of an intervention designed to address inconsistent exercise, leaving the

possibility that findings do not apply to individuals with long-term high or low activity levels. Our

sample, like the adult cohort of the type 1 diabetes exchange (38), had limited racial or ethnic

diversity, though it was diverse regarding HbA1c, blood pressure, anthropometrics, and

occupational physical activity (Table 1). Additionally, the qualitative nature of the study may

introduce biases, as participants could have altered their responses due to social desirability. To

address these limitations, future research should focus on larger, longitudinal qualitative and

quantitative studies to examine the stability of motivation states over time and their long-term

impacts on movement and sedentary behaviors.

Despite these limitations, it is important to recognize the strengths of this study. The qualitative

methodology employed allowed for an in-depth exploration of participants' lived experiences, with

semi-structured interviews yielding rich, nuanced data that captures the complexity of motivation

states. Furthermore, to our knowledge, this is the first study to explore motivation states in middle-

aged adults with T1DM using the WANT model. This approach not only sheds light on the

motivational dynamics within this population but also lays the groundwork for further research

and targeted interventions tailored to their specific needs and desires.

Another theme that emerged from the data, though not explicitly delineated in the paper, was the

recurring language of "have to" (obligation) reflecting a sense of necessity in participants'

motivation. This dimension permeated throughout the results and may represent an important

complement to the "want to/don't want to" motivational perceptions emphasized in the WANT

model. For future studies, it may be useful to explore how a sense of obligation interacts with

"want to" or "don't want to" motivations to influence behavior and how it's further modulated by

motivational conflicts (e.g., "want to move" vs concurrent "want to rest"), a finding corroborated

in our previous qualitative study with 17 adults without any chronic disease (17).

Moving forward, future researchers should focus on larger, longitudinal studies to examine the

stability of motivation states over time and their long-term impacts on movement and sedentary

behaviors/rest. Additionally, investigating how varying motivational states can inform targeted

intervention designs aimed at improving physical activity among individuals with T1DM is

essential. It is also crucial to explore the role of healthcare providers in facilitating conversations

about motivation and offering ongoing support tailored to individual experiences and challenges.

By addressing both the limitations and strengths of this study, we can pave the way for more

effective strategies to enhance T1DM self-management support.

5. References

1. Colberg SR, Sigal RJ, Yardley JE, et al. Physical activity/exercise and diabetes: A position statement of the American Diabetes Association. *Diabetes Care.* 2016;39(11):2065-2079. doi:10.2337/dc16-1728

2. McCarthy MM, Whittemore R, Grey M. Physical activity in adults with type 1 diabetes. *Diabetes Educ.* 2016;42(1):108-115. doi:10.1177/0145721715620021

3. Dollman J. Social and environmental influences on physical activity behaviours. *Int J Environ Res Public Health.* 2018;15(1):169. doi:10.3390/ijerph15010169

4. Adu MD, Malabu UH, Malau-Aduli AE, Malau-Aduli BS. Enablers and barriers to effective diabetes self-management: A multi-national investigation. *PLoS One.* 2019;14(6):e0217771. doi:10.1371/journal.pone.0217771

5. Lascar N, Kennedy A, Hancock B, et al. Attitudes and barriers to exercise in adults with type 1 diabetes (T1DM) and how best to address them: a qualitative study. *PLoS One.* 2014;9(9):e108019. doi:10.1371/journal.pone.0108019

6. Finn M, Sherlock M, Feehan S, Guinan EM, Moore KB. Adherence to physical activity recommendations and barriers to physical activity participation among adults with type 1 diabetes. *Ir J Med Sci.* 2022;191(4):1639-1646. doi:10.1007/s11845-021-02741-w

7. Cockcroft EJ, Narendran P, Andrews RC. Exercise-induced hypoglycaemia in type 1 diabetes. *Exp Physiol.* 2020;105(4):590-599. doi:10.1113/EP088219

8. Yardley JE, Sigal RJ. Exercise strategies for hypoglycemia prevention in individuals with type 1 diabetes. *Diabetes Spectr.* 2015;28(1):32-38. doi:10.2337/diaspect.28.1.32

9. Young-Hyman D, de Groot M, Hill-Briggs F, et al. Psychosocial care for people with diabetes: A position statement. *Diabetes Care.* 2016;39(12):2126-2140. doi:10.2337/dc16-2053

10. Qin W, Blanchette JE, Yoon M. Self-efficacy and diabetes self-management in middle-aged and older adults in the United States: A systematic review. *Diabetes Spectr.* 2020;33(4):315-323. doi:10.2337/ds19-0051

11. Rhodes RE, McEwan D, Rebar AL. Theories of physical activity behaviour change: a history and synthesis of approaches. *Psychol Sport Exerc.* 2019;42:100-109. doi:10.1016/j.psychsport.2018.11.010

12. Conroy DE, Berry TR. Automatic affective evaluations of physical activity. *Exerc Sport Sci Rev.* 2017;45(4):230-237. doi:10.1249/JES.0000000000000120

13. Alarcón-Gómez J, Chulvi-Medrano I, Martin-Rivera F, Calatayud J. Effect of high-intensity interval training on quality of life, sleep quality, exercise motivation and

enjoyment in sedentary people with type 1 diabetes mellitus. *Int J Environ Res Public Health.* 2021;18(23):12612. doi:10.3390/ijerph182312612

14. Mak SS, Nally LM, Montoya J, et al. An exergames program for adolescents with type 1 diabetes: qualitative study of acceptability. *JMIR Diabetes.* 2025;10:e65665. doi:10.2196/65665

15. Kennedy A, Narendran P, Andrews RC, Daley A, Greenfield SM; EXTOD Group. Attitudes and barriers to exercise in adults with a recent diagnosis of type 1 diabetes: a qualitative study of participants in the Exercise for Type 1 Diabetes (EXTOD) study. *BMJ Open.* 2018;8(1):e017813. doi:10.1136/bmjopen-2017-017813

16. Vlcek C, Greenberg D, Yardley JE, et al. "How we do it": A qualitative study of strategies for adopting an exercise routine while living with type 1 diabetes. *Front Endocrinol (Lausanne)*. 2023;13:1063859. Published 2023 Jan 5. doi:10.3389/fendo.2022.1063859

17. Stults-Kolehmainen MA, Gilson TA, SantaBarbara N, et al. Qualitative and quantitative evidence of motivation states for physical activity, exercise and being sedentary from university student focus groups. *Front Sports Act Living.* 2023;5:1190027. doi:10.3389/fspor.2023.1033619

18. Stults-Kolehmainen MA, Blacutt M, Bartholomew JB, et al. Motivation states for physical activity and sedentary behavior: Desire, urge, wanting, and craving. *Front Psychol.* 2020;11:568861. doi:10.3389/fpsyg.2020.568390

19. Stults-Kolehmainen MA, Blacutt M, Fogelman N, et al. Measurement of motivation states for physical activity and sedentary behavior: development and validation of the CRAVE Scale. *Front Psychol.* 2021;12:568286. doi:10.3389/fpsyg.2021.568286

20. Ash GI, Jeon S, Lukasik J, Griggs S, Stults-Kolehmainen M, Fucito LM. Motivation States for Physical Activity Among Middle-Aged Adults With Type 1 Diabetes Mellitus: An Ecological Momentary Assessment Study. *SportrXiv* [preprint]. Posted September 29, 2025. doi:10.51224/SRXIV.626

21. Noyes J, Booth A, Moore G, et al. Synthesising quantitative and qualitative evidence to inform guidelines on complex interventions: clarifying the purposes, designs, and outlining some methods. *BMJ Glob Health.* 2019;4(Suppl 1):e000893. doi:10.1136/bmjgh-2018-000893

22. Sandelowski M. Whatever happened to qualitative description? *Res Nurs Health.* 2000;23(4):334-340. doi:10.1002/1098-240X(200008)23:4<334::AID-NUR9>3.0.CO;2-G

23. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res.* 2005;15(9):1277-1288. doi:10.1177/1049732305276687

24. Smit B, Scherman V. Computer-Assisted Qualitative Data Analysis Software for Scoping Reviews: A Case of Atlas.ti. *Int J Qual Methods.* 2021;20:16094069211019140. doi:10.1177/16094069211019140

25. Stults-Kolehmainen MA, Blacutt M, Bartholomew JB, et al. Urges to Move and Other Motivation States for Physical Acivity in Clinical and Healthy Populations: A Scoping Review Protocol. *Front Psychol.* 2022;13:901272. doi:10.3389/fpsyg.2022.901272

26. Casper RC. Restlessness and an Increased Urge to Move (Drive for Activity) in Anorexia Nervosa May Strengthen Personal Motivation to Maintain Caloric Restriction and May Augment Body Awareness and Proprioception: A Lesson From Leptin Administration in Anorexia Nervosa. *Front Psychol.* 2022;13:885274. doi:10.3389/fpsyg.2022.885274

27. Cheng Z, Taylor A, Stults-Kolehmainen MA, et al. Validation of the CRAVE-C scale in Chinese adults... *Front Psychol.* 2024;15:1467949. doi:10.3389/fpsyg.2024.1467949

28. Filgueiras A, Stults-Kolehmainen MA, Boullosa D, et al. The CRAVE and ARGE scales for motivation states for physical activity and sedentarisim: Brazilian Portugese translation and single-item versions. *Front Psychol.* 2023;14:1106571. doi:10.3389/fpsyg.2023.1106571

29. Budnick CJ, Stults-Kolehmainen MA, Dadina C, et al. Motivation states to move, be physically active and sedentary vary like circadian rhythms. *Front Sports Act Living.* 2023;5:1094288. doi:10.3389/fspor.2023.1094288

30. Verhulst CEM, Fabricius TW, Nefs G, et al. Consistent Effects of Hypoglycemia on Cognitive Function in People With or Without Diabetes. *Diabetes Care.* 2022 Jul 25;45(9):2103–2110. doi:10.2337/dc21-2502

31. Kodl CT, Seaquist ER. Cognitive dysfunction and diabetes mellitus. *Endocr Rev.* 2008;29(4):494-511. doi:10.1210/er.2007-0034

32. McCarthy MM, Whittemore R, Gholson G, Grey M. Self-management of physical activity in adults with type 1 diabetes. *Appl Nurs Res.* 2017;35:18-23. doi:10.1016/j.apnr.2017.02.010

33. Chatwin H, Broadley M, de Galan B, et al. Effectiveness of educational and behavioural interventions for reducing fear of hypoglycaemia. *Diabet Med.* 2023;40(5):e15071. doi:10.1111/dme.15071

34. Martyn-Nemeth P, Duffecy J, Quinn L, et al. FREE: A randomized controlled feasibility trial of a cognitive behavioral therapy and technology-assisted intervention to reduce fear

of hypoglycemia in young adults with type 1 diabetes. *J Psychosom Res.*

2024;181:111679. doi:10.1016/j.jpsychores.2024.111679

35. Brazeau AS, Rabasa-Lhoret R, Strychar I, Mircescu H. Barriers to physical activity

among patients with type 1 diabetes. *Diabetes Care.* 2008;31(11):2108-2109.

doi:10.2337/dc08-0720

36. Alobaid AM, Zulyniak MA, Ajjan RA, et al. Barriers to Exercise in Adults With Type 1

Diabetes and Insulin Resistance. *Can J Diabetes.* 2023;47(6):503-508.

doi:10.1016/j.jcjd.2023.04.016

37. Aysola J, Werner RM, Keddem S, et al. Asking the Patient About Patient-Centered

Medical Homes: A Qualitative Analysis. *J Gen Intern Med.* 2015;30(10):1461-1467.

doi:10.1007/s11606-015-3312-8

38. Foster NC, Beck RW, Miller KM, et al. State of Type 1 Diabetes Management and

Outcomes from the T1D Exchange in 2016-2018. *Diabetes Technol Ther.* 2019;21(2):66-

72. doi:10.1089/dia.2018.0384

6. Figure Legends

FIGURE 1. Conceptual model of analysis based on WANT Model. Highlighted colors

correspond to motivational states as described by participants. Flow = Yellow; Animation =

Purple; Agitation = Orange; Rest = Blue. T1DM = type 1 diabetes mellitus.

7. Tables

Table 1. Participant demographics

Table 2. Self-reported occupation and impact on activity and rest.

Supplemental Material

Table S1. COREQ Checklist.

Table S2. Interview Guide.

Table S3. Literature Review.

FIGURE 1: Conceptual model of analysis based on WANT Model. Highlighted colors correspond to motivational states as described by participants. Flow = Yellow; Animation = Purple; Agitation = Orange; Rest = Blue. T1DM = type 1 diabetes mellitus.

Table 1. Participant demographics (n = 23) [a]

Variables	Mean	SD
Age (years)	48	11 years
Duration Diabetes Diagnosis (years)	27	13 years
HbA1c		
%	6.8	0.9
Mmol/mol	51	10
Body mass index (kg/m^2)	22.7[a]	1.5
	31.5[b]	6.1
Systolic blood pressure (mmHg)	119	12
Diastolic blood pressure (mmHg)	79	8
	N	**%**
Diagnosis		
Type 1 diabetes mellitus	21	91.3
Latent autoimmune diabetes of adulthood	2	8.7
Sex		
Male	5	21.7
Female	18	78.3
Race X Ethnicity		
Non-Hispanic White	21	91.3
Non-Hispanic Black	1	4.3
Hispanic White	1	4.3
Marital Status		
Single	3	13.0
Married	17	73.9
Divorced	2	8.7
N/A	1	4.3
Household Income[c]		
Less than $60,000	1	4.3
$20,000 - $39,999	2	8.7
$40,000 - $59,999	2	8.7
$60,000 - $79,999	2	8.7
$80,000 - $90,999	1	4.3
More than or equal to $100,000	14	60.9

Generational Status		
Self and parent/s USA-born	20	87.0
Self USA-born, ≥1 parent outside USA	3	13.0
Insurance		87.0
Private	20	13.0
Public	3	
Diabetes Management	4	17.4
Injections	1	4.3
Sensor-Augmented Pump		
Automated Insulin Delivery (AID)	18	78.3
Pump Brand		
Omnipod Dash	3	13.0
Omnipod 5	4	17.4
Tandem t:slim X2	9	39.1
Medtronic 770G	3	13.0
[a]N=9, [b]N=14		
[c]One participant dropped out before reaching the exit interview for this sub-study due to competing time demands. [b]One participant declined to respond to this question		

Table 2. Self-reported occupation and impact on activity and rest.

Profession	Job Title	N	Activity Level	Disrupts Rest
Healthcare & Public Social Services	Nurse Practitioner	2	Active / Sedentary	No / Yes
	Ultrasound Tech	1	Mixed	No
	Social Worker	1	Sedentary	Yes
	Firefighter Paramedic	1	Active	Yes
	Care Staff	1	Mixed	Yes
	Police Officer	1	Mixed	Yes
Education & Training	Teacher	2	Active	No / Yes
	School Faculty	1	Sedentary	No
	Software Trainer	1	Sedentary	No
Agricultural	Arborist	1	Mixed	Yes
	Wildlife Biologist	1	Mixed	No
Creative Business	Lighting Designer	1	Mixed	N/A
	Marketing Consultant	1	Sedentary	Sometimes
	Small Business Owner	2	Sedentary	No / No
Miscellaneous	Stay-At-Home Parent	1	Active	No
	Facilities Coordinator	1	Mixed	No
	Unemployed	1	Mixed	No
	Retired	1	Mixed	No
	Unknown	1	Mixed	N/A