Title: Clinical and imaging characteristics of Parkinson's disease with negative alpha-synuclein seed amplification assay

Authors: Sarah M. Brooker MD, PhD[1], Jacopo Pasquini MD[2,3], Seung Ho Choi MS[4], David-Erick Lafontant PhD[4], Seyed-Mohammad Fereshtehnejad MD, PhD[5,6,7], Yashar Zeighami PhD[8,9], Piergiorgio Grillo MD[10,11,12], Giulietta M. Riboldi MD, PhD[13], Houman Azizi BSc[8, 9,14], Roqaie Moqadam MSc[8], Un Jung Kang MD[10,13,15,16], Andrew Siderowf MD[17], Caroline M. Tanner MD, PhD[18], Thomas F. Tropea DO[17,19], Tatiana Foroud PhD[20], Lana M. Chahine MD[21], Brit Mollenhauer MD[22,23], Kalpana M. Merchant PhD[1], Douglas Galasko MD[24], Christopher S. Coffey PhD[4], Roseanne D. Dobkin PhD[25], Ethan G. Brown MD[18], Roy N. Alcalay MD[26,27], Daniel Weintraub MD[17,28,29], Kenneth Marek MD[19], Tanya Simuni MD[1], Paulina Gonzalez Latapi MD[1], Nicola Pavese MD, PhD[2], Kathleen L. Poston MD[30,31,32,33] on behalf of the Parkinson's Progression Markers Initiative*

Affiliations
1. Northwestern University Feinberg School of Medicine, Department of Neurology, Chicago, IL, United States
2. Clinical Ageing Research Unit, Newcastle University, Newcastle upon Tyne, UK
3. Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy
4. Department of Biostatistics, College of Public Health, University of Iowa, Iowa City, IA, United States
5. Division of Neurology, Toronto Western Hospital, University of Toronto, Toronto, ON, Canada
6. Edmond J. Safra Program in Parkinson's Disease and the Morton and Gloria Shulman Movement Disorders Clinic, Toronto Western Hospital, UHN, Toronto, ON, Canada
7. Department of Neurobiology, Care Sciences and Society (NVS), Karolinska Institute, Stockholm, Sweden
8. Douglas Mental Health University Institute, Department of Psychiatry, McGill University, Montreal, Quebec, Canada
9. Integrated Program in Neuroscience, McGill University, Montreal, Quebec, Canada
10. The Marlene and Paolo Fresco Institute for Parkinson's and Movement Disorders, Department of Neurology, NYU Langone Health, NY, United States
11. Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy
12. IRCCS Mondino Foundation, Pavia, Italy
13. Department of Neurology, NYU Grossman School of Medicine, NY, United States
14. Neuro (Montreal Neurological Institute-Hospital), McGill University, Montreal, Quebec, Canada;
15. Departments of Neuroscience, NYU Grossman School of Medicine, NY, United States
16. Parekh Center for interdisciplinary Neurology, NYU Langone Health, NY, United States
17. Department of Neurology, University of Pennsylvania, Philadelphia, PA, United States

18. Department of Neurology, Weill Institute for Neurosciences, University of California San Francisco, San Francisco, California, United States
19. Institute for Neurodegenerative Disorders, New Haven, CT, United States
20. Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN, United States
21. Department of Neurology University of Pittsburgh, Pittsburgh, PA, United States
22. Department of Neurology, University Medical Center Göttingen, Göttingen, Germany
23. Paracelsus-Elena-Klinik Kassel, Kassel, Germany
24. Department of Neurosciences, UC San Diego, La Jolla, CA, United States
25. Department of Psychiatry, Rutgers University, Robert Wood Johnson Medical School, Piscataway, NJ, United States
26. Division of Movement Disorders, Department of Neurology, Tel Aviv Soursaky Medical Center, Tel Aviv, Israel.
27. Columbia University Irving Medical Center, NY, NY.
28. Departments of Psychiatry, University of Pennsylvania School of Medicine, Philadelphia, PA, United States
29. Parkinson's Disease Research, Education and Clinical Center, Philadelphia Department of Veterans Affairs
30. Department of Neurology and Neurological Sciences, Stanford University, Stanford CA, United States
31. Department of Neurosurgery Stanford University, Stanford CA, United States
32. Wu Tsai Neuroscience Institute, Stanford University, Stanford CA, United States
33 Knight Initiative for Brain Resilience, Stanford University, Stanford CA, United States

Corresponding author:
Sarah Brooker, MD, PhD
303 East Chicago Avenue, Ward 12-140, Chicago, IL, 60611
Phone: 312-695-7950
Email: sarah-brooker@northwestern.edu

Word count: 3700

Running Title: PD with negative alpha-synuclein aggregation

Financial Disclosure/Conflict of Interest pertaining to this manuscript:
S.M.B., S.H.C., D.E.L, A.S., C.M.T., T.F.T, T.F., L.M.C., B.M., K.M.M., D.G., C.S.C., R.D.D., E.G.B., R.N.A., D.W., K.M., T.S., P.G.L, N.P., and K.L.P. receive research support and/or consulting fees from the Michael J. Fox Foundation. U.J.K. and K.L.P are on the Scientific Advisory Board of Amprion, Inc. B.M. has received honoraria for consultancy from Amprion. The other co-authors do not report any conflict of interest related to this article.

Funding Sources
PPMI – a public-private partnership – is funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners, including 4D Pharma, Abbvie, AcureX, Allergan, Amathus Therapeutics, Aligning Science Across Parkinson's, AskBio, Avid

Radiopharmaceuticals, BIAL, BioArctic, Biogen, Biohaven, BioLegend, BlueRock Therapeutics, Bristol-Myers Squibb, Calico Labs, Capsida Biotherapeutics, Celgene, Cerevel Therapeutics, Coave Therapeutics, DaCapo Brainscience, Denali, Edmond J. Safra Foundation, Eli Lilly, Gain Therapeutics, GE HealthCare, Genentech, GSK, Golub Capital, Handl Therapeutics, Insitro, Jazz Pharmaceuticals, Johnson & Johnson Innovative Medicine, Lundbeck, Merck, Meso Scale Discovery, Mission Therapeutics, Neurocrine Biosciences, Neuron23, Neuropore, Pfizer, Piramal, Prevail Therapeutics, Roche, Sanofi, Servier, Sun Pharma Advanced Research Company, Takeda, Teva, UCB, Vanqua Bio, Verily, Voyager Therapeutics, the Weston Family Foundation and Yumanity Therapeutics. This study is also funded by the NIH/NINDS

Abstract

Background: The CSF alpha-synuclein (a-syn) seed amplification assay (CSFasynSAA) detects a-syn aggregation in over 90% of individuals with sporadic PD (sPD). However, the clinical characteristics of sPD with negative CSFasynSAA have not been well defined.

Objectives: Describe clinical and neuroimaging characteristics of individuals with sPD enrolled in the Parkinson's Progression Markers Initiative (PPMI) who have a negative CSFasynSAA.

Methods: Data were from the PPMI. We compared individuals in the sPD cohort who had a negative CSFasynSAA (SAA-, n=79) to a group with positive CSFasynSAA (SAA+, n=237) propensity-score matched on age, sex, and time since clinical diagnosis at baseline assessment. Clinical parameters, DAT-SPECT, and MRI brain volumetric analyses were analyzed.

Results: The SAA- and matched SAA+ groups had similar motor performance on the MDS-UPDRS part III and similar cognitive performance on the MoCA at baseline. The proportion with severe hyposmia was 12% for SAA- versus 73% for SAA+ participants ($p < 0.001$). The SAA- group showed a higher degree of atrophy in subcortical brain regions including substantia nigra. Longitudinally, the SAA- group had faster time to reach disease progression milestones, and 14.3% had a change in diagnosis, most commonly to multiple system atrophy.

Conclusions: At baseline, SAA- sPD PPMI participants have a substantially lower rate of hyposmia, but they otherwise cannot be readily distinguished from SAA+ participants based on clinical characteristics. However, SAA- participants have a greater degree of subcortical brain atrophy, faster time to reach disease milestones, and approximately 1 out of 6 received a change in research diagnosis.

Introduction

Pathologically, Parkinson's disease (PD) is characterized by intra-neuronal accumulation of alpha-synuclein (a-syn) within Lewy bodies. Until recently, confirmation

of a-syn aggregation was only feasible via pathologic assessment of post-mortem brain tissue. As such, clinical trials use established clinical diagnostic criteria,[1] and when enrolling newly diagnosed PD may supplement with a biomarker of presynaptic dopaminergic neuronal dysfunction such as dopamine transport (DAT)-SPECT.[2] While DAT-SPECT increases diagnostic accuracy, DAT-SPECT abnormalities are non-specific for PD,[3] and clinical trials using these criteria still may enroll a subset of individuals without Lewy body pathology.

We have recently introduced a research framework for a biological definition of Neuronal Synuclein Disease (NSD) that includes biomarkers specific for identifying people with Lewy body pathology.[4] One use of this framework could be to increase the positive predictive value of the clinical diagnosis for presence of Lewy body pathology early in the disease course when the clinical diagnosis may be less accurate. This framework leverages advances in a-syn biomarker development, specifically the a-syn seed amplification assay from cerebrospinal fluid (CSFasynSAA).[5] CSFasynSAA has emerged as a reliable biomarker for detection of predominantly neuronal a-syn aggregation, and CSFasynSAA has been validated as a highly sensitive and specific technique for differentiating those with a clinical diagnosis of PD from healthy control individuals.[6-8] Across multiple cohorts, including the Parkinson's Progression Markers Initiative (PPMI), over 90% of people with sporadic PD (sPD) have positive CSFasynSAA.[9, 10]

However, some individuals that meet sPD clinical diagnostic criteria do not have aggregated a-syn on the CSFasynSAA. It remains unknown how those with clinical sPD with a negative CSFasynSAA (SAA-), differ from those with positive CSFasynSAA (SAA+). Answering this question could clarify the potential utility of the CSFasynSAA assay, and the application of the NSD framework, to clinical trial design. Specifically, if the progression of SAA- individuals differs from those who are SAA+, inclusion of SAA- individuals into clinical trials could introduce heterogeneity that is unrelated to Lewy body pathology.

To address this knowledge gap, we aimed to determine whether sPD participants in the PPMI cohort who are SAA- at baseline have distinct clinical and neuroimaging

characteristics compared to SAA+ participants. We also sought to evaluate longitudinal disease progression amongst SAA- sPD participants and determine whether the diagnosis of PD in these individuals was revised over time.

Methods

Study population

Data were from PPMI, a multicenter prospective cohort study. The PPMI study protocol, including biofluid collection techniques and storage processes, have been published elsewhere in detail.[11] Participants included in the study were those enrolled in the sPD PPMI cohort prior to March 10, 2025, who had CSFasynSAA performed at baseline enrollment. The inclusion criteria for enrollment in the sPD PPMI cohort were: 1) clinical diagnosis of PD, 2) disease duration of 2 years or less, and 3) abnormal DAT-SPECT based on study criteria. Participants were excluded from this analysis if they had: 1) inconclusive or type II CSFasynSAA result at baseline analysis (see below for details), 2) were PPMI participants with a pathogenic variant in PD-associated genes (see below for details).

CSFasynSAA

CSF analysis was performed at the baseline visit for all participants. Assessment of aggregated a-syn was performed via a CSFasynSAA completed by Amprion (Amprion Inc, San Diego, CA) as previously described.[9, 12] There are two protocol variations that have been used in PPMI, the Amprion 150hr assay and the newer Amprion 24hr assay. The following assay parameters were used to define CSFasynSAA results: Fmax (highest fluorescence intensity), T50 (time to 50% of Fmax), and TTT (time to reach a target relative fluorescence unit threshold). Based on interpretation of these parameters, the following possible results were obtained: negative CSFasynSAA (SAA-), positive CSFasynSAA (SAA+), and inconclusive. The positive 24hr CSFasynSAA shows a distinct type I pattern in PD subjects and type II pattern in individuals with a pathologically confirmed diagnosis of multiple system atrophy (MSA).[12] Therefore, for participants for whom both the 150hr and 24hr CSFasynSAA were performed, results of

the 24hr CSFasynSAA were used to determine CSFasynSAA status. The number of SAA- participants who had the 150hr and 24hr CSFasynSAA protocol, and the concordance between assay results is detailed in Table S1. Regarding blinding, an elective return of research information (RORI) PPMI initiative was initiated in December 2023 so some participants and investigators may have been informed of CSFasynSAA results. Future work is needed to determine whether RORI has impacted investigator diagnosis classifications in PPMI.

Clinical Assessments

Demographic data was collected per the PPMI protocol as previously reported, including age at enrollment, sex, race, Hispanic or Latino ethnicity, and education.[11] Time since diagnosis at date of enrollment and family history of PD was recorded. The PPMI study includes clinical measures assessing motor performance, non-motor symptoms, and cognitive function conducted annually. This study included data on Hoehn & Yahr Stage, Movement Disorder Society-Unified Parkinson's Disease Rating Scale (MDS-UPDRS)-Parts I, II, III, and IV scores, percentile score on the University of Pennsylvania Smell Identification Test (UPSIT), Modified Schwab & England Activities of Daily Living score, REM Sleep Behavior Disorder (RBD) screen questionnaire score (RBDSQ), Geriatric Depression Scale (GDS), and Scales for Outcomes in Parkinson's disease-Autonomic (SCOPA-AUT), including total SCOPA-AUT score and subscores in each autonomic domain. For cognitive function, the Montreal Cognitive Assessment (MoCA) score was used as well as investigator diagnosis of cognitive status (normal, mild cognitive impairment, or dementia). To assess motor asymmetry, a motor symptom asymmetry index was calculated as follows: lateralized items on MDS-UPDRS-part III were assessed (items 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.15, 3.16, and 3.17) and the following calculation was made: absolute value of (right side – left side)/(right side + left side). At annual visits PPMI site investigator makes a clinical determination of most likely diagnosis. For this study Dementia with Lewy Bodies was not considered a change in diagnosis as it is a neuronal synuclein disease.

Genetic testing

Genotyping methods for the PPMI study are described at ppmi-info.org. Participants with known pathogenic variants identified in the *LRRK2*, *GBA1*, *PRKN*, or *SNCA* genes were excluded from analysis. Genetic testing for the *LRRK2* G2019S variant was available for 76/80 SAA- sPD participants. Assessment for all known pathogenic *LRRK2* variants was available on 28/80 of SAA- participants at time of analysis.

Dopamine transporter imaging analysis

DAT-SPECT at baseline visit was used to evaluate dopamine dysfunction in SAA- and SAA+ groups. Dopamine dysfunction was assessed quantitatively by the following parameters: lowest putamen specific binding ratio (SBR), percent expected for age and sex, mean striatum binding, mean caudate binding, mean putamen binding, and the DAT binding asymmetry index, calculated as previously described.[13]

Magnetic Resonance Imaging (MRI) analysis

MRI sequences for PPMI are described at [ppmi-info.org](https://www.ppmi-info.org) (https://www.ppmi-info.org/sites/default/files/docs/PPMI2.0_002_MRI_TOM_Final_v4.0_20221118_FE.pdf). T1-weighted MRI at baseline visit was used to evaluate atrophy patterns between SAA- and SAA+ sPD groups. Similar to our previous work, deformation-based morphometry (DBM) was used as the main measure of regional brain atrophy.[14-16] Briefly, all T1-weighted images were pre-processed including denoising,[17] non-uniformity correction,[18] and intensity normalization. All images were then registered linearly and nonlinearly to MNI-ICBM-2009c template. All steps underwent visual quality control and participant scans that didn't pass the quality control in any step were excluded from analysis (67/516 scans ~ 12.98%), consistent with the PPMI dataset failure rate at 13.10%. Voxel-wise DBM maps were generated by calculating the determinant of the Jacobian based on the deformation field based on the nonlinear transformations from subject T1-weighted to the template. These voxel-wise DBM maps were then averaged using an accurate atlas of 22 subcortical brain regions based on integration of BigBrain and MNI-ICBM-2009c template to calculate the regional DBM values for each participant scan for the statistical analysis.[19] MRI analysis was carried out with the matched subset of subjects based on propensity-score matching as well as the complete unmatched sample.

Longitudinal outcomes

Disease progression was assessed using time to domain-based milestones as described.[20] This methodology assesses 25 disease milestone metrics across six domains: "walking and balance," "motor complications," "cognition," "autonomic dysfunction," "functional dependence," and "activities of daily living." Other outcomes assessed include Hoehn & Yahr stage ≥ 3, moderate to severe non-dyskinesia motor complications defined as a score of ≥ 3 on MDS-UPDRS items 4.3 or 4.4, and research diagnosis.

Neuropathologic assessment

Autopsy evaluation of post-mortem brain tissue was performed on one SAA- sPD participant. Neuropathologic assessment was performed as described previously.[21]

Statistical Analysis

Baseline demographic, clinical, and DAT-SPECT characteristics are summarized for SAA- and SAA+ sPD groups, using median (interquartile range, IQR) and/or mean (standard deviation, SD) for continuous measures and frequency (percentage) for categorical measures. Differences in baseline characteristics between groups were assessed using the Wilcoxon rank sum test for continuous variables, and Chi-Square (or Fisher's exact test when at least one expected cell count is below 5) for categorical variables.

For longitudinal survival analyses, Kaplan-Meier curves were generated for each endpoint as defined by Brumm et al.[20]: any progression milestone, 6 domain milestones; Hoehn & Yahr stage ≥ 3; MDS-UPDRS item 4.3 or 4.4 ≥ 3). These curves are presented by group over a 3-year period from enrollment, with at-risk tables and log-rank p-value. Participants who did not experience an event during the follow-up period were censored at the time of their last assessment within 3 years. Participants who had already reached any progression milestone or either of the two other survival outcomes at the baseline visit were excluded from analysis.

To account for age differences at enrollment between SAA- and SAA+ sPD groups, a propensity-score matching approach was employed where SAA+ sPD participants were

matched to the SAA- sPD group based on age, sex, and time since diagnosis. The matching process utilized a caliper width of 0.25 times the standard deviation of the logit of the propensity score with an optimal fixed-ratio matching procedure designed to maximize the number of matched units while minimizing the standardized mean differences.

Statistical analyses were performed using SAS v9.4 (SAS Institute Inc., Cary,NC, USA; sas.com; RRID:SCR_008567). Sankey diagrams were prepared using the "ggsankey" package in R (version 4.1.1; R Core Team 2024).

Data Sharing

Data used in preparation of this article were obtained on March 10, 2025 from the PPMI database (www.ppmi-info.org/access-data-specimens/download-data), RRID:SCR_006431. For up-to-date information on PPMI, visit www.ppmi-info.org. This analysis was conducted by the PPMI Statistics Core and used actual dates of activity for participants, a restricted data element not available to public users of PPMI data. Statistical analysis codes used to perform the analyses in this article are shared on Zenodo (10.5281/zenodo.15238103). Protocol information for The Parkinson's Progression Markers Initiative (PPMI) Clinical - Establishing a Deeply Phenotyped PD Cohort AM 3.2. can be found on protocols.io or by following this link: https://dx.doi.org/10.17504/protocols.io.n92ldmw6ol5b/v2.

Results

Sample Characteristics

Of the total sPD PPMI participants 80 were SAA- at baseline compared to 856 who were SAA+ at baseline. Of the 80 SAA- participants, repeat CSFasynSAA data at follow-up visits was available for 26. Twenty-two (84.6%) of these participants remained SAA- on follow-up while 4 (15.4%) had a type II result on follow-up. None of the SAA- participants had a positive type I SAA on follow-up.

Baseline Demographic Characteristics of SAA- *sPD participants*

The median age at enrollment for the SAA- group was 66.8 years (IQR 61.7-73.1) compared to 63.9 years (IQR 57.2-69.9) in the SAA+ group, p=0.001 (Table S2). The median time since diagnosis and distribution of sex was comparable between groups (Table S2).

Due to the statistically significant between-group age difference, subsequent analyses used a reduced dataset matched for age, sex, and time since diagnosis, as described in the Methods section. Of note, one SAA- participant was excluded from matched analysis due to missing data on disease duration.

Baseline Clinical Characteristics

Baseline demographic and clinical characteristics of the matched SAA- (n=79) and SAA+ (n=237) sPD groups are summarized in Table 1 and Table S3. Notably, the median percentile score on the UPSIT was 55.0 (IQR 26.0-78.5) for the SAA- sPD group compared to 8.0 (IQR 4.0-16.0) for SAA+. The percentage of participants with an UPSIT score less than or equal to the 15[th] percentile was 12% for SAA- versus 73% for SAA+. The SAA- group had higher scores on the MDS-UPDRS part II (p = 0.003), but there was no significant difference in MDS-UPDRS Part III scores in the OFF state (p = 0.648) (Fig. 1A). Analysis of tremor subscores demonstrated a lower rest tremor subscore in the SAA- group compared to SAA+ (p = 0.002). Regarding non-motor features, there was no significant difference between groups in cognitive performance on the MoCA (p = 0.531) (Fig. 1A) or autonomic features as assessed by median SCOPA-AUT score (p=0.151).

Baseline DAT-SPECT metrics

There were no significant differences between matched SAA- and SAA+ sPD groups when comparing DAT-SPECT imaging metrics at baseline assessment including lowest putamen ratio, mean striatum binding, mean caudate binding, mean putamen binding, and the DAT binding asymmetry index (Table 1). The association between dopamine transporter binding, olfaction, and CSFasynSAA results is shown in Fig.1B.

Baseline Structural MRI analysis

Compared to the propensity-score matched SAA+ group, SAA- sPD participants had a significantly higher degree of atrophy in 9 out of 22 subcortical regions including bilateral substantia nigra, subthalamic nucleus, globus pallidus interna and externa, and left red nucleus (FDR corrected p-values < 0.05) (Fig. 2, Table S4). While not significant, the remaining 13 regions except for the right caudate showed a trend towards higher atrophy. To ensure the robustness of our findings, we repeated the analysis on the full unmatched sample, including Age and Sex as covariates. The results were consistent, except for the left putamen, which reached statistical significance in the full-sample analysis (Table S5).

Longitudinal Analysis

Median follow-up from baseline was 1.9 years for the SAA- group. At 2 years there was data available for 53% of SAA- sPD participants. 39% of SAA- participants had not yet reached the 2-year enrollment timepoint, 4% were lost to follow-up or were overdue for their 2-year visit, and 5% withdrew from the study (Table S6).

Compared to the propensity-score matched SAA+ sPD group, the SAA- group had a faster time to reach any domain-based disease milestones (Fig. 3). Stratification of the SAA+ participants by NSD stage demonstrated that the SAA- group progressed at a rate similar to those SAA+ participants who were in NSD stage 4 at enrollment (Fig. S1). The most notable difference in attainment of milestones was in the proportion of participants who met the walking and balance milestones, which includes complications such as freezing of gait and postural instability. The walking and balance milestones were met by 13% of SAA- participants within 2 years compared to 3% of SAA+. For those SAA- participants who reached walking and balance milestones the most common milestone met was postural instability (Table S7).

Diagnostic change over time

Amongst the 80 SAA- sPD participants, 77 had at least one follow-up visit and of these participants 11 (14.3%) had a change in research diagnosis at a follow-up visit (Fig. 4). By contrast, of the 237 matched SAA+ sPD group there were 231 who had at least one follow up visit, and only 2/231 (0.9%) had a change in diagnosis.

In the SAA- group, 6 participants (7.8%) had their research diagnosis changed to MSA [11]. The other changed diagnoses were progressive supranuclear palsy (PSP), corticobasal syndrome (CBS), primary progressive freezing of gait, and prodromal synucleinopathy, each in one individual participant. In addition, one participant was deemed not to have PD or another neurologic disorder. Amongst the six participants for whom the diagnosis was changed to MSA, repeat CSFasynSAA data was available for two of these participants one of whom remained SAA- and the other had a type II MSA-like result at 1-year follow-up. To determine whether participants with a change in diagnosis impacted analysis of longitudinal disease progression we performed assessment of time to reach domain-based milestones with exclusion of the 11 SAA-participants who had a change in research diagnosis. With these participants excluded, the SAA- group still had a faster time to reach any domain-based disease milestone (p = 0.0023), and a faster time to walking and balance milestones (p = 0.0007) (Fig. S2).

Pathologic Assessment

Post-mortem pathologic assessment of brain tissue was available for one of the SAA-sPD participants. The research diagnosis for this individual was changed from PD to MSA at year 4 of follow-up. The participant died 7 years following enrollment in the study. Post-mortem brain tissue examination demonstrated glial cytoplasmic inclusions of a-syn consistent with a diagnosis of MSA-P.

Discussion

In this study, we assessed clinical and neuroimaging characteristics of sPD participants in the PPMI cohort who lacked evidence of a-syn aggregation in the CSF and compared these participants to SAA+ sPD participants matched by key demographic characteristics. We found that there were several clinical features that differed between the two groups. Foremost amongst these was hyposmia with 73% of SAA+ participants performing at or less than the 15[th] percentile on the UPSIT scale while only 12% of SAA- sPD participants were hyposmic by this parameter. This finding is consistent with prior reports of high correlation between hyposmia and a-syn pathology.[9, 22] However, the SAA- and SAA+ sPD groups had similar baseline motor features and cognitive

assessments and thus the data suggest that aside from hyposmia, standard clinical assessments alone at the time of clinical diagnosis are inadequate to identify individuals who may lack a-syn pathology. Using MRI deformation-based morphometry, the SAA- participants showed significantly greater atrophy at a group level in multiple subcortical regions, including the substantia nigra, subthalamic nucleus, globus pallidus, putamen, and red nucleus. These differences were significant after matching for age, suggesting that the observed atrophy is not solely age-related.

On longitudinal assessment, progression to clinically meaningful milestones was faster in SAA- individuals, taking into account the limitations of our small number of participants with long-term follow up data. The SAA- individuals had a notably high rate of diagnostic change as well. This suggests that CSFasynSAA status could be important to consider in clinical trial design, particularly for a-syn targeting therapies.

When interpreting the study results, we consider potential causes of negative CSFasynSAA, including: 1) a true negative in an individual with a clinical syndrome indistinguishable from sPD, 2) a false negative in an individual with sPD and Lewy body pathology, and 3) a true negative in an individual who was erroneously diagnosed with sPD but actually has an alternative pathology, such as a 4-repeat tauopathy or glial cytoplasmic inclusions. To this last consideration, 14.3% of SAA- sPD participants had a change in research diagnosis on follow-up, most commonly to MSA. However, extended longitudinal assessment is needed for those who are still within a few years of diagnosis. In regard to possible false negative results, for the subset of SAA- sPD participants for whom repeat CSF analysis was available on follow-up 85% remained SAA-, 15% had a change to the type II MSA-like CSFasynSAA result and no participants had a change to type I CSFasynSAA positive. Therefore the consistency of the results is evidence in support of a true negative result. The one SAA- participant with autopsy confirmation of glial cytoplasmic inclusions, and no Lewy body pathology, also supports this conclusion. However, additional neuropathological concordance data is necessary to interpret a SAA- result in sPD as our data do not confirm that SAA- status in parkinsonian individuals negates the possibility that they have Lewy bodies at autopsy. Therefore, we do not purport that this study should be extrapolated to the clinical setting and do not suggest that SAA- parkinsonian patients be told they do not

have a Lewy body disease based on a biomarker alone. Rather, clinicians should continue to use all of the clinical and biological data, including biomarkers when available, to make the best diagnosis.

The median age at disease onset for the SAA- sPD group was significantly higher than the SAA+ group and we performed age-matching to account for the potentially confounding effects of age. However, the older age in the SAA- group may in itself be biologically relevant as older individuals may be more likely to have non-Lewy body pathologies including 4-repeat tauopathies such as PSP or corticobasal degeneration.[23, 24] In addition to identifying individuals with pathologic a-syn, there is an urgent need for development of biomarkers for other pathologies to improve diagnostic accuracy, assess co-pathology, and improve clinical trial design.

In this study, individuals with pathogenic variants in *LRRK2*, *GBA1*, *PRKN*, and *SNCA* were excluded as they were not classified as sporadic. Notably, amongst people with LRRK2-PD approximately one third lack a-syn containing Lewy body pathology in post-mortem brain tissue,[25, 26] and 32.5% have negative CSFasynSAA results.[9] Similarly, PD patients with pathogenic *PRKN* variants often lack neuronal a-syn on brain autopsy.[27, 28] Since some genetic forms of PD lack pathologic a-syn, it is possible that a subset of the SAA- participants in this study may have an as yet unrecognized genetic cause of PD resulting in a clinically indistinguishable, yet biologically distinct disease process.

Limitations and future studies

A limitation of this study is the limited long-term follow-up of SAA- sPD participants. The PPMI study will continue longitudinal follow-up to characterize symptom progression and diagnostic change in the SAA- sPD group. Another factor impacting interpretation of this study is the relatively small number of SAA- sPD participants, which is reflective of the high rate of CSFasynSAA detection in sPD. One additional factor impacting study interpretation is that the SAA- sPD group is heterogeneous with a subset of individuals likely falling into distinct disease categories, including those with MSA, PSP, or CBD, who were mis-diagnosed as sPD. An additional limitation is that genetic testing for the *LRRK2* G2019S variant was only available for 95% of participants and full variant assessment was only available for 35%. However, the rate of *LRRK2* positivity amongst

individuals with PD is approximately 2.4%,[29] and thus amongst the SAA- participants without genetic testing it is unlikely that any individuals have a pathogenic *LRRK2* variant. Future work will involve detailed genetic analysis of SAA- sPD participants as well as comprehensive assessment of additional fluid biomarkers in this cohort.

Conclusion

SAA- sPD PPMI participants have a substantially lower rate of hyposmia compared to SAA+ participants, but otherwise cannot be readily distinguished by baseline clinical characteristics. However, SAA- participants had a faster time to reach disease milestones and a substantial proportion of SAA- participants had a change in diagnosis. Together, this study confirms the utility of determining CSFasynSAA results in sPD clinical trials where inclusion of individuals unlikely to have Lewy body pathology may confound trial data.

Acknowledgements

PPMI – a public-private partnership – is funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners, including 4D Pharma, Abbvie, AcureX, Allergan, Amathus Therapeutics, Aligning Science Across Parkinson's, AskBio, Avid Radiopharmaceuticals, BIAL, BioArctic, Biogen, Biohaven, BioLegend, BlueRock Therapeutics, Bristol-Myers Squibb, Calico Labs, Capsida Biotherapeutics, Celgene, Cerevel Therapeutics, Coave Therapeutics, DaCapo Brainscience, Denali, Edmond J. Safra Foundation, Eli Lilly, Gain Therapeutics, GE HealthCare, Genentech, GSK, Golub Capital, Handl Therapeutics, Insitro, Jazz Pharmaceuticals, Johnson & Johnson Innovative Medicine, Lundbeck, Merck, Meso Scale Discovery, Mission Therapeutics, Neurocrine Biosciences, Neuron23, Neuropore, Pfizer, Piramal, Prevail Therapeutics, Roche, Sanofi, Servier, Sun Pharma Advanced Research Company, Takeda, Teva, UCB, Vanqua Bio, Verily, Voyager Therapeutics, the Weston Family Foundation and Yumanity Therapeutics.

Authors' Roles

(1) Research Project: A. Conception, B. Organization, C. Execution; (2) Statistical Analysis: A. Design, B. Execution, C. Review and Critique; (3) Manuscript Preparation: A. Writing of the First Draft, B. Review and Critique.

S.M.B. 1A, 1B, 1C, 2A, 2C, 3A, 3B

J.P. 1A, 1B, 1C, 2A, 2C, 3B

S.H.C. 1B, 1C, 2A, 2B, 2C, 3B

D.E.L.: 1B, 1C, 2A, 2B, 2C, 3B

S.M.F. 1A, 1C, 2B, 2C, 3B

Y.Z. 1A, 1C, 2B, 2C, 3B

P.G. 1C, 2C, 3B

G.M.R. 1C, 2C, 3B

H.A. 1C, 2C, 3B

R.M. 1C, 2C, 3B

U.J.K. 1C, 2C, 3B

A.S. 1A, 2C, 3B

C.M.T. 1A, 2C, 3B

T.F.T. 1A, 2C, 3B

T.F. 1A, 2C, 3B

L.M.C. 1A, 1B 2C, 3B

B.M. 1A, 2C, 3B

K.M.M 1A, 2C, 3B

D.G. 1A, 2C, 3B

C.S.C. 1A, 2A, 2C, 3B

R.D.D. 1A, 2C, 3B

E.G.B. 1A, 2C, 3B

R.N.A. 1A, 2C, 3B

D.W. 1A, 2C, 3B

K.M. 1A, 1B, 2C, 3B

T.S. 1A, 1B, 2C, 3B

P.G.L 1A, 2C, 3B

N.P. 1A, 1B, 2A, 2C, 3B

K.L.P. 1A, 1B, 2A, 2C, 3B

Financial Disclosures of all authors (for the preceding 12 months)

S.M.B receives funding from the NIH/NINDS and the Michael J. Fox Foundation. She has received travel expense reimbursement from the Parkinson Study Group and The Michael J Fox Foundation.

J.P. has no disclosures.

S.H.C, is funded by grants from the Michael J Fox Foundation for Parkinson's Research.

D.E.L. is funded by grants from the Michael J Fox Foundation for Parkinson's Research.

S.M.F. has no disclosures

Y.Z. has received research funding from the Fonds de recherche du Québec—Santé Chercheurs boursiers et chercheuses boursières en Intelligence artificielle, Natural Sciences and Engineering Research discovery grant, and Canadian Institutes of Health Research.

P.G. has no disclosures.

G.M.R. has no disclosures.

H.A. has received a PhD scholarship from Fonds de la Recherche du Québec—Santé and Parkinson's Canada.

R.M. has no disclosures

U.J.K. is supported by research grants from NIH, Parekh Center for Interdisciplinary Neurology. U.JK. is on SAB for Amprion, Inc. and NurrON. He is a consultant for UCB, Lundbeck, and HanAll.

A.S. has been a consultant to the following companies in the past year: Acadia, Eli Lilly and Co, Neurocrine, Theravance, Cerevance, Spark/Roche, Boerhinger-Ingelheim, Wave Life Sciences, Inhibikase, Prevail, Mitzubishi and Alertity Therapeutics. He has served on DSMBs for the Huntington Study Group and The Healey ALS Consortium (Massachusetts General Hospital).   He has received grant funding from the Michael J. Fox Foundation, NIA and NINDS.

C.M.T. declares consultancies for Neurocrine, Praxis (advisory board via the International Parkinson's and Movement Disorder Society), Jazz/Cavion, Roche Genentech and Bial . CMT also declares grant support to her institution from The Michael J. Fox Foundation, National Institutes of Health, Gateway LLC, Department of Defense, Roche Genentech and Parkinson Foundation.

T.F.T. has received research support from NIH, The Michael J Fox Foundation, the Parkinson Foundation, and Eli Lilly and Company. Dr Tropea serves as a clinical trial advisory board member for Bial, has received travel expense reimbursement from the Parkinson Study Group, The Michael J Fox Foundation and the Parkinson Foundation, and speakers fees from Catalyst Medical Education.

T.F. receives funding from the NIH and Michael J. Fox Foundation and serves on several advisory committees for NIH funded studies (with < $1,000 each).

L.M.C. Financial disclosure: consulting fees and research support from the Michael J Fox Foundation. COI: none

B.M. has received honoraria for consultancy and/or educational presentations from GE, Bial, Roche, Biogen, AbbVie, Desitin and Amprion. BM is member of the executive steering committee of the Parkinson Progression Marker Initiative of the Michael J. Fox Foundation for Parkinson's Research and has received research funding from Aligning

Science Across Parkinson's disease (ASAP, CRN), the German Parkinsonstiftung and the Michael J. Fox Foundation.COI: none

K.M.M.: Consultant for Axial Therapeutics, Asceneuron, Calico, Hanall, JnJ, Michael J. Fox Foundation, Nitrase Therapeutics, NuraBio, NRG Therapeutics, Rome Therapeutics, Schrodinger, Ventyx, private equity companies. Serves on the BoD for Envisagenics, Retromer Tx; serves on the SAB for Axial, Nitrase, NRG, Sinopia, Vanqua; Received research Funding support from Michael J. Fox Foundation and Honoraria from ASAP.

D.G. has received grant support from the NIH and Michael J Fox Foundation, and is a consultant for Eisai, Lilly, Artery Therapeutics and Cognition Therapeutics.

C.S.C. receives funding from the NIH/NINDS, MJFF, and PCORI.

R.D.D. receives research funding from the Michael J. Fox foundation and the VA office of rural health.

E.G.B. has received grant support for research from the Michael J. Fox Foundation, the National Institutes of Health, the Department of Defense, and Gateway Institute for Brain Research, Inc. He has received consulting fees from Guidepoint Inc.

R.N.A. research is funded by the Michael J. Fox Foundation, the Silverstein Foundation, the Parkinson's Foundation and the Aufzien Family Center for the Prevention and Treatment of Parkinson's Disease. He received consultation fees from Bexxion, Biogen, Biohaven, Capsida, Gain Therapeutics, Genzyme/Sanofi, Janssen, SK Biopharmaceuticals, Takeda and Vanqua Bio.

D.W. has received research funding or support from Michael J. Fox Foundation for Parkinson's Research, International Parkinson and Movement Disorder Society (IPMDS), National Institute on Health (NIH) and the U.S. Department of Veterans Affairs; honoraria for consultancy from AbbVie, Boehringer Ingelheim, CHDI Foundation, Citrus Health Group, Negev Labs, Otsuka, Parkinson Study Group, Sage Therapeutics, Signant Health and Vanqua Bio; and license fee payments from the University of Pennsylvania for the QUIP, QUIP-RS and PDAQ.

K.M.:Consultant for Michael J Fox Foundation, CPP, Roche, Biohaven, Neuron23, Prothena, Lilly, Calico, ABli, Mitro, BMS, Novartis, Teva.

T.S. declares consultancies for AcureX, Adamas, AskBio, Amneal, Blue Rock Therapeutics, Critical Path for Parkinson's Consortium, Denali, The Michael J. Fox Foundation, Neuroderm, Roche, Sanofi, Sinopia, Takeda, and Vanqua Bio; on advisory boards for AcureX, Adamas, AskBio, Biohaven, Denali, GAIN, Neuron23 and Roche; on scientific advisory boards for Koneksa, Neuroderm, Sanofi and UCB; and received research funding from Amneal, Biogen, Roche, Neuroderm, Sanofi, Prevail and UCB and an investigator for NINDS, MJFF, Parkinson's Foundation.

P.G.L declares an early investigator award from The Michael J. Fox Foundation

N.P. has received consultancy honoraria form: Hoffmann-La Roche, Britannia, Bial, AbbVie, Teva; Speaker fees from: Bial, Britannia, AbbVie, GE Healthcare, Boston Scientific; Research Funding from Independent Research Fund Denmark, Danish Parkinson's disease Association, Parkinson's UK, Center of Excellence in Neurodegeneration (CoEN) network award, GE Healthcare Grant, Multiple System Atrophy Trust, Weston Brain Institute, EU Joint Program Neurodegenerative Disease Research (JPND), EU Horizon 2020 research, The Michael J. Fox Foundation, F. Hoffmann-La Roche, Medtronic, Symbyx.

K.L.P. is funded by grants from the NIH, Michael J Fox Foundation for Parkinson's Research, the Knight Initiative for Brain Resilience, the Wu Tsai Neuroscience Institute, Lewy Body Dementia Association, Parkinson's Foundation, American Parkinson's Disease Association, and the Sue Berghoff LBD Research Fellowship. She has been on the Scientific Advisory Board for Amprion and has been a consultant for Novartis, Lilly, BioArctic, Biohaven, Curasen and Neuron23.

References

1.   Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. Mov Disord 2015;30(12):1591-1601.

2.   Bega D, Kuo PH, Chalkidou A, et al. Clinical utility of DaTscan in patients with suspected Parkinsonian syndrome: a systematic review and meta-analysis. NPJ Parkinsons Dis 2021;7(1):43.

3.   Quintas S, Sanles-Falagan R, Berbis MA. I(123)-FP-CIT (DaTSCAN) SPECT beyond the Most Common Causes of Parkinsonism: A Systematic Review. Mov Disord Clin Pract 2024;11(6):613-625.

4.   Simuni T, Chahine LM, Poston K, et al. A biological definition of neuronal alpha-synuclein disease: towards an integrated staging system for research. Lancet Neurol 2024;23(2):178-190.

5.   Concha-Marambio L, Pritzkow S, Shahnawaz M, Farris CM, Soto C. Seed amplification assay for the detection of pathologic alpha-synuclein aggregates in cerebrospinal fluid. Nat Protoc 2023;18(4):1179-1196.

6.   Rossi M, Candelise N, Baiardi S, et al. Ultrasensitive RT-QuIC assay with high sensitivity and specificity for Lewy body-associated synucleinopathies. Acta Neuropathol 2020;140(1):49-62.

7.   Berg D, Klein C. alpha-synuclein seed amplification and its uses in Parkinson's disease. Lancet Neurol 2023;22(5):369-371.

8.   Grossauer A, Hemicker G, Krismer F, et al. alpha-Synuclein Seed Amplification Assays in the Diagnosis of Synucleinopathies Using Cerebrospinal Fluid-A Systematic Review and Meta-Analysis. Mov Disord Clin Pract 2023;10(5):737-747.

9.   Siderowf A, Concha-Marambio L, Lafontant DE, et al. Assessment of heterogeneity among participants in the Parkinson's Progression Markers Initiative cohort using alpha-synuclein seed amplification: a cross-sectional study. Lancet Neurol 2023;22(5):407-417.

10.   Orru CD, Vaughan DP, Vijiaratnam N, et al. Diagnostic and prognostic value of alpha-synuclein seed amplification assay kinetic measures in Parkinson's disease: a longitudinal cohort study. Lancet Neurol 2025;24(7):580-590.

11.   Marek K, Chowdhury S, Siderowf A, et al. The Parkinson's progression markers initiative (PPMI) - establishing a PD biomarker cohort. Ann Clin Transl Neurol 2018;5(12):1460-1477.

12. Ma Y, Farris CM, Weber S, et al. Sensitivity and specificity of a seed amplification assay for diagnosis of multiple system atrophy: a multicentre cohort study. Lancet Neurol 2024;23(12):1225-1237.

13. Kaasinen V. Ipsilateral deficits of dopaminergic neurotransmission in Parkinson's disease. Ann Clin Transl Neurol 2016;3(1):21-26.

14. Fereshtehnejad SM, Moqadam R, Azizi H, et al. Distinct Longitudinal Clinical-Neuroanatomical Trajectories in Parkinson's Disease Clinical Subtypes: Insight toward Precision Medicine. Mov Disord 2025.

15. Zeighami Y, Fereshtehnejad SM, Dadar M, Collins DL, Postuma RB, Dagher A. Assessment of a prognostic MRI biomarker in early de novo Parkinson's disease. Neuroimage Clin 2019;24:101986.

16. Zeighami Y, Ulla M, Iturria-Medina Y, et al. Network structure of brain atrophy in de novo Parkinson's disease. Elife 2015;4.

17. Coupe P, Yger P, Prima S, Hellier P, Kervrann C, Barillot C. An optimized blockwise nonlocal means denoising filter for 3-D magnetic resonance images. IEEE Trans Med Imaging 2008;27(4):425-441.

18. Sled JG, Zijdenbos AP, Evans AC. A nonparametric method for automatic correction of intensity nonuniformity in MRI data. IEEE Trans Med Imaging 1998;17(1):87-97.

19. Xiao Y, Lau JC, Anderson T, et al. An accurate registration of the BigBrain dataset with the MNI PD25 and ICBM152 atlases. Sci Data 2019;6(1):210.

20. Brumm MC, Siderowf A, Simuni T, et al. Parkinson's Progression Markers Initiative: A Milestone-Based Strategy to Monitor Parkinson's Disease Progression. J Parkinsons Dis 2023;13(6):899-916.

21. Bukhari SA, Nudelman KNH, Rumbaugh M, et al. Parkinson's Progression Markers Initiative brain autopsy program. Parkinsonism Relat Disord 2022;101:62-65.

22. Marek K, Russell DS, Concha-Marambio L, et al. Evidence for alpha-synuclein aggregation in older individuals with hyposmia: a cross-sectional study. EBioMedicine 2025;112:105567.

23.   Savica R, Grossardt BR, Bower JH, Ahlskog JE, Rocca WA. Incidence and pathology of synucleinopathies and tauopathies related to parkinsonism. JAMA Neurol 2013;70(7):859-866.

24.   Robinson JL, Lee EB, Xie SX, et al. Neurodegenerative disease concomitant proteinopathies are prevalent, age-related and APOE4-associated. Brain 2018;141(7):2181-2193.

25.   Kalia LV, Lang AE, Hazrati LN, et al. Clinical correlations with Lewy body pathology in LRRK2-related Parkinson disease. JAMA Neurol 2015;72(1):100-105.

26.   Zimprich A, Biskup S, Leitner P, et al. Mutations in LRRK2 cause autosomal-dominant parkinsonism with pleomorphic pathology. Neuron 2004;44(4):601-607.

27.   Farrer M, Chan P, Chen R, et al. Lewy bodies and parkinsonism in families with parkin mutations. Ann Neurol 2001;50(3):293-300.

28.   Kluge A, Borsche M, Streubel-Gallasch L, et al. alpha-Synuclein Pathology in PRKN-Linked Parkinson's Disease: New Insights from a Blood-Based Seed Amplification Assay. Ann Neurol 2024;95(6):1173-1177.

29.   Cook L, Verbrugge J, Schwantes-An TH, et al. Parkinson's disease variant detection and disclosure: PD GENEration, a North American study. Brain 2024;147(8):2668-2679.

Figure Legends

Figure 1: Baseline characteristics of SAA- and SAA+ sPD participants

A) Violin plots of selected clinical characteristics measured at the baseline assessment are presented for SAA- and SAA+ participants. The number of SAA- and SAA+ participants for whom data is available is indicated on the x-axis. For each plot the point indicates the median and the range indicates the interquartile range. (A) Median score on the MDS-UPDRS Part II, p = 0.003. (B) Median score on the MDS-UPDRS Part III in the off state, p = 0.648. (C) median score on the Montreal Cognitive Assessment (MOCA), p = 0.531. (D) median score on the geriatric depression scale (GDS), p = 0.025. B) The association between olfaction, dopamine transporter binding, and CSFasynSAA results is shown. Olfaction as indicated by percentile score on the UPSIT

is shown on the x-axis and DATscan lowest putamen specific binding ratio is shown on the y-axis. SAA- participants are represented with an open circle and SAA+ participants are represented with a closed circle.

Figure 2: Cross-sectional brain atrophy map of the SAA- sPD group.

T-statistic maps (FDR-corrected p < 0.05) show regions where the SAA- group exhibited significantly greater atrophy compared to the matched SAA+ group. Results are overlaid on axial slices of the MNI template, ranging from z = 58 (top left) to z = 91 (bottom right) in 3 mm increments. DBM: deformation based morphometry. The colormap indicates the t-values from the SAA-group contrast between SAA- and SAA+ participants from the linear model controlling for age and sex

Figure 3: Time to reach disease progression milestones in matched SAA- and SAA+ sPD

Kaplan-Meier curves are presented for time to reach disease progression milestones in the SAA- and SAA+ matched groups over a 3-year time period. The number of SAA- and SAA+ participants for whom data is available at each annual time point is indicated. The endpoints analyzed include: (A) Time to reach any of the 6 domain progression milestones defined by Brumm et al., (B) Walking and balance domain milestones, (C) Motor complications domain milestones, (D) cognitive domain milestones, (E) autonomic dysfunction domain milestones, (F) functional independence domain milestones, (G) activities of daily living domain milestones, (H) Hoehn & Yahr stage ≥ 3, and (I) moderate to severe non-dyskinesia motor complications defined as a score of ≥ 3 on the MDS-UPDRS item 4.3 or 4.4 which measure time in the off state and functional impact of motor fluctuations respectively.

Figure 4 Sankey diagram of primary research diagnosis in SAA- sporadic PD participants at annual visits.

A sankey diagram is presented indicating the primary research diagnosis recorded at each annual follow-up visit for the SAA- sPD group. Yellow indicates a diagnosis of Parkinson's disease (PD) and purple indicates any other diagnosis that is not PD. The height of the bar at each year is proportional to the number of participants in each

diagnosis category. For each annual timepoint the number of SAA- participants for whom research diagnosis information was available is indicated on the x-axis.

Table 1: Demographic and baseline characteristics of matched SAA- and SAA+ sporadic PD participants

Variable	SAA- sPD (N = 79[a])	SAA+ sPD (N = 237)	p-value[b]
Age at enrollment, years, median (IQR)	66.7 (61.3–73.3)	67.4 (62.1–72.5)	0.959
Male sex, n (%)	50 (63%)	150 (63%)	1.000
Family history of PD, n (%)			0.752
1st-degree family w/ PD	10 (13%)	37 (16%)	
Non-1st-degree family w/ PD	9 (11%)	30 (13%)	
No family w/ PD	60 (76%)	170 (72%)	
Disease duration at enrollment, median (IQR)	0.5 (0.3–0.8)	0.5 (0.3–0.8)	0.369
Duration of follow-up from BL, years, median (IQR)	1.9 (1.0–3.0)	2.1 (1.0–6.0)	0.060
Hoehn & Yahr stage, n (%)			0.201[c]
1	19 (24%)	75 (32%)	
2	59 (75%)	161 (68%)	
3	1 (1%)	1 (<1%)	
UPSIT percentile, median (IQR)	55.0 (26.0–78.5)	8.0 (4.0–16.0)	<.001
UPSIT ≤ 15th %ile, n (%)	9 (12%)	171 (73%)	<.001
MDS-UPDRS Part I, median (IQR)	6.0 (4.0–11.0)	5.0 (3.0–8.0)	0.069
MDS-UPDRS Part II, median (IQR)	7.0 (4.0–11.0)	5.0 (3.0–8.0)	0.003
MDS-UPDRS Part III (OFF), median (IQR)	22.5 (18.0–27.0)	22.0 (15.5–29.0)	0.648
Motor symptom asymmetry index (OFF), median (IQR)	0.40 (0.20–0.60)	0.52 (0.32–0.81)	0.001
Tremor score (OFF), median (IQR)	4.0 (2.0–7.0)	5.0 (3.0–7.0)	0.075
Rest tremor subscore (OFF), median (IQR)	2.0 (0.0–4.0)	4.0 (2.0–5.0)	0.002
MDS-UPDRS Total Score (OFF), median (IQR)	39.0 (28.0–48.0)	32.0 (25.0–43.0)	0.020
MoCA, median (IQR)	27.0 (25.0–29.0)	27.0 (25.0–29.0)	0.531
Cognitive categorization, n (%)			0.244
Normal	54 (83%)	150 (89%)	
Mild cognitive impairment	11 (17%)	19 (11%)	
GDS, median (IQR)	2.0 (1.0–5.0)	2.0 (0.0–3.0)	0.025
SCOPA-AUT, median (IQR)	11.0 (7.0–14.0)	9.0 (6.0–13.0)	0.151
Lowest putamen ratio, median (IQR)	0.35 (0.23–0.60)	0.35 (0.29–0.43)	0.736
Mean striatum binding, median (IQR)	1.29 (1.03–1.83)	1.40 (1.18–1.67)	0.557
Mean caudate binding, median (IQR)	1.87 (1.45–2.33)	1.93 (1.66–2.35)	0.188
Mean putamen binding, median (IQR)	0.84 (0.56–1.39)	0.84 (0.68–1.05)	0.704
DAT binding asymmetry index, median (IQR)	0.12 (0.07–0.20)	0.15 (0.08–0.25)	0.169

UPSIT = University of Pennsylvania Smell Identification Test; MDS-UPDRS = Movement Disorder Society-Unified Parkinson's Disease Rating Scale; MoCA = Montreal Cognitive Assessment; GDS = Geriatric Depression Scale; SCOPA-AUT = Scales for Outcomes in Parkinson's Disease-Autonomic

Missing data (if more than 5): Cognitive categorization n = 82 (consistent data collection on this variable began after 2020); DaTscan measures (lowest putamen ratio, mean binding, asymmetry index) n = 6.

[a]One SAA- sPD participant excluded from matched analysis due to missing disease duration.
[b]Comparisons by SAA status used Chi-Square or Fisher's Exact tests for categorical variables and Wilcoxon rank sum tests for continuous variables.
[c]For the purposes of comparisons, Hoehn & Yahr stage was dichotomized as stage 1 vs. ≥ 2.

Figure 1: Baseline characteristics of SAA- and SAA+ sporadic PD participants

Note: The point (range) within each violin plot indicates the median (interquartile range) for each group.

Figure 2: Cross-sectional brain atrophy map of the SAA- sporadic PD group

Figure 3: Time to reach disease progression milestones in matched SAA- and SAA+ sporadic PD

Figure 4. Sankey diagram of primary research diagnosis in SAA- sporadic PD participants at annual visits.

*Ns reflect the number of participants with non-missing research diagnosis at each timepoint.
In cases of intermittent missingness, the Sankey carries forward the diagnosis (PD or non-PD) from the last available visit.