Widespread transmission in diverse ecotypes challenges visceral leishmaniasis control in East Africa

Eva Iniguez[1†], Mercy Tuluso[2†], Steve Kiplagat[3], Araya Gebresilassie[4], Esayas Aklilu[5], Olivia Battistoni[6], Johnstone Ingonga[2], John Mark Makwatta[3], Mohamed Alamin[7], Osman Dakien[7,8], Alphine Chebet[2], Esther Kaunda[2], Patrick Huffcutt[1], Serena Doh[1], Pedro Cecilio[9], Galgallo Bonaya[3], Claudio Meneses[1], Tiago D. Serafim[1], Myrthe Pareyn[10], Mohamed Osman[11], Eltahir A. G. Khalil[7], Omran F. Osman[7], Brima M. Younis[7], Guofa Zhou[12], Jesus G. Valenzuela[1], Dan K. Masiga[3], Ahmed M. Musa[13], Asrat Hailu[14], Abhay Satoskar[6], Shaden Kamhawi[1]*, Damaris Matoke-Muhia[2]*

[1]Vector Molecular Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA

[2]Centre for Biotechnology Research and Development, Kenya Medical Research Institute, Nairobi, Kenya

[3]International Centre of Insect Physiology and Ecology (*icipe*), Nairobi, Kenya

[4]Department of Zoological Sciences, College of Natural and Computational Sciences, Addis Ababa University, Addis Ababa, Ethiopia

[5]Vector Biology and Control Research Unit, Aklilu Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia

[6]Departments of Pathology and Microbiology, Wexner Medical Center, The Ohio State University, Columbus, OH, USA

[7]Department of Zoology, Faculty of Science, University of Khartoum, Khartoum, Sudan

[8]Kala Zara Research Center, Faculty of Medicine and Health Science, University of

Gedaref, Gedaref, Sudan

[9]Vector Biology Section, Laboratory of Malaria and Vector Research, National Institute

of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, USA

[10]Department of Clinical Sciences, Institute of Tropical Medicine Antwerp, Belgium

[11]York Biomedical Research Institute, University of York, York, UK

[12]Program in Public Health, University of California at Irvine, Irvine, CA, USA

[13]Department of Clinical Pathology and Immunology, Institute of Endemic Diseases,

University of Khartoum, Khartoum, Sudan

[14]College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

[†]Equal contribution

*Corresponding authors email address: skamhawi@nih.niaid.gov and

DMatoke@kemri.go.ke

Abstract

East Africa is emerging as the global hot spot of visceral leishmaniasis (VL), yet efforts

to eliminate it are hindered by substantial knowledge gaps in its ecoepidemiology. Here,

we report on the high prevalence of *Leishmania* infection in *Phlebotomus orientalis* in

Marsabit county, Kenya (3.9%), and Gedaref state, Sudan (3.6%), where this species

comprised 99.8% (n = 1185) and 100% (n = 1350) of captured *Phlebotomus* females,

respectively. In Aba Roba, Ethiopia, *Ph. martini* accounted for 99% of 184 collected

Phlebotomus females and had a lower infection rate of 1.5%. *Ph. orientalis* and *Ph.*

martini exhibited different habitat and feeding preferences. While *Ph. orientalis* was

abundant in diverse peridomestic and sylvatic microhabitats, *Ph. martini* was

predominantly collected from termite hills. Moreover, *Ph. orientalis* primarily fed on

humans and less on domestic and sylvatic animals. In contrast, *Ph. martini* exhibited

zoophagic behavior, mostly feeding on cows and ovis. Widespread transmission of

Leishmania in our study sites is supported by high rK39 seroprevalence in both Kenya

(20.88%) and Sudan (7.36%). An observed greater prevalence of antibodies to rK39 in

individuals living near than away from VL cases in both Kenya (21.68% versus 16.51%,

$P = 0.03$) and Sudan (10.03% versus 2.05%, $P = 0.02$) demonstrated that proximity to a

VL case carries an increased risk of infection. Our findings highlight the need for a

targeted site-specific elimination strategy that accounts for the intensity, diversity, and

complexity of VL transmission in today's East Africa.

Keywords: East Africa, visceral leishmaniasis, rK39 seroprevalence, ecotype,

microhabitat, breeding sites, blood meal analysis, *Leishmania*-infected sand flies,

transmission.

Introduction

East Africa is currently the epicenter of global transmission of visceral leishmaniasis

(VL), a fatal neglected disease ranked second after malaria in mortality rates[1]. VL in

East Africa is epidemiologically and clinically diverse, caused by genetically distinct

Leishmania donovani strains[2-4], and transmitted by several sand fly vector species

including *Phlebotomus orientalis, Ph. martini,* and *Ph. celiae* that have divergent

ecologies and bionomics[5-8]. VL cases have recently surged in East Africa, emerging in

new foci that disproportionally affected Kenya, Sudan, and Ethiopia, accounting for

more than 65% of cases worldwide, out of which ~50% are children under 15 years

old[1, 9].

VL in East Africa had initially been targeted for control rather than elimination due to

significant knowledge gaps concerning disease transmission[10]. In June 2024, the

World Health Organization launched a framework to eliminate VL as a public health

concern from East Africa by 2030[11]. Elimination is proposed through early diagnosis

and treatment, integrated vector control management, and disease surveillance[11]. At

the country level, the target for elimination was defined as <1% case-fatality rate due to

primary VL, 90% decline in new VL cases, absence of death due to VL in children by

2030, and 100% detection and treatment referral of VL-HIV and PKDL patients[11].

However, East Africa faces major challenges to VL elimination. Treatment differs

between countries, and it is often toxic and lengthy, with low compliance and poor

access to care facilities[5, 11-13]. Moreover, the serological test for *Leishmania* diagnosis,

the rK39 rapid test, has a decreased sensitivity and specificity in regions of East Africa,

particularly in HIV-VL co-infected patients[14-16]. In addition, significant knowledge gaps

in ecoepidemiology, vector biology, and nature of reservoirs need to be addressed for a

successful implementation of an elimination campaign in East Africa,[5, 12] where there

are no current effective integrated vector control programs[11]. Though *Ph.* orientalis, *Ph.*

martini, and *Ph. celiae* sand flies have been found in both peridomestic and sylvatic

niches in strong association with *Acacia* and *Balanites* woodlands, vertisol (black cotton

soil), and termite hills[17-20], linking these microhabitats to VL transmission in the

region[19, 21-24], confirmation of their epidemiological relevance and their potential as

resting and breeding sites for these vectors[20, 25] needs to be demonstrated.

Additionally, while VL caused by *L. donovani* is considered anthroponotic, the

opportunistic feeding behavior reported for *Ph. orientalis* together with high transmission

rates reported in relatively scarcely populated areas of East Africa, points to the

potential presence of an animal reservoir[17, 24, 26-28]. Additionally, we need to determine

the role of asymptomatic individuals as infectious reservoirs for sand flies[29-31]. Further

hindering control of VL in East Africa is the presence of semi-nomadic populations, such

as pastoralists and farmers living in temporary shelters[12], and populations displaced by

conflict[32, 33], that are at a higher risk of contracting VL[32-34], complicating the

epidemiological landscape.

In this study, we report the prevalence of *Leishmania*-infected sand flies from diverse

microhabitats in endemic study sites in Kenya, Sudan, and Ethiopia. We also establish

widespread transmission through the high rate of rK39 seroprevalence in Marsabit county,

Kenya, and Gedaref state, Sudan, where *Ph. orientalis* is the primary vector. Altogether, this

study provides current information on VL transmission dynamics in East Africa.

Methods

Ethics approval

The study was approved by the Scientific Research Unit, KEMRI, and licensed by The

National Commission for Science, Technology and Innovation (clinical protocol #

KEMRI/SERU/CBRD/249/4634) and by the State Ministry of Health and Social

Development at Gedaref state, Sudan ethical committee (clinical protocol #

UG/EC/3/2023) in Sudan. Both clinical protocols were translated into local languages.

Communal consent was obtained by meeting with community leaders. Written consent

was obtained from adults or guardians of minors before blood collection. Verbal consent

was acquired from homeowners before entomological collections.

Study sites

We selected study sites with stable VL transmission in Kenya, Sudan, and Ethiopia. The

study sites were chosen to encompass features common to VL ecotypes in East Africa.

<u>Kenya</u>

Marsabit county has a population of 459,785 people[35], and has experienced sporadic

outbreaks since 2014[36, 37]. It has an arid and semi-arid tropical savanna climate with an

average temperature ranging between 15°C and 26°C. Average precipitation is about

128.24 mm per annum. Marsabit county has three climatic seasons: two dry seasons in

January to February and June to July, short rains in March to May, and long rains in

August to December [38]. The communal free-range land is dominated by diverse *Acacia*

and related trees, a wide variety of shrubs, and seasonal grass.

Inhabitants of Marsabit county are mainly nomadic pastoralists belonging to one of

fourteen Cushitic-speaking ethnic groups. Camel, goat, sheep, donkeys, and cattle are

major domestic animals kept as a source of income. The community moves with their

domestic animals from semi-permanent (manyatta) houses to temporary houses (foras)

for a period of up to six months in search of pasture and water. We investigated three

locations: Laisamis, Logo Logo, and Karare, in Laisamis sub-county, Marsabit county.

Sudan

Gedaref state has an urban structure with well-established villages. The region is a flat plain, with small, scattered hills and flowing watercourses, and the climate is tropical. The rainy season spans June to October with an annual rainfall of 400-1400 mm. The natural vegetation is a savanna woodland with *Acacia* and *Balanites* trees, and the principal soil type is black cotton vertisol. Agriculture is the main livelihood, with sorghum, sesame, and millet as major crops. Goats, sheep, donkeys, and cattle are major domestic animals kept as a source of income. Most people are subsistence farmers or engage in small animal husbandry. High demand for manual labor attracts seasonal workers from within Sudan or bordering Ethiopia during the rainy season. In a recent study, a high VL prevalence of 27.6% was reported among Dinder National Park wildlife soldiers[39]. Dinder National Park falls between the Sahel of Gedaref state and Ethiopian Highlands and is home to several species of large and small mammals, such as lions, gazelle, monkeys, bats and reptiles. We investigated the villages of Kunaynah, bordering north Ethiopia, and Umslala-Houng, closest to Dinder National Park, in Gedaref state.

Ethiopia

The Konso zone is a semi-arid rugged terrain with many hills, lowlands, and riverbeds. The area is generally semi-arid, with an annual rainfall of about 551 mm per year. The long rains last from February to May, followed by short rains in September and October. The mean annual temperature is 24°C. The predominant soil type is sandy. Dispersed homesteads and animal enclosures are scattered in lowlands or valleys, while villages are on top of large hills or highlands whose slopes lead to the valleys. Segen river flows

through the valley providing water for irrigation and creating flat plains suitable for

farming and animal herding. Even though there are no permanent human settlements,

many families own farms in the valleys and spend a considerable number of nights in

temporary shelters (foras) herding cattle, goats and sheep. The vegetation comprises a

mixture of shrub and forests with diverse trees, including species of *Acacia* and

Balanites. Castellated active or eroded shrinking termite hills[7] are abundant in the area.

Aba Roba communities mostly depend on subsistence farming on the cultivated

terraced hillsides and flat plains of the Segen River. Sorghum, maize, sesame and

sunflower are major crops. They also raise goats, sheep, and cattle, kept in enclosures

at night. We investigated the villages of Galga and Goinada, 1180-1250 m above sea

level, and Maira and Lahalaha, 830-885 m above sea level, in the Konso zone.

174   Sand fly collection

In each country, sand fly sampling ecotypes included peridomestic (associated with

humans) and sylvatic (away from humans) sampling microhabitats (Supplementary

Figure 1). Ecotypes included indoors (inside manyatta/hut) and outdoors (outside

manyatta/hut), temporary shelters or foras, animal enclosures, vegetation in

association or not with animal burrows and vertisols, and termite hills. The study sites

and the total number of CDC light traps used per collection varied in each country, but

the same number of traps were placed per ecotype (Supplementary Table 1).

183   Species identification of *Phlebotomus* sand flies

The head and the last three segments of each *Phlebotomus* sand fly were wet mounted

for morphological identification. Spermatheca and the pharynx teeth were examined to

distinguish the female species, while external genitalia were examined for identification

of males using a combination of keys[40, 41].

Rotation of *Phlebotomus* male genitalia

During the first 16-24 h of the adult life of a male sand fly the external genitalia undergo

a rotation to orient the claspers during mating. Thus, unrotated or partially rotated male

genitalia can be used to identify newly emerged juveniles captured during their first

night of activity as adults, as a proxy to identify sand fly breeding sites[17, 20, 22]. Under

the microscope, we recorded the number of males with rotated, unrotated or partially

rotated genitalia.

Mitigation of sample contamination from the field to the bench

In the field, all female sand flies were dissected using dissecting pins that were first

bleached (10%) and rinsed twice with water before each sand fly dissection. This step

was critical to avoid contamination caused by carry over DNA from sample to sample. In

the laboratory, three separate rooms were used to process and run the field samples by

qPCR. Room one was used only for preparation of buffer and primer aliquots and

loading of the qPCR master mix into the plate. Room two was used for extraction of

nucleic acids of field specimens. Room three was used to load the DNA samples into

the qPCR plate. Each plate included a standard curve, and DNA from an uninfected

midgut and non-template negative controls. To avoid contamination all the instruments,

benches and pipettes were continuously bleached (10%) and rinsed with water.

Extraction of DNA from field-collected *Phlebotomus* female sand flies

Unfed and gravid midgut samples were individually collected in Ethiopia, and individual

or pools of 5 midguts in Kenya. For Sudan, pools of 10 sand flies (whole bodies or

midguts) were collected. Samples were stored in 50 μl of DNA/RNA Shield buffer

(Zymo), and stored at room temperature, or 4°C if available. DNA was extracted from

individual specimens of unfed and gravid sand flies using the Quick-DNA/RNA Miniprep

Plus kits (Zymo) tissue protocol, with a proteinase K and tissue lysis digestion step of 1

h 56°C. DNA was extracted from pools of unfed and gravid sand flies using the Tissue

DNA kit (Omega) according to the manufacturer's instructions. For whole body pooled

samples, we first macerated each using a plastic pestle, followed by chemical digestion

with proteinase K and tissue lysis buffer for 2-6 h or until fully digested at 56°C.

Blood-fed sand flies were preserved individually as midguts/whole bodies on Whatman

903 Protein Saver cards (Cytica), or in up to 50 μl of DNA/RNA Shield buffer (Zymo).

Briefly, field-collected blood fed midguts were individually punched from the Whatman

cards using a 3 mm punch biopsy, and Proteinase K (20 μl), Lysis buffer (20 μl), and

DNA/RNA shield (30 μl) were added to each punched sample and digested at 56°C for

2 to 4 h, with brief vortexing preformed every hour. Blood fed samples stored in

DNA/RNA Shield buffer as midguts/whole bodies were first macerated using ceramic

bashing beads (Roche) using the MagNAlyzer (Roche), followed by digestion for 24 h.

DNA was extracted using the Quick-DNA/RNA Miniprep Plus kits (Zymo). All samples

were eluted to a final volume of 100 µL and stored at -20/-80°C until further processing.

Blood meal analysis by multiplex PCR

Standardization of a blood meal multiplex custom PCR panel

Lutzomyia longipalpis (Jacobina colony) was reared at the Laboratory of Malaria and

Vector Research, NIAID, NIH insectary. Female sand flies (4-6 day old) were allowed to

feed for 2 h on whole blood from various species through an artificial chicken skin

membrane[42] and collected 24 h post-feeding as positive controls. Blood meals included

domestic and sylvatic hosts such as mouse and chicken (LMVR4 animal protocol);

human (leftover samples collected for another study under IRB human protocol 000331-

I); cow (Charles River Laboratories); dog, goat, sheep, and donkey (Innovative

Research); hyrax and camel (donated by Dan Masiga, *icipe*).

A universal reverse primer previously reported[43], with custom forward primers

developed based on the mitochondrial *cytochrome b* region, highly conserved among

vertebrates[44], were used for several species of interest (Supplementary Table 2) . A

human primer set based on the cytochrome c oxidase I mitochondrial region[45], and the

rodent (*Muroidea* superfamily) primer set targeting the mitochondrial d-loop region were

used from previous published studies[46]. Briefly, DNA (40 ng, 10 µL) was mixed in a

single tube as a multiplex reaction with all forward and reverse primers and the

DreamTaq[TM] Hot Start Green PCR Master Mix (Thermo Scientific) to a final volume of

30 µL (Supplementary Table 2). PCR conditions were 95°C initial denaturation for 5

minutes, 35 cycles of 20 seconds at 95°C, 20 seconds at 58°C, 60 seconds at 72°C,

and a final extension of 7 minutes at 72°C. PCR products were separated on a 2%

agarose gel electrophoresis ran at 8 v/cm for 60 minutes. Two microliters of a 50 to

1000 base-pair ladder (Thermo Scientific) was used. Bands were visualized on a c600

Azure Biosystems imager.

Bloodmeal analysis for field-collected samples

PCR of DNA from field samples was performed following the same procedure and

conditions described above and in Supplementary Table 2. The PCR products were run

on a 2% agarose gel, cut and purified (Thermo Scientific). The PCR conditions were

94°C initial denaturation for 5 min, 35 cycles of 30 s at 94°C, 30 s at 55°C, 60 s at 72°C,

and a final extension of 7 min at 72°C. For field samples that did not amplify against any

of our custom-made targets, the DNA was amplified using a PCR against generic

vertebrate primers (0.2 µM) for the *cytochrome b* gene region[47]. The resulting products

were again purified and reamplified under the same conditions. Five microliters of each

reamplified product were separated on a 2% agarose gel. The remaining product was

purified using GeneJet PCR purification kit (Thermo Scientific) and sequenced (Eurofins

Genomics). Sequences were aligned with reference genomes from the NCBI database

to determine blood meal identity.

Leishmania detection and quantification by qPCR

A standard curve was created for each different sample condition by spiking and

uninfected sand fly midgut, a whole body, or a pool of whole bodies, with 10^5 *L.*

donovani (MHOM/SD/62/1S) parasites followed by DNA extraction, followed by a 10-

fold serial dilution.

For unfed and gravid samples (processed individually or as a pool of 5-10 sand flies),

we used qPCR to amplify the kinetoplast minicircle from *Leishmania* parasites using

JW11 and JW12 primers[42] in a 25 µL reaction volume containing SYBR green

(PerfeCTA SYBR Green Fast Mix or FIREPol Master Mix Ready to Load). PCR

conditions were 95°C initial denaturation for 3 min with 49 cycles of 10 s at 95°C and 30

s at 55°C. A probe-based qPCR was used for individually blood fed sand flies as

described[42]. All field samples were run in duplicate (unfed/gravid) or triplicate (blood

fed) using 40 ng of DNA template. A standard curve (of serially diluted parasite-spiked

midguts), and negative (uninfected laboratory-reared sand flies) and positive

(*Leishmania donovani*-infected laboratory-reared sand flies) controls were included in

each plate. Melting curve analysis of field-samples were compared to the melting curve

from positive and negative controls. Samples considered positive for *Leishmania* kDNA

by qPCR were re-run to confirm positive amplification.

Human dried blood spot collection

All reported VL cases to the Ministry of Health were diagnosed and treated at reference

hospitals according to national guidelines[11, 48]. Males and females were included in our

study. A total of 752 participants were enrolled from Laisamis and Karare, Marsabit

county, Kenya, and 440 participants were enrolled from Kunaynah village, Gedaref

state, Sudan. Blood was collected by finger prick using a lancet, spotted on a Whatman

903 protein saver card (Cytiva), and preserved at room temperature.

Leishmania rK39 ELISA from dried blood spots

ELISA plates (Thermo Fisher Scientific) were coated overnight at 4°C with 25 ng/well of rK39 antigen (donated by Steve Reed from the Infectious Disease Research Institute, now the Access to Advanced Health Institute). Plates were blocked with 200 μl diluted in Tis-Bris Saline 0.05% Tween 20 (TBST) with 20% donor equine serum (DES; Cytiva), and incubated for 1 h at room temperature. In parallel, dried blood spots (DBSs) were eluted with 300 μl of PBS 0.05% Tween 20 (Sigma), and incubated for 4 h shaking at room temperature or overnight at 4°C. After blocking, 25 μl of the eluted DBS sample was diluted in TBST/5% DES (1:4 dilution) to a total volume of 100 μl/well in duplicates. Plates were incubated for 1 h at room temperature, followed by 6 washes with TBST. One hundred microliters of secondary antibody, [Goat anti-human IgG (H+L) Alkaline phosphatase conjugated; Sigma], diluted in TBST/5% DES was added to each well. After 1 h of incubation, the plates were washed 12 times with TBST. Lastly, 100 μl of substrate-p-nitrophenyl phosphate liquid (Sigma) was added to each well and data was collected 1 h after addition of the substrate at a 405 nm wavelength. Non-endemic negative controls obtained from individuals living in Nairobi, Kenya were included in each plate.

Statistical analysis

Chi-square goodness of fit test was performed for entomology analysis, followed by Post-hoc z-score analysis with Bonferroni correction to compare ecotypes within each country. A z-score ≥ 2.69 was considered significant. ELISA data was normalized using

the plate cutoff calculated by the mean optical density (O.D.) of non-endemic negative

controls + 2 SD, followed by log transformation. After plate normalization, samples with

an O.D. ≤ 0.04 were considered negative and statistical comparisons were calculated

by the Mann-Whitney test. A *P* value of ≤ 0.05 was considered significant.

Results

Selection of study sites in Kenya, Sudan, and Ethiopia

We selected endemic study sites in Kenya, Ethiopia and Sudan (Figure 1A) with stable

VL transmission (Figure 1B) and distinct ecological features common to several VL foci

in East Africa. Marsabit county in Kenya experienced sporadic outbreaks[36, 37], reporting

an incidence of 1 case per 10,0000 inhabitants over the past seven years excluding a

notable increase in incidence during 2019. The Konso zone in Ethiopia, reported 2.66

cases per 10,000 inhabitants per year since 2018. In Sudan, Gedaref state is a highly

endemic VL region that borders Ethiopia to the east and reports a steady 10 cases per

10,000 inhabitants per year since 2017 (Figure 1B).

Vector species have distinct microhabitat and breeding preferences

Entomological surveys of phlebotomine sand flies were carried out periodically from

September 2023 to September 2024 (Supplementary Table 1). Trapping sites were

selected from representative peridomestic and sylvatic ecotypes (Supplementary Figure

1). Our collections determined that *Ph. orientalis* was the predominant VL vector

species in Marsabit, Kenya, and Gedaref, Sudan, comprising 99.75% (n = 1185) and

100% (n = 1350) of *Phlebotomus* females, respectively (Figure 2A). In contrast, *Ph.*

martini/Ph. celiae females (morphologically indistinguishable) were the most prevalent

of five *Phlebotomus* species in Konso, Ethiopia, accounting for 99% of the collection (n

= 184; Figure 2A). Other female *Phlebotomus* species captured included one *Ph.*

martini/Ph. celiae and two *Ph. alexandri* in Kenya; and six *Ph. orientalis*, four *Ph.*

saevus, and one *Ph. alexandri* in Ethiopia.

Comparing ecotype productivity of *Ph. orientalis* and *Ph. martini/Ph. celiae* revealed a

significant difference in habitat distribution across all study sites in the three countries

(c^2 = 1352.89, d.f. = 12, *P* <0.0001). The majority of *Ph. orientalis* females in Marsabit,

Kenya, were collected from outdoor habitats associated with peridomestic and sylvatic

vegetation (45.01%, *P* <0.0001), mostly *Acacia* and *Balanites* trees, followed by animal

enclosures (25.92%, *P* <0.0001; Figure 2B). In Gedaref, Sudan, *Ph. orientalis* was

primarily captured in vegetation (31.78%, *P* <0.0001) and inside houses (27.19%, *P*

<0.0001) whose floors commonly consist of black cotton soil, a risk factor for VL[19, 24].

The remaining specimens were collected from outside houses (12.15%) and in animal

enclosures (15.93%; Figure 2B). Worth noting, the majority of *Ph. orientalis* females

(39.34%) were collected near human dwellings. In Konso, Ethiopia, *Ph. martini/Ph.*

celiae females were mainly captured from termite hills (74.46%, *P* <0.0001) that were

prevalent in both peridomestic and sylvatic ecotypes, followed by temporary shelters

locally known as "foras" (8.15%; Figure 2B).

We used *Phlebotomus* males to verify species prevalence and to indicate breeding

sites. *Ph. orientalis* represented 100% of all collected males in Kenya (n = 260) and

Sudan (n = 4714) (Supplementary Figure 2A). Like females, *Ph. orientalis* males in

Marsabit, Kenya, were collected from vegetation (46.92%, *P* <0.0001), outside houses

near *Acacia* and *Balanites* trees (18.85%), and in animal enclosures (19.23%) (Supplementary Figure 2B). Preliminary evidence of breeding sites, identified by finding unrotated or partially rotated male genitalia (n = 12), implicated outdoor ecotypes (Supplementary Figure 2C). In Gedaref, Sudan, *Ph. orientalis* males were mainly collected from vegetation (33.64%, *P* <0.0001) and indoors (22.70%, *P* <0.0001) (Supplementary Figure 2B), similar to females. Of the captured males with unrotated or partially rotated genitalia (n = 33), 51.52% (*P* <0.0001) were collected from vegetation and 21.21% from animal enclosures (Supplementary Figure 2C), potentially indicating that *Ph. orientalis* may not be breeding indoors in either Kenya or Sudan.

In Konso, Ethiopia, *Ph. martini* was the main male species collected (Supplementary Figure 2A), mostly from termite hills (72.35%, *P* <0.0001) (Supplementary Figure 2B), again similar to females. Other male species included *Ph. celiae* (5.80%) and *Ph. orientalis* (1.02%) (Supplementary Figure 2A). Some 72.92% (n = 35, *P* <0.0001) of *Ph. martini* males with unrotated or partially rotated genitalia were collected from termite hills (Supplementary Figure 2C), implicating this microhabitat as a significant breeding site for *Ph. martini*. Of note, we collected one *Ph. celiae* male with unrotated/partially rotated genitalia from a domestic animal enclosure. The relative abundance of *Ph. martini* over *Ph. celiae* males (94.14%) indicates predominance of the former species in our female collections.

Vector species exhibit different feeding preferences

We performed host blood meal analysis on field-collected individual blood fed *Phlebotomus* females using a custom-made multiplex PCR tailored to amplify blood

from humans and nine animals abundant in our study sites (Supplementary Figure 3,

Supplementary Table 2). The multiplex PCR panel identified 63.8% (n = 270/423) of

blood meals. After *cytochrome b* (*cytb*) gene sequencing of unknowns this increased to

96.9% (n = 410/423). Interestingly, the diversity of identified blood meal sources was

considerably higher in Marsabit, Kenya, (13 hosts) compared to either Gedaref, Sudan,

(5 hosts) or Konso, Ethiopia, (4 hosts) (Figure 3A-C). Of interest, the observed number

of sand flies with multiple blood meals were higher than expected for Marsabit, Kenya,

(c^2 = 44.09, d.f. = 15, *P* <0.0001) and Gedaref, Sudan (c^2 = 17.83, d.f. = 6, *P* <0.0067).

Our results also establish that several females fed on humans and domestic or sylvatic

animals such as hyraxes and mongooses (Figure 3A-C).

In Marsabit, Kenya, we collected 226 *Ph. orientalis* and two *Ph. alexandri* blood fed

females from animal enclosures (32.30%, *P* <0.0001), vegetation (31.86%, *P* <0.0001),

and temporary shelters (23.89%, *P* = 0.001), following a similar pattern to unfed/gravid

sand flies (Figure 3D). Humans were identified as the dominant host for *Ph. orientalis*,

representing 37.76% (*P* <0.0001) of single blood meals, followed by rodents (15.89%, *P*

<0.0001), dogs (9.27%) and other sylvatic animals like gazelles (4.64%) (Figure 3A).

Mixed blood meals from two or three hosts were identified in 63 *Ph. orientalis*

specimens including human/ovis (22.22%), and human/cow (17.46%) (Figure 3A). In

Gedaref, Sudan, 29.51% (*P* <0.0001) of the 183 collected blood fed *Ph. orientalis*

specimens were captured inside the house, and 33.33% (*P* <0.0001) in vegetation

(Figure 3D). Similar to Marsabit, Kenya, humans (74.83%, *P* <0.0001) were the

predominant host, while 14.97% fed on donkey (Figure 3B). Only seven specimens

contained mixed blood meals and fed on human plus other domestic and sylvatic

animals such as mongooses and rabbits (Figure 3B). In Ethiopia, only 12 *Ph.*

martini/Ph. celiae blood fed specimens were collected from termite hills (50%, *P* <0.001)

and animal enclosures (25%) (Figure 3D). Contrary to the anthropophilic feeding

behavior observed for *Ph. orientalis*, *Ph. martini/Ph. celiae* seemed to preferentially feed

on cows (37.5%), and ovis (25.0%), and five sand flies fed on multiple hosts such as

human/cow and hyrax/cow (Figure 3C). Of note, the two *Ph. alexandri* collected from

Kenya fed on humans, while in Ethiopia, one *Ph. alexandri* fed on an Ovis.

Leishmania-infected vector species were recovered from diverse microhabitats

Using qPCR (Supplemental Figure 4) on individual (n_i) or pooled (n_p) sand flies, we

detected infected specimens in both peridomestic and sylvatic niches across all study

sites (Figure 4A), and from distinct ecotypes (Figure 4B-C). The minimal infection rates

of *Ph. orientalis* with *Leishmania* were similar in Marsabit, Kenya, (26/648, 3.9%) and

Gedaref, Sudan, (10/281, 3.6%). Ethiopia had a lower rate of infection of 1.5% (3/183)

for *Ph. martini/Ph. celiae* females (Figure 4A). Of note, none of the other sand fly

species collected throughout this study were infected.

In Marsabit, Kenya, 4.3% (18/422) unfed/gravid and 3.5% (8/226) blood fed *Ph.*

orientalis were *Leishmania* positive (Figure 4B-C). Infected *Ph. orientalis* were

associated with vegetation (30.77%), where we captured ten infected unfed/gravid and

four infected blood fed females in sylvatic sites, and seven unfed/gravid and three blood

fed *Ph. orientalis* from peridomestic sites (Supplementary Tables 1 and 3). Interestingly,

a cluster of six infected sand flies was collected outside and inside the house of a

relapsed VL patient in Laisamis, Kenya (Supplementary Tables 1 and 3), alluding to

how hotbeds of transmission form. The patient, negative for the human

immunodeficiency virus, received treatment in 2021 and refused treatment after

relapsing in January 2024 (Supplementary Table 3), highlighting the need for better

surveillance and treatment. Though four of the eight (50%) infected blood fed *Ph.*

orientalis fed on humans, confirming the vector's anthropophilic behavior, finding three

and one infected sand flies that fed on gazelles and a rodent, respectively (Figure 4C,

Supplementary Tables 3 and 4), stresses the need to investigate whether there is a role

for animal reservoirs in VL transmission in East Africa.

For Gedaref, Sudan, one of the 98 (1.0%) screened unfed/gravid specimens collected

from inside a house was positive for *Leishmania* (Figure 4B). Importantly, we found nine

(9/183, 4.9%) blood fed *Leishmania*-infected *Ph. orientalis* females, seven from

peridomestic sites, and two from vegetation and a termite hill in sylvatic sites within

Dinder Park (Figure 4C, Supplementary Table 3). All nine infected sand flies fed on

humans, further emphasizing the anthropophilic behavior of *Ph. orientalis* in this region

(Figure 4C, Supplementary Tables 3 and 4). The focal nature of transmission was again

revealed by finding three infected blood fed sand flies, two inside the house of a VL

patient, collected in consecutive days, and a third that was captured from inside the

house of the patient's neighbor (Supplementary Tables 1 and 3).

In Ethiopia, three (1.8%, n=3/171) unfed/gravid *Ph. martini/Ph. celiae* were positive for

Leishmania, all captured in termite hills (Figure 4B). Two of the three were captured 9

months apart from the same castellated termite hill located in Galga near the house of a

recent VL case (Supplementary Tables 1 and 3), again highlighting the focality of VL.

The prevalence of *Ph. martini/Ph. celiae* females in termite hills and the exclusive

collection of *Leishmania*-infected specimens from this microhabitat implicates it in VL transmission in Ethiopia.

Leishmania rK39 seroprevalence reveals an increased risk of infection for individuals living near VL cases

As *Ph. orientalis* is the vector of *L. donovani* in both Marsabit, Kenya, and Gedaref, Sudan, we wanted to assess seroprevalence of *Leishmania* rK39 antibodies in these study foci. We conducted a case-control cross-sectional survey in household members of VL cases and five neighboring households for comparison against controls (households located >4000 meters from confirmed cases). DBSs were collected from household members of 48 VL cases diagnosed during December 2023 to May 2025 in Laisamis and Karare villages, Marsabit county, Kenya, and from five VL cases diagnosed during August 2023 to August 2024 in Kunaynah village, Gedaref state, Sudan (Supplementary Table 5). In Kenya, we enrolled 752 subjects (male to female ratio of 1:1.54) with a median age of 14 years (range, 2 – 85 years). In Sudan, we enrolled 440 participants (male to female ratio of 1:1.3) with a median age of 15 years (range, 5 months – 85 years). Thirty-two and ten control households were screened from control HHs in Kenya and Sudan, respectively.

The rK39 titers of individuals living near cases were significantly higher than those of control households in both Kenya and Sudan (Figure 5). In Kenya, 21.68% (129/595) of screened asymptomatic participants living with or near VL cases exhibited significantly higher rK39 antibodies, compared to 16.51% (18/109) of those living farther away (Figure 5A). Similarly, 10.03% (29/289) of asymptomatic people living with or near VL

cases in Sudan were seropositive, versus only 2.05% (3/146) of those living away from

cases (Figure 5B). Of note, at the time of serum collection, only 23/48 (Kenya) and 2/5

(Sudan) of the diagnosed and treated VL patients had detectable antibodies by rK39

ELISA (Supplementary Table 5), considered the gold standard VL serodiagnostic test,

even though DBSs were collected within 1-4 months of treatment.

Discussion

The WHO has recently targeted visceral leishmaniasis in East Africa for elimination as a

public health concern by 2030[11]. In the present study, we provide a comprehensive

account of sand fly species composition, habitat preference, and *Leishmania*-infection

rates within distinct VL endemic foci in three East African countries. We also compare

the prevalence of asymptomatic cases near and away from recently treated VL cases in

endemic foci. Our findings provide evidence of widespread and intense VL transmission

in diverse settings across our study sites.

Consistent with earlier reports from the region[6, 8, 49], *Ph. orientalis* was the most

prevalent *Phlebotomus* species in Sudan and northern Kenya*,* while *Ph. martini*

predominated in southern Ethiopia. *Ph. orientalis* was predominant in *Acacia* and

Balanites forests in association with vertisol and animal burrows, microhabitats reported

to be suitable breeding sites for this vector[8, 17, 20, 50]. In comparison, termite hills were

the most productive site for *Ph. martini*, reported previously[7, 51-53], though the collection

of over 72% of newly emerged males from this microhabitat provides evidence of its

significance as a breeding site.

Several studies have reported infection rates in field-collected specimens from Kenya

(0.3%)[54], Sudan (1.6% and 1.4%)[19, 55], and southern (5.8%)[7] and northern Ethiopia

(23%)[23]. In our study, the infection rates were higher than previously reported for

Kenya (3.9%) and Sudan (3.6%) and lower for southern Ethiopia (1.5%). More studies

targeted towards establishing the longevity, population structure, and feeding behavior

of vectors from field-captured specimens, particularly for *Leishmania*-infected females,

are needed to fully understand vectorial competence and VL transmission in East

Africa.

We recovered *Leishmania*-infected *Ph. orientalis* from different microhabitats within our

study sites in Marsabit, Kenya, and Gedaref, Sudan, in both sylvatic and peridomestic

niches, and frequently from the same location months apart. This indicates that *Ph.*

orientalis has adapted to diverse ecological microhabitats that drive focal transmission

of VL by providing sugar and blood meal sources, and breeding habitats[17, 18, 20]. In

addition, *Ph. orientalis* likely shifts its behavior to adapt to climatic conditions.

Collections from Helat-Belo, Gedaref state, Sudan, conducted during the dry season in

March to June 2016-2018 found infected blood fed *Ph. orientalis* mainly outside houses

and in sylvatic ecotypes, and reported that they fed on human, cattle, and donkeys[19].

In contrast, our sand fly collections from Kunaynah and Umslala-Houg were performed

during the rainy season (June to September), when people tend to sleep inside their

houses. Consequently, *Leishmania*-infected *Ph. orientalis* females were collected

mainly indoors and had fed soley on humans, suggesting a potential shift in vector

behavior, with sand flies likely resting inside the house after feeding on humans. While

the majority of juvenile *Ph. orientalis* males were collected from vegetation and animal

enclosures, we also collected juvenile males indoors in Sudan, possibly due to the widespread nature of black cotton soil in the village, including inside houses, serving as a breeding site for this vector[20], and creating an ideal environment for focal and active VL transmission indoors. Further confirmation of this shift in vector behavior and transmission pattern is needed to inform control measures that are currently focused on outdoor applications.

In Konso, Ethiopia, *Ph. martini/Ph. celiae* were predominantly collected from termite hills, both in sylvatic and peridomestic settings, pointing to a restricted habitat preference compared to *Ph. orientalis*. Additionally, all three *Leishmania*-infected sand flies were recovered from termite hills reinforcing findings of a previous study[7]. Capturing two *Leishmania*-infected females nine months apart from a peridomestic castellated termite hill reinforces the focality of infection sources and highlights the importance of this microhabitat in sustaining transmission by *Ph. martini/Ph. celiae*. Notably, this termite hill was located only 20 meters away from a household that reported a VL case.

Finding *Leishmania*-infected sand flies near VL cases from all our study sites affirms the importance of human reservoirs in maintaining VL transmission, and the need for continued surveillance and rapid treatment of cases. This is further reinforced by rK39 serological surveys where participants living near VL cases exhibited significantly higher antibody levels compared to those living farther away from cases in both Marsabit, Kenya, and Gedaref, Sudan, highlighting the risk associated with proximity to VL cases. Overall, the high prevalence of rK39 antibodies in asymptomatic participants provides

evidence of prevalent and efficient transmission, and stresses the urgency of

determining their involvement as potential parasite reservoirs[56].

Our study and others[7, 19, 21, 23, 54, 55] have reported infected sand flies from both sylvatic

and domestic/peri-domestic habitats. Host attractiveness to vector sand flies,

seroprevalence of *Leishmania* antibodies, and PCR detection of *L. donovani* DNA have

been reported for various domestic and wild animal species in East Africa[27, 28, 57-59], but

live parasites have not been isolated from any thus far. Irrespective of whether domestic

or wild animals are reservoirs, VL endemicity in relatively scarcely populated areas

points to the existence of a zoonotic transmission cycle in East Africa[19, 24, 60].

Establishing whether animal reservoirs are involved in sustaining VL transmission in

East Africa is a major knowledge gap that needs to be urgently addressed to inform on

the best strategy for VL elimination in this region.

A limitation of our study is the use rK39 to assess seroprevalence. The sensitivity of

rK39 is East Africa is suboptimal[14-16]. Diagnostic tools with better specificity and

sensitivity to detect *Leishmania* infections in East Africa are needed. For studies of

vector sand flies, our aim was to identify their preferred habitats through a broad

investigation of distinct foci within three East African countries. Moving forward, in-depth

longitudinal studies focused on identified hotbeds of infection will be needed to better

understand transmission dynamics throughout the season. Another study limitation is

the pooling of non-fed specimens which provided a minimal infection rate. A high

throughput in-depth analysis of infection status in individual specimens, conducted in

defined microhabitats, will yield original data vital to elucidation of vector competence in

VL foci in East Africa.

Our exploratory ecoepidemiological study conducted in three East African countries

reinforces widespread and intense VL transmission in the region. Our results also

pinpoint habitat preference of vector species and transmission hotspots in distinct

microhabitats within endemic foci that emphasize both the focality and complexity of VL

transmission in East Africa. Concurrent longitudinal clinical and entomological studies of

several disease foci throughout East Africa are needed to galvanize an effective

elimination campaign.

Acknowledgements

We thank the community, administrative officers, and health care workers from all our

study sites. We are also grateful to Marsabit County, Kenya, for supporting us with

fieldwork implementation and logistics. We thank Professor EL-Hassan Center for

Tropical Diseases staff, community leaders and the population in the Kunaynah village,

Gedaref state, Sudan. We thank the Karat Zone Health Bureau, Ethiopia, for supporting

this work and the Arbaminch University for hosting the project in Southern Ethiopia. We

also acknowledge Rogers Asamba (KEMRI), and David Mbuvi (ICIPE) for their valuable

support during fieldwork; Dr. Geyeto Garra (Ethiopia) for support given to this project;

Mr. Haile Gebremariam (Ethiopia) for technical assistance.

Funding

Funding support for this study was obtained from the NIH under the Tropical Medicine

Research Centers (TMRC) grant (U01AI168619). This research was supported in part

by the Intramural Research Program of the National Institutes of Health (NIH).

The contributions of the NIH authors were made as part of their official duties as NIH

federal employees, and are in compliance with agency policy requirements, and

are considered Works of the United States Government. However, the findings and

conclusions presented in this paper are those of the authors and do not necessarily

reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author contributions

E.I. performed the experiments, analyzed the data, conceived the study, supervised the

work, and wrote the manuscript. M.T., Steve Kiplagat, A.G., E.A., O.B., J.I., J.M.M.,

M.A., O.D., A.C., E.K., P.H., S.D., P.C., G.B., C.M., T.D.S., M.P., M.O., E.A.G.K.,

O.F.O., and B.M.Y. and D.M.M. undertook the field work, performed the experiments,

analyzed the data, and edited the manuscript. G.Z. performed the sand fly statistical

analysis. J.G.V., D.K.M., A.M.M., A.H., A.S., E.I. and Shaden Kamhawi and D.M.M

wrote the manuscript. E.I., J.G.V., D.K.M., A.M.M., A.H., A.S., Shaden Kamhawi., and

D.M.M. conceived the study and supervised the work.

Competing interest

The authors declared no competing interest.

Data availability

All data produced in the present work are contained in the manuscript

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used GPT-4o (OpenAI, NIAID GenAI Toolkit) in order to improve readability and language of the manuscript. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

1. World Health Organization. WHO: Leishmaniasis Fact Sheet. 2024 [cited 2024 May, 17, 2024]; Available from: https://www.who.int/news-room/fact-sheets/detail/leishmaniasis

2. Baleela R, Llewellyn MS, Fitzpatrick S, Kuhls K, Schonian G, Miles MA, et al. Leishmania donovani populations in Eastern Sudan: temporal structuring and a link between human and canine transmission. Parasit Vectors. 2014 Nov 20;7:496.

3. Zackay A, Cotton JA, Sanders M, Hailu A, Nasereddin A, Warburg A, et al. Genome wide comparison of Ethiopian Leishmania donovani strains reveals differences potentially related to parasite survival. PLoS Genet. 2018 Jan;14(1):e1007133.

4. Zackay A, Nasereddin A, Takele Y, Tadesse D, Hailu W, Hurissa Z, et al. Polymorphism in the HASPB repeat region of East African Leishmania donovani strains. PLoS Negl Trop Dis. 2013;7(1):e2031.

5. Alvar J, den Boer M, Dagne DA. Towards the elimination of visceral leishmaniasis as a public health problem in east Africa: reflections on an enhanced control strategy and a call for action. Lancet Glob Health. 2021 Dec;9(12):e1763-e9.

6.   Elnaiem DE. Ecology and control of the sand fly vectors of Leishmania donovani

in East Africa, with special emphasis on Phlebotomus orientalis. J Vector Ecol. 2011

Mar;36 Suppl 1:S23-31.

7.   Gebre-Michael T, Lane RP. The roles of Phlebotomus martini and P.celiae

(Diptera: Phlebotominae) as vectors of visceral leishmaniasis in the Aba Roba focus,

southern Ethiopia. Med Vet Entomol. 1996 Jan;10(1):53-62.

8.   Gebre-Michael T, Malone JB, Balkew M, Ali A, Berhe N, Hailu A, et al. Mapping

the potential distribution of Phlebotomus martini and P. orientalis (Diptera:

Psychodidae), vectors of kala-azar in East Africa by use of geographic information

systems. Acta Trop. 2004 Mar;90(1):73-86.

9.   José Antonio Ruiz-Postigo aSJ, a Serge Madjou,a Ana Nilce Maia-Elkhoury,b

Samantha Valadas,b Supriya Warusavithana,c Mona Osman,c Aya Yajima,d Zaw Lin,d

Abate Beshahe and Sunghye Kimf,. Global leishmaniasis surveillance: 2021, assessing

the impact of the COVID-19 pandemic. 2021.

10.   World Health Organization. Ending the neglect to attain the Sustainable

Development Goals: A road map for neglected tropical diseases 2021–2030; 2021.

11.   World Health Organization. Strategic framework for the elimination of visceral

leishmaniasis as a public health problem in eastern Africa 2023–2030. 2024;Licence:

CC BY-NC-SA 3.0 IGO.

12.   Alvar J, Beca-Martinez MT, Argaw D, Jain S, Aagaard-Hansen J. Social

determinants of visceral leishmaniasis elimination in Eastern Africa. BMJ Glob Health.

2023 Jun;8(6).

13. Wasunna M, Musa A, Hailu A, Khalil EA, Olobo J, Juma R, et al. The

Leishmaniasis East Africa Platform (LEAP): strengthening clinical trial capacity in

resource-limited countries to deliver new treatments for visceral leishmaniasis. Trans R

Soc Trop Med Hyg. 2016 Jun;110(6):321-3.

14. Hagos DG, Schallig H, Kiros YK, Abdulkadir M, Wolday D. Performance of rapid

rk39 tests for the diagnosis of visceral leishmaniasis in Ethiopia: a systematic review

and meta-analysis. BMC Infect Dis. 2021 Nov 17;21(1):1166.

15. Iniguez E, Kamhawi S. Discovery of new antigens for serodiagnosis of visceral

leishmaniasis in humans and dogs adds valuable tools to our arsenal. mBio. 2024 Aug

14;15(8):e0107924.

16. Kassa M, Abdellati S, Cnops L, Bremer Hinckel BC, Yeshanew A, Hailemichael

W, et al. Diagnostic accuracy of direct agglutination test, rK39 ELISA and six rapid

diagnostic tests among visceral leishmaniasis patients with and without HIV coinfection

in Ethiopia. PLoS Negl Trop Dis. 2020 Dec;14(12):e0008963.

17. Gebresilassie A, Kirstein OD, Yared S, Aklilu E, Moncaz A, Tekie H, et al.

Species composition of phlebotomine sand flies and bionomics of Phlebotomus

orientalis (Diptera: Psychodidae) in an endemic focus of visceral leishmaniasis in

Tahtay Adiyabo district, Northern Ethiopia. Parasit Vectors. 2015 Apr 25;8:248.

18. Hassaballa IB, Torto B, Sole CL, Tchouassi DP. Exploring the influence of

different habitats and their volatile chemistry in modulating sand fly population structure

in a leishmaniasis endemic foci, Kenya. PLoS Negl Trop Dis. 2021 Feb;15(2):e0009062.

19. Khogali A, Elnaiem DA, Diaz-Reganon R, Jibreel T, Nour BYM, Abdelrahman

SH, et al. Infection of Leishmania donovani in Phlebotomus orientalis Sand Flies at

Different Microhabitats of a Kala-Azar Endemic Village in Eastern Sudan. Trop Med

Infect Dis. 2024 Feb 2;9(2).

20. Moncaz A, Kirstein O, Gebresellassie A, Lemma W, Yared S, Gebre-Michael T,

et al. Characterization of breeding sites of Phlebotomus orientalis - the vector of visceral

leishmaniasis in northwestern Ethiopia. Acta Trop. 2014 Nov;139:5-14.

21. Elnaiem DE, Osman OF. Evidence for active transmission of visceral

leishmaniasis within a village in eastern Sudan. Acta Trop. 1998 Nov 30;71(3):305-9.

22. Hailu A, Balkew M, Berhe N, Meredith SE, Gemetchu T. Is Phlebotomus

(Larroussius) orientalis a vector of visceral leishmaniasis in south-west Ethiopia? Acta

Trop. 1995 Sep;60(1):15-20.

23. Kirstein OD, Skrip L, Abassi I, Iungman T, Horwitz BZ, Gebresilassie A, et al. A

fine scale eco-epidemiological study on endemic visceral leishmaniasis in north

ethiopian villages. Acta Trop. 2018 Jul;183:64-77.

24. Zijlstra EE, el-Hassan AM. Leishmaniasis in Sudan. Visceral leishmaniasis.

Trans R Soc Trop Med Hyg. 2001 Apr;95 Suppl 1:S27-58.

25. Votypka J, Pavlasova M, Volfova V, Volf P. Rotation of male genitalia in various

species of phlebotomine sandfly. Med Vet Entomol. 2015 Dec;29(4):355-60.

26. Elaagip A, Ahmed A, Wilson MD, Boakye DA, Abdel Hamid MM. Studies of host

preferences of wild-caught Phlebotomus orientalis and Ph. papatasi vectors of

leishmaniasis in Sudan. PLoS One. 2020;15(7):e0236253.

27. Elnaiem DA, Hassan MM, Maingon R, Nureldin GH, Mekawi AM, Miles M, et al.

The Egyptian mongoose, Herpestes ichneumon, is a possible reservoir host of visceral

leishmaniasis in eastern Sudan. Parasitology. 2001 May;122(Pt 5):531-6.

28. Hassan MM, Osman OF, El-Raba'a FM, Schallig HD, Elnaiem DE. Role of the

domestic dog as a reservoir host of Leishmania donovani in eastern Sudan. Parasit

Vectors. 2009 Jun 17;2(1):26.

29. Mondal D, Bern C, Ghosh D, Rashid M, Molina R, Chowdhury R, et al.

Quantifying the Infectiousness of Post-Kala-Azar Dermal Leishmaniasis Toward Sand

Flies. Clin Infect Dis. 2019 Jul 2;69(2):251-8.

30. Singh OP, Chaubey R, Kushwaha AK, Fay MP, Sacks D, Sundar S. Visceral

Leishmaniasis-Human Immunodeficiency Virus-Coinfected Patients Are Highly

Infectious to Sandflies in an Endemic Area in India. J Infect Dis. 2024 May 7.

31. Singh OP, Tiwary P, Kushwaha AK, Singh SK, Singh DK, Lawyer P, et al.

Xenodiagnosis to evaluate the infectiousness of humans to sandflies in an area

endemic for visceral leishmaniasis in Bihar, India: a transmission-dynamics study.

Lancet Microbe. 2021 Jan;2(1):e23-e31.

32. Al-Salem W, Herricks JR, Hotez PJ. A review of visceral leishmaniasis during the

conflict in South Sudan and the consequences for East African countries. Parasit

Vectors. 2016 Aug 22;9(1):460.

33. Berry I, Berrang-Ford L. Leishmaniasis, conflict, and political terror: A spatio-

temporal analysis. Soc Sci Med. 2016 Oct;167:140-9.

34. Boodman C, van Griensven J, Gupta N, Diro E, Ritmeijer K. Anticipating visceral

leishmaniasis epidemics due to the conflict in Northern Ethiopia. PLoS Negl Trop Dis.

2023 Mar;17(3):e0011188.

35. Kenya National Bureau of Statistics. Kenya Population and Housing Census.

2019;II.

36.   de Souza DK, Picado, A., Bessell, P. R., Liban, A., Wachira, D., Mwiti, D.,

Matendechero, S. H., Bosch, M., Ndung'u, J. M., & Cruz, I.,. Strengthening Visceral

Leishmaniasis Diagnosis Capacity to Improve Access to Care in Kenya: The Example of

Marsabit County. Frontiers in Tropical Diseases. 2021.

37.   Mewara A, Gudisa R, Padhi BK, Kumar P, Sah R, Rodriguez-Morales AJ.

Visceral leishmaniasis outbreak in Kenya-a setback to the elimination efforts. New

Microbes New Infect. 2022 Nov-Dec;49-50:101060.

38.   Department KM. Kenya Meterological deperment, . 2024 [cited; Available from:

https://meteo.go.ke

39.   Ibrahim B, Elmadani M, Sahar H, Gadallh M, Abdallh A, Ahmed AEE. Prevalence

of Visceral Leishmaniasis among Wildlife Rangers in Dinder National Park, Sudan.

medRxiv. 2024:2024.06.24.24309386.

40.   Abonnenc E, Minter D. Bilingual keys for the identification of the sandflies of the

Ethiopian Region (Frenceh and English). Cahier, Office de la Recherche Scintifique et

Technique d'Outre, Mer Entomologie Medicale. 1965;5(1).

41.   Lewis D.J. The phlebotomine sandflies (Diptera: Psychodidae) of the Oriental

Region. Bull Br Museum Nat Hist 1978;37:217–343. .

42.   Dey R, Joshi AB, Oliveira F, Pereira L, Guimaraes-Costa AB, Serafim TD, et al.

Gut Microbes Egested during Bites of Infected Sand Flies Augment Severity of

Leishmaniasis via Inflammasome-Derived IL-1beta. Cell Host Microbe. 2018 Jan

10;23(1):134-43 e6.

43.   Kent RJ, Norris DE. Identification of mammalian blood meals in mosquitoes by a

multiplexed polymerase chain reaction targeting cytochrome B. Am J Trop Med Hyg.

2005 Aug;73(2):336-42.

44.   Irwin DM, Kocher TD, Wilson AC. Evolution of the cytochrome b gene of

mammals. J Mol Evol. 1991 Feb;32(2):128-44.

45.   Parodi B, Aresu O, Bini D, Lorenzini R, Schena F, Visconti P, et al. Species

identification and confirmation of human and animal cell lines: a PCR-based method.

Biotechniques. 2002 Feb;32(2):432-4, 6, 8-40.

46.   Collini M, Albonico F, Hauffe HC, Mortarino M. Identifying the last bloodmeal of

questing sheep tick nymphs (Ixodes ricinus L.) using high resolution melting analysis.

Vet Parasitol. 2015 Jun 15;210(3-4):194-205.

47.   Abbasi I, Cunio R, Warburg A. Identification of blood meals imbibed by

phlebotomine sand flies using cytochrome b PCR and reverse line blotting. Vector

Borne Zoonotic Dis. 2009 Feb;9(1):79-86.

48.   Kenya Ministry of Health. Prevention, Diagnosis and Treatment of Visceral

Leishmaniasis (Kala azar) in Kenya: National Guidelines. 2017 [cited August 27,

2025];   Available   from: https://www.afrikadia.org/wp-

content/uploads/2018/08/VL_Guidelines_Kenya_2017.pdf

49.   Anjili CO, Ngumbi PM, Kaburi JC, Irungu LW. The phlebotomine sandfly fauna

(Diptera: Psychodidae) of Kenya. J Vector Borne Dis. 2011 Dec;48(4):183-9.

50.   Thomson MC, Elnaiem DA, Ashford RW, Connor SJ. Towards a kala azar risk

map for Sudan: mapping the potential distribution of Phlebotomus orientalis using digital

data of environmental variables. Trop Med Int Health. 1999 Feb;4(2):105-13.

51.   Basimike M MM, Kumar R..,. Habitat preference and seasonal variations of

phlebotomine sandflies. Insect Science and Its Application 1992;13(3):307-314. .

52.   Minter D. The distribution of sandflies (Diptera, Psychodidae) in Kenya. Bulletin

of Entomological Research. 1964; 55, 205-217

53.   Wijers DJ, Minter DM. Studies on the vector of kala-azar in Kenya. V. The

outbreak in Meru district. Ann Trop Med Parasitol. 1966 Mar;60(1):11-21.

54.   Perkins PV, Githure JI, Mebrahtu Y, Kiilu G, Anjili C, Ngumbi PS, et al. Isolation

of Leishmania donovani from Phlebotomus martini in Baringo district, Kenya. Trans R

Soc Trop Med Hyg. 1988;82(5):695-700.

55.   Hassan MM, Elraba'a FM, Ward RD, Maingon RD, Elnaiem DA. Detection of high

rates of in-village transmission of Leishmania donovani in eastern Sudan. Acta Trop.

2004 Sep;92(1):77-82.

56.   Singh OP, Hasker E, Sacks D, Boelaert M, Sundar S. Asymptomatic Leishmania

infection: a new challenge for Leishmania control. Clin Infect Dis. 2014

May;58(10):1424-9.

57.   Dereure J, El-Safi SH, Bucheton B, Boni M, Kheir MM, Davoust B, et al. Visceral

leishmaniasis in eastern Sudan: parasite identification in humans and dogs; host-

parasite relationships. Microbes Infect. 2003 Oct;5(12):1103-8.

58.   Gebresilassie A, Yared S, Aklilu E, Kirstein OD, Moncaz A, Tekie H, et al. Host

choice of Phlebotomus orientalis (Diptera: Psychodidae) in animal baited experiments: a

field study in Tahtay Adiyabo district, northern Ethiopia. Parasit Vectors. 2015 Mar

31;8:190.

59.   Kassahun A, Sadlova J, Benda P, Kostalova T, Warburg A, Hailu A, et al. Natural

infection of bats with Leishmania in Ethiopia. Acta Trop. 2015 Oct;150:166-70.

60.   Elnaiem DA, Hassan HK, Ward RD. Phlebotomine sandflies in a focus of visceral

leishmaniasis in a border area of eastern Sudan. Ann Trop Med Parasitol. 1997

Apr;91(3):307-18.

Figure legends

Figure 1. Study sites and incidence of visceral leishmaniasis

(A,B) Map of East Africa showing the location of study sites. 1, Marsabit ; 2, Konso; 3,

Gedaref state. (B) Visceral leishmaniasis (VL) cases per 10,000 population over the

past seven years in Marsabit county, Kenya; Konso, Ethiopia; and Gedaref state,

Sudan. Incidence data are based on Ministry of Health reports from each country. Maps

at QGIS software QGIS shapefile, https://open.africa/dataset/?tags=Shapefiles.

Figure 2. Distribution of different sand fly species across distinct microhabitats in

endemic foci in Kenya, Sudan and Ethiopia study sites

(A) Relative abundance of collected *Phlebotomus* females by species. (B) Relative

abundance of *Phlebotomus* females by ecotype.

Figure 3. A mosaic of blood meal sources observed in leishmaniasis vectors

across endemic foci in Kenya, Sudan and Ethiopia study sites

(A-C) Host blood meal identity of blood fed females in Kenya (A), Sudan (B), and Ethiopia (C) with a single or multiple sources of blood detected. (D) Relative abundance of collected blood fed *Phlebotomus* females by ecotype.

Figure 4. Distribution of *Leishmania*-infected sand flies from distinct and diverse microhabitats reveals hotspots of active transmission near and away human habitation

(A) Number of *Leishmania*-infected sand flies collected from peridomestic or sylvatic habitats. (B) Habitat productivity of infected unfed/gravid female *Ph. orientalis* sand flies in Kenya and Sudan, and *Ph. martini/Ph. celiae* in Ethiopia. (C) Host blood meal identity of infected blood fed *Ph. orientalis and Ph. martini/Ph. celiae* females. Specimens were processed individually or as pools of up to 5-10 sand flies (B) or individually (C).

Figure 5. *Leishmania* rK39 antibody prevalence in our study sites indicate high risk of infection in Marsabit county, Kenya, and Gedaref state, Sudan

(A,B) *Leishmania* rK39 antibody prevalence measured from dry blood spots (DBS) collected from individuals living with or near and away recently reported VL cases in Laisamis and Karare, Marsabit county (A) and Kunaynah, Gedaref state (B). Data was normalized using the plate cutoff calculated by the mean optical density (O.D.) of non-endemic negative controls + 2 SD. Non-endemic negative controls were obtained from individuals living in Nairobi, Kenya. Black dotted line, samples with O.D. ≤ 0.04 were considered negative after normalization. A P value of ≤ 0.05 was considered significant by Mann-Whitney test (A,B); *$P < 0.05$.