Widespread transmission in diverse ecotypes challenges visceral leishmaniasis control in East Africa

Eva Iniguez[1†*], Mercy Tuluso[2†], Steve Kiplagat[3], Araya Gebresilassie[4], Esayas Aklilu[5], Olivia Battistoni[6], Johnstone Ingonga[2], John Mark Makwatta[3], Mohamed Alamin[7], Osman Dakien[7,8], Alphine Chebet[2], Esther Kaunda[2], Patrick Huffcutt[1], Serena Doh[1], Pedro Cecilio[9], Galgallo Bonaya[3], Claudio Meneses[1], Tiago D. Serafim[1], Myrthe Pareyn[10], Mohamed Osman[11], Eltahir A. G. Khalil[7], Omran F. Osman[7], Brima M. Younis[7], Sithar Dorjee[12], Guofa Zhou[13], Jesus G. Valenzuela[1], Dan K. Masiga[3], Ahmed M. Musa[14], Asrat Hailu[15], Abhay Satoskar[6], Shaden Kamhawi[1*], Damaris Matoke-Muhia[2*]

[1]Vector Molecular Biology Section, Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD 20852, USA

[2]Centre for Biotechnology Research and Development, Kenya Medical Research Institute, Nairobi 00200, Kenya

[3]International Centre of Insect Physiology and Ecology (*icipe*), Nairobi 30772-00100, Kenya

[4]Department of Zoological Sciences, College of Natural and Computational Sciences, Addis Ababa University, Addis Ababa 1176, Ethiopia

[5]Vector Biology and Control Research Unit, Aklilu Lemma Institute of Pathobiology, Addis Ababa University, Addis Ababa 1176, Ethiopia

[6]Departments of Pathology and Microbiology, Wexner Medical Center, The Ohio State

University, Columbus, OH 43210, USA

[7]Department of Zoology, Faculty of Science, University of Khartoum, Khartoum 11111,

Sudan

[8]Kala Zara Research Center, Faculty of Medicine and Health Science, University of

Gedaref, Gedaref 32210, Sudan

[9]Vector Biology Section, Laboratory of Malaria and Vector Research, National Institute

of Allergy and Infectious Diseases, National Institutes of Health, Rockville 20852, MD,

USA

[10]Department of Clinical Sciences, Institute of Tropical Medicine Antwerp 2000, Belgium

[11]York Biomedical Research Institute, University of York, York YO10 5DD, UK

[12]Faculty of Postgraduate Medicine, Khesar Gyalpo University of Medical Sciences of

Bhutan, Thimphu 11001, Bhutan

[13]Program in Public Health, University of California at Irvine, Irvine, CA 92697, USA

[14]Department of Clinical Pathology and Immunology, Institute of Endemic Diseases,

University of Khartoum, Khartoum 11111, Sudan

[15]College of Health Sciences, Addis Ababa University, Addis Ababa 1000, Ethiopia

[†]Equal contribution

*Corresponding authors: Eva Iniguez; Vector Molecular Biology Section, Laboratory of

Malaria and Vector Research, National Institute of Allergy and Infectious Diseases,

National Institutes of Health; Rockville, MD 20852, USA.; Phone: 1.301.761.7357;

Email: eva.iniguez@nih.gov. Shaden Kamhawi; Vector Molecular Biology Section,

Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious

Diseases, National Institutes of Health; Rockville, MD 20852, USA.; Phone:

1.301.761.5081; Email: skamhawi@niaid.nih.gov. Damaris Matoke-Muhia; Centre for

Biotechnology Research and Development, Kenya Medical Research Institute, Nairobi

00200, Kenya; Phone: 54.722.205.901; Email: DMatoke@kemri.go.ke.

53 **Abstract**

East Africa is emerging as the global hot spot of visceral leishmaniasis (VL), yet efforts

to eliminate it are hindered by substantial knowledge gaps in its ecoepidemiology. Here,

we report on the high prevalence of *Leishmania* infection in *Phlebotomus orientalis* in

Marsabit county, Kenya (3.9%), and Gedaref state, Sudan (3.6%), where this species

comprised 99.8% (n = 1185) and 100% (n = 1350) of captured *Phlebotomus* females,

respectively. In Aba Roba, Ethiopia, *Ph. martini* accounted for 99% of 184 collected

Phlebotomus females and had a lower infection rate of 1.5%. *Ph. orientalis* and *Ph.*

martini exhibited different habitat and feeding preferences. While *Ph. orientalis* was

abundant in diverse peridomestic and sylvatic microhabitats, *Ph. martini* was

predominantly collected from termite hills. Moreover, *Ph. orientalis* primarily fed on

humans and less on domestic and sylvatic animals. In contrast, *Ph. martini* exhibited

zoophagic behavior, mostly feeding on cows and Ovis. Widespread transmission of

Leishmania in our study sites is supported by high rK39 seroprevalence in both Kenya

(17.9%) and Sudan (6.22%). An observed greater prevalence of antibodies to rK39 in

individuals living near than away from VL cases in both Kenya (19.8% versus 7.41%, *P*

= 0.0015) and Sudan (8.4% versus 2.1%, *P* = 0.0105) demonstrated that proximity to a

VL case carries an increased risk of infection. Our findings highlight the need for a risk-

based targeted site-specific elimination strategy that accounts for the intensity, diversity,

and complexity of VL transmission in today's East Africa.

Keywords: East Africa; visceral leishmaniasis; Sand fly ecology; *Leishmania*-infected

sand flies; rK39 seroprevalence.

Introduction

East Africa is currently the epicenter of global transmission of visceral leishmaniasis

(VL), a fatal neglected disease ranked second after malaria in mortality rates[1, 2]. VL in

East Africa is epidemiologically and clinically diverse, caused by genetically distinct

Leishmania donovani strains[3-5], and transmitted by several sand fly vector species

including *Phlebotomus orientalis, Ph. martini,* and *Ph. celiae* that have divergent

ecologies and bionomics[6-9]. VL cases have recently surged in East Africa, emerging in

new foci that disproportionally affected Kenya, Sudan, and Ethiopia where they account

for more than 65% of cases worldwide, and where ~50% are children under 15 years

old[1, 10].

VL in East Africa had initially been targeted for control rather than elimination due to

significant knowledge gaps concerning disease transmission[11]. In June 2024, the

World Health Organization launched a framework to eliminate VL as a public health

concern from East Africa by 2030[12]. Elimination is proposed through early diagnosis

and treatment, integrated vector control management, and disease surveillance[12]. At

the country level, the target for elimination was defined as <1% case-fatality rate due to

primary VL, 90% decline in new VL cases, absence of death due to VL in children by

2030, and 100% detection and treatment referral of VL-HIV and PKDL patients[12].

However, East Africa faces major challenges to VL elimination. Treatment differs

between countries, and it is often toxic and lengthy, with low compliance and poor

access to care facilities[6, 12-14]. In addition, significant knowledge gaps in

ecoepidemiology, vector biology, and nature of reservoirs need to be addressed for a

successful implementation of an elimination campaign in East Africa,[6, 13] where there

are no current effective integrated vector control programs[12]. Though *Ph.* orientalis, *Ph.*

martini, and *Ph. celiae* sand flies have been found in both peridomestic and sylvatic

niches in strong association with *Acacia* and *Balanites* woodlands, vertisol (black cotton

soil), and termite hills[15-18], linking these microhabitats to VL transmission in the

region[17, 19-22], confirmation of their epidemiological relevance and their potential as

resting and breeding sites for these vectors[18, 23] needs to be demonstrated.

Additionally, we need to determine the prevalence of asymptomatic individuals and their

role as infectious reservoirs for sand flies[24-26]. Further hindering control of VL in East

Africa is the presence of semi-nomadic populations, such as pastoralists and farmers

living in temporary shelters[13], and populations displaced by conflict[27, 28], that are at a

higher risk of contracting VL[27-29], complicating the epidemiological landscape.

In this study, we provide a comprehensive account of sand fly species composition,

habitat preference, and report the prevalence of *Leishmania*-infected sand flies from

diverse microhabitats in endemic study sites in Kenya, Sudan, and Ethiopia. We also

compare the prevalence of asymptomatic cases near and away from recently treated VL

cases establishing widespread transmission through the high rate of rK39

seroprevalence in Marsabit county, Kenya, and Gedaref state, Sudan, where *Ph.*

orientalis is the primary vector. Altogether, this study provides current information on VL

transmission dynamics in East Africa.

Methods

Ethics approval

The study was approved by the Scientific Research Unit, KEMRI, and licensed by The

National Commission for Science, Technology and Innovation (clinical protocol #

KEMRI/SERU/CBRD/249/4634) and by the State Ministry of Health and Social

Development at Gedaref state, Sudan ethical committee (clinical protocol #

UG/EC/3/2023) in Sudan. Both clinical protocols were translated into local languages.

Communal consent was obtained by meeting with community leaders. Written consent

was obtained from adults or guardians of minors before blood collection. Verbal consent

was acquired from homeowners before entomological collections.

Study sites

We selected study sites with stable VL transmission in Kenya, Sudan, and Ethiopia. The

study sites were chosen to encompass features common to VL ecotypes in East Africa.

<u>Kenya</u>

Marsabit county has a population of 459,785 people[30], and has experienced sporadic

outbreaks since 2014[31, 32]. It has an arid and semi-arid tropical savanna climate with an

average temperature ranging between 15°C and 26°C. Average precipitation is about

128.24 mm per annum. The long rainy season spans April to May while the short rains

occur from November to December; January to March are dry and hot, and June to

October are dry cold months [33]. The communal free-range land is dominated by

diverse *Acacia* trees and related species, a wide variety of shrubs, and seasonal grass.

Inhabitants of Marsabit county are mainly nomadic pastoralists belonging to several

tribes, along with traders and farmers, and smaller communities. Camel, goat, sheep,

donkeys, and cattle are major domestic animals kept as a source of income. The

community moves with their domestic animals from semi-permanent (manyatta) houses

to temporary homesteads (foras) for a period of up to six months in search of pasture

and water. We investigated Laisamis, Logo Logo, and Karare, sub-counties of Marsabit

county.

Sudan

Gedaref state has an urban structure with well-established villages. The region is a flat

plain, with small, scattered hills and flowing watercourses, and the climate is tropical.

The rainy season spans June to October with an annual rainfall of 400-1400 mm. The

natural vegetation is a savanna woodland with *Acacia* and *Balanites* trees, and the

principal soil type is black cotton vertisol. Agriculture is the main livelihood, with

sorghum, sesame, and millet as major crops. Goats, sheep, donkeys, and cattle are

major domestic animals kept as a source of income. Most people are subsistence

farmers or engage in small animal husbandry. High demand for manual labor attracts

seasonal workers from within Sudan or bordering Ethiopia during the rainy season.

Dinder National Park falls between the Sahel of Gedaref state and Ethiopian Highlands

and is home to several species of large and small mammals, such as lions, gazelle,

monkeys, bats and reptiles. A high VL prevalence of 27.6% was recently reported

among Dinder National Park wildlife soldiers[34]. We investigated the villages of

Kunaynah, Sabouni, bordering north Ethiopia, and Umslala-Houng, closest to Dinder

National Park, in Gedaref state.

Ethiopia

The Konso zone is a semi-arid rugged terrain with many hills, lowlands, and riverbeds.

The area is generally semi-arid, with an annual rainfall of about 551 mm per year. The

long rains last from February to May, followed by short rains in September and October.

The mean annual temperature is 24°C. The predominant soil type is sandy. Dispersed

homesteads and animal enclosures are scattered in lowlands or valleys, while villages

are on top of large hills or highlands whose slopes lead to the valleys. Segen river flows

through the valley providing water for irrigation and creating flat plains suitable for

farming and animal herding. Even though there are no permanent human settlements,

many families own farms in the valleys and spend a considerable number of nights in

temporary shelters (foras) herding cattle, goats, and sheep. The vegetation comprises a

mixture of shrub and forests with diverse trees, including species of *Acacia* and

Balanites. Castellated active or eroded shrinking termite hills[8] are abundant in the area.

Aba Roba communities mostly depend on subsistence farming on the cultivated

terraced hillsides and flat plains of the Segen River. Sorghum, maize, sesame and

sunflower are major crops. Inhabitants also raise goats, sheep, and cattle, kept in

enclosures at night. We investigated the villages of Galga and Goinada, 1180-1250 m

above sea level, and Maira and Lahalaha, 830-885 m above sea level, in the Konso

zone.

Sand fly collection

In each country, sand fly sampling ecotypes included peridomestic (associated with humans) and sylvatic (away from humans) sampling microhabitats (Supplementary Figure 1). Ecotypes included indoors (inside manyatta/hut) and outdoors (outside manyatta/hut), temporary shelters or foras, animal enclosures, vegetation in association or not with animal burrows and vertisols, and termite hills. The study sites and the total number of CDC light traps used per collection varied in each country, but the same number of traps were placed per ecotype (Supplementary Table 1).

Identification of *Phlebotomus* sand fly species

The head and the last three segments of each *Phlebotomus* sand fly were wet mounted for morphological identification. Spermatheca and the pharynx teeth were examined to distinguish the female species, while external genitalia were examined for identification of males using a combination of keys[35, 36].

Rotation of *Phlebotomus* male genitalia

During the first 16-24 h of the adult life of a male sand fly, the external genitalia undergo a rotation to orient the claspers for mating. Thus, unrotated or partially rotated male genitalia can be used to identify newly emerged juveniles, captured during their first night of activity as adults, as a proxy to identify sand fly breeding sites[15, 18, 20]. Under the microscope, we recorded the number of males with rotated, unrotated or partially rotated genitalia.

Mitigation of sample contamination from the field to the bench

In the field, all female sand flies were dissected using dissecting pins that were bleached (10%) and rinsed twice with water before each sand fly dissection. This step was critical to avoid contamination caused by carry-over DNA from sample to sample. In the laboratory, three separate rooms were used to process and run the field samples by qPCR. Room one was used only for preparation of buffer and primer aliquots and loading of the qPCR master mix into the plate. Room two was used for extraction of nucleic acids of field specimens. Room three was used to load the DNA samples into the qPCR plate. Each plate included a standard curve, and DNA from an uninfected midgut and non-template negative controls. To avoid contamination all the instruments, benches and pipettes were continuously bleached (10%) and rinsed with water.

Extraction of DNA from field-collected *Phlebotomus* female sand flies

Unfed and gravid midgut samples were individually collected in Ethiopia. Individual or pools of 5 midguts were collected in Kenya. For Sudan, pools of 10 sand flies (whole bodies or midguts) were collected. Samples were stored in 50 ml of DNA/RNA Shield buffer (Zymo), and stored at room temperature, or 4°C if available. DNA was extracted from individual specimens of unfed and gravid sand flies using the Quick-DNA/RNA Miniprep Plus kit (Zymo) tissue protocol, with a digestion step of 1 h at 56°C with proteinase K and tissue lysis. DNA was extracted from pools of unfed and gravid sand flies using the Tissue DNA kit (Omega) according to the manufacturer's instructions. For whole body pooled samples, we first macerated each using a plastic pestle, followed by

chemical digestion with proteinase K and tissue lysis buffer at 56°C for 2-6 h or until

fully digested.

Blood-fed sand flies were preserved individually as midguts/whole bodies on Whatman

903 Protein Saver cards (Cytica), or in up to 50 ml of DNA/RNA Shield buffer (Zymo).

Briefly, field-collected blood fed midguts were individually punched from the Whatman

cards using a 3 mm punch biopsy, and Proteinase K (15 ml), Lysis buffer (30 ml), and

DNA/RNA shield (30 ml) were added to each punched sample and digested at 56°C for

2 to 4 h, with brief vortexing preformed every hour. Blood fed samples stored in

DNA/RNA Shield buffer as midguts/whole bodies were first macerated using ceramic

bashing beads (Roche) using the MagNAlyzer (Roche), followed by digestion for 24 h.

DNA was extracted using the Quick-DNA/RNA Miniprep Plus kits (Zymo). All samples

were eluted to a final volume of 100 µL and stored at -20/-80°C until further processing.

Blood meal analysis by multiplex PCR

Standardization of a blood meal multiplex custom PCR panel

Lutzomyia longipalpis sand flies (Jacobina) were reared at the Laboratory of Malaria

and Vector Research, NIAID, NIH insectary. Female sand flies (4-6 day old) were

allowed to feed for 2 h on whole blood from various species through an artificial chicken

skin membrane[37] and collected 24 h post-feeding as positive controls. Blood meals

included domestic and sylvatic hosts such as mouse and chicken (LMVR4 animal

protocol); human (leftover samples collected for another study under IRB human

protocol 000331-I); cow (Charles River Laboratories); dog, goat, sheep, and donkey

(Innovative Research); hyrax and camel (donated by Dan Masiga, *icipe*).

A universal reverse primer previously reported[38] was used together with custom

forward primers designed in this study based on the mitochondrial *cytochrome b* region,

highly conserved among vertebrates[39], to identify several species of interest

(Supplementary Table 2). A human primer set based on the cytochrome c oxidase I

mitochondrial region[40], and the rodent (*Muroidea* superfamily) primer set targeting the

mitochondrial d-loop region reported in previous studies were also used[41]. Briefly, DNA

(40 ng, 10 µL) was mixed in a single tube as a multiplex reaction with all forward and

reverse primers (Supplementary Table 2) and the DreamTaq^{TM} Hot Start Green PCR

Master Mix (Thermo Scientific) to a final volume of 30 µL. PCR conditions were 94°C

initial denaturation for 5 min, 35 cycles of 30 s at 94°C, 30 s at 55°C, and 60 s at 72°C,

and a final extension of 7 min at 72°C. PCR products were separated on a 2% agarose

gel electrophoresis ran at 8 v/cm for 60 minutes. Two microliters of a 50 to 1000 base-

pair ladder (Thermo Scientific) were used. Bands were visualized on a c600 Azure

Biosystems imager.

Bloodmeal analysis of field-collected samples

PCR of DNA from field samples was performed using the primers listed in

Supplementary Table 2 and following the same procedure and conditions described

above. The PCR products were run on a 2% agarose gel. For field samples that did not

amplify against any of our custom-made targets, the DNA was amplified using a PCR

against generic vertebrate primers (0.2 µM) for the *cytochrome b* gene region[42]. The

products were separated on a 2% agarose gel and the leftover product was purified

(Thermo Scientific) using the GeneJet PCR purification kit (Thermo Scientific), and

reamplified under the same conditions. After confirming amplification by agarose gel,

the remaining reamplified product was purified and sequenced (Eurofins Genomics).

Sequences were aligned with reference genomes from the NCBI database to determine

blood meal identity.

Leishmania detection and quantification by qPCR

A standard curve was created for each different sample condition by spiking and

uninfected sand fly midgut, a whole body, or a pool of whole bodies, with 10^5 _L._

donovani (MHOM/SD/62/1S) parasites followed by DNA extraction, followed by a 10-

fold serial dilution into corresponding uninfected samples.

For unfed and gravid samples (processed individually or as a pool of 5-10 sand flies),

we used qPCR to amplify the kinetoplast minicircle from _Leishmania_ parasites using

JW11 and JW12 primers[37] in a 25 µL reaction volume containing SYBR green

(Perfecta SYBR Green Fast Mix or FIREPol Master Mix Ready to Load). PCR

conditions were 95°C initial denaturation for 3 min with 49 cycles of 10 s at 95°C and 30

s at 55°C. A probe-based qPCR was used for individually blood fed sand flies as

described[37]. All field samples were run in duplicate (unfed/gravid) or triplicate (blood

fed) using 40 ng of DNA template. A standard curve of serially diluted parasite-spiked

midguts, negative uninfected laboratory-reared sand flies, and positive _Leishmania_

donovani-infected laboratory-reared sand flies were included as controls in each plate.

The melting curve of field-samples was compared to the melting curve of positive and

negative controls. Samples considered positive for _Leishmania_ kDNA by qPCR were re-

run to confirm positive amplification.

Human dried blood spot collection

All reported VL cases to the Ministry of Health were diagnosed and treated at reference hospitals according to national guidelines[12, 43]. Males and females were included in our study. A total of 752 participants were enrolled from Laisamis and Karare villages, Marsabit county, Kenya, and 435 participants were enrolled from Kunaynah and Sabouni villages, Gedaref state, Sudan. Blood was collected by finger prick using a lancet, spotted on a Whatman 903 protein saver card (Cytiva), and preserved at room temperature. For Kenya, non-neighboring controls lived in a neighborhood located within 5 km from areas of index VL cases within Laisamis but with no reports of VL cases. In Sudan, controls households were located in Sabouni, an adjacent village at a distance of 80 km from Kunaynah, with no reports of VL cases, a similar ecology, and where sand flies are present.

***Leishmania* rK39 ELISA from dried blood spots**

ELISA plates (Thermo Fisher Scientific) were coated overnight at 4°C with 25 ng/well of rK39 antigen (donated by Steve Reed from the Infectious Disease Research Institute, now the Access to Advanced Health Institute). In parallel, dried blood spots (DBSs) were eluted with 300 µl of PBS 0.05% Tween 20 (Sigma) and incubated for 4 h shaking at room temperature or overnight at 4°C. Plates were blocked with 200 µl diluted in Tis-Bris Saline 0.05% Tween 20 (TBST) with 20% donor equine serum (DES; Cytiva), and incubated for 1 h at room temperature. After blocking, 25 µl of the eluted DBS sample was diluted in TBST/5% DES (1:4 dilution) to a total volume of 100 µl/well in duplicates.

Plates were incubated for 1 h at room temperature, followed by 6 washes with TBST.

One hundred microliters of secondary antibody, [Goat anti-human IgG (H+L) Alkaline

phosphatase conjugated; Sigma], diluted in TBST/5% DES was added to each well.

After 1 h of incubation, the plates were washed 12 times with TBST. Lastly, 100 µl of

substrate-p-nitrophenyl phosphate liquid (Sigma) was added to each well and data was

collected 1 h after addition of the substrate at a 405 nm wavelength. Non-endemic

negative controls obtained from individuals living in Nairobi, Kenya were included in

each plate for normalization.

Statistical analysis

A chi-square goodness-of-fit test was performed for the entomology data, followed by

post-hoc pairwise comparisons using standardized residual (Z-score) analysis with

Bonferroni correction to identify differences among ecotypes within each country. A

Bonferroni-adjusted critical value of |Z| 2.69 (corresponding to P value ≤ 0.007) was

considered significant. ELISA data was normalized using the plate cutoff calculated by

the mean optical density (O.D.) of non-endemic negative controls + 5 SD, followed by

log transformation. After normalization, samples were statistically compared using the

Mann-Whitney test. Odds ratio was used to assess the risk for having rK39 antibodies

(categorized as positive or negative) between endemic index case household members

and neighborhood households versus non-neighboring households (Kenya), or non-

neighborhood controls (Sudan), and its significance was tested using Fisher's exact

test. A univariable binary logistic regression was performed to assess the effect of

distance in meters from index case households to neighboring versus non-neighboring

households for Kenya, or distance in meters from a VL case in Kunaynah to households

a non-endemic adjacent village for Sudan, on rK39 seropositivity (risk by log-odds of

positive outcome after normalization). The 'margins' command of Stata was used to

compute adjusted log-odds of seropositivity at selected distances from index case

households. A *P* value of ≤ 0.05 was considered significant.

Results

Study sites in Kenya, Sudan, and Ethiopia

Figure 1 depicts our selected endemic study sites in Kenya, Ethiopia and Sudan (Figure

1A) and their stable VL transmission (Figure 1B). Marsabit county in Kenya experienced

sporadic outbreaks[31, 32], reporting an incidence of 1 case per 10,0000 inhabitants over

the past seven years excluding a notable increase in incidence during 2019. The Konso

zone in Ethiopia, reported 2.66 cases per 10,000 inhabitants per year since 2018. In

Sudan, Gedaref state is a highly endemic VL region that borders Ethiopia to the east

and reports a steady 10 cases per 10,000 inhabitants per year since 2017 (Figure 1B).

Distinct microhabitat and breeding preferences of vector species

Entomological surveys of phlebotomine sand flies were carried out periodically from

September 2023 to September 2024 (Supplementary Table 1). Trapping sites were

selected from representative peridomestic and sylvatic ecotypes (Supplementary Figure

1). Our collections determined that *Ph. orientalis* was the predominant VL vector

species in Marsabit, Kenya, and Gedaref, Sudan, comprising 99.75% (n = 1185) and

100% (n = 1350) of *Phlebotomus* females, respectively (Figure 2A). In contrast, *Ph.*

martini/Ph. celiae females (morphologically indistinguishable) were the most prevalent

of five *Phlebotomus* species in Konso, Ethiopia, accounting for 99% of the collection (n

= 184; Figure 2A). Other female *Phlebotomus* species captured included one *Ph.*

martini/Ph. celiae and two *Ph. alexandri* in Kenya; and six *Ph. orientalis*, four *Ph.*

saevus, and one *Ph. alexandri* in Ethiopia.

Comparing ecotype productivity of *Ph. orientalis* and *Ph. martini/Ph. celiae* revealed a

significant difference in habitat distribution across all study sites in the three countries

(c^2 = 1352.89, d.f. = 12, *P* <0.0001). The majority of *Ph. orientalis* females in Marsabit,

Kenya, were collected from outdoor habitats associated with peridomestic and sylvatic

vegetation (45.01%, *P* <0.0001), mostly *Acacia* trees, followed by animal enclosures

(25.92%, *P* <0.0001; Figure 2B). In Gedaref, Sudan, *Ph. orientalis* was primarily

captured in vegetation (31.78%, *P* <0.0001) and inside houses (27.19%, *P* <0.0001)

whose floors commonly consist of black cotton soil, a risk factor for VL[17, 22]. The

remaining specimens were collected from outside houses (12.15%) and in animal

enclosures (15.93%; Figure 2B). Worth noting that in Sudan, the majority of *Ph.*

orientalis females (39.34%) were collected near human dwellings. In Konso, Ethiopia,

Ph. martini/Ph. celiae females were mainly captured from termite hills (74.46%, *P*

<0.0001) that were prevalent in both peridomestic and sylvatic ecotypes, followed by

temporary shelters locally known as "foras" (8.15%; Figure 2B).

We used *Phlebotomus* males to verify species prevalence and to indicate breeding

sites. *Ph. orientalis* represented 100% of all collected males in Kenya (n = 260) and

Sudan (n = 4714) (Supplementary Figure 2A). Like females, *Ph. orientalis* males in

Marsabit, Kenya, were collected from vegetation (46.92%, *P* <0.0001), outside houses

near *Acacia* and *Balanites* trees (18.85%), and in animal enclosures (19.23%) (Supplementary Figure 2B). Preliminary evidence of breeding sites, identified by finding unrotated or partially rotated male genitalia (n = 12), implicated outdoor ecotypes (Supplementary Figure 2C). In Gedaref, Sudan, *Ph. orientalis* males were mainly collected from vegetation (33.64%, *P* <0.0001) and indoors (22.70%, *P* <0.0001) (Supplementary Figure 2B), similar to females. Of the captured males with unrotated or partially rotated genitalia (n = 33), 51.52% (*P* <0.0001) were collected from vegetation and 21.21% from animal enclosures (Supplementary Figure 2C), potentially indicating that *Ph. orientalis* may not be breeding indoors in either Kenya or Sudan.

In Konso, Ethiopia, *Ph. martini* was the main male species collected (Supplementary Figure 2A), mostly from termite hills (72.35%, *P* <0.0001) (Supplementary Figure 2B), again similar to females. Other male species included *Ph. celiae* (5.80%) and *Ph. orientalis* (1.02%) (Supplementary Figure 2A). Some 72.92% (n = 35, *P* <0.0001) of *Ph. martini* males with unrotated or partially rotated genitalia were collected from termite hills (Supplementary Figure 2C), implicating this microhabitat as a significant breeding site for *Ph. martini*. Of note, we collected one *Ph. celiae* male with unrotated/partially rotated genitalia from a domestic animal enclosure. The relative abundance of *Ph. martini* (94.14%) over *Ph. celiae* males (5.80 %) indicates predominance of the former species in our female collections. As such, we will use *Ph. martini* instead of *Ph. martini/Ph. celiae* for females from here on.

Vector species exhibit different feeding preferences

We performed host blood meal analysis on field-collected individual blood fed

Phlebotomus females using a custom-made multiplex PCR tailored to amplify blood

from humans and seven animals present in our study sites (Supplementary Figure 3,

Supplementary Table 2). The multiplex PCR panel identified 63.8% (n = 270/423) of

blood meals. After *cytochrome b* (*cytb*) gene sequencing of unknowns this increased to

96.9% (n = 410/423). Interestingly, the diversity of identified blood meal sources was

considerably higher in Marsabit, Kenya, (13 hosts) compared to either Gedaref, Sudan,

(5 hosts) or Konso, Ethiopia, (4 hosts) (Figure 3A-C). Of interest, the observed number

of sand flies with multiple blood meals were higher than expected for Marsabit, Kenya,

(c^2 = 44.09, d.f. = 15, *P* <0.0001) and Gedaref, Sudan (c^2 = 17.83, d.f. = 6, *P* <0.0067).

Our results also establish that several females contained mixed blood meals from

humans and domestic or sylvatic animals such as hyraxes and mongooses (Figure 3A-

C).

In Marsabit, Kenya, we collected 226 *Ph. orientalis* and two *Ph. alexandri* blood fed

females from animal enclosures (32.30%, *P* <0.0001), vegetation (31.86%, *P* <0.0001),

and temporary shelters (23.89%, *P* = 0.001), following a similar pattern to unfed/gravid

sand flies (Figure 3D). Humans were identified as the dominant host for *Ph. orientalis*,

representing 37.76% (*P* <0.0001) of single blood meals, followed by rodents (15.89%, *P*

<0.0001), dogs (9.27%) and other sylvatic animals like gazelles (4.64%) (Figure 3A).

Mixed blood meals from two or three hosts were identified in 63 *Ph. orientalis*

specimens including human/Ovis (22.22%), and human/cow (17.46%) (Figure 3A). In

Gedaref, Sudan, 29.51% (*P* <0.0001) of the 183 collected blood fed *Ph. orientalis*

specimens were captured inside the house, and 33.33% (*P* <0.0001) in vegetation

(Figure 3D). Similar to Marsabit, Kenya, humans (74.83%, *P* <0.0001) were the

predominant host, while 14.97% fed on donkey (Figure 3B). Only seven specimens

contained mixed blood meals that fed on human plus other domestic and sylvatic

animals such as mongooses and rabbits (Figure 3B). In Ethiopia, only 12 *Ph. martini*

blood fed specimens were collected from termite hills (50%, *P* <0.001) and animal

enclosures (25%) (Figure 3D). Contrary to the anthropophilic feeding behavior observed

for *Ph. orientalis*, *Ph. martini* seemed to preferentially feed on cows (37.5%), and Ovis

(25.0%), and five sand flies fed on multiple hosts such as human/cow and hyrax/cow

(Figure 3C). Of the few collected *Ph. alexandri* blood fed females, the two from Kenya

fed on humans, while in Ethiopia, the one specimen fed on an Ovis.

***Leishmania*-infected vector species were recovered from diverse microhabitats**

Using qPCR (Supplemental Figure 4) on individual (n_i) or pooled (n_p) sand flies, we

detected *Leishmania*-infected specimens in both peridomestic and sylvatic niches

across all study sites (Figure 4A), and from distinct ecotypes (Figure 4B-C). The

minimal infection rates of *Ph. orientalis* with *Leishmania* were similar in Marsabit,

Kenya, (26/648, 3.9%) and Gedaref, Sudan, (10/281, 3.6%). Ethiopia had a lower rate

of infection of 1.5% (3/183) for *Ph. martini* females (Figure 4A). Of note, none of the

other sand fly species collected throughout this study were infected.

In Marsabit, Kenya, 4.3% (18/422) unfed/gravid and 3.5% (8/226) blood fed *Ph.*

orientalis were *Leishmania* positive (Figure 4B-C). Infected *Ph. orientalis* were

associated with vegetation (30.77%), where we captured ten infected unfed/gravid and

four infected blood fed females in sylvatic sites, and seven unfed/gravid and three blood

fed specimens from peridomestic sites (Supplementary Tables 1 and 3). Interestingly, a cluster of six infected sand flies was collected outside and inside the house of a relapsed VL patient in Laisamis, Kenya (Supplementary Tables 1 and 3), alluding to how hotbeds of transmission form. The patient, negative for the human immunodeficiency virus, received treatment in 2021 and refused treatment after relapsing in January 2024 (Supplementary Table 3), highlighting the need for better surveillance and treatment. Though four of the eight (50%) infected blood fed *Ph. orientalis* fed on humans, confirming the vector's anthropophilic behavior, finding three and one infected sand flies that fed on gazelles and a rodent, respectively (Figure 4C, Supplementary Tables 3 and 4), stresses the need to investigate whether there is a role for animal reservoirs in VL transmission in East Africa.

For Gedaref, Sudan, one of the 98 (1.0%) screened unfed/gravid specimens collected from inside a house was positive for *Leishmania* (Figure 4B). Importantly, we found nine (9/183, 4.9%) blood fed *Leishmania*-infected *Ph. orientalis* females, seven from peridomestic sites, and two from vegetation and a termite hill in sylvatic sites within Dinder Park (Figure 4C, Supplementary Table 3). All nine infected sand flies fed on humans, further emphasizing the anthropophilic behavior of *Ph. orientalis* in this region (Figure 4C, Supplementary Tables 3 and 4). The focal nature of transmission was again revealed by finding three infected blood fed sand flies, two inside the house of a VL patient, collected in consecutive days, and a third that was captured from inside the house of the patient's neighbor (Supplementary Tables 1 and 3).

In Ethiopia, three (1.8%, n = 3/171) unfed/gravid *Ph. martini* were positive for *Leishmania,* all captured in termite hills (Figure 4B). Two of the three were captured 9

months apart from the same castellated termite hill located in Galga near the house of a

recent VL case (Supplementary Tables 1 and 3), again highlighting the focality of VL.

The prevalence of *Ph. martini* females in termite hills and the exclusive collection of

Leishmania-infected specimens from this microhabitat implicates it in VL transmission in

Ethiopia, and potentially elsewhere in East Africa.

Leishmania rK39 seroprevalence reveals an increased risk of infection for

individuals living near VL cases

As *Ph. orientalis* is the vector of *L. donovani* in both Marsabit, Kenya, and Gedaref,

Sudan, we assessed seroprevalence of *Leishmania* rK39 antibodies in these study foci.

We conducted a case-control study of household members of VL cases and five

neighboring households for comparison against non-neighboring controls. These

comprised households located approximately 1.5 to 10 km from respective index VL

cases within the endemic villages (Kenya) or households located in an adjacent non-

endemic village 80 km away from Kunaynah (Sudan). DBSs were collected from

household members of 48 VL cases diagnosed during December 2023 to May 2025 in

Laisamis and Karare villages, Marsabit county, Kenya, and household members of five

VL cases diagnosed during August 2023 to August 2024 in Kunaynah village, Gedaref

state, Sudan (Supplementary Table 5). In Kenya, we enrolled a total of 752 subjects

(male to female ratio of 1:1.54) with a median age of 14 years (range, 2 – 85 years). In

Sudan, we enrolled a total of 435 participants (male to female ratio of 1:1.3) with a

median age of 15 years (range, 5 months – 85 years). Thirty-two (n = 108 household

members) and ten (n = 145 household members) control households were screened in

Kenya and Sudan, respectively.

rK39 titers were significantly higher for individuals living with or near cases (neighboring

households) than for those of non-neighboring controls in both Kenya and Sudan

(Figure 5 A-B). In Kenya, 19.80% (118/596) of screened asymptomatic participants

living with or near VL cases exhibited antibodies against rK39, compared to 7.41%

(8/108) of those living farther away. Similarly, 8.42% (24/285) of asymptomatic people

living with or near VL cases in Sudan were seropositive, versus only 2.07% (3/145) of

those living away from cases. The odds ratio calculated by Fisher's test for those living

with or near a VL case was 3.08 (P = 0.0015) and 4.35 (P = 0.0105) for Kenya and

Sudan, respectively. Additionally, logistic regression demonstrated that the likelihood of

developing rK39 antibodies decreases significantly with distance from a VL case for

both Kenya (Figure 5C; Likelihood ratio = 19.8805, P < 0.0001; z = -3.1344, P = 0.0017)

and Sudan (Figure 5D; Likelihood ratio = 7.2247, P = 0.0072; z = -2.2891, P = 0.0221).

Importantly, the same outcome was observed when the analysis excluded VL index

case household members for both Kenya (Supplemental Figure 5 A; Likelihood ratio =

16.5773, P < 0.0001; z = -2.95592, P = 0.0031) and Sudan (Supplemental Figure 5B;

Likelihood ratio, 5.71154, P = 0.0169; z = -2.08351, P = 0.0372). Of note, at the time of

serum collection, only 22/48 (Kenya) and none (Sudan) of the diagnosed and treated VL

patients had detectable antibodies by rK39 ELISA (Supplementary Table 5), considered

the gold standard VL serodiagnostic test, even though DBSs were collected within 1-4

months of treatment.

Discussion

The WHO has recently targeted visceral leishmaniasis in East Africa for elimination as a public health concern by 2030[12]. Our findings provide evidence of widespread and intense VL transmission in diverse settings across our study sites to inform control efforts.

Consistent with earlier reports from the region[7, 9, 44], *Ph. orientalis* was the most prevalent *Phlebotomus* species in Sudan and northern Kenya, while *Ph. martini* predominated in southern Ethiopia. We established that *Ph. orientalis* was predominant in *Acacia* forests in association with vertisol and animal burrows, with a high proportion of juvenile males collected outside, in vegetation and animal enclosures, demonstrating that these microhabitats are suitable breeding sites for this vector as previously suggested[9, 15, 18, 45]. In comparison, termite hills were the most productive site for *Ph. martini*, as reported previously[8, 46-48], and the collection of over 72% of newly emerged males, provides direct evidence of this ecotype as a preferential breeding site for *Ph. martini.*

We recovered *Leishmania*-infected *Ph. orientalis* from different microhabitats within our study sites in Marsabit, Kenya, and Gedaref, Sudan, in both sylvatic and peridomestic niches, and frequently from the same location months apart. This indicates that *Ph. orientalis* has adapted to diverse ecological microhabitats that provide sugar and blood meal sources, and breeding habitats[15, 16, 18], driving widespread yet focal transmission of VL. Supporting this observation, our data indicates that *Ph. orientalis* likely shifts its behavior to adapt to climatic conditions. Collections from Helat-Belo, Gedaref state, Sudan, conducted during the dry season in March to June 2016-2018 found infected

blood fed *Ph. orientalis* mainly outside houses and in sylvatic ecotypes, indicating they

are exophilic, and reported that they fed on human, cattle, and donkeys[17]. In contrast,

our sand fly collections from Kunaynah and Umslala-Houg were performed during the

rainy season (June to September), when people tend to sleep inside their houses.

Consequently, *Leishmania*-infected *Ph. orientalis* females were collected mainly indoors

and had fed soley on humans, suggesting a potential shift in vector behavior, with sand

flies likely resting inside the house after feeding on humans. While the majority of

juvenile *Ph. orientalis* males were collected from vegetation and animal enclosures, we

also collected juvenile males indoors in Sudan, possibly due to the widespread nature of

vertisols (black cotton soil) in the village, including inside houses, serving as a breeding

site for this vector[18], and creating an ideal environment for focal and active VL

transmission indoors. Further confirmation of this shift in vector behavior and

transmission pattern is needed to inform control measures that are currently focused on

outdoor applications.

In Konso, Ethiopia, *Ph. martini* were predominantly collected from termite hills, both in

sylvatic and peridomestic settings, pointing to a restricted habitat preference compared

to *Ph. orientalis*. Additionally, all three *Leishmania*-infected sand flies were recovered

from termite hills reinforcing findings of a previous study[8]. Capturing two *Leishmania*-

infected females nine months apart from a peridomestic castellated termite hill

reinforces the focality of infection sources and highlights the importance of this

microhabitat in sustaining transmission by *Ph. martini*. Notably, this termite hill was

located only 20 meters away from a household that reported a VL case. In-depth

studies focused on termite hills are needed to fully elucidate the involvement of this

microhabitat in VL transmission.

Several studies have reported infection rates in field-collected sand flies from Kenya

(0.3%), Sudan (1.6% and 1.4%)[17, 49], and southern (5.8%)[8] and northern Ethiopia

(23%)[21]. In our study, infection rates were higher than was previously reported for

Kenya (3.9%) and Sudan (3.6%), and were lower for southern Ethiopia (1.5%).

Additionally, reinforcing previous findings, we recovered infected specimens from both

sylvatic and domestic/peri-domestic habitats[8, 17, 19, 21, 49, 50].

Host attractiveness to vector sand flies, seroprevalence of *Leishmania* antibodies, and

PCR detection of *L. donovani* DNA have been reported for various domestic and wild

animal species in East Africa[51-55], but live parasites have not been isolated from any

thus far. Irrespective of whether domestic or wild animals are reservoirs, VL endemicity

in scarcely populated areas supports the existence of a zoonotic transmission cycle in

East Africa[17, 22, 56]. Establishing whether animal reservoirs are involved in sustaining VL

transmission in East Africa is a major knowledge gap that needs to be urgently

addressed to inform on the best strategy for VL elimination in this region.

Finding *Leishmania*-infected sand flies near VL cases from all our study sites affirms the

importance of human reservoirs in maintaining VL transmission, and the need for

continued surveillance and rapid treatment of cases. This is further reinforced by rK39

serological surveys where participants living near VL cases exhibited a significantly

higher proportion of positive rK39 antibodies and an increased risk associated with

proximity to VL cases (log-odds of positivity decreased with increasing distance from the

index households) in both Marsabit, Kenya, and Gedaref, Sudan. Importantly, the high

prevalence of rK39 antibodies in asymptomatic participants provides further evidence of

prevalent and efficient transmission, and stresses the urgency of determining their

involvement as potential parasite reservoirs[57].

A limitation of our study is the use rK39 to assess seroprevalence and the small sample

size of VL index cases in Sudan. The sensitivity of rK39 in East Africa is known to be

suboptimal[58-60]. In our study, this is reinforced by the loss of rK39 positivity in recently

diagnosed VL cases emphasizing the need for diagnostic tools with better specificity

and sensitivity to detect *Leishmania* infections in the region. Although we observed a

significant decrease in rK39 seropositivity with increasing distance from index VL case

households in both Kenya and Sudan, future studies with a larger sample size are

needed to assess the increased risk of living in proximity to VL cases. For studies of

vector sand flies, our aim was to identify their preferred habitats through a broad

investigation of distinct foci within three East African countries. Moving forward, we

need in-depth longitudinal studies focused on identified hotbeds of infection to better

understand the requisites supporting transmission in specific microhabitats.

Our exploratory ecoepidemiological study conducted in three East African countries

reports widespread VL transmission in the region and emphasizes both its focality and

complexity. The information we provide on habitat preference and transmission hotspots

of vector species in distinct microhabitats will galvanize further studies across disease

foci throughout East Africa, needed for an effective elimination campaign.

Acknowledgements

We thank the community, administrative officers, and health care workers from all our

study sites. We are also grateful to Marsabit County, Kenya, for supporting us with

fieldwork implementation and logistics. We thank Professor EL-Hassan Center for

Tropical Diseases staff, community leaders and the population in the Kunaynah village,

Gedaref state, Sudan. We thank the Karat Zone Health Bureau, Ethiopia, for supporting

this work and the Arbaminch University for hosting the project in Southern Ethiopia. We

also acknowledge Rogers Asamba (KEMRI), and David Mbuvi (ICIPE) for their valuable

support during fieldwork; Dr. Geyeto Garra (Ethiopia) for support given to this project;

Mr. Haile Gebremariam (Ethiopia) for technical assistance.

Funding

Funding support for this study was obtained from the NIH under the Tropical Medicine

Research Centers (TMRC) grant (U01AI168619). This research was supported in part

by the Intramural Research Program of the National Institutes of Health (NIH).

The contributions of the NIH authors were made as part of their official duties as NIH

federal employees, and are in compliance with agency policy requirements, and

are considered Works of the United States Government. However, the findings and

conclusions presented in this paper are those of the authors and do not necessarily

reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author contributions

E.I. performed the experiments, analyzed the data, conceived the study, supervised the

work, and wrote the manuscript. M.T., Steve Kiplagat, A.G., E.A., O.B., J.I., J.M.M.,

M.A., O.D., A.C., E.K., P.H., S.D., P.C., G.B., C.M., T.D.S., M.P., M.O., E.A.G.K.,

O.F.O., and B.M.Y. and D.M.M. undertook the field work, performed the experiments,

analyzed the data, and edited the manuscript. G.Z. and S.D. performed the sand fly

statistical analysis. J.G.V., D.K.M., A.M.M., A.H., A.S., E.I. and Shaden Kamhawi wrote

the manuscript. E.I., J.G.V., D.K.M., A.M.M., A.H., A.S., Shaden Kamhawi, and D.M.M.

conceived the study and supervised the work.

Competing interest

The authors declared no competing interest.

Data availability

All data produced in the present work are contained in the manuscript

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used GPT-4o (OpenAI, NIAID GenAI

Toolkit) in order to improve readability and language of the manuscript. After using this

tool/service, the authors reviewed and edited the content as needed and take full

responsibility for the content of the publication.

References

1.   World Health Organization. WHO: Leishmaniasis Fact Sheet. 2024 [cited 2024

May, 17, 2024]; Available from: https://www.who.int/news-room/fact-

sheets/detail/leishmaniasis

2.   eBioMedicine. Leishmania: an urgent need for new treatments. eBioMedicine.

2023;87.

3.   Baleela R, Llewellyn MS, Fitzpatrick S, Kuhls K, Schonian G, Miles MA, et al.

Leishmania donovani populations in Eastern Sudan: temporal structuring and a link

between human and canine transmission. Parasit Vectors. 2014 Nov 20;7:496.

4.   Zackay A, Cotton JA, Sanders M, Hailu A, Nasereddin A, Warburg A, et al.

Genome wide comparison of Ethiopian Leishmania donovani strains reveals differences

potentially related to parasite survival. PLoS Genet. 2018 Jan;14(1):e1007133.

5.   Zackay A, Nasereddin A, Takele Y, Tadesse D, Hailu W, Hurissa Z, et al.

Polymorphism in the HASPB repeat region of East African Leishmania donovani strains.

PLoS Negl Trop Dis. 2013;7(1):e2031.

6.   Alvar J, den Boer M, Dagne DA. Towards the elimination of visceral

leishmaniasis as a public health problem in east Africa: reflections on an enhanced

control strategy and a call for action. Lancet Glob Health. 2021 Dec;9(12):e1763-e9.

7.   Elnaiem DE. Ecology and control of the sand fly vectors of Leishmania donovani

in East Africa, with special emphasis on Phlebotomus orientalis. J Vector Ecol. 2011

Mar;36 Suppl 1:S23-31.

8.   Gebre-Michael T, Lane RP. The roles of Phlebotomus martini and P.celiae

(Diptera: Phlebotominae) as vectors of visceral leishmaniasis in the Aba Roba focus,

southern Ethiopia. Med Vet Entomol. 1996 Jan;10(1):53-62.

9.   Gebre-Michael T, Malone JB, Balkew M, Ali A, Berhe N, Hailu A, et al. Mapping

the potential distribution of Phlebotomus martini and P. orientalis (Diptera:

Psychodidae), vectors of kala-azar in East Africa by use of geographic information
systems. Acta Trop. 2004 Mar;90(1):73-86.

10. José Antonio Ruiz-Postigo aSJ, a Serge Madjou,a Ana Nilce Maia-Elkhoury,b
Samantha Valadas,b Supriya Warusavithana,c Mona Osman,c Aya Yajima,d Zaw Lin,d
Abate Beshahe and Sunghye Kimf,. Global leishmaniasis surveillance: 2021, assessing
the impact of the COVID-19 pandemic. 2021.

11. World Health Organization. Ending the neglect to attain the Sustainable
Development Goals: A road map for neglected tropical diseases 2021–2030; 2021.

12. World Health Organization. Strategic framework for the elimination of visceral
leishmaniasis as a public health problem in eastern Africa 2023–2030. 2024;Licence:
CC BY-NC-SA 3.0 IGO.

13. Alvar J, Beca-Martinez MT, Argaw D, Jain S, Aagaard-Hansen J. Social
determinants of visceral leishmaniasis elimination in Eastern Africa. BMJ Glob Health.
2023 Jun;8(6).

14. Wasunna M, Musa A, Hailu A, Khalil EA, Olobo J, Juma R, et al. The
Leishmaniasis East Africa Platform (LEAP): strengthening clinical trial capacity in
resource-limited countries to deliver new treatments for visceral leishmaniasis. Trans R
Soc Trop Med Hyg. 2016 Jun;110(6):321-3.

15. Gebresilassie A, Kirstein OD, Yared S, Aklilu E, Moncaz A, Tekie H, et al.
Species composition of phlebotomine sand flies and bionomics of Phlebotomus
orientalis (Diptera: Psychodidae) in an endemic focus of visceral leishmaniasis in
Tahtay Adiyabo district, Northern Ethiopia. Parasit Vectors. 2015 Apr 25;8:248.

16.   Hassaballa IB, Torto B, Sole CL, Tchouassi DP. Exploring the influence of

different habitats and their volatile chemistry in modulating sand fly population structure

in a leishmaniasis endemic foci, Kenya. PLoS Negl Trop Dis. 2021 Feb;15(2):e0009062.

17.   Khogali A, Elnaiem DA, Diaz-Reganon R, Jibreel T, Nour BYM, Abdelrahman

SH, et al. Infection of Leishmania donovani in Phlebotomus orientalis Sand Flies at

Different Microhabitats of a Kala-Azar Endemic Village in Eastern Sudan. Trop Med

Infect Dis. 2024 Feb 2;9(2).

18.   Moncaz A, Kirstein O, Gebresellassie A, Lemma W, Yared S, Gebre-Michael T,

et al. Characterization of breeding sites of Phlebotomus orientalis - the vector of visceral

leishmaniasis in northwestern Ethiopia. Acta Trop. 2014 Nov;139:5-14.

19.   Elnaiem DE, Osman OF. Evidence for active transmission of visceral

leishmaniasis within a village in eastern Sudan. Acta Trop. 1998 Nov 30;71(3):305-9.

20.   Hailu A, Balkew M, Berhe N, Meredith SE, Gemetchu T. Is Phlebotomus

(Larroussius) orientalis a vector of visceral leishmaniasis in south-west Ethiopia? Acta

Trop. 1995 Sep;60(1):15-20.

21.   Kirstein OD, Skrip L, Abassi I, Iungman T, Horwitz BZ, Gebresilassie A, et al. A

fine scale eco-epidemiological study on endemic visceral leishmaniasis in north

ethiopian villages. Acta Trop. 2018 Jul;183:64-77.

22.   Zijlstra EE, el-Hassan AM. Leishmaniasis in Sudan. Visceral leishmaniasis.

Trans R Soc Trop Med Hyg. 2001 Apr;95 Suppl 1:S27-58.

23.   Votypka J, Pavlasova M, Volfova V, Volf P. Rotation of male genitalia in various

species of phlebotomine sandfly. Med Vet Entomol. 2015 Dec;29(4):355-60.

24. Mondal D, Bern C, Ghosh D, Rashid M, Molina R, Chowdhury R, et al.
Quantifying the Infectiousness of Post-Kala-Azar Dermal Leishmaniasis Toward Sand
Flies. Clin Infect Dis. 2019 Jul 2;69(2):251-8.

25. Singh OP, Chaubey R, Kushwaha AK, Fay MP, Sacks D, Sundar S. Visceral
Leishmaniasis-Human Immunodeficiency Virus-Coinfected Patients Are Highly
Infectious to Sandflies in an Endemic Area in India. J Infect Dis. 2024 May 7.

26. Singh OP, Tiwary P, Kushwaha AK, Singh SK, Singh DK, Lawyer P, et al.
Xenodiagnosis to evaluate the infectiousness of humans to sandflies in an area
endemic for visceral leishmaniasis in Bihar, India: a transmission-dynamics study.
Lancet Microbe. 2021 Jan;2(1):e23-e31.

27. Al-Salem W, Herricks JR, Hotez PJ. A review of visceral leishmaniasis during the
conflict in South Sudan and the consequences for East African countries. Parasit
Vectors. 2016 Aug 22;9(1):460.

28. Berry I, Berrang-Ford L. Leishmaniasis, conflict, and political terror: A spatio-
temporal analysis. Soc Sci Med. 2016 Oct;167:140-9.

29. Boodman C, van Griensven J, Gupta N, Diro E, Ritmeijer K. Anticipating visceral
leishmaniasis epidemics due to the conflict in Northern Ethiopia. PLoS Negl Trop Dis.
2023 Mar;17(3):e0011188.

30. Kenya National Bureau of Statistics. Kenya Population and Housing Census.
2019;II.

31. de Souza DK, Picado, A., Bessell, P. R., Liban, A., Wachira, D., Mwiti, D.,
Matendechero, S. H., Bosch, M., Ndung'u, J. M., & Cruz, I.,. Strengthening Visceral

Leishmaniasis Diagnosis Capacity to Improve Access to Care in Kenya: The Example of

Marsabit County. Frontiers in Tropical Diseases. 2021.

32.   Mewara A, Gudisa R, Padhi BK, Kumar P, Sah R, Rodriguez-Morales AJ.

Visceral leishmaniasis outbreak in Kenya-a setback to the elimination efforts. New

Microbes New Infect. 2022 Nov-Dec;49-50:101060.

33.   Department KM. Kenya Meterological deperment, . 2024 [cited; Available from:

https://meteo.go.ke

34.   Ibrahim B, Elmadani M, Sahar H, Gadallh M, Abdallh A, Ahmed AEE. Prevalence

of Visceral Leishmaniasis among Wildlife Rangers in Dinder National Park, Sudan.

medRxiv. 2024:2024.06.24.24309386.

35.   Abonnenc E, Minter D. Bilingual keys for the identification of the sandflies of the

Ethiopian Region (Frenceh and English). Cahier, Office de la Recherche Scintifique et

Technique d'Outre, Mer Entomologie Medicale. 1965;5(1).

36.   Lewis D.J. The phlebotomine sandflies (Diptera: Psychodidae) of the Oriental

Region. Bull Br Museum Nat Hist 1978;37:217–343. .

37.   Dey R, Joshi AB, Oliveira F, Pereira L, Guimaraes-Costa AB, Serafim TD, et al.

Gut Microbes Egested during Bites of Infected Sand Flies Augment Severity of

Leishmaniasis via Inflammasome-Derived IL-1beta. Cell Host Microbe. 2018 Jan

10;23(1):134-43 e6.

38.   Kent RJ, Norris DE. Identification of mammalian blood meals in mosquitoes by a

multiplexed polymerase chain reaction targeting cytochrome B. Am J Trop Med Hyg.

2005 Aug;73(2):336-42.

39.   Irwin DM, Kocher TD, Wilson AC. Evolution of the cytochrome b gene of

mammals. J Mol Evol. 1991 Feb;32(2):128-44.

40.   Parodi B, Aresu O, Bini D, Lorenzini R, Schena F, Visconti P, et al. Species

identification and confirmation of human and animal cell lines: a PCR-based method.

Biotechniques. 2002 Feb;32(2):432-4, 6, 8-40.

41.   Collini M, Albonico F, Hauffe HC, Mortarino M. Identifying the last bloodmeal of

questing sheep tick nymphs (Ixodes ricinus L.) using high resolution melting analysis.

Vet Parasitol. 2015 Jun 15;210(3-4):194-205.

42.   Abbasi I, Cunio R, Warburg A. Identification of blood meals imbibed by

phlebotomine sand flies using cytochrome b PCR and reverse line blotting. Vector

Borne Zoonotic Dis. 2009 Feb;9(1):79-86.

43.   Kenya Ministry of Health. Prevention, Diagnosis and Treatment of Visceral

Leishmaniasis (Kala azar) in Kenya: National Guidelines.   2017   [cited August 27,

2025];   Available   from: https://www.afrikadia.org/wp-

content/uploads/2018/08/VL_Guidelines_Kenya_2017.pdf

44.   Anjili CO, Ngumbi PM, Kaburi JC, Irungu LW. The phlebotomine sandfly fauna

(Diptera: Psychodidae) of Kenya. J Vector Borne Dis. 2011 Dec;48(4):183-9.

45.   Thomson MC, Elnaiem DA, Ashford RW, Connor SJ. Towards a kala azar risk

map for Sudan: mapping the potential distribution of Phlebotomus orientalis using digital

data of environmental variables. Trop Med Int Health. 1999 Feb;4(2):105-13.

46.   Basimike M MM, Kumar R..,. Habitat preference and seasonal variations of

phlebotomine sandflies. Insect Science and Its Application 1992;13(3):307-314. .

47.   Minter D. The distribution of sandflies (Diptera, Psychodidae) in Kenya. Bulletin

of Entomological Research. 1964; 55, 205-217

48.   Wijers DJ, Minter DM. Studies on the vector of kala-azar in Kenya. V. The

outbreak in Meru district. Ann Trop Med Parasitol. 1966 Mar;60(1):11-21.

49.   Hassan MM, Elraba'a FM, Ward RD, Maingon RD, Elnaiem DA. Detection of high

rates of in-village transmission of Leishmania donovani in eastern Sudan. Acta Trop.

2004 Sep;92(1):77-82.

50.   Perkins PV, Githure JI, Mebrahtu Y, Kiilu G, Anjili C, Ngumbi PS, et al. Isolation

of Leishmania donovani from Phlebotomus martini in Baringo district, Kenya. Trans R

Soc Trop Med Hyg. 1988;82(5):695-700.

51.   Dereure J, El-Safi SH, Bucheton B, Boni M, Kheir MM, Davoust B, et al. Visceral

leishmaniasis in eastern Sudan: parasite identification in humans and dogs; host-

parasite relationships. Microbes Infect. 2003 Oct;5(12):1103-8.

52.   Elnaiem DA, Hassan MM, Maingon R, Nureldin GH, Mekawi AM, Miles M, et al.

The Egyptian mongoose, Herpestes ichneumon, is a possible reservoir host of visceral

leishmaniasis in eastern Sudan. Parasitology. 2001 May;122(Pt 5):531-6.

53.   Gebresilassie A, Yared S, Aklilu E, Kirstein OD, Moncaz A, Tekie H, et al. Host

choice of Phlebotomus orientalis (Diptera: Psychodidae) in animal baited experiments: a

field study in Tahtay Adiyabo district, northern Ethiopia. Parasit Vectors. 2015 Mar

31;8:190.

54.   Hassan MM, Osman OF, El-Raba'a FM, Schallig HD, Elnaiem DE. Role of the

domestic dog as a reservoir host of Leishmania donovani in eastern Sudan. Parasit

Vectors. 2009 Jun 17;2(1):26.

55.   Kassahun A, Sadlova J, Benda P, Kostalova T, Warburg A, Hailu A, et al. Natural

infection of bats with Leishmania in Ethiopia. Acta Trop. 2015 Oct;150:166-70.

56.   Elnaiem DA, Hassan HK, Ward RD. Phlebotomine sandflies in a focus of visceral

leishmaniasis in a border area of eastern Sudan. Ann Trop Med Parasitol. 1997

Apr;91(3):307-18.

57.   Singh OP, Hasker E, Sacks D, Boelaert M, Sundar S. Asymptomatic Leishmania

infection: a new challenge for Leishmania control. Clin Infect Dis. 2014

May;58(10):1424-9.

58.   Hagos DG, Schallig H, Kiros YK, Abdulkadir M, Wolday D. Performance of rapid

rk39 tests for the diagnosis of visceral leishmaniasis in Ethiopia: a systematic review

and meta-analysis. BMC Infect Dis. 2021 Nov 17;21(1):1166.

59.   Iniguez E, Kamhawi S. Discovery of new antigens for serodiagnosis of visceral

leishmaniasis in humans and dogs adds valuable tools to our arsenal. mBio. 2024 Aug

14;15(8):e0107924.

60.   Kassa M, Abdellati S, Cnops L, Bremer Hinckel BC, Yeshanew A, Hailemichael

W, et al. Diagnostic accuracy of direct agglutination test, rK39 ELISA and six rapid

diagnostic tests among visceral leishmaniasis patients with and without HIV coinfection

in Ethiopia. PLoS Negl Trop Dis. 2020 Dec;14(12):e0008963.

Figure legends

Figure 1. Study sites and incidence of visceral leishmaniasis

(A,B) Map of East Africa showing the location of study sites. 1, Marsabit; 2, Konso; 3,

Gedaref state. (B) Visceral leishmaniasis (VL) cases per 10,000 population over the

past seven years in Marsabit county, Kenya; Konso, Ethiopia; and Gedaref state,

Sudan. Incidence data are based on Ministry of Health reports from each country. Maps

at QGIS software QGIS shapefile, https://open.africa/dataset/?tags=Shapefiles.

Figure 2. Distribution of different sand fly species across distinct microhabitats in

endemic foci in Kenya, Sudan and Ethiopia study sites

(A) Relative abundance of collected *Phlebotomus* females by species. (B) Relative

abundance of *Phlebotomus* females by ecotype.

Figure 3. A mosaic of blood meal sources observed in leishmaniasis vectors

across endemic foci in Kenya, Sudan and Ethiopia study sites

(A-C) Host blood meal identity of blood fed females in Kenya (A), Sudan (B), and

Ethiopia (C) with a single or multiple sources of blood detected. (D) Relative abundance

of collected blood fed *Phlebotomus* females by ecotype.

Figure 4. Distribution of *Leishmania*-infected sand flies from distinct and diverse

microhabitats reveals hotspots of active transmission near and away human

habitation

(A) Number of *Leishmania*-infected sand flies collected from peridomestic or sylvatic

habitats. (B) Habitat productivity of infected unfed/gravid female *Ph. orientalis* sand flies

in Kenya and Sudan, and *Ph. martini/Ph. celiae* in Ethiopia. (C) Host blood meal identity

of infected blood fed *Ph. orientalis and Ph. martini/Ph. celiae* females. Specimens were

processed individually or as pools of up to 5-10 sand flies (B) or individually (C).

Figure 5. Prevalence of *Leishmania* rK39 antibodies in our study sites indicate high risk of infection in Marsabit county, Kenya, and Gedaref state, Sudan

(A,B) *Leishmania* rK39 antibody prevalence measured from dry blood spots (DBS) collected from individuals living with or near and away from recently reported VL cases in Laisamis and Karare, Marsabit county (A) and Kunaynah, Gedaref state (B). Data was normalized using the plate cutoff calculated by the mean optical density (O.D.) of non-endemic negative controls + 5 SD. Non-endemic negative controls were obtained from individuals living in Nairobi, Kenya. Samples with O.D. ≥ 0.00 after normalization were considered positive. (C,D) Univariable binary logistic regression analysis was performed to assess the effect of distance (meters) from index case households to neighboring and control households, on rK39 positivity in Marsabit, Kenya (C), and Gedaref, Sudan (D). The 'margins' command of Stata was used to compute adjusted log-odds of seropositivity at selected distances from the index households. Analysis includes index household members. A *P* value of ≤ 0.05 was considered significant by Mann-Whitney test (A,B), or univariable binary logistic regression (C,D); *$P < 0.05$, **$P < 0.01$.

Number of observations = 704
Number of positive outcomes = 126
Log likelihood (model) = −1.35940
Likelihood ratio (G^2) = 19.8805, $P<0.0001$
Slope: Z factor = −3.1344, $P = 0.0017$
R^2 = 0.02029
Equation log(odds) = -1.35940-0.000302236*X

Number of observations = 430
Number of positive outcomes = 26
Log likelihood (model) = −91.888
Likelihood ratio (G^2) = 7.2247, $P = 0.0072$
Slope: Z factor = −2.2891, $P = 0.0221$
R^2 = 0.01666
Equation log(odds) = $-2.43047 - 1.76834\times10^{-5}*X$