PANCDetect: Early Detection of Pancreatic Cancer from Multimodal EHR data with LLM Embeddings

Zicheng Jin[1*], Xuhui Guo[1*], Zehua Wang[1,2*], Qiang Yang[3], Xiaotong Yang[1], Xinyu Zhang[4], Rui Yin[3], Lana X. Garmire[1$]

[1]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

[2]Current Address: Department of Computer Science, University of Virginia, Charlottesville, VA, USA

[3]Department of Health Outcomes & Biomedical Informatics, University of Florida, Gainesville, FL, USA

[4]Department of Mathematics, University of Michigan, Ann Arbor, MI, USA

[*] These authors contributed equally to this work

[$] Corresponding author email: lgarmire@umich.edu

Abstract

Background

Pancreatic cancer (PANC) is often diagnosed at late stages due to the absence of specific early symptoms, resulting in one of the highest cancer mortality rates. While imaging modalities such as MRI and CT offer high diagnostic accuracy, their population-wide application is however impractical due to the cost. Electronic health records (EHRs) provide a routine, easily accessible, longitudinal and scalable data source for risk prediction, particularly for diseases with no specific symptom such as PANC.

Method

We introduce PANCDetect, a multimodal framework that leverages large language model (LLM)-derived embeddings of diagnoses, procedures, medications, and laboratory tests, and integrates these data modalities through a Transformer-based architecture. We train the model on MarketScan (≈250M patients) and validate it externally on additional large real-world EHR datasets of University of Michigan Precision Health, or UMPH data (n≈6M patients) and OneFlorida+ data (n≈26M patients). We then fine-tuned the general model on UMPH EHR data. We evaluated the performance of both models using metrics including area under the receiver operating characteristic curve (AUROC) and area under the precision-recall-gain curve (AUPRG). We assessed the top predictive features with integrated gradients (IG).

Result

In the MarketScan cohort, PANCDetect achieved an AUROC of 0.812 and AUPRG of 0.851 at the 6-month prediction window, and an AUROC of 0.735 and AUPRG of 0.629 for 60-month prediction, significantly outperforming CancerRiskNet. External validation on UMPH and OneFlorida+ demonstrated good generalizability, with 6-months AUROC scores of 0.711 and 0.793, respectively. Fine-tuning on UMPH with laboratory data further improved performance,

reaching an AUROC of 0.927 and an AUPRG of 0.979 at 6 months. Even at the 60-month horizon,

the refined PANCDetect model maintained strong performance, with an AUROC of 0.927 and

AUPRG of 0.979. Attribution analysis highlighted type 2 diabetes, pancreatic diseases, personal

and family cancer history as the most important risk factors.

Conclusion

PANCDetect is the state-of-the-art method integrating multimodal EHR data with LLM

embeddings for accurate, interpretable, and generalizable early prediction of pancreatic cancer.

This framework holds promise for precision screening of high-risk patients, with the potential to

improve survival outcomes without increasing healthcare costs.

Introduction

Pancreatic cancer (PANC), is characterized by the lack of specific early clinical manifestations.[1] In many cases, individuals may experience vague or nonspecific symptoms that precede diagnosis by years.[2] Based on a global analysis of data from 48 countries, incidence and mortality of PANC are rising globally, particularly among women and individuals aged 50 and older, with concerning increases also seen in younger populations.[3] The absence of validated screening protocols, lack of reliable early biomarkers, combined with the retroperitoneal and anatomically inaccessible location of the pancreas, hinders timely detection. As a result, less than 20% patients are diagnosed at an early, resectable stage, resulting in a persistently high mortality rate of this malignancy.[4] Additionally, delayed diagnosis of PANC imposes substantial financial burdens on patients, families, and healthcare systems, as late-stage treatments are far more costly than early surgery with adjuvant therapy.[5]

Routine MRI and CT can be an effective screening method for detecting PANC with high sensitivity and specificity,[6] but applying cross-sectional imaging at a population scale is impractical due to prohibitive costs and resource burden.[7] Electronic health record (EHR) is a much more economical, accessible and scalable data resource. EHR data include valuable baseline information such as demographics, as well as the longitudinal information including disease diagnoses, laboratory results, medication, and procedures, making them well-suited for early detection and prognosis modeling.[8] Moreover, the standardized EHR frameworks allow them to be more AI-ready, and comparable across different healthcare systems, which enhances the generalizability of predictive models across health centers.[9–11]

Previous transformer-based pretrained models demonstrated the feasibility of applying self-attention architectures to structured EHR data for the general prediction of future diagnoses of patients. Models such as BEHRT[12], Hi-BEHRT[13], and Med-BERT[14] adapted BERT-style encoders

to longitudinal claims and clinical coding sequences, effectively capturing temporal dynamics,

disease progression, and patient heterogeneity. Building on this trend, researchers have applied

Transformers to pancreatic cancer risk prediction, leveraging self-attention mechanisms to better

model temporal dependencies and disease evolution. Moreover, recent advances in large

language models (LLMs) offer new opportunities to extract clinically meaningful patterns hidden

among the millions of patients' EHR data. By treating EHRs as sequences of "clinical tokens",

such as diagnoses (ICD/SNOMED), procedures (CPT), medications (RxNorm/NDC), and

discretized lab values, pretrained LLMs can learn rich contextualized embeddings that capture

clinical semantics and longitudinal continuity across care settings.[15,16] General-purpose LLMs

such as GPT have been adapted to biomedical corpora (e.g., BioGPT[17]) and even to institution-

scale EHR data (e.g., NYUTron[18]), enabling predictive modeling, free-text understanding, and

zero or few-shot adaptation to diverse downstream tasks.

A few recent studies have leveraged EHR data to build models tailored for predicting PANC risk,

enabling identification of high-risk patient cohorts[19–22]. Among these efforts, CancerRiskNet[21]

stands out as a representative study that demonstrated the feasibility of using Transformers in

this domain. However, its multi-window prediction design, which does not exclude disease events

occurring shortly before cancer diagnosis, carries the risk of data leakage, potentially inflating

accuracy estimates. Moreover, CancerRiskNet relies solely on diagnosis codes, and it is

acknowledged that incorporating other clinically informative modalities could further improve

prediction accuracy with the real-world availability of data. Many existing models are trained on

relatively small datasets, restricting generalizability across diverse populations and health

systems[22]. To address these limitations and take advantage of LLMs, we propose PANCDetect,

a Transformer-based multimodal prediction framework trained on the MarketScan dataset, which

contains over 110,000 pancreatic cancer patients, making it the largest resource of its kind to date

(Fig. 1). PANCDetect not only incorporates diagnostic information but also procedures,

medications, and laboratory tests into LLM-based embeddings. We also employed a more

rigorous exclusion strategy to minimize data leakage in modeling and improve reliability. To

validate the generalizability of our approach, we tested and fine-tuned the model across two

independent healthcare systems. Consequently, PANCDetect substantially enhances prediction

accuracy and clinical applicability for early pancreatic cancer detection as compared to other

earlier efforts.

Methods

Study Cohorts

This study is conducted on three different retrospective datasets: Merative MarketScan

Databases,[23] University of Michigan Precision Health (UMPH),[24] and OneFlorida+ Clinical

Research Network.[25] MarketScan is a commercial database, which we have access to through

the University of Michigan. UMPH and OneFlorida+ data usage was approved by the Institutional

Review Board (IRB) of the University of Michigan Medical School (HUM#00259964) and the IRB

of the University of Florida (#IRB202400752), with a waiver of informed consent for secondary

use of de-identified EHR data. No data from UMPH or OneFlorida+ was shared outside the

respective organizations, ensuring compliance with data security and confidentiality.

MarketScan Database contains records from approximately 250 million patients from 2009–2022.

We identified 113,518 individuals diagnosed with PANC based on the diagnosis codes listed in

Supplementary Table 1. UMPH includes approximately six million patients from 2006 to 2025,

among whom 8,257 individuals are recorded as PANC.. OneFlorida+ Clinical Research Network

encompasses around 26 million patients from 72 hospitals across Florida, Alabama, Georgia, and

Arkansas. We identified 10,500 individuals diagnosed with PANC during 2013–2024. For each

cohort, we identified corresponding matched control patients, by applying a 1:5 matching strategy

based on sex, age, and geographic location (Supplementary Fig. 1) to mitigate the substantial

class imbalance.

We then used the MarketScan cohort as the primary training set given its largest PANC patient

size. We performed external validation on UMPH and OneFlorida+, and fine-tuned the model on

the UMPH cohort for better adaptation to the local population characteristics.

Feature Embedding

As shown in Fig. 2A, the input features comprise time-varying information and time-invariant

features. Time-varying information includes EHR clinical events such as diagnoses, procedures,

and medications, each annotated with temporal markers such as visit date and patient age at that

visit. Time-invariant attributes include sex and geographic location. For the time-variant features,

standardized textual descriptions of medical codes are encoded using the OpenAI text-

embedding-3-small model to generate dense semantic representations (without patient private

information). These embeddings are combined with positional embeddings to capture temporal

order and irregular intervals after linear projection, thereby enhancing the representation of

longitudinal patient trajectories. During fine-tuning on the UMPH cohort, we integrated laboratory

test results as a new input modality, using the same embedding generation method to maintain

consistency while enabling exploration of richer multi-modal inputs in a real-world clinical setting.

Other time-invariant features (e.g., sex and geographic location) are mapped into dense

representations using a feed-forward layer before being concatenated with the longitudinal

sequence representations.

Multi-modal Fusion Strategy

The proposed framework integrates multi-modal information within a Transformer-based

architecture (Fig. 2A). Here, three modalities, namely diagnoses, procedures, and medications,

are modeled through self-attention blocks to capture intra-modality dependencies. Cross-

attention mechanisms are then applied, using diagnosis events as queries and procedure and

medication events as key–value inputs, enabling the model to learn complementary information

across modalities. During fine-tuning on the UMPH cohort, additional modalities, such as

laboratory tests, can be incorporated in the same manner. The resulting multimodal

representations are concatenated and passed through a fusion multilayer perceptron (MLP) to

construct longitudinal patient representations, which are subsequently combined with time-

invariant features to predict pancreatic cancer risk (Fig. 2A). This framework is modular and can

be readily extended to accommodate further modalities as they become available, such as in the

refined model using UMPH data.

Diagnosis Prediction Intervals

Compared with prior studies[21,22], we adopted a single-interval prediction design. When predicting

"PANC diagnosis within N months", all EHR information on visits occurring within the N-month

interval before the PANC diagnosis was masked and only earlier EHR information was retained

to train the model (Fig. 2B). Specifically, for each PANC patient, we extracted all diagnoses,

procedures, and medication records occurring prior to the PANC diagnosis date. We truncated

the clinical trajectories according to different predefined prediction intervals N (N=6-, 12-, 24-, 36-,

60- month). For non-PANC controls, the reference date was defined as the end of their

observation period, and their records were truncated backward from this time point using the

same prediction intervals. This strategy is essential to reduce the risk of data leakage from

prodromal symptoms very close to the diagnosis date and provides a more realistic assessment

for early detection potential.

Implementation Details & Evaluation Metrics

Patients with valid longitudinal trajectories after preprocessing are randomly split into training, validation, and test sets in an 8:1:1 ratio. The model was trained using the binary cross-entropy loss with logits, optimized with AdamW (initial learning rate of 10^{-4}, weight decay applied). All training experiments were performed on 4 NVIDIA A40 GPUs for 10 epochs with a batch size of 128. Model performance was primarily assessed using several metrics: (1) the area under the receiver operating characteristic curve (AUROC), (2) the area under the precision-recall-gain curve (AUPRG),[26] normalized adjusted area under the precision–recall curve (AUPRC) with precision-gain and recall-gain scores, which handles the imbalances between case vs control samples. For each metric, we used the 95% confidence interval (CI) estimated via 200 bootstrap resamples. To provide a more comprehensive assessment, we also reported precision, recall, F1 score, and specificity (Supplementary Table 2-5).

Feature Interpretation

To enhance model interpretability, we applied integrated gradients (IG)[27] to perform attribution analysis on the input embeddings. IG is a path-integrated, gradient-based attribution method that quantifies feature contributions to the prediction outcome (see Supplementary Appendix for formulation). Specifically, it computes feature attributions by accumulating gradients of the prediction score with respect to the input along a straight-line path in the embedding space, interpolating from a baseline input to the actual input. We defined the baseline as a zero embedding vector, representing the absence of any clinical information. Based on the attribution results, we aggregated importance scores for all inputs and identified the clinical features most relevant to PANC risk prediction.

Results

Overview of Study cohorts

This study includes 3 large real-world patient clinical databases: (1) MarketScan Database which contains records from approximately 250 million patients from 2009–2022. (2) UMPH which includes approximately six million patients from 2006-2025. (3) OneFlorida+ Clinical Research Network which encompasses around 26 million patients from 72 hospitals across Florida, Alabama, Georgia, and Arkansas from 2013–2024. We employed 1:5 matching for PANC patients and controls, based on sex, age, and geographic location. As shown in Table 1, we identified 113,518, 8,257 and 10,500 PANC individuals in MartketScan, UMPH and OneFlorida+ respectively, among which the males account for 50.8%, 53.3% and 52.7%, similar among the datasets. The highest proportion of diagnoses is observed between ages 55 and 65, 65-70 and 65-75 (Fig. 3A, 3C and 3E). For MarketScan, most PANC patients exhibit trajectory lengths of 0–6 years with 50–400 accumulated diagnosis records (Fig. 3B), reflecting intensive healthcare utilization and multiple comorbidities in the years preceding diagnosis. Compared to the MarketScan database, PANC patients in UMPH generally have shorter trajectory lengths with most of them spanning 2 years or less, reflecting the high volume of referred high-risk patients in a research hospital (Fig. 3D). The OneFlorida+ cohort showed that PANC patients had trajectory lengths of ≤4 years (Fig. 3F) and accumulated a median of 96 diagnosis records (Table 1). In contrast, non-PANC patients generally have shorter trajectories in all three datasets, consistent with lower healthcare utilization.

Model Building on MarketScan Cohort

For each prediction interval setting, we obtained the model that achieved the minimum validation loss on the validation set, and then we evaluated its predictive ability on the MarketScan hold-out test set. As shown in Figs. 4A-B, the 6-month prediction window achieves the best performance, with an AUROC of 0.812 (95% CI, 0.807–0.818) and an AUPRG of 0.851 (95% CI, 0.843–0.860). The AUROC and AUPRG decrease as the prediction period gets longer, as expected. Still, the 60-month prediction window reaches the AUROC of 0.735 (95% CI, 0.724–0.745) and AUPRG of 0.629 (95% CI, 0.600–0.656). The shorter prediction window also has a larger training patient size, likely contributing to better model performance (Supplementary Table 6). To show where PANCDetect stands, we compared our model with CancerRiskNet[21], with the same data preprocessing and the same training, validation, and testing cohorts. PANCDetect consistently outperforms CancerRiskNet across all prediction intervals (averaged AUC 0.633-0.765), particularly on long-term predictions over 36 and 60 months (Supplementary Fig. 2).

Tests on the technical rigor of PANCDetect

We further evaluated several LLMs pretrained on biomedical corpora, such as BioGPT[17], which is trained on PubMed abstracts, and GatorTron[28], which is trained on large-scale de-identified clinical notes of University of Florida (UF) Health, PubMed articles, and Wikipedia. These domain-specific models were compared against the general-purpose embedding model OpenAI text-embedding-3-small, with predictive performance evaluated by AUROC and AUPRG on the test set. Experimental results show that the OpenAI text-embedding-3-small model achieves the performance that is comparable to, and in most cases better than, these specialized biomedical domain-specific models (Supplementary Table 7). This aligns with some of the existing literature showing that specialized models, whether trained from scratch or fine-tuned, often underperform compared to more general large-scale models across multiple tasks.[29,30]

We also performed another ablation study on multimodal fusion strategies by sequentially

removing procedure and/or drug medication modalities (Supplementary Table 8). The model

using diagnosis codes alone provided the baseline performance (AUROC of 0.779 and AUPRG

of 0.810 for 6-month prediction). Adding procedures or drugs yielded consistent gains, with

AUROC of 0.802 and AUPRG of 0.841, and AUROC of 0.801 and AUPRG of 0.828, for 6-month

prediction, respectively. The best results were obtained when all three modalities were combined

(AUROC of 0.812 and AUPRG of 0.851 for 6-month prediction). The results show that

incorporating both procedure and drug information led to a significant improvement on predictive

accuracy, revealing the complementary nature of these modalities.

External Validation Results on UMPH and OneFlorida+ Cohorts

To further evaluate the generalizability of our models, we tested them on two additional cohorts

from the OneFlorida+ research network and the UMPH (Fig. 4C-F)**.** The model achieves

comparable results on the OneFlorida+ cohort (Fig. 4C-D)**,** particularly in short-term (6-12 months)

prediction with AUROCs of 0.793 and 0.729, and AUPRGs of 0.917 and 0.859. The performance

declines with longer exclusion intervals as expected, as fewer patients are remaining for training

the model and the inherent difficulty for prediction over longer intervals . On the other hand, for

the UMPH Cohort, the model shows decent but lower predictive AUROC scores, ranging from

0.711 at 6-month prediction to 0.584 at 60-month prediction (Fig. 4E). AUPRG, however,

decreases as the exclusion interval lengthens, from 0.891 at 6-month prediction to 0.598 at 60

months (Fig. 4F).

Fine-tuning results on the UMPH Cohort

The performance decline observed on the two EHR cohorts (particularly UMPH data), as

compared to that of MarketScan hold-out data, is primarily attributable to the difference in EHR

data structure as compared to the MarketScan claims-based data. This includes differences in

coding systems, variable coverage, patterns of missingness,[31] as well as the underlying patient

populations, age distributions, and care-seeking behaviors, etc. Collectively, these factors

contribute to input feature distribution shifts that limit the generalizability of the models trained on

the MarketScan cohort originally. Therefore, it is necessary to fine-tune the model on the EHR

cohort to improve adaptability and healthcare-specific predictive performance.

Thus, we fine-tuned the model previously trained from the MarketScan cohort, using the training

subset of the UMPH hospital EHR data. We incorporated laboratory test results as an additional

Lab modality. We kept the data processing pipeline and model architecture unchanged. As shown

in Fig. 4G-H, the fine-tuning results on UM hold-out testing data drastically improved the model's

performance on UMPH test data. The AUROC reached 0.927 (95% CI, 0.899–0.951) and the

AUPRG reached 0.979 (95% CI, 0.968–0.986) for the 6-month prediction window, whereas the

60-month prediction window achieved an AUROC of 0.835 (95% CI, 0.783–0.879) and an AUPRG

of 0.911 (95% CI, 0.849–0.955).

Top risk predictor for pancreatic cancer

Based on IG attribution analysis, we summarized the top ten risk factors identified across different

prediction windows (Fig.5). In the MarketScan cohort, the model consistently highlighted type 2

diabetes mellitus and other diseases of pancreas (except acute pancreatitis) as the top 1 and top

2 most important risk factors across all prediction windows, respectively. Even though their

absolute scores decrease with the increase of prediction interval, these abnormalities maintained

high importance scores throughout, indicating that they are the most stable and significant signals

preceding pancreatic cancer.[32–34] Abdominal & pelvic pain and acute pancreatitis are additional

top risk factors as well, though their importance decreases as the prediction interval increases.

By 60-month prediction, abdominal & pelvic pain falls below the top 10 risk factors.

In addition, personal or family history of malignant neoplasms repeatedly appear among the top
10 risk factors (Fig. 5), reflecting their importance in the preclinical phase of pancreatic cancer
and potential comorbidity mechanism. In particular, neoplasm in different organs, such as colon
and other GI tract organs, prostate and breast are also identified as important risk factors across
different prediction windows, suggesting possible shared risk pathways or genetic susceptibility
mechanisms with pancreatic cancer. Previous epidemiological studies have confirmed that these
factors indeed elevate the risk of pancreatic cancer, thereby providing strong external evidence
of our attribution result.[35–39]

Discussion

This study leveraged the large-scale MarketScan database to develop a predictive model using
multi-modal EHR information, including diagnoses, procedures and medications. Incorporating
laboratory test data during fine-tuning on UM hospital validation cohort further improved model
performance.

The strengths of this study lie in three areas. First, at the data level, the MarketScan database—
covering a large and diverse population—provided a solid foundation for robust model
development. We also conducted rigorous testing and validation on two regional hospital-based
datasets, UMPH and OneFlorida+, to ensure the comprehensiveness of the study design and the
reliability of the experimental results. Moreover, we show that inclusion of multimodal attributes in
the EHR data can substantially improve the accuracy of cancer risk prediction through the ablation
study. In particular, the inclusion of laboratory test results in hospital datasets significantly
increased sensitivity and precision in the fine-tuned model. It is expected that with the inclusion
of additional data modalities, such as unstructured clinical notes, the model performance will
improve further. Second, at the methodological level, the proposed LLM-based embedding plus

multimodal fusion framework also contributes to superior performance. This is demonstrated by

better performance of PANCDetect in comparison to CancerRiskNet, using the same training,

validation, and testing cohort. Interestingly, our embedding comparison revealed that

representations derived from general LLMs outperformed those trained exclusively on biomedical

text, underscoring the cross-domain transferability and robustness of general-purpose models.

Third, at the study design level, the integration of IG-based interpretability revealed clinically

meaningful drivers of predictions.

Nevertheless, several limitations remain. First, the generalizability of the model requires further

improvement, as performance may decline when applied to external institutions or different

populations. Second, discrepancies in coding systems across healthcare settings have not been

resolved, posing challenges for real-world deployment and cross-system adaptation. Additionally,

using ICD codes as diagnostic criteria is not perfectly accurate, as miscoding or incomplete

documentation may introduce noise into the data. Thirdly, longer exclusion intervals tend to

exclude a much larger number of patients and truncate the available trajectories, resulting in

shorter and potentially less informative input sequences. This increases the difficulty of the

prediction task over a long period of time. Federated learning frameworks could be utilized to

facilitate more powerful long-term risk prediction, while preserving patient privacy, thereby

improving generalizability[40]. Lastly, future efforts should focus on testing PANCDetect in

prospective studies, before being deployed in real-world clinical environments for clinical decision

support.

In conclusion, we present a multimodal disease risk prediction framework based on large-scale

EHR data and applied it to pancreatic cancers, which demonstrated superior performance and

interpretability compared with existing methods. Our model shows promise for potential

applications in high-risk pancreatic cancer screening, early diagnosis, and personalized

interventions. Ultimately, it may improve pancreatic cancer survival outcomes while reducing
healthcare costs.

Data Availability

This study analyzed de-identified EHR and claims data from three sources: the MarketScan
Research databases (licensed from Merative), the University of Michigan Precision Health
(UMPH), and the OneFlorida+ Clinical Research Network. Due to licensing restrictions,
institutional regulations, and patient privacy considerations, these data are not publicly available.
Access to MarketScan requires a data use agreement and paid license from Merative. Access to
UMPH and OneFlorida+ data may be provided to qualified researchers upon reasonable request
to the corresponding author and is subject to approval by the respective institutional review boards
and data custodians.

Acknowledgement

336   We would like to thank Dr. Anirban Maitra for his suggestion and encouragement to pursue this
337   research, Patrick Brady and Institute for HealthCare Policy and Innovation (IHPI) of University of
338   Michigan for helping us to access the MarketScan data. LXG would like to dedicate this work to
339   her father Jianwen Xia, who passed away from pancreatic cancer on 09/28/2024.

Source of Funding

341   LXG is supported by grants from NLM R01 LM012373 and NIH R03 OD039978. ZJ is supported
342   by a summer student grant from the E-HAIL initiative of University of Michigan. XY is supported
343   by NIGMS T32GM141746. QY and RY are supported by the NIH National Center for Advancing
344   Translational Sciences under award number UL1TR001427 and UL1TR000064.

Author Contribution

346 LXG envisioned and supervised this project. ZJ, XG and ZW built and tested the model, conducted

347 the analysis and wrote the manuscript. QY and RY conducted validation analysis using

348 FloridaOne+ cohort. XY and XZ helped with analysis. XY helped to access UMPH data. All

349 authors have read, revised and approved the final manuscript.

Competing Interest

351 The authors declare no conflict of interest.

Reference

1. Al-Shaheri, F. N. *et al.* Blood biomarkers for differential diagnosis and early detection of pancreatic cancer. *Cancer Treat. Rev.* **96**, 102193 (2021).

2. Pereira, S. P. *et al.* Early detection of pancreatic cancer. *Lancet Gastroenterol. Hepatol.* **5**, 698–710 (2020).

3. Huang, J. *et al.* Worldwide Burden of, Risk Factors for, and Trends in Pancreatic Cancer. *Gastroenterology* **160**, 744–754 (2021).

4. Yang, J. *et al.* Early screening and diagnosis strategies of pancreatic cancer: a comprehensive review. *Cancer Commun.* **41**, 1257–1274 (2021).

5. Søreide, K., Ismail, W., Roalsø, M., Ghotbi, J. & Zaharia, C. Early Diagnosis of Pancreatic Cancer: Clinical Premonitions, Timely Precursor Detection and Increased Curative-Intent Surgery. *Cancer Control* **30**, 10732748231154711 (2023).

6. Elbanna, K. Y., Jang, H.-J. & Kim, T. K. Imaging diagnosis and staging of pancreatic ductal adenocarcinoma: a comprehensive review. *Insights Imaging* **11**, 58 (2020).

7. Wittram, R., Kreis, L., König, H.-H. & Brettschneider, C. Economic evaluations of early detection strategies for pancreatic cancer: a systematic review. *Eur. J. Health Econ.* (2025).

8. Mishra, A. K., Chong, B., Arunachalam, S. P., Oberg, A. L. & Majumder, S. Machine Learning Models for Pancreatic Cancer Risk Prediction Using Electronic Health Record Data—A Systematic Review and Assessment. *Am. J. Gastroenterol.* **119**, 1466–1482 (2024).

9. Guo, L. L. *et al.* A multi-center study on the adaptability of a shared foundation model for electronic health records. *Npj Digit. Med.* **7**, 171 (2024).

10. Xian, S. *et al.* Transformer patient embedding using electronic health records enables patient stratification and progression analysis. *Npj Digit. Med.* **8**, 521 (2025).

11. Zhang, S., Cornet, R. & Benis, N. Cross-Standard Health Data Harmonization using

Semantics of Data Elements. *Sci. Data* **11**, 1407 (2024).

12. Li, Y. *et al.* BEHRT: Transformer for Electronic Health Records. *Sci. Rep.* **10**, 7155

(2020).

13. Li, Y. *et al.* Hi-BEHRT: Hierarchical Transformer-Based Model for Accurate Prediction of

Clinical Events Using Multimodal Longitudinal Electronic Health Records. *IEEE J. Biomed.*

Health Inform. **27**, 1106–1117 (2023).

14. Rasmy, L., Xiang, Y., Xie, Z., Tao, C. & Zhi, D. Med-BERT: pretrained contextualized

embeddings on large-scale structured electronic health records for disease prediction. *Npj*

Digit. Med. **4**, 86 (2021).

15. Su, X. *et al.* Multimodal Medical Code Tokenizer. Preprint at

https://doi.org/10.48550/ARXIV.2502.04397 (2025).

16. Lee, S. A., Wu, A. & Chiang, J. N. Clinical ModernBERT: An efficient and long context

encoder for biomedical text. Preprint at https://doi.org/10.48550/ARXIV.2504.03964 (2025).

17. Luo, R. *et al.* BioGPT: Generative Pre-trained Transformer for Biomedical Text

Generation and Mining. *Brief. Bioinform.* **23**, bbac409 (2022).

18. Jiang, L. Y. *et al.* Health system-scale language models are all-purpose prediction

engines. *Nature* **619**, 357–362 (2023).

19. Jia, K. *et al.* A pancreatic cancer risk prediction model (Prism) developed and validated

on large-scale US clinical data. *eBioMedicine* **98**, 104888 (2023).

20. Appelbaum, L. *et al.* Development and validation of a pancreatic cancer risk model for

the general population using electronic health records: An observational study. *Eur. J.*

Cancer **143**, 19–30 (2021).

21. Placido, D. *et al.* A deep learning algorithm to predict risk of pancreatic cancer from

disease trajectories. *Nat. Med.* **29**, 1113–1122 (2023).

22. Park, J. *et al.* Enhancing EHR-based pancreatic cancer prediction with LLM-derived

embeddings. *Npj Digit. Med.* **8**, 465 (2025).

23. Butler, A. M., Nickel, K. B., Overman, R. A. & Brookhart, M. A. IBM MarketScan

Research Databases. in *Databases for Pharmacoepidemiological Research* (eds

Sturkenboom, M. & Schink, T.) 243–251 (Springer International Publishing, Cham, 2021).

doi:10.1007/978-3-030-51455-6_20.

24. Zawistowski, M. *et al.* The Michigan Genomics Initiative: A biobank linking genotypes

and electronic clinical records in Michigan Medicine patients. *Cell Genomics* **3**, 100257

(2023).

25. Shenkman, E. *et al.* OneFlorida Clinical Research Consortium: Linking a Clinical and

Translational Science Institute With a Community-Based Distributive Medical Education

Model. *Acad. Med.* **93**, 451–455 (2018).

26. Flach, P. & Kull, M. Precision-Recall-Gain Curves: PR Analysis Done Right. *In Proc.*

28th Conference on Neural Information Processing Systems (NeurIPS) (2015).

27. Sundararajan, M., Taly, A. & Yan, Q. Axiomatic Attribution for Deep Networks. *In Proc.*

34th International Conference on Machine Learning (ICML) (2017).

28. Yang, X. *et al.* A Large Clinical Language Model to Unlock Patient Information from

Unstructured Electronic Health Records. *Npj Digit. Med.* **5**, 194 (2022).

29. Excoffier, J.-B. *et al.* Generalist embedding models are better at short-context clinical

semantic search than specialized embedding models. Preprint at

https://doi.org/10.48550/ARXIV.2401.01943 (2024).

30. Hegselmann, S. *et al.* Large Language Models are Powerful Electronic Health Record

Encoders. Preprint at https://doi.org/10.48550/ARXIV.2502.17403 (2025).

31. Laws, M. B., Michaud, J., Shield, R., McQuade, W. & Wilson, I. B. Comparison of

Electronic Health Record–Based and Claims-Based Diabetes Care Quality Measures:

Causes of Discrepancies. *Health Serv. Res.* **53**, 2988–3006 (2018).

32. Cho, J., Scragg, R. & Petrov, M. S. Postpancreatitis Diabetes Confers Higher Risk for

Pancreatic Cancer Than Type 2 Diabetes: Results From a Nationwide Cancer Registry.

Diabetes Care **43**, 2106–2112 (2020).

33. Huxley, R., Ansary-Moghaddam, A., Berrington de González, A., Barzi, F. & Woodward,

M. Type-II diabetes and pancreatic cancer: a meta-analysis of 36 studies. *Br. J. Cancer* **92**,

2076–2083 (2005).

34. Lai, S.-W. & Liao, K.-F. Pancreatitis, Type 2 Diabetes Mellitus, and Pancreatic Cancer

(Diabetes Metab J 2025;49:252-63). *Diabetes Metab. J.* **49**, 518–519 (2025).

35. Price, S. J., Gibson, N., Hamilton, W. T., King, A. & Shephard, E. A. Intra-abdominal

cancer risk with abdominal pain: a prospective cohort primary care study. *Br. J. Gen. Pract.*

J. R. Coll. Gen. Pract. **72**, e361–e368 (2022).

36. Matsubayashi, H. *et al.* Surveillance of Individuals with a Family History of Pancreatic

Cancer and Inherited Cancer Syndromes: A Strategy for Detecting Early Pancreatic

Cancers. *Diagn. Basel Switz.* **9**, 169 (2019).

37. Seppälä, T. T., Burkhart, R. A. & Katona, B. W. Hereditary colorectal, gastric, and

pancreatic cancer: comprehensive review. *BJS Open* **7**, zrad023 (2023).

38. Mocci, E. *et al.* Risk of pancreatic cancer in breast cancer families from the breast

cancer family registry. *Cancer Epidemiol. Biomark. Prev. Publ. Am. Assoc. Cancer Res.*

Cosponsored Am. Soc. Prev. Oncol. **22**, 803–811 (2013).

39. Jacobs, E. J. *et al.* Family history of cancer and risk of pancreatic cancer: a pooled

analysis from the Pancreatic Cancer Cohort Consortium (PanScan). *Int. J. Cancer* **127**,

1421–1428 (2010).

40. Sadilek, A. *et al.* Privacy-first health research with federated learning. *Npj Digit. Med.* **4**,

132 (2021).

Figure Legend

Fig 1. Overall workflow of this study.

(1) Cohort selection: The MarketScan cohort was used for model training, while two independent cohorts (UMPH and OneFlorida+) served as external validation datasets. (2) Feature Engineering: During model training, diagnoses, procedures, and medications were extracted as inputs. Standardized textual descriptions of these medical codes were encoded into embeddings using a pretrained LLM. (3) Model Building and Validation: The generated embeddings were passed into a Transformer-based model, with the best model selected based on minimized validation loss. Evaluation was conducted on the held-out test set and external validation cohorts. Attribution analysis using Integrated Gradients (IG) was applied to identify key risk factors. (4) Model Refinement: Fine-tuning was further performed on the UMPH cohort to include additional laboratory tests into the pretrained model using MarketScan data. UMPH: University of Michigan Precision Health; PANC: pancreatic cancer.

Fig 2, Model structure of PANCDetect and single-interval prediction design.

Panel A shows the overall model structure of PANCDetect. Firstly, time-varying EHR events (diagnoses, procedures, medications) are encoded with LLM-based embeddings and temporal markers to represent longitudinal patient trajectories and then processed by a Transformer with masked self- and cross-attention. The resulting sequence representation is pooled into a patient vector and late-fused by concatenation with time-invariant attributes (e.g., sex, geographic location); a final MLP maps the fused vector to a continuous pancreatic cancer risk score. Panel B illustrates the single-interval prediction design, where EHR events within N months before cancer diagnosis are masked to reduce data leakage, and trajectories are truncated according to predefined prediction windows (6, 12, 24, 36, and 60 months). LLM: large language model; MLP: multi-layer perceptron; PANC: pancreatic cancer.

Fig 3. Characteristics of the MarketScan, UMPH and OneFlorida+ cohorts.

Panels A,C, and E show the age-sex distribution of pancreatic cancer patients on MarketScan (Panel A), UMPH (Panel C) and OneFlorida+ (Panel E) cohorts, respectively. Panels B,D, and F show heatmaps of the joint distribution of diagnosis trajectory length and the number of distinct disease codes for PANC and non-PANC patients in the MarketScan (Panel B), UMPH (Panel D), and OneFlorida+ (Panel F) cohorts. Color intensity reflects the number of patients in each bin of age group, with darker shades indicating larger patient counts. UMPH: University of Michigan Precision Health; PANC: pancreatic cancer.

Fig 4. Model performance across MarketScan, UMPH, and OneFlorida+ cohorts.

Panels A and B show results of the model trained on MarketScan training set tested on MarketScan hold-out test data; Panels C and D show OneFlorida+ external validation results using the model trained on MarketScan training data; Panels E and F show UMPH external validation results on the model trained on MarketScan training data; Panels G and H show fine-tuning results on UMPH hold-out data, after adding laboratory test data to the model built on MarketScan data. Panels A,C,E, and G: Results in AUROC; Panels B,D,F, and H: Results in AUPRG. Shaded areas represent 95% confidence intervals. AUROC: area under the receiver operating characteristic curve; AUPRG: area under the precision-recall-gain curve; UMPH: University of Michigan Precision Health; PANC: pancreatic cancer.

Fig. 5. Attribution analysis of top EHR risk predictors for pancreatic cancer.

Top 10 features from the model trained on MarketScan data. The top predictive clinical features across different prediction windows are ranked by the IG attribution scores.

496 **Tables**

General dataset information	MarketScan	UMPH	OneFlorida+
Dataset timeline	2009-2022	2006-2025	2013-2024
Total n patients	680,821	48,318	54,853
Male (%)	346,084(50.8%)	25,751(53.3%)	27,596 (50.3%)
Female (%)	334,737(49.2%)	22,567(46.7%)	27,257 (49.7%)
Median n disease codes per patient	101	298	77
Median length of trajectory in years	3	5	2
Pancreatic cancer cohort information			
Total n patients	113,518	8,257	10,500
Male (%)	57,713(50.8%)	4,406(53.4%)	5,537 (52.7%)
Female (%)	55,805(49.2%)	3,851(46.6%)	4,963 (47.3%)
Median age at PC diagnosis	62	67	67
Median n disease codes per patient	242	193	96
Median length of trajectory in years	3	1	1
Smoker (%)	24,965(22.0%)	891(10.8%)	4,319 (41.1%)
Alcohol user (%)	5,520 (4.9%)	239(2.9%)	693 (6.6%)
Illegal drug user (%)	3,819 (3.4%)	328(4.0%)	712 (6.8%)

497 **Table 1**: Summary of patient statistics in the three datasets of this study

| Cohort Composition | Feature Engineering | Model Building | Model Refinement |

Cohort Composition

MarketScan (2009-2022) PANC patients (n = 113,518)

Training Cohort

UMPH (2006-2025) PANC patients (n = 8,257)

OneFlorida+ (2013-2024) PANC patients (n = 10,500)

External Validation Cohort

Feature Engineering

DIAGNOSIS ICD CODES — Diagnosis

Demographics

Rx — Drug

PROCEDURE CPT CODES — Procedure

LLM

LLM embeddings

Model Building

Model training Minimized loss

Train Valid

Validation & Interpretation

AUROC AUPRG

Validation results Risk factor discovery

Model Refinement

On UMPH cohort

Add lab test features

Finetuning

A Overall model architecture of PANCDetect

B Timepoints and prediction intervals

Test results on MarketScan holdout test set

Validation results on OneFlorida+ cohort

Validation results on UMPH cohort

Finetuing results on UMPH holdout test set

Feature attribution results

6-month Prediction	Score	12-month Prediction	Score	24-month Prediction	Score	36-month Prediction	Score	60-month Prediction	Score
Type 2 diabetes mellitus	2.070	Type 2 diabetes mellitus	1.118	Type 2 diabetes mellitus	0.777	Type 2 diabetes mellitus	0.358	Type 2 diabetes mellitus	0.259
Other diseases of pancreas	1.169	Other diseases of pancreas	0.749	Other diseases of pancreas	0.395	Other diseases of pancreas	0.183	Other diseases of pancreas	0.081
Abdominal and pelvic pain	0.773	Abdominal and pelvic pain	0.214	Benign neoplasm of colon, rectum, anus and anal canal	0.230	Benign neoplasm of colon, rectum, anus and anal canal	0.087	Benign neoplasm of colon, rectum, anus and anal canal	0.041
Acute pancreatitis	0.444	Acute pancreatitis	0.207	Abdominal and pelvic pain	0.207	Family history of malignant neoplasm	0.082	Family history of malignant neoplasm	0.034
Malignant neoplasm of prostate	0.277	Malignant neoplasm of prostate	0.157	Personal history of malignant neoplasm	0.182	Abdominal and pelvic pain	0.070	Personal history of malignant neoplasm	0.027
Benign neoplasm of colon, rectum, anus and anal canal	0.258	Personal history of malignant neoplasm	0.138	Acute pancreatitis	0.163	Acute pancreatitis	0.063	Malignant neoplasm of prostate	0.026
Malignant neoplasm of colon	0.216	Benign neoplasm of other digestive organs	0.106	Malignant neoplasm of prostate	0.158	Personal history of malignant neoplasm	0.061	Malignant neoplasm of kidney, except renal pelvis	0.024
Personal history of malignant neoplasm	0.198	Malignant neoplasm of colon	0.101	Malignant neoplasm of breast	0.098	Malignant neoplasm of breast	0.052	Acute pancreatitis	0.023
Benign neoplasm of other digestive organs	0.178	Genetic susceptibility to malignant neoplasm	0.100	Malignant neoplasm of colon	0.090	Malignant neoplasm of prostate	0.044	Other disorders of kidney and ureter	0.022
Other diseases of biliary tract	0.153	Family history of malignant neoplasm	0.098	Family history of malignant neoplasm	0.087	Benign neoplasm of other digestive organs	0.040	Malignant neoplasm of breast	0.019

ICD-10 chapters

II Neoplasms

IV Endocrine, nutritional and metabolic diseases

XI Diseases of the digestive system

XIV Diseases of the genitourinary system

XVIII Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified

XXI Factors influencing health status and contact with health services