TITLE

Incorporating Dietary Information to Enhance Polygenic Prediction Models with Applications to Body Mass Index and Type 2 Diabetes

AUTHORS

Eunice Y. Lee[1], Bryan L. Dinh[1,2], Ji Tang[1], Samantha Streicher[3], Subarna Biswas[1], He Tian[1], Gertraud Maskarinec[3], Lynne R. Wilkens[3], Song-Yi Park[3], Charleston W.K. Chiang[1,2]*

AFFILIATIONS

1. Center for Genetic Epidemiology, Department of Population and Public Health Sciences, University of Southern California, California

2. Department of Quantitative and Computational Biology, University of Southern California, California

3. Epidemiology Program, Cancer Center, University of Hawai□i, Honolulu

CORRESPONDING AUTHOR

Charleston W.K. Chiang*

ABSTRACT

Polygenic predictors can enhance screening for biomedical conditions, such as metabolism-related traits and diseases, but explain limited phenotypic variance and face implementation challenges in non-European populations. On the other hand, dietary quality and other sociocultural factors are well established metabolic risk factors that remain under-investigated in risk stratification models. In this study, we developed and evaluated risk stratification model combining polygenic predictors and diet-based models for body mass index (BMI) and type 2 diabetes (T2D). Using 5,368 Native Hawaiians from the Multiethnic Cohort (MEC-NH) with genetic data, we integrated large-scale cross-ancestry GWAS summary statistics to develop polygenic score (PGS) models with better prediction accuracies (partial-R^2 [SE] = 0.12 [0.04] for BMI; liability-R^2 [SE] = 0.09 [0.04] for T2D) than using GWAS information from single ancestry (partial-R^2 = 0.07-0.09 for BMI and liability-R^2 = 0.05-0.07 for T2D) or in combination with GWAS from MEC-NH (partial-R^2 = 0.05-0.06 for BMI and liability-R^2 = 0.01-0.05 for T2D). Moreover, machine learning models trained on 520 dietary variables available from 14,346 MEC-NH individuals substantially explained BMI variation (partial-R^2 [SE] = 0.12 [0.01]) and enhanced prediction when combined with PGS (adjusted-R^2 = 0.29). The best performing diet score model for BMI was associated with multiple chronic diseases in the same cohort, potentially mediated via inflammatory and lipid pathways. Both PGS and dietary scores provided significant, complementary information for predicting BMI and T2D. For populations with limited genetic studies like Native Hawaiians, integrating external GWAS data or non-genetic dietary information can improve risk stratification models.

Word count: 248 (excluding headers)

INTRODUCTION

Metabolic syndrome (MetS), also known as insulin resistance syndrome, is defined by the presence of three or more co-occurring conditions: abdominal obesity, hypertension, hyperglycemia (1-2). These conditions increase the risk of developing cardiovascular disease (CVD), nephropathy, and type 2 diabetes (T2D), potentially leading to premature death (3-5). Currently, nearly 2.5 billion adults (43% of the world's population) are overweight or obese (6). According to the CDC, obesity and T2D prevalence in both adults and children in the United States continues to increase and is projected to rise further (7-8). Early identification of risk factors can help prevent future incidence and reduce disease prevalence.

Genetic factors influence individuals' responses to environmental stressors, causing them to react differently to the same health challenges. The within-family heritability estimates, which are relatively free of bias due to shared environment, are 0.43 for T2D and 0.49 for BMI across major ethnic groups in the United States (9). Hundreds to thousands of genetic variants discovered in recent large-scale genome-wide association studies of metabolic phenotypes across diverse populations demonstrate that these traits are polygenic. Rather than focusing on single genes or small gene sets, aggregating cumulative effects across the genome provides better risk prediction and stratification. Furthermore, polygenic scores (PGS) offer additional benefits, as adding PGS as covariates significantly increases disease prediction accuracy alongside other clinical factors (10-12). However, PGS face transferability issues across populations because their performance depends heavily on the underlying GWAS, which have been predominantly conducted in European-ancestry individuals (13-14). This results in low prediction accuracy for non-European populations, hindering clinical implementation and presenting an ongoing challenge (15-16).

Besides genetic contributions, MetS results from multiple environmental factors, with diet being one of the primary contributors. Research highlights that chronic low-grade inflammation, insulin resistance, and adipose tissue dysfunction are key pathogenic mechanisms in MetS. The intake of specific food components plays an important role in MetS development, prevention, and treatment (17-19). Unhealthy dietary habits such as consuming excess energy, sugar, saturated fats, and salt are major sources of low-density lipoprotein accumulation, promoting inflammation and altering glucose metabolism, thereby increasing diabetes risk and atherosclerotic plaque formation that can lead to cardiovascular conditions. A meta-analysis of 50 prospective and randomized controlled studies demonstrated that Mediterranean diet adherence reduced MetS

103 incidence by 50%, indicating that healthy dietary patterns such as Mediterranean, DASH,
104 vegetarian, or high-Healthy Eating Index diets can improve immune and cardiometabolic
105 function while reducing chronic disease risk (20-22).
106
Similar to PGS, rather than examining individual food items or nutrients, dietary quality indices
have been developed to assess overall dietary effects on chronic disease risk. However, these
dietary indices are typically based on specific dietary recommendations and therefore do not
capture the entirety of a person's food choices and eating patterns, which are interactive and
synergistic. Developing quantitative methods to detect eating behaviors and patterns that
comprehensively reflect a person's overall diet remains a major challenge. Moreover, these
dietary indices usually tend to be developed to capture the overall health and well-being, rather
than focusing on any specific disease or conditions. Therefore, individual dietary indices are
likely to show weaker correlations with specific clinical outcomes.
116
In this study, we explored the role of genetic and non-genetic risk factors in MetS-related
phenotypes including body mass index (BMI) and T2D by developing polygenic risk and diet
scores in Native Hawaiians, a population with high prevalence of obesity and T2D (23-24).
Moreover, there are limited available genetic data in this population, making polygenic
predictions more challenging; non-genetic information such as dietary information is thus more
pertinent in risk stratification in a clinical setting. For PGS development, we applied two recently
developed Bayesian methods that integrate GWAS summary statistics from multiple
populations, which have been reported to outperform existing PGS construction methods and
improve cross-population polygenic prediction. For our diet score, we applied a data-driven
approach to derive dietary patterns from all available food and nutrition consumption data using
three different machine learning algorithms. Unlike traditional methods such as factor and
cluster analysis, our data-driven approach considers both linear and non-linear relationships
without making arbitrary decisions about factor or cluster structure. Our hypothesis-free method,
requiring no prior knowledge, represents the totality of foods and drinks that individuals
consume regularly and captures the synergistic effects of foods eaten together that may affect
MetS outcomes. Our study has significant implications for disease prevention and precision
medicine.
134
135 **METHODS**

136 Overview of the methods, input data, and phenotypes for this study are described in
137 **Supplement Figure S1**. The institutional review board of the University of Southern California
138 approved the study protocol. All participants signed an informed consent form.
139
Study Subjects: Multiethnic Cohort
The Multiethnic Cohort (MEC) is a large-scale cohort designed to study diet and cancer among
ethnic minorities in the United States. Study participants comprise 215,251 adult men and
women aged 45 to 75 years at baseline who resided in Hawaii and California (primarily Los
Angeles County). The cohort includes five racial and ethnic groups: Japanese Americans
(28.5%), whites (24.8%), Latinos/Hispanic (22.7%), African Americans (16.8%), and Native
Hawaiians (7.2%). Recruitment occurred from 1993 to 1996. Each participant completed a self-
administered 26-page questionnaire that was mailed to them. A main portion of the
questionnaire was a quantitative food frequency questionnaire (QFFQ) with more than 180 food
items to assess habitual diet during the previous year (25). For each food item, the QFFQ
queried consumption frequency using eight or nine categories ("neve or hardly ever" to "≥2
times/day" or "≥4 times/day") and portion size (three serving size choices). A calibration study
demonstrated a good validity of the QFFQ against three 24-hour dietary recalls based on
correlations in nutrient estimates (26). Beyond dietary information, the questionnaire also
captured demographic factors, personal behaviors (smoking, solar exposure, physical activity,
sleep), occupation, medical conditions, medication use, reproductive history for women, as well
as family cancer history. Genetic data were collected from a subset of volunteers who are
slightly younger, less likely to smoke, and slightly more educated, but otherwise similar to the
subset who did not donate biospecimen for genetics (**Table 1**). More information regarding the
study design, questionnaire, and exclusion and inclusion criteria can be found elsewhere (25-
27).
161
The labels and groupings of the MEC individuals are based on self-report at baseline using
terminology that was in practice in the early 1990s when the surveys were distributed to
participants. In the survey, participants were asked to provide self-reported ethnic or racial
background, marking all that applies, with the options of "Black or African-American", "Chinese",
"Filipino", "Hawaiian", "Japanese (includes Okinawan)", "Korean", "Mexican or other Hispanic",
"White or Caucasian", or "Other". Most MEC analyses categorize participants into one of the five
major racial/ethnic groups mentioned above that were targeted during recruitment, prioritizing
group memberships in the following order: "African-American", "Hawaiian", "Latino", "Japanese",

"White". Thus, if an individual reports "Chinese" and "Hawaiian" in the survey, they would be
classified as Native Hawaiian for these analyses. Among the 0.3% of individuals where no
response was given for the participant, but existed for the mother and father, a race/ethnicity
was created from the parent's information; otherwise the individual is labeled as "Other."
While the MEC subcohorts represent multiple levels of self-reported ethnicities, nationality, and
genetic ancestries, we continue to use these labels for consistency with other MEC studies and
for ease of recognizing and interpreting the historical experience of health disparity for some of
these populations, including potential benefits in genomic health through polygenic predictions.
We recognize these labels are imperfect proxies for genetic ancestry, but note that our study
focuses on a cohort of individuals from a community that emphasize self-identity or genealogical
record for group membership (in place of estimated genetic ancestry), who have experienced
significant disparity due to social determinants of health, who feel strongly of this identity, and
who encompass multiple ancestries on a continuum that would be difficult and unethical to
delineate or recategorize.
184
185 ***Definition of Metabolic Health Phenotypes:***
186
187 *Obesity/BMI*
188
189 BMI was calculated using height and weight reported on the questionnaire at the entry of cohort
190 for each participant. The raw BMI was first stratified by sex and regressed against the age. The
191 residuals of the sex stratified regression models were inverse normalized and combined across
192 sexes for the analyses. This residualized and inverse-normalized BMI is used throughout the
193 study, and for brevity we simply refer to it as BMI.
194
195 *Type 2 Diabetes*
196
T2D status was ascertained from a few major sources of information for the MEC. First, self-
reports were available from 6 questionnaires, in response to the question "Has your doctor ever
told you that you had diabetes?" Second, as part of the biorepository, cohort members who
reported T2D medication at time of blood draw were classified as cases. Third, a one-time
linkage to Hawai'i insurance databases as of Jan. 1, 2008, also indicated T2D cases for
participants recruited in Hawai'i. Finally, a linkage to Medicare Chronic Condition Warehouse for
diabetes for Medicare claims between the years 1999-2019 (the latest linkage). Individuals

indicated on any of these sources are considered a case, with the remaining individuals
considered as controls. Our only information of prevalent diabetes is based on self-reports on
the baseline questionnaire, and thus cases who did not report diabetes in the baseline
questionnaire is considered incident T2D. For Native Hawaiians, we identified 6566 total
individuals with T2D (2561 with genetic and dietary data, 4005 with dietary data only). Of these,
4380 are defined as incident T2D (1947 with genetic and dietary data, 2799 with dietary data
only). 8984 are defined as controls (2433 with genetic and dietary data, 4979 with dietary data
only).

Genotype Data Generation, Processing, and Quality Control
In total, 5,388 self-identified Native Hawaiians from the MEC were genotyped on two separate
GWAS arrays: Illumina MEGA (n = 4144) and Illumina Global Diversity Array (GDA; n = 1238).
Each dataset was processed through the same quality control pipeline: we removed variants
that were genotyped in fewer than 95% of people in the sample, as well as variants out of
Hardy-Weinberg equilibrium ($p < 10^{-6}$). We also applied a filter for people with more than 2%
missing genotypes but removed no one with this filter. The datasets were then lifted to hg38
using triple-LiftOver (v.1.33) (28) to ensure that alleles in inverted sequences between reference
genome builds were properly lifted. We phased the genotypes with EAGLE (v.2.4.1) (29) using
a Native-Hawaiian specific genetic map (30), and then imputed using the TOPMed imputation
server against the TOPMed r3 (31).

Global ancestry estimates for the MEC-NH cohort were previously calculated (32) using
ADMIXTURE (v.1.3.0) in unsupervised mode, in combination with other MEC ethnic groups and
the 1000 Genome Project reference populations. We found MEC-NH individuals at k = 4
exhibited known components of European (EUR)-like, East-Asian (EAS)-like, and African
(AFR)-like ancestries (interpreted based on 1000 Genomes reference samples), as well as a
component of ancestry that is unique to the MEC-NH, presumed to be Polynesian (PNS)-like in
origin.

We performed GWAS for BMI and T2D (prevalent + incident T2D cases) based on the subset of
4144 individuals genotyped on the MEGA array, leaving out the 1238 individuals genotyped on
the GDA array as validation dataset. Specifically, we implemented the UW-GAC TOPMed
GENESIS analysis pipeline (https://github.com/UW-GAC/analysis_pipeline) and inferred a
Genetic Relationship Matrix (GRM) and Principal Components (PCs) using array data for MEC-

NH. In brief, this pipeline filtered variants (MAF < 1%, missingness > 1%) before LD pruning (R^2
> 0.1, 10Mb window). Initial kinship and population divergence estimates are inferred with
KING(33). These estimates were used as input for PC-AiR to infer PCs that in turn served as
input into PC-Relate to infer a genetic-relatedness matrix (34-35). An second iteration of PC-AiR
and PC-Relate was run with the PC-Relate GRM as the input. Association testing is done with
the GENESIS package (36). The null model is fit using the phenotype, GRM, and 20 PCs.
Association testing is done with imputed dosages.

Trans-ethnic Polygenic Score (PGS)
The PGS were developed for each phenotype in MEC-NH participants applying two Bayesian
approaches, namely PRS-CSx and BridgePRS, which are described below (37-38). For the
development of BMI PGS, GWAS summary statistics conducted in individuals from the Taiwan
Biobank and Biobank Japan (n=256,450) representing EAS-like ancestry, Genetic Investigation
of Anthropometric Traits (GIANT) consortium and U.K. BioBank (n=~700,000) representing
EUR-like ancestry, and Native Hawaiians in MEC study (n=4,156 genotyped on MEGA array
only) representing largely PNS-like ancestry (**Supplement Figure S2)** were used. For the
development of T2D PGS, we used ancestry-specific GWAS summary statistics from EAS-like
populations (n=427,504) and EUR-like ancestry populations (n=1,812,017) as released by prior
trans-ancestry GWAS efforts for T2D (39). We conducted GWAS for T2D in 4,164 MEC-NH in-
house.

Pairs of these GWAS summary statistics were integrated and trained in both PRS-CSx and
BridgePRS models separately. PGS were calculated based on the resulting model for validation
in held-out MEC-NH (those genotyped on GDA array) for each phenotype (n_{BMI} = 1226; n_{T2D} =
1218); scores were standardized to have mean = 0 and standard deviation = 1. Following
previous study (32), we measured performance of a PGS model based on the partial R^2 (for
continuous outcome, calculated by R package rsq v2.5) or the Nagelkerke pseudo-R^2
(calculated by R package DescTools v0.99.43) transformed to the liability scale assuming an
estimated lifetime risk of 60% (40). Depending on the relative performance between PRS-CSx
and BridgePRS models, the better-performing PGS by partial R2 was then used in the
downstream analyses for the respective phenotype.

PRScsx

PRS-CSx utilizes a high-dimensional Bayesian regression framework to perform cross-
population polygenic prediction by integrating GWAS summary statistics from multiple
populations (37). For effect size estimation, information is shared between summary statistics
while considering genetic diversity across discovery samples through population-specific allele
frequencies and LD patterns. PRS-CSx applies a shared continuous shrinkage prior, which
allows varying degrees of shrinkage to be applied to SNP effect sizes based on the strength of
their GWAS associations.

First, summary statistics and ancestry-matched LD reference panels were used to estimate
population-specific posterior SNP effect sizes using PRS-CSx in auto mode, with default
Strawderman-Berger prior parameters (a=1, b=0.5) for local shrinkage and the global shrinkage
parameter ($\square$) learned from data using a fully Bayesian approach. The population-specific
posterior effect estimates were meta-analyzed using inverse-variance weighting within the
Gibbs sampler to produce combined SNP effects. The polygenic score for each study subject
was generated using PLINK commands (--score and sum modifier that combines chromosome-
specific scores into genome-wide scores).

We used pre-compiled 1000 Genomes reference panels provided by PRS-CSx for EAS-like and
EUR-like ancestry summary statistics (~1.2 million SNPs across multiple populations), and
constructed a Native Hawaiian reference panel that conforms to the pre-compiled reference
panel format. For the Native Hawaiian reference panel construction, we extracted the imputed
genotypes at HapMap3 variants from a randomly chosen subset of 500 Native Hawaiian
samples, restricted to variants with imputation quality filtering ($R^2 > 0.8$), MAF filtering (>1%),
and converted the genome build to hg19 to be consistent with the pre-compiled 1000 Genomes
references. We computed LD correlation matrices using PLINK and organized them into LD
blocks using Asian population boundaries derived from population-specific LD breakpoints
estimated by LDetect (https://bitbucket.org/nygcresearch/ldetect-data/src/master/). The final
reference panel includes population-specific allele frequencies and strand-flipping indicators
integrated into the standard PRS-CSx multi-population framework, with LD matrices stored in
compressed HDF5 format for computational efficiency.

BridgePRS
While PRS-CSx constructs PGS using continuous shrinkage priors applied at individual causal
SNPs, BridgePRS operates at the locus level, combining SNP effects within genomic loci
through Bayesian posterior mean estimation rather than simple averaging (38). SNP effects

within each locus are modeled by a multivariate Gaussian distribution, where a locus is defined
as variants with R² > 0.01 and within 1Mb of a lead variant (the SNP with the lowest p-value in
that genomic region). We constructed three different trans-ancestry models for each phenotype:
1) EAS-like population as discovery and Native Hawaiian as target, 2) EUR-like population as
discovery and Native Hawaiian as target, and 3) EAS-like population as discovery and EUR-like
population as target. Single-ancestry or population-specific PRS models were trained and
optimized using default parameters (α = 0.5 for allele frequency dependency, $\lambda^{(□)}$ = 0.5 for
baseline shrinkage), with zero-centered Gaussian prior distributions assigned to SNP effects at
each locus. In the two-stage cross-ancestry models, posterior SNP effect sizes estimated in the
discovery populations were used as informative priors for the target population (τ = 100 for
cross-ancestry shrinkage). The final PRS combined all single-ancestry and two-stage polygenic
scores using optimal weights determined through cross-validation ridge regression in the
validation dataset (n_BMI=1,226; n_T2D=1,218). For EAS-like and EUR-like populations, we
constructed LD reference based on the EAS and EUR populations from 1000 Genomes,
respectively, using variants with MAF > 5%. For Native Hawaiians, we constructed the LD
reference based on 500 randomly selected individuals from the MEC-NH (the same set of
individuals used above to construct the reference panel for PRS-CSx models).

Dietary Variables and Phenotype Specific Diet Scores

Based on the QFFQ responses, participants' food and nutrient intakes were computed using
comprehensive food composition database developed and maintained for the MEC, including a
large recipe database (25). As part of the Dietary Patterns Methods Project that the MEC and
two other large US cohort studies participated, four commonly used indices (HEI, AHEI, aMED,
DASH) for overall diet quality were calculated using a standardized methodology across the
three cohorts (41). Later, 27 more dietary indices were additionally calculated. MEC dietary data
includes nutrients (absolute intake), individual foods (grams), food groups (grams), MyPyramid
Equivalents Database (MPED), and dietary patterns (component scores and total scores), which
resulted in a total of 520 diet variables that were used to train and predict diet scores in MEC-
NH subjects (n=13,595). For the phenotype-specific diet score, we applied an elastic net
regularization, random forest, and neural network machine learning algorithms to prioritize diet
variables or build regression models with dietary variables, from which we can compute a diet
score that was then standardized within cohort to have a mean of 0 and standard deviation of 1.
Across all diet score prediction models in Native Hawaiians, subjects without genetics data
served as the training set and subjects with genetics data served as the validation set.

For the development and selection of the best-performing diet score, we applied three different machine learning algorithms. First, an elastic net regularization was implemented with a two-step nested cross-validation approach (5-fold cross validation) that identified the optimal L2 penalty parameter by evaluating cross-validation error across the lambda2 grid which ranged from 0.00001 to 0.01 with increments of 0.001, then the corresponding optimal L1 parameter for the selected L2 value was determined. After training the elastic net model on BMI (n=8,372), a single composite score was calculated for each participant in the validation dataset (n=5,223) based on the learned coefficients of the selected dietary variables. For the prediction of the diabetes outcome, a cross-validated elastic net logistic regression model was developed using the optimal lambda value identified through cross-validation, keeping the alpha fixed at 0.3 (blend of L1 and L2 penalties: 30% Lasso penalty for variable selection and 70% Ridge penalty for handling correlated predictors) in the training dataset (n= 7,026; 2,294 cases, 4,732 controls). Individual dietary scores were computed based on the elastic net model coefficients for the validation dataset (n= 4,632). We used the 'gcdnet' (v.1.0.6) in R for the elastic net models

Second, a random forest regression model was trained to predict BMI using all dietary variables in the training dataset, configured with 500 decision trees. The trained random forest model was then applied to predict BMI values for each subject in the validation dataset. For diabetes status, a random forest classification model was trained to predict incident T2D status using all dietary variables in the training dataset. We applied the trained random forest model to the validation dataset to predict the probability of each individual having diabetes rather than providing binary classifications, and used the predicted probabilities as diabetes risk scores. For the random forest models, we used randomForest (v.4.7-1.2) in R.

Third, we applied a neural network diet score model. Multilayer perceptrons (MLPs) possess sufficient representational capacity to model the complex nonlinear relationships in terms of the Universal Approximation Theorem (42), and it is a common practice to use MLPs to fit a classifier and regressor in the machine learning field. The predictions of T2D and BMI based on dietary variables are technically a classification task and a regression-fitting task, respectively. Therefore, we employed MLPs to build neural network models for predicting T2D and BMI. Since the training is fast (one epoch takes seconds), we used grid-search to find the optimal values of the hyperparameters. Specifically, we defined multiple candidate values for the

hyperparameters (**Supplement Table S5**); the combination of these candidate values results in
more than 1,500 candidate models for predicting BMI and T2D, respectively. We trained all the
candidate models in parallel and chose the ones with the lowest validation loss as the final
models. The key details for the training include: (a) 80% of the training data was used to
optimize the weights and bias in the neurons, and the rest 20% of the training data was used as
validation set to evaluate the validation loss; (b) The initial Learning rate was 0.01, and was
multiplied by 0.5 if validation loss was not improved for 10 epochs; (c) Early stop was used to
stop training after no improvement in validation loss for 15 epochs; (d) At most 200 epochs of
training were executed; (e) Batch Normalization was used to normalize the outputs of each
hidden layer; (f) Mini-batch training was used; (g) Dropout regularization with a rate of 0.1 was
applied to hidden layers. As (c), (e), and (g) work together to effectively avoid overfitting, making
it unnecessary to use the L1 and L2 regularizations further. The Keras within Tensorflow (43)
was used to implement the building and training of the models.
386

Mediation Analysis (Biomarkers)
We tested the association of the best-performing BMI diet score with the incidence of different
chronic diseases ascertained from the Medicare Chronic Condition Warehouse in MEC-NH,
adjusting for age, sex, BMI, total energy intake (kcal/day), and the first 10 genetic PCs. These
chronic conditions included Alzheimer's disease, anemia, asthma, atrial fibrillation, cataracts,
chronic kidney disease, chronic obstructive pulmonary disease (COPD), depression, gout,
hypertension and related phenotypes, ischemic heart disease, osteoporosis, rheumatoid
arthritis, and stroke. For disease outcomes that were significantly associated with diet score, we
conducted causal mediation analysis to examine whether levels of four different biomarkers
mediate the relationship between diet score and disease status. We employed a two-stage
regression approach within the counterfactual framework for causal mediation analysis using
the mediation R package (version 4.5.1). In the first stage, we fitted a linear regression model to
estimate the mediator model, regressing in turn each biomarker on diet score while controlling
for BMI, daily energy intake, and the first 10 PCs. In the second stage, we fitted a probit
regression model for the binary disease outcome, regressing disease phenotype on both the
biomarker (mediator) and diet score (treatment), while adjusting for age, sex, BMI, daily energy
intake, and the same 10 PCs. We used nonparametric bootstrap resampling with 100
simulations to generate confidence intervals for the mediation effects. The analysis estimated
the Average Causal Mediation Effect (ACME), Average Direct Effect (ADE), and total effect,
representing the indirect effect through the biomarker, the direct effect of diet score, and the

overall effect, respectively. Biomarkers measured in a subset of MEC-NH individuals included

C-reactive protein (CRP; n=1,929), low-density lipoprotein (LDL; n=1,886), high-density

lipoprotein (HDL; n=1,898), and triglycerides (n=1,894). Phenotype transformation for these

biomarkers were performed as previously described (44).

RESULTS

Baseline characteristics for the Native Hawaiian participants are shown in **Table 1**. The total

number of subjects with dietary data was 14,344 and a subset of these individuals had their

DNA samples genotyped (n=5,360). There were more female participants than males. Subjects

with genetics data were slightly younger and had higher education levels than the overall group.

All study subjects had their BMI information. The incidence of T2D from the enrollment to the

latest follow-up was 4380 and the subject with genetics data had significantly higher rate of T2D

(36.3%).

Variable	Dietary data (n=14,344)	Genotype Data (n=5,360)	P-value*
BMI (mean ± sd)	29.1 ± 6.3	28.9 ± 5.9	0.28
T2D Incidence (n,%)	4380 (30.5)	1947 (36.3)	1.09×10^{-14}
AGE (mean ± sd)	56.6 ± 8.7	54.9 ± 7.9	$<2.20 \times 10^{-16}$
SEX (n,%)			
Female	8117 (56.6)	3028 (56.5)	0.92
Male	6227 (43.4)	2332 (43.5)	0.92
Smoking (n,%)			
Never Smoker	5499 (38.3)	2309 (43.1)	1.55×10^{-9}
Former Smoker	5481 (38.2)	2087 (39.0)	0.36
Current Smoker	3214 (22.4)	923 (17.2)	2.11×10^{-15}
Physical Activity (mean ± sd)			
AC-METS	1.7 ± 0.3	1.6 ± 0.3	0.04
Total energy intake (mean ± sd)	2601.9 ± 1342.6	2533.1 ± 1234.4	0.001
Education (n,%)			
Missing	135 (0.9)	42 (0.8)	0.34
< Middle School	100 (0.7)	14 (0.3)	4.92×10^{-4}
Middle School	577 (4.0)	101 (1.9)	3.25×10^{-13}
High School	7023 (49.0)	2206 (41.2)	$<2.20 \times 10^{-16}$
Vocational School	1122 (7.8)	459 (8.6)	0.09
Some College	2991 (20.9)	1325 (24.7)	2.06×10^{-7}
Graduated College	1366 (9.5)	701 (13.1)	5.15×10^{-13}
Graduate/professional School	1030 (7.2)	512 (9.6)	4.11×10^{-8}
Ancestry (mean ± sd)			
EUR-like		0.28±0.26	
EAS-like		0.26±0.26	

AFR-like		0.01±0.03	
PNS-like		0.45±0.23	

Table 1. Descriptive statistics for Native Hawaiians in the Multiethnic Cohort (MEC). Never smoker, an adult who has never smoked or has smoked < 100 cigarettes in his or her lifetime; former smoker, an adult who has smoked ≥ 100 cigarettes in his or her lifetime but who had quit smoking at the time of interview; AC-METS, average of daily metabolic equivalent of physical activity during the last year (continuous); total energy intake, kcal/day. * Unpaired two-sample t-test and two-proportion z-test were conducted for continuous and categorical variables, respectively.

Developing the trans-ancestry PGS for BMI and T2D

To improve upon single-population PGS models previously developed for Native Hawaiians (32), we used two approaches (PRS-CSx and BridgePRS) to develop trans-ancestry PGS for BMI and T2D based on pairwise combinations of GWAS summary statistics from three continental populations: East Asian (n= 256,450 for BMI, 427,504 for T2D), European (n = 700,000 for BMI, 1,812,017 for T2D), and Native Hawaiians (n = 4,156 for BMI, 4,164 for T2D) **(Methods).**

For BMI, the best-performing trans-ancestry model with the highest accuracy in held-out validation MEC-NH (n = 1,226) individuals is one derived by PRS-CSx, combining GWAS summary statistics from European and East-Asian populations (927,557 common SNPs included in the model, with partial R^2 = 0.119; **Figure 1; Supplementary Table S1**). This model outperformed single-population models previously constructed using pruning-and-threshold or LDPred2 (32), publicly available models (32), as well as models estimated by PRS-CS here (partial R^2 = 0.029-0.093; **Figure 1; Supplementary Table S1**). Notably, in single-population PGS models using Native Hawaiian GWAS did not perform better than single-population PGS using East-Asian or European GWAS (partial R^2 = 0.029 vs. 0.074-0.093; **Figure 1; Supplementary Table S1**), presumably due to its small size. In trans-ancestry PGS models, using Native Hawaiian GWAS in combination with East-Asian or European GWAS actually appeared to introduce more noise and lowered the accuracy of PGS models, compared to single-population PGS using either GWAS (partial R^2 = 0.036-0.065 vs. 0.074-0.093; **Figure 1; Supplementary Table S1**). In fact, when we compared the two-population trans-ancestry PGS models to a three-population model, we found that the inclusion of Native Hawaiian GWAS continue to negatively impact the efficacy of the model (partial R^2 = 0.034; **Supplement Figure S3**).

Despite the improved performance of a trans-ancestry PGS combining EAS and EUR GWAS,
because information from Native Hawaiian or Polynesian-ancestry cohorts could not be
integrated effectively, we expect this model to continue to show the disparity in performance
accuracy across population as previously observed for single-population PGS models (33).
Indeed, we found the performance of these trans-ancestry PGS model to be the lowest among
Native Hawaiians most enriched with Polynesian ancestry (**Figure 2**) (45).

For T2D, we observed much of the similar trend (**Figure 1; Supplementary Table S1**). A trans-
ancestry PGS model combining information from EAS and EUR GWAS tend to perform better
(liability R^2 = 0.092) than single-population PGS (liability R^2 = 0.013-0.074) model or trans-
ancestry models including Native Hawaiians (liability R^2 = 0.013-0.048), though in this case
BridgePRS model slightly outperforms the PRS-CSx model (liability R^2 = 0.090 vs. 0.092;
Figure 1; Supplementary Table S1).

Figure 1. Trance-ethnic and single-population polygenic scores in Native Hawaiians. Multiple polygenic scores were
estimated for each phenotype integrating information from single-population GWAS (European [EUR], East Asian
[EAS], or Native Hawaiian [PNS] alone) and combination of GWAS from two populations using either PRS-CSx or
BridgePRS. (a) For BMI, the partial R^2 represents the proportion of variance in BMI that is explained by PGS after
adjusting for the first 10 PCs in validation dataset. (b) The liability R^2 was used to evaluate the predictive ability of
each PGS for T2D in Native Hawaiians after adjusting for age, sex, BMI, and first 10 PCs in validation data. Black

dashed lines represent partial R^2 and liability R^2 of zero and red dotted lines crossing the lowest average variance
explained among the three PGS models per each method.

478
479
480

482 **Figure 2**. Performance of the trans-ethnic PGS models for BMI. Different approaches to combine GWAS summary
483 statistics from multiple ethnic populations using PRS-CSx was evaluated and compared across strata by PNS-like
484 ancestries. The PGS accuracy as measured by partial R^2 was evaluated in different strata of Native Hawaiians based
485 on estimated PNS-like ancestry proportions: < 30%, between 30% to 60%, and > 60%.

Dietary characteristics across the Multiethnic Cohort

The correlation matrix (**Supplementary Figure S4**) displays relationships between the final list
of 520 variables that were used to train our models. We provide the full list of these diet
variables in supplement.

To visualize individual level variation in this high-dimensional space of dietary variables, we first
applied principal components analysis (PCA) to the data after pruning the data for related
variables (**Supplementary Figure S4**), and then projected the top 10 PCs down to two-
dimensions with uniform manifold approximation and projection (UMAP). We performed this
procedure across all five ethnic groups of MEC individuals (n = 191,183) to visualize and
examine differential and shared dietary patterns between racial/ethnic groups. Nearly all Native

Hawaiians in MEC were recruited from Hawaii and most of African-American and Latinos
participants in MEC were recruited in Los Angeles. Japanese and Whites are more evenly
distributed in terms of their recruitment locations (25). When we labeled individuals based on
area of recruitment, it is clear that there may be shared food consumption behaviors among
those recruited in Los Angeles (African Americans, Latinos, and some of the Whites), as well as
those recruited in Hawaiʻi (Native Hawaiians, Japanese, and some of the Whites), indicating a
pattern perhaps driven by food type accessibility that differed by geographical regions (**Figure
3a**). A notable exception may be the Japanese in the MEC, who were more similar to each other
regardless of the area they were recruited from, in contrast to, say, the Whites (**Figure 3a**),
consistent with a stronger influence of consistent cultural practices in food choices.

Restricting our analysis to only participants recruited from Hawaiʻi so to lessen the influence of
geographical constraints of food access, we repeated the UMAP analysis (**Figure 3b**). In this
case, we observed that Japanese and Whites tend to differentiate on the 2-dimensional UMAP
space, suggesting that ethnic- or culturally-driven influences on food choices. Furthermore,
there may be shared food consumption behaviors among the Native Hawaiians with both the
Japanese and Whites, as the pattern from Native Hawaiians span the UMAP space that
overlapped with both the Japanese and Whites (**Figure 3b**). Indeed, by stratifying Native
Hawaiians by estimated EAS-like or PNS-like ancestry, we found a closer affinity in UMAP
space between Native Hawaiians with EAS-like ancestry and Japanese (Kullback-Leibler
divergence, $KLD_{UMAP1}=0.08$. $KLD_{UMAP2}=0.12$) than the Native Hawaiians with PNS-like ancestry
and Japanese ($KLD_{UMAP1}=0.23$, $KLD_{UMAP2}=0.17$) (**Supplementary Figure S5**). We observed a
similar pattern when stratified by EUR-like ancestry as well (**Supplementary Figure S5**).
Together, our findings suggest that there may be cultural effects on dietary choices, which is in
part captured by genetic ancestry.

Subjects by Geographical Location: ▲ Subjects in Hawaii ▲ Subjects in Los Angeles

Subjects by Race/ethnicity in Hawaii: ▲ Native Hawaiian ▲ Black ▲ Japanese ▲ White ▲ Latino/Hispanic

Figure 3. Dietary patterns/behaviors across MEC participants by geographical location and race/ethnicity. We applied
uniform manifold approximation and projection (UMAP) to the top 10 principal components of the dietary data to
visualize the diet/nutrition variability. UMAP is performed once across ethnic groups, but colored and visualized
separately to aid visualization. UMAP was performed either (a) in all MEC participant, colored by recruitment area
(Hawaii vs Los Angeles) or (b) in MEC participants recruited from Hawai'I, colored by their self-identified
race/ethnicity.

Developing a risk stratification model based on dietary information for BMI and T2D

We used three different machine learning algorithms to train and predict diet scores in Native

Hawaiians (n=8,372). Performance of these diet scores were evaluated for BMI and T2D in the

held-out validation dataset (n=5223 for BMI, n=4632 for T2D) (**Table 2**). Diet score models

applying random forest algorithm outperformed scores trained by elastic net and neural network

for both BMI and T2D. For BMI, the model evaluation metrics suggest that a standard deviation

(sd) increase in diet score was significantly associated with 0.38 sd (s.e of ±0.01) increase in

BMI and explained approximately 12% of the variability after adjusting for daily energy intake

(kcal/day) (**Table 2**). Moreover, a sd unit increase in T2D diet score was significantly associated

with 21% increase in odds of having the disease after adjusting for age, sex, BMI, and daily

energy intake (kcal/day). However, T2D diet scores estimated based on these three algorithms

had relatively poor predictability (AUC ranged from 0.57 to 0.58) and explained relatively small

variance of T2D, regardless of whether it is broadly defined incident T2D (Liability R^2 ranged

from 6.04×10^{-3} to 0.012), or incident T2D in 5 years or 10 years from baseline (data not shown).

BMI (n test = 5,223)			
	Elastic	*Random Forest*	*Neural Network*
P-value	5.21×10^{-91}	8.66×10^{-148}	1.04×10^{-127}
Beta ± SE	0.29±0.01	0.38±0.01	0.34±0.01
R^2 [95% CI]	0.096 [0.081 to 0.113]	0.140 [0.123 to 0.158]	0.125 [0.109 to 0.142]
Partial-R^2 [95% CI]	0.075 [0.062 to 0.089]	0.121 [0.104 to 0.138]	0.105 [0.089 to 0.121]
T2D (n test = 4,632)			
	Elastic	*Random Forest*	*Neural Network*
P-value	1.65×10^{-4}	1.96×10^{-7}	1.47×10^{-5}
OR [95% CI]	1.13 [1.06 to 1.21]	1.21 [1.12 to 1.30]	1.16 [1.08 to 1.25]
AUC	0.568 [0.565 to 0.571]	0.582 [0.578 to 0.586]	0.567 [0.564 to 0.570]
Liability R^2 [95% CI]	6.04×10^{-3} [1.20×10^{-3} to 0.014]	0.012 [4.14×10^{-3} to 0.022]	8.14×10^{-3} [2.48×10^{-3} to 0.018]

Table 2. Diet scores trained and validated on BMI and T2D in Native Hawaiians. Performance of the predicted diet

scores were evaluated and presented by phenotype and machine learning algorithm. Partial R^2 for BMI was adjusted

for daily energy intake (kcal/day). Odd ratio and liability R^2 for T2D were adjusted for age, sex, BMI, and daily energy

intake (kcal/day).

For the BMI, the diet score models developed here are much more strongly associate with BMI

and explained much greater proportion of the variation in BMI than established dietary indices

(**Supplementary Table S2**). This is perhaps not surprising as information of BMI was directly

utilized in the development of the diet score model. In the Native Hawaiian cohort, we also find

that the diet scores are significantly associated with future BMI, taken approximately 10 years
after baseline (n=3492; partial R^2 [95%CI] = 0.12 [0.10 to 0.14]), though adult BMI tend to stay
stable over time. While T2D diet scores generally explained little of the phenotypic variation on
the liability scale, it still appeared to be more strongly associated with the outcome than other
dietary indices, including outperforming empirical lifestyle or dietary indices for insulin resistance
(**Supplementary Table S3**).

Finally, we examined how well the diet score models trained and validated in Native Hawaiians
would perform in other MEC populations, focusing on the BMI diet score model as it appeared
to be most efficacious. As is the case with PGS models, we also found that diet score models
generally performed less well in explaining variation in BMI in other ethnic groups, particularly
for Latinos (partial R^2 = 0.027 – 0.056 across diet scores; **Supplementary Figure S6**). Our
results thus suggest that diet score models currently are not well-transferred across populations,
even if the same level of dietary information is available.

Combining PGS and diet score in risk stratification model
As we expect the genetic and non-genetic predictors would capture complementary information
in the variation of the phenotype of interest, we fitted regression models with the best-
performing trans-ancestry PGS and diet score as the main predictors on each of BMI and T2D
to assess their individual and joint effects adjusting for potential confounding variables (**Table 3**
and 4). For BMI, the multivariate model containing both PGS and diet score (along with other
covariates) is a significantly better predictor than models with either of the scores (adjusted R^2 in
Model 3 = 0.29 vs. adjusted R^2 in Model 1 or 2 = 0.21; P = 4.95×10^{-117} and P = 4.89×10^{-120},
respectively by likelihood ratio test; **Table 3.a and 3.b**), suggesting that adding diet score to PGS
model significantly improved the model fit. Both the PGS and diet scores capture
complementary and nearly additive information, as the partial R^2 barely diminished in the joint
model (0.12 or 0.11 in Models 1 or 2, respectively, to 0.10 in Model 3). We did not observe a
significant interaction between PGS and diet score, whether treating the scores as continuous
variables (Model 4) or as categorical variables (Model 5 and 6), though the effect coefficients of
diet score in Model 5 and 6 may hint at a non-linear effect with upper quartiles than the first
quartile of the diet score (**Table 3.a**).

(a). BMI					
n=5,225	**Variables**	**Beta**	**P-value**	**Partial R^2**	**Adjusted R^2**
Model 1	*PGS*	0.39	6.80 x10^{-134}	0.12	0.21
Model 2	*Diet Score*	0.36	1.36 x10^{-131}	0.11	0.21
Model 3	*PGS*	0.35	3.98 x10^{-119}	0.10	0.29
	Diet Score	0.32	5.12 x10^{-116}	0.10	
Model 4	*PGS*	0.35	4.20 x10^{-119}	0.10	0.29
	Diet Score	0.32	5.46 x10^{-116}	0.10	
	PGS :Diet Score	-6.25	0.96	5.53 x10^{-7}	
Model 5	*PGS*	0.36	2.41 x10^{-120}	0.10	0.28
	Diet Score Q2	0.27	1.89 x10^{-14}	1.18 x10^{-2}	
	Diet Score Q3	0.51	5.35 x10^{-47}	4.12 x10^{-2}	
	Diet Score Q4	0.77	3.69 x10^{-88}	7.73 x10^{-2}	
Model 6	*PGS*	0.36	3.88 x10^{-42}	0.04	0.28
	Diet Score Q2	0.27	3.94 x10^{-14}	0.01	
	Diet Score Q3	0.51	2.59 x10^{-46}	0.04	
	Diet Score Q4	0.77	1.50 x10^{-86}	0.08	
	PGS : Diet Score Q2	1.66x10^{-3}	0.96	4.76 x10^{-7}	
	PGS : Diet Score Q3	-1.22x10^{-2}	0.73	2.49 x10^{-5}	
	PGS : Diet Score Q4	-3.12x10^{-3}	0.93	1.64 x10^{-6}	

(b). BMI					
Comparison	**Models Compared**	**df**	**χ^2**	**p-value**	**Δ R^2**
Model 1 vs Model3	PGS vs PGS + diet	1	528.88	4.95×10^{-117}	0.0795
Model 2 vs Model3	Diet vs PGS + diet	1	542.69	4.89×10^{-120}	0.0817
Model 1 vs Model4	PGS vs PGS + diet + PGS×diet	2	528.89	1.42×10^{-115}	0.0795
Model 2 vs Model4	Diet vs PGS + diet + PGS×diet	2	542.70	1.42×10^{-118}	0.0817
Model 3 vs Model4	Additive vs Interaction	1	0.0085	0.93	1.21×10^{-6}
Model 1 vs Model5	PGS vs PGS + diet_Qs	3	438.77	8.83×10^{-95}	0.0665
Model 2 vs Model5	Diet vs PGS + diet_ Qs	3	452.59	8.97×10^{-98}	0.0688
Model 1 vs Model6	PGS vs PGS + diet_ Qs + PGS×diet_Qs	6	438.98	1.16×10^{-91}	0.0666
Model 2 vs Model6	Diet vs PGS + diet_Qs + PGS×diet_Qs	6	452.79	1.23×10^{-94}	0.0688
Model 4 vs Model6	Continuous vs Categorical	4	89.91	1.37×10^{-18}	-0.0129
Model 5 vs Model6	Additive vs Interaction	3	0.2045	0.98	2.97×10^{-5}

Table 3. Regression models incorporating both polygenic score and diet score in Native Hawaiians for BMI. a).
Relationship and predictive power of both PGS and diet score were assessed by estimating their individual effects on
BMI as the main predictor or jointly in the same model while controlling for covariates and potential confounding
(smoking status, physical activity, education level, daily energy intake (kcal/day) and first 10 PCs). Diet variable was
treated as both continuous and categorical variables. Potential interaction effects of PGS and diet score on BMI was
also evaluated by adding a multiplicative interaction term in regression model; b). Model comparisons were
performed using the log-likelihood ratio test.

Overall, the T2D PGS appeared to have a stronger association and better predictability than diet

score based on OR, liability R^2 and AUC as an independent predictor and when included in the

same model with diet score (**Table 4.a**). The log-likelihood ratio tests suggest that the joint

effects of PGS and diet score in Model 3 statistically improved the model fit compared to single

predictor models, though the improvement was marginal (ΔR^2 =0.0028 for Model 1 vs 3 and ΔR^2

=0.032 for Model 2 vs 3 shown in **Table 4.b**). Similar trends were observed when diet score was

treated as a categorical variable. The odds of having T2D increased by 16%, 23%, and 52% for

the 2nd, 3rd, and 4th quartiles, respectively, when compared to the 1ST quartile (Model 5 in **Table**

4.a). Interactions between PGS and diet score (both as continuous and categorical) were

reported that the interaction between two significantly decrease the odds of having T2D.

(a). T2D					
n=4,383	**Variables**	**OR [95% CI]**	**P-value**	**Liability R^2 [95% CI]**	**AUC**
Model 1	PGS	1.71 [1.59 to 1.86]	4.29 x10^{-41}	0.0830 [0.061 to 0.108]	0.7295
Model 2	Diet Score	1.17 [1.09 to 1.26]	3.19 x10^{-5}	0.008 [0.002 to 0.017]	0.7042
Model 3	PGS	1.72 [1.59 to 1.86]	5.19 x10^{-41}	0.0825 [0.060 to 0.107]	0.7312
	Diet Score	1.03 [1.02 to 1.03]	3.81 x10^{-5}		
Model 4	PGS	1.72 [1.59 to 1.87]	3.02 x10^{-41}	0.083 [0.060 to 0.107]	0.7322
	Diet Score	1.18 [1.09 to 1.27]	3.73 x10^{-5}		
	PGS :Diet Score	0.92 [0.85 to 0.98]	1.75 x10^{-2}		
Model 5	PGS	1.72 [1.59 to 1.86]	2.64 x10^{-41}	0.083 [0.061 to 0.108]	0.7312
	Diet Score Q2	1.16 [0.96 to 1.40]	0.12		
	Diet Score Q3	1.23 [1.02 to 1.49]	0.03		
	Diet Score Q4	1.52 [1.23 to 1.88]	8.26 x10^{-5}		
Model 6	PGS	2.04 [1.79 to 2.38]	3.17 x10^{-19}	0.038 [0.026 to 0.054]	0.7320
	Diet Score Q2	1.15 [0.95 to 1.38]	0.14		
	Diet Score Q3	1.22 [1.01 to 1.48]	0.04		
	Diet Score Q4	1.55 [1.27 to 1.91]	1.00 x10^{-4}		
	PGS : Diet Score Q2	0.83 [0.68 to 1.02]	0.06		
	PGS : Diet Score Q3	0.79 [0.65 to 0.97]	0.02		
	PGS : Diet Score Q4	0.84 [0.68 to 1.03]	0.04		

(b). T2D					
Comparison	**Models Compared**	**df**	**χ^2**	**p-value**	**ΔR^2**
Model 1 vs Model3	PGS vs PGS + diet	1	16.89	3.97x10^{-5}	0.0028
Model 2 vs Model3	Diet vs PGS + diet	1	192.79	7.83x10^{-44}	0.0324
Model 1 vs Model4	PGS vs PGS + diet + PGS×diet	2	22.71	1.17x10^{-5}	0.0038
Model 2 vs Model4	Diet vs PGS + diet + PGS×diet	2	198.61	7.44x10^{-44}	0.0334
Model 3 vs Model4	Additive vs Interaction	1	5.83	0.0158	0.001
Model 1 vs Model5	PGS vs PGS + diet_Qs	3	15.92	0.0012	0.0027
Model 2 vs Model5	Diet vs PGS + diet_ Qs	3	191.82	2.47x10^{-41}	0.0323
Model 1 vs Model6	PGS vs PGS + diet_ Qs + PGS×diet_Qs	6	23.09	7.68x10^{-4}	0.0039
Model 2 vs Model6	Diet vs PGS + diet_Qs + PGS×diet_Qs	6	198.99	3.11x10^{-40}	0.0335
Model 4 vs Model6	Continuous vs Categorical	4	0.38	0.9844	1.00 x10^{-4}

Model 5 vs Model6	Additive vs Interaction	3	7.17	0.0667	0.0012

Table 4. Logistic regression models for polygenic risk score and diet score in Native Hawaiians on T2D. a). The relationship and predictive power of both PGS and diet score were assessed by estimating their individual effects on T2D as the main predictor or independent variable and as a covariate to each other controlling for other covariates and potential confounding. Diet variable was treated as both continuous and categorical variables. The combined effect of PGS and diet score on T2D was evaluated by adding a multiplicative interaction term in logistic regression model; b). Each logistic regression model was compared to each other applying the log-likelihood test. All models for T2D were adjusted for age, sex, BMI, smoking status, physical activity, education level, daily energy intake (kcal/day), and first 10 PCs.

Prediction of chronic diseases using BMI diet score (mediation analysis)

Because BMI is a risk factor for a number of chronic diseases, we tested the association of the BMI diet score on the incidence of multiple chronic diseases and conditions in MEC participants. We found that the BMI diet score was associated with 21% increase in odds of developing T2D, 20% increase in odds of having stroke and chronic kidney disease, 17% higher odds of having ischemic heart disease, and 12% increase in having anemia (**Table 5, Supplementary Figure S8**). These associations were found even after adjusting BMI directly in the model, suggesting that the effect due to the diet might be mediated through other mechanisms in addition to BMI. We conducted mediation analysis for each of these five diseases with four different biomarkers that are available to us in the cohort, including CRP (as an inflammatory marker), HDL, LDL, and Triglyceride (reflecting different types of lipids) (**Table 6**). While after correcting for a total of 20 tests we do not generally have significant findings (Bonferroni *p*-value threshold = 0.0025), we did observe several suggestive associations for mediation effects. For example, the average causal mediation effect (ACME) of CRP on T2D (ACME =0.0022, p-value=0.06) suggests that the BMI-related diet could lead to incident T2D through the inflammatory pathway. We also found that BMI-related diet may affect the levels of healthy lipids (HDL) in the system that could play a role in developing chronic kidney disease (ACME=0.0028, p-value=0.02) or stroke (ACME=0.0020, p-value=0.06). For T2D, approximately 13% of the overall relationship between BMI-associated diet and phenotype was attributed by triglycerides (p-value=0.04); there were 23% increased odds of T2D for each unit increase in triglycerides (p-value=4.35×10^{-10}).

Phenotype	P-value	N_{case}	$N_{control}$	OR [95% CI]
Anemia	3.76×10^{-3}	1760	1800	1.12 [1.03 to 1.21]
Chronic Kidney Disease	3.01×10^{-6}	1556	1989	1.20 [1.12 to 1.31]
Ischemic Heart Disease	1.33×10^{-4}	1627	1944	1.17 [1.08 to 1.27]
Stroke	7.14×10^{-4}	561	2921	1.20 [1.08 to 1.34]
T2D	1.17×10^{-7}	1898	2734	1.21 [1.13 to 1.30]

648 **Table 5.** The association of BMI diet score with other disease endpoints. Only diseases with significant association

649 after correcting for multiple testing burden (0.05 / 24 traits = 2.1×10^{-3}) were shown. Anemia was included as its p-

650 value was very close to the threshold. For the complete list of results please refer to **Supplementary Figure S8**.

Mediator: CRP				
Disease	*N*		*Estimate [95% CI]*	*P-value*
T2D	1879 (n_{case}=505)	*Total Effect*	0.0319 [0.0092 to 0.0604]	<0.0001
		ACME	0.0022 [5.9208×10^{-5} to 0.005]	0.06
		ADE	0.0297 [0.0059 to 0.0593]	0.02
		Prop. Mediated	0.0068 [0.0282 to 0.3688]	0.06
Mediator: HDL				
Disease	*N*		*Estimate [95% CI]*	*P-value*
Chronic Kidney Disease	1274 (n_{case}=502)	*Total Effect*	0.0467 [0.0206 to 0.0771]	<0.001
		ACME	0.0028 [0.0003 to 0.0062]	0.02
		ADE	0.0439 [0.0192 to 0.0748]	<0.001
		Prop. Mediated	0.0603 [0.0076 to 0.1533]	0.02
Stroke	1253 (n_{case}=197)	*Total Effect*	0.0406 [0.0163 to 0.0649]	0.02
		ACME	0.0020 [-5.8239×10^{-6} to 0.0054]	0.06
		ADE	0.0387 [0.0118 to 0.0637]	0.02
		Prop. Mediated	0.0482 [-0.0044 to 0.1770]	0.08
Mediator: Triglyceride				
Disease	*N*		*Estimate [95% CI]*	*P-value*
T2D	1814 (n_{case}=507)	*Total Effect*	0.0336 [0.0124 to 0.0615]	0.02
		ACME	0.0042 [0.0004 to 0.0082]	0.02
		ADE	0.0294 [0.0093 to 0.0575]	0.02
		Prop. Mediated	0.1256 [0.0095 to 0.3515]	0.04

Disease	N	Variable	OR [95% CI]	P-value
T2D	1879 (n_{case}=505)	*CRP*	1.1394 [1.0693 to 1.2145]	5.87×10^{-5}
T2D	1814 (n_{case}=507)	*Triglyceride*	1.2330 [1.1547 to 1.3173]	4.35×10^{-10}
Chronic Kidney Disease	1274 (n_{case}=502)	*HDL*	0.9186 [0.8488 to 0.9938]	0.04
Stroke	1253 (n_{case}=197)	*HDL*	0.9216 [0.8364 to 1.0150]	0.10

Table 6. Potential mediators in Native Hawaiians. For the outcome phenotypes in which a significant association was

observed between the BMI diet score and outcome, we tested the potential indirect effects of BMI diet score on

disease phenotypes through mediation analysis with potential mediators. Specifically, we explored biomarkers that

may reflect inflammation (C-reactive protein) and lipid pathways (low- and high-density lipoproteins, and

triglycerides). The mediation analysis estimated total effect, which sums up direct and indirect effect of diet score,

average causal mediated/indirect effect (ACME) of biomarkers, average direct effect (ADE) of diet score, and

proportion of mediated effect. The marginal effects of mediator on phenotype were also reported as odds ratios after

accounting for covariates (diet score, age, sex, BMI, daily energy intake (kcal/day), and first 10 PCs).

DISCUSSION

In the present study we developed comprehensive risk stratification models integrating genetic and dietary predictors for BMI and type-2 diabetes in Native Hawaiians. Our trans-ancestry polygenic score approach showed improved prediction accuracy over single-ancestry models, demonstrating that leveraging large-scale cross-ancestry GWAS summary statistics can improve prediction accuracy in populations with limited genetic studies. Beyond genetic factors, dietary intake emerged as a powerful complementary predictor, with our hypothesis-free diet-based models explaining substantial variance in BMI. The combination of polygenic scores and dietary predictors achieved markedly improved prediction performance compared to either approach alone. Furthermore, our best-performing BMI diet score showed associations with multiple chronic diseases, even when BMI was included as a covariate, suggesting that BMI-associated dietary patterns may influence broader health outcomes through inflammatory and lipid-mediated pathways.

Building on these predictive capabilities, our analysis revealed that dietary intake explains phenotypic variation beyond genetic factors. Our best-performing diet score accounted for approximately 12% of adult BMI variance, demonstrating substantially higher explanatory power than Mediterranean diet scores (~0.6-3%), Healthy Eating Index (~3.9%), and other established dietary pattern indices (1-3%) (45-49). In contrast, the best-performing diet model trained on T2D incidence, though significantly associated with the outcome in held-out validation cohort, explained only ~1% of phenotypic variation on the liability scale. This limited predictive ability for T2D likely reflects several factors: dietary patterns vary considerably over time, making past diet a poor predictor of future disease onset; and individuals with pre-diabetic conditions may have already modified their diets toward healthier patterns, confounding the diet-disease relationship.

Overall, the effect sizes of elastic net-selected dietary variables were relatively small for T2D incidence compared to BMI (**Supplement Data**). As we anticipated, there were shared dietary patterns between BMI and T2D given that BMI is an established T2D risk factor. For example, established risk factors including higher red meat consumption and diet soda intake were positively associated with both BMI and T2D (50-53). Interestingly, more non-Western dietary patterns (ramen/saimin, refried beans, enchiladas) were associated with reduced T2D (54), suggesting population-specific dietary-metabolic relationships that warrant further investigation.

One current limitation of our diet score models is that it is not transferable to other cohorts. Unlike genetic data which is more standardized or can be imputed to the same set of genetic markers for calculating the polygenic scores, our diet score models currently require having the same set of diet information available to calculate the score. Indeed, we have examined the possibility of harmonizing the available diet phenotype in the MEC with that from UK Biobank through manual curation but found that variables can only match ~30-57% of the times depending on the strictness of interpretation. Therefore, to utilize diet scores in risk stratification in different cohorts will require future research on variable harmonization of dietary surveys or assessment tools.

Another limitation is that while our composite scores effectively predict metabolic phenotypes, establishing a causal role of diet remains challenging. Unlike polygenic scores, which are determined at conception and remain constant throughout life, dietary information in the MEC was collected in adulthood, when participants may already have been in the preliminary stages of disease but not yet diagnosed. As a step toward establishing causality, we tested whether our best-performing BMI diet score could predict BMI measured approximately 10 years after baseline, finding significant predictive ability (n=3492; partial R^2 [95%CI] = 0.12 [0.10 to 0.14]), although we should note that adult BMI is generally stable and our two measurements of BMI, 10 years apart, are highly correlated (R^2 = 0.7358). Future longitudinal study designs that collect detailed dietary data, particularly in adolescent or young adults, could greatly benefit efforts to establish causal relationships between diet and metabolic conditions (55-59).

Despite the challenges to establish causality with diet scores, we found BMI diet score to be strongly associated with incidence of some chronic disease status in Native Hawaiians, such as T2D, chronic kidney disease, ischemic heart disease and stroke. These associations were observed even after adjusting for BMI, suggesting that dietary effects involve mechanisms independent of body weight. To begin probing the mechanism through which BMI-related diet is associated with incident chronic diseases, we turned to mediation analysis. Our mediation analysis is preliminary and limited given the relatively small number of individuals with available molecular biomarkers. Given this caveat, our analysis revealed that the markers we tested explain less than 13% of the association between the dietary scores and chronic disease risk, indicating that the majority of diet's health effects operate through unmeasured biological pathways beyond inflammatory markers and lipid metabolism. Future studies should consider emerging pathways such as gut microbiome interactions, metabolomic alterations, epigenetic

mechanisms, and direct cellular mechanisms (including mitochondrial and endothelial function) to capture the full impact of diet on chronic disease risk (60-64). Population-specific mechanisms should also be investigated, such as genetic variants affecting metabolic enzyme function, population-specific genetic adaptations in Native Hawaiians, and traditional versus Western diet interactions (65-68). Nevertheless, due to this apparent 'pleiotropic' effect of dietary patterns associated with BMI, these BMI-prone diet scores could be incorporated into clinical risk assessment tools to improve chronic disease risk prediction.

Finally, irrespective of the clinical utilities of the diet score models, at least for BMI and to a lesser extent T2D, it is clear that diet information explains part of the phenotypic variation that are not captured by the best available polygenic predictors. This also means when the diet information is available for a cohort in a genetic study, computing and adjusting for diet scores will help remove a part of the phenotypic variation explained by the non-genetic factors, thereby improving the statistical power of the genetic mapping study. Such an increase in power will be welcomed when one is trying to map population-specific associations with a relatively small GWAS cohort.

CONCLUSION

For populations with limited genetic data, integrating large-scale GWAS data from other ancestries substantially improved polygenic score performance compared to single-ancestry approaches in Native Hawaiians. Similarly, data-driven diet scores outperformed established dietary indices in explaining BMI variance (12% vs. 0.6-3.9%). Importantly, diet scores trained on shared risk factors such as BMI can be applied across different chronic diseases, though effectiveness may vary by the disease, the population and dietary patterns. However, our mediation analysis revealed that traditional biomarkers (inflammation and lipid pathways) explain less than 13% of the association between BMI-associated dietary patterns and chronic diseases. Future research should more robustly characterize the mediation effect of molecular biomarkers, as well as investigate emerging pathways including gut microbiome, metabolomics, and epigenetic mechanisms to fully understand diet-health relationships. Overall, polygenic scores and diet scores provide complementary and substantial predictive information for BMI and, with more limited success, for T2D in this population.

REFERENCES

1. Samson, S.L. & Garber, A.J. (2024). Metabolic Syndrome. In: StatPearls. StatPearls Publishing

2. Beltrán-Sánchez H, Harhay MO, Harhay MM, McElligott S. Prevalence and trends of metabolic syndrome in the adult U.S. population, 1999-2010. J Am Coll Cardiol. 2013 Aug 20;62(8):697-703. doi: 10.1016/j.jacc.2013.05.064. Epub 2013 Jun 27. PMID: 23810877; PMCID: PMC3756561.

3. American Heart Association (2024). What is Metabolic Syndrome? Retrieved from https://www.heart.org/en/health-topics/metabolic-syndrome/about-metabolic-syndrome
4. Prasad GV. Metabolic syndrome and chronic kidney disease: Current status and future directions. World J Nephrol. 2014 Nov 6;3(4):210-9. doi: 10.5527/wjn.v3.i4.210. PMID: 25374814; PMCID: PMC4220353.

5. Li W, Chen D, Peng Y, Lu Z, Kwan MP, Tse LA. Association Between Metabolic Syndrome and Mortality: Prospective Cohort Study. JMIR Public Health Surveill. 2023 Sep 5;9:e44073. doi: 10.2196/44073. PMID: 37669100; PMCID: PMC10509744.

6. World Health Organization (2025). Obesity and overweight. Fact Sheet.

7. CDC (2024). National Diabetes Statistics Report. Centers for Disease Control and Prevention

8. CDC (2025). Diabetes in Young People Is on the Rise. Centers for Disease Control and Prevention.

9. Yengo, L., Liang, Y., 23andMe Research Team, Wang, X., Granka, J. M., Evans, D. M., Sidorenko, J., & Visscher, P. M. (2025). Within-family heritability estimates for behavioural and disease phenotypes from 500,000 sibling pairs of diverse ancestries. *medRxiv*. https://doi.org/10.1101/2025.09.17.25336022

10. Xiang, R., Kelemen, M., Xu, Y. *et al.* Recent advances in polygenic scores: translation, equitability, methods and FAIR tools. *Genome Med* **16**, 33 (2024). https://doi.org/10.1186/s13073-024-01304-9

11. **Truong, B. et al. (2024).** Integrative polygenic risk score improves the prediction accuracy of complex traits and diseases. Cell Genomics, 4(3), 100504

12. Slunecka, J.L., van der Zee, M.D., Beck, J.J. *et al.* Implementation and implications for polygenic risk scores in healthcare. *Hum Genomics* **15**, 46 (2021). https://doi.org/10.1186/s40246-021-00339-y

13. Duncan, L., Shen, H., Gelaye, B. *et al.* Analysis of polygenic risk score usage and performance in diverse human populations. *Nat Commun* **10**, 3328 (2019). https://doi.org/10.1038/s41467-019-11112-0

14. Wang, Y., Guo, J., Ni, G. *et al.* Theoretical and empirical quantification of the accuracy of polygenic scores in ancestry divergent populations. *Nat Commun* **11**, 3865 (2020). https://doi.org/10.1038/s41467-020-17719-y

15. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current polygenic risk scores may exacerbate health disparities. Nat Genet. 2019 Apr;51(4):584-591. doi: 10.1038/s41588-019-0379-x. Epub 2019 Mar 29. Erratum in: Nat Genet. 2021 May;53(5):763. doi: 10.1038/s41588-021-00797-z. PMID: 30926966; PMCID: PMC6563838.

16. Roberts MC, Khoury MJ, Mensah GA. Perspective: The Clinical Use of Polygenic Risk Scores: Race, Ethnicity, and Health Disparities. Ethn Dis. 2019 Jul 18;29(3):513-516. doi: 10.18865/ed.29.3.513. PMID: 31367172; PMCID: PMC6645721.

17. Ambroselli D, Masciulli F, Romano E, Catanzaro G, Besharat ZM, Massari MC, Ferretti E, Migliaccio S, Izzo L, Ritieni A, Grosso M, Formichi C, Dotta F, Frigerio F, Barbiera E, Giusti AM, Ingallina C, Mannina L. New Advances in Metabolic Syndrome, from Prevention to Treatment: The Role of Diet and Food. Nutrients. 2023 Jan 26;15(3):640. doi: 10.3390/nu15030640. PMID: 36771347; PMCID: PMC9921449.

18. Clemente-Suárez VJ, Beltrán-Velasco AI, Redondo-Flórez L, Martín-Rodríguez A, Tornero-Aguilera JF. Global Impacts of Western Diet and Its Effects on Metabolism and Health: A Narrative Review. Nutrients. 2023 Jun 14;15(12):2749. doi: 10.3390/nu15122749. PMID: 37375654; PMCID: PMC10302286.
19. Castro-Barquero S, Ruiz-León AM, Sierra-Pérez M, Estruch R, Casas R. Dietary Strategies for Metabolic Syndrome: A Comprehensive Review. Nutrients. 2020 Sep 29;12(10):2983. doi: 10.3390/nu12102983. PMID: 33003472; PMCID: PMC7600579.

20. Kastorini CM, Milionis HJ, Esposito K, Giugliano D, Goudevenos JA, Panagiotakos DB. The effect of Mediterranean diet on metabolic syndrome and its components: a meta-analysis of 50 studies and 534,906 individuals. J Am Coll Cardiol. 2011 Mar 15;57(11):1299-313. doi: 10.1016/j.jacc.2010.09.073. PMID: 21392646.

21. Klapp R, Laxamana JA, Shvetsov YB, Park SY, Kanehara R, Setiawan VW, Danquah I, Le Marchand L, Maskarinec G. The EAT-Lancet Diet Index Is Associated with Lower Obesity and Incidence of Type 2 Diabetes in the Multiethnic Cohort. J Nutr. 2024 Nov;154(11):3407-3415. doi: 10.1016/j.tjnut.2024.06.018. Epub 2024 Jul 15. PMID: 39019161; PMCID: PMC11600087.

22. Park SY, Boushey CJ, Wilkens LR, Haiman CA, Le Marchand L. High-Quality Diets Associate With Reduced Risk of Colorectal Cancer: Analyses of Diet Quality Indexes in the Multiethnic Cohort. Gastroenterology. 2017 Aug;153(2):386-394.e2. doi: 10.1053/j.gastro.2017.04.004. Epub 2017 Apr 17. PMID: 28428143; PMCID: PMC5526717.

23. CDC/HHS Office of Minority Health data. https://minorityhealth.hhs.gov/diabetes-and-native-hawaiianspacific-islanders

24. Karter AJ, Schillinger D, Adams AS, Moffet HH, Liu J, Adler NE, Kanaya AM. Elevated rates of diabetes in Pacific Islanders and Asian subgroups: The Diabetes Study of Northern California (DISTANCE). Diabetes Care. 2013 Mar;36(3):574-9. doi: 10.2337/dc12-0722. Epub 2012 Oct 15. PMID: 23069837; PMCID: PMC3579366.

25. Kolonel LN, Henderson BE, Hankin JH, Nomura AM, Wilkens LR, Pike MC, Stram DO, Monroe KR, Earle ME, Nagamine FS. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. Am J Epidemiol. 2000 Feb 15;151(4):346-57. doi: 10.1093/oxfordjournals.aje.a010213. PMID: 10695593; PMCID: PMC4482109.

26. Stram DO, Hankin JH, Wilkens LR, Pike MC, Monroe KR, Park S, Henderson BE, Nomura AM, Earle ME, Nagamine FS, Kolonel LN. Calibration of the dietary questionnaire for a multiethnic cohort in Hawaii and Los Angeles. Am J Epidemiol. 2000 Feb 15;151(4):358-70. doi: 10.1093/oxfordjournals.aje.a010214. PMID: 10695594; PMCID: PMC4482461.

27. Bogumil D, Sheng X, Wan P, Xia L, Pooler L, Cheng I, Streicher S, Huang BZ, Chen F, Stram D, Shen S, King G, Chiang CWK, Ongaco C, Adams M, McMullen I, Zhang P, Ling H, Mawhinney M, Doheny KF, Marchand LL, Wilkens LR, Haiman CA, Conti DV. The Multiethnic Cohort: A Resource for the study of Genetic and non-Genetic Cancer Risk Across Populations. medRxiv [Preprint]. 2025 Jun 11:2025.06.09.25328993. doi: 10.1101/2025.06.09.25328993. PMID: 40585109; PMCID: PMC12204407.

28. Sheng X, Xia L, Cahoon JL, Conti DV, Haiman CA, Kachuri L, Chiang CWK. Inverted genomic regions between reference genome builds in humans impact imputation accuracy and decrease the power of association testing. HGG Adv. 2022 Nov 11;4(1):100159. doi: 10.1016/j.xhgg.2022.100159. PMID: 36465187; PMCID: PMC9709082.

29. Loh, P.R., Danecek, P., Palamara, P.F., Fuchsberger, C., Reshef, Y.A., Finucane, H.K., Schoenherr, S., Forer, L., McCarthy, S., Abecasis, G.R., Durbin, R., & Price, A.L. (2016). Reference-based phasing using the Haplotype Reference Consortium panel. *Nature Genetics*, 48(11), 1443-1448. https://doi.org/10.1038/ng.3679

30. Dinh, B.L., Tang, E., Taparra, K., Nakatsuka, N., Chen, F., & Chiang, C.W.K. (2024). Recombination map tailored to Native Hawaiians may improve robustness of genomic scans for positive selection. *Human Genetics*, 143(1), 85-99. https://doi.org/10.1007/s00439-023-02625-2

31. Taliun, D., Harris, D.N., Kessler, M.D., Carlson, J., Szpiech, Z.A., Torres, R., Taliun, S.A.G., Corvelo, A., Gogarten, S.M., Kang, H.M., et al. (2021). Sequencing of 53,831 diverse genomes from the NHLBI TOPMed Program. *Nature*, 589(7845), 290-299. https://doi.org/10.1038/s41586-021-03205-y

32. Lo, YC., Tian, H., Chan, T.F. *et al.* The accuracy of polygenic score models for BMI and Type II diabetes in the Native Hawaiian population. *Commun Biol* **8**, 651 (2025). https://doi.org/10.1038/s42003-025-08050-7

33. Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM (2010) Robust relationship inference in genome-wide association studies. Bioinformatics 26(22):2867-2873

34. Conomos MP, Miller MB, Thornton TA. Robust inference of population structure for ancestry prediction and correction of stratification in the presence of relatedness. Genet Epidemiol. 2015 May;39(4):276-93. doi: 10.1002/gepi.21896. Epub 2015 Mar 23. PMID: 25810074; PMCID: PMC4836868.

35. Conomos MP, Reiner AP, Weir BS, Thornton TA. Model-free Estimation of Recent Genetic Relatedness. Am J Hum Genet. 2016 Jan 7;98(1):127-48. doi: 10.1016/j.ajhg.2015.11.022. PMID: 26748516; PMCID: PMC4716688.

36. Stephanie M Gogarten, Tamar Sofer, Han Chen, Chaoyu Yu, Jennifer A Brody, Timothy A Thornton, Kenneth M Rice, Matthew P Conomos, Genetic association testing using the GENESIS R/Bioconductor package, *Bioinformatics*, Volume 35, Issue 24, December 2019, Pages 5346–5348, https://doi.org/10.1093/bioinformatics/btz567

37. Y Ruan, YF Lin, YCA Feng, CY Chen, M Lam, Z Guo, Stanley Global Asia Initiatives, L He, A Sawa, AR Martin, S Qin, H Huang, T Ge. Improving polygenic prediction in ancestrally diverse populations. *Nature Genetics*, 54:573-580, 2022.

38. Hoggart, C.J., Choi, S.W., García-González, J. *et al.* BridgePRS leverages shared genetic effects across ancestries to increase polygenic risk score portability. *Nat Genet* **56**, 180–186 (2024). https://doi.org/10.1038/s41588-023-01583-9

39. Suzuki K, Hatzikotoulas K, Southam L, Taylor HJ, Yin X, Lorenz KM, Mandla R, Huerta-Chagoya A, Melloni GEM, Kanoni S, Rayner NW, Bocher O, Arruda AL, Sonehara K, Namba S, Lee SSK, Preuss MH, Petty LE, Schroeder P, Vanderwerff B, Kals M, Bragg F, Lin K, Guo X, Zhang W, Yao J, Kim YJ, Graff M, Takeuchi F, Nano J, Lamri A, Nakatochi M, Moon S, Scott RA, Cook JP, Lee JJ, Pan I, Taliun D, Parra EJ, Chai JF, Bielak LF, Tabara Y, Hai Y, Thorleifsson G, Grarup N, Sofer T, Wuttke M, Sarnowski C, Gieger C, Nousome D, Trompet S, Kwak SH, Long J, Sun M, Tong L, Chen WM, Nongmaithem SS, Noordam R, Lim VJY, Tam CHT, Joo YY, Chen CH, Raffield LM, Prins BP, Nicolas A, Yanek LR, Chen G, Brody JA, Kabagambe E, An P, Xiang AH, Choi HS, Cade BE, Tan J, Broadaway KA, Williamson A, Kamali Z, Cui J, Thangam M, Adair LS, Adeyemo A, Aguilar-Salinas CA, Ahluwalia TS, Anand SS, Bertoni A, Bork-Jensen J, Brandslund I, Buchanan TA, Burant CF, Butterworth AS, Canouil M, Chan JCN, Chang LC, Chee ML, Chen J, Chen SH, Chen YT, Chen Z, Chuang LM, Cushman M, Danesh J, Das SK, de Silva HJ, Dedoussis G, Dimitrov L, Doumatey AP, Du S, Duan Q, Eckardt KU, Emery LS, Evans DS, Evans MK, Fischer K, Floyd JS, Ford I, Franco OH, Frayling TM, Freedman BI, Genter P, Gerstein HC, Giedraitis V, González-Villalpando C, González-Villalpando ME, Gordon-Larsen P, Gross M, Guare LA, Hackinger S, Hakaste L, Han S, Hattersley AT, Herder C, Horikoshi M, Howard AG, Hsueh W, Huang M, Huang W, Hung YJ, Hwang MY, Hwu CM, Ichihara S, Ikram MA, Ingelsson M, Islam MT, Isono M, Jang HM, Jasmine F, Jiang G, Jonas JB, Jørgensen T, Kamanu FK, Kandeel FR, Kasturiratne A, Katsuya T, Kaur V, Kawaguchi T, Keaton JM, Kho AN, Khor CC, Kibriya MG, Kim DH, Kronenberg F, Kuusisto J, Läll K, Lange LA, Lee KM, Lee MS, Lee NR, Leong A, Li L, Li Y, Li-Gao R, Ligthart S, Lindgren CM, Linneberg A, Liu CT, Liu J, Locke AE, Louie T, Luan J, Luk AO, Luo X, Lv J, Lynch JA, Lyssenko V, Maeda S, Mamakou V, Mansuri SR, Matsuda K, Meitinger T, Melander O, Metspalu A, Mo H, Morris AD, Moura FA, Nadler JL, Nalls MA, Nayak U, Ntalla I, Okada Y, Orozco L, Patel SR, Patil S, Pei P, Pereira MA, Peters A, Pirie FJ, Polikowsky HG, Porneala B, Prasad G, Rasmussen-Torvik LJ, Reiner AP, Roden M, Rohde R, Roll K, Sabanayagam C, Sandow K, Sankareswaran A, Sattar N, Schönherr S, Shahriar M, Shen B, Shi J, Shin DM, Shojima N, Smith JA, So WY, Stančáková A, Steinthorsdottir V, Stilp AM, Strauch K, Taylor KD, Thorand B, Thorsteinsdottir U, Tomlinson B, Tran TC, Tsai FJ, Tuomilehto J, Tusie-Luna T, Udler MS, Valladares-Salgado A, van Dam RM, van Klinken JB, Varma R, Wacher-Rodarte N, Wheeler E, Wickremasinghe AR, van Dijk KW, Witte DR, Yajnik CS, Yamamoto K, Yamamoto K, Yoon K, Yu C, Yuan JM, Yusuf S, Zawistowski M, Zhang L, Zheng W; VA Million Veteran Program; Raffel LJ, Igase M, Ipp E, Redline S, Cho YS, Lind L, Province MA, Fornage M, Hanis CL, Ingelsson E, Zonderman AB, Psaty BM, Wang YX, Rotimi CN, Becker DM, Matsuda F, Liu Y, Yokota M, Kardia SLR, Peyser PA, Pankow JS, Engert JC, Bonnefond A, Froguel P, Wilson JG, Sheu WHH, Wu JY, Hayes MG, Ma RCW, Wong TY, Mook-Kanamori DO, Tuomi T, Chandak GR, Collins FS, Bharadwaj D, Paré G, Sale MM, Ahsan H, Motala AA, Shu XO, Park

KS, Jukema JW, Cruz M, Chen YI, Rich SS, McKean-Cowdin R, Grallert H, Cheng CY, Ghanbari M, Tai ES, Dupuis J, Kato N, Laakso M, Köttgen A, Koh WP, Bowden DW, Palmer CNA, Kooner JS, Kooperberg C, Liu S, North KE, Saleheen D, Hansen T, Pedersen O, Wareham NJ, Lee J, Kim BJ, Millwood IY, Walters RG, Stefansson K, Ahlqvist E, Goodarzi MO, Mohlke KL, Langenberg C, Haiman CA, Loos RJF, Florez JC, Rader DJ, Ritchie MD, Zöllner S, Mägi R, Marston NA, Ruff CT, van Heel DA, Finer S, Denny JC, Yamauchi T, Kadowaki T, Chambers JC, Ng MCY, Sim X, Below JE, Tsao PS, Chang KM, McCarthy MI, Meigs JB, Mahajan A, Spracklen CN, Mercader JM, Boehnke M, Rotter JI, Vujkovic M, Voight BF, Morris AP, Zeggini E. Genetic drivers of heterogeneity in type 2 diabetes pathophysiology. Nature. 2024 Mar;627(8003):347-357. doi: 10.1038/s41586-024-07019-6. Epub 2024 Feb 19. PMID: 38374256; PMCID: PMC10937372.

40. Lee SH, Goddard ME, Wray NR, Visscher PM. A better coefficient of determination for genetic profile analysis. Genet Epidemiol. 2012 Apr;36(3):214-24. doi: 10.1002/gepi.21614. PMID: 22714935.

41. Liese AD, Krebs-Smith SM, Subar AF, George SM, Harmon BE, Neuhouser ML, Boushey CJ, Schap TE, Reedy J. The Dietary Patterns Methods Project: synthesis of findings across cohorts and relevance to dietary guidance. J Nutr. 2015 Mar;145(3):393-402. doi: 10.3945/jn.114.205336. Epub 2015 Jan 21. PMID: 25733454; PMCID: PMC4336525.

42. Castro JL, Mantas CJ, Benítez JM. Neural networks with a continuous squashing function in the output are universal approximators. Neural Netw. 2000 Jul;13(6):561-3. doi: 10.1016/s0893-6080(00)00031-9. PMID: 10987509.

43. arXiv:1603.04467v2

44. Wojcik GL, Graff M, Nishimura KK, Tao R, Haessler J, Gignoux CR, Highland HM, Patel YM, Sorokin EP, Avery CL, Belbin GM, Bien SA, Cheng I, Cullina S, Hodonsky CJ, Hu Y, Huckins LM, Jeff J, Justice AE, Kocarnik JM, Lim U, Lin BM, Lu Y, Nelson SC, Park SL, Poisner H, Preuss MH, Richard MA, Schurmann C, Setiawan VW, Sockell A, Vahi K, Verbanck M, Vishnu A, Walker RW, Young KL, Zubair N, Acuña-Alonso V, Ambite JL, Barnes KC, Boerwinkle E, Bottinger EP, Bustamante CD, Caberto C, Canizales-Quinteros S, Conomos MP, Deelman E, Do R, Doheny K, Fernández-Rhodes L, Fornage M, Hailu B, Heiss G, Henn BM, Hindorff LA, Jackson RD, Laurie CA, Laurie CC, Li Y, Lin DY, Moreno-Estrada A, Nadkarni G, Norman PJ, Pooler LC, Reiner AP, Romm J, Sabatti C, Sandoval K, Sheng X, Stahl EA, Stram DO, Thornton TA, Wassel CL, Wilkens LR, Winkler CA, Yoneyama S, Buyske S, Haiman CA, Kooperberg C, Le Marchand L, Loos RJF, Matise TC, North KE, Peters U, Kenny EE, Carlson CS. Genetic analyses of diverse populations improves discovery for complex traits. Nature. 2019 Jun;570(7762):514-518. doi: 10.1038/s41586-019-1310-4. Epub 2019 Jun 19. PMID: 31217584; PMCID: PMC6785182.

45. Minster, R.L., et al. (2020). Evolutionary history of modern Samoans. *Proceedings of the National Academy of Sciences*, 117(17), 9458-9465. https://doi.org/10.1073/pnas.1913157117

46. Maskarinec G, Novotny R, Tasaki K. Dietary patterns are associated with body mass index in multiethnic women. J Nutr. 2000 Dec;130(12):3068-72. doi: 10.1093/jn/130.12.3068. PMID: 11110871.

47. Martínez-González MA, García-Arellano A, Toledo E, Salas-Salvadó J, Buil-Cosiales P, Corella D, Covas MI, Schröder H, Arós F, Gómez-Gracia E, Fiol M, Ruiz-Gutiérrez V, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pintó X, Muñoz MA, Wärnberg J, Ros E, Estruch R;

PREDIMED Study Investigators. A 14-item Mediterranean diet assessment tool and obesity indexes among high-risk subjects: the PREDIMED trial. PLoS One. 2012;7(8):e43134. doi: 10.1371/journal.pone.0043134. Epub 2012 Aug 14. PMID: 22905215; PMCID: PMC3419206.

48. Hurley KM, Oberlander SE, Merry BC, Wrobleski MM, Klassen AC, Black MM. The healthy eating index and youth healthy eating index are unique, nonredundant measures of diet quality among low-income, African American adolescents. J Nutr. 2009 Feb;139(2):359-64. doi: 10.3945/jn.108.097113. Epub 2008 Dec 11. PMID: 19074210; PMCID: PMC2646206.

49. Tande DL, Magel R, Strand BN. Healthy Eating Index and abdominal obesity. Public Health Nutr. 2010 Feb;13(2):208-14. doi: 10.1017/S1368980009990723. Epub 2009 Aug 4. PMID: 19650960.

50. Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Willett WC, Hu FB. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. Am J Clin Nutr. 2011 Oct;94(4):1088-96. doi: 10.3945/ajcn.111.018978. Epub 2011 Aug 10. PMID: 21831992; PMCID: PMC3173026.

51. Li C, Bishop TRP, Imamura F, Sharp SJ, Pearce M, Brage S, Ong KK, Ahsan H, Bes-Rastrollo M, Beulens JWJ, den Braver N, Byberg L, Canhada S, Chen Z, Chung HF, Cortés-Valencia A, Djousse L, Drouin-Chartier JP, Du H, Du S, Duncan BB, Gaziano JM, Gordon-Larsen P, Goto A, Haghighatdoost F, Härkänen T, Hashemian M, Hu FB, Ittermann T, Järvinen R, Kakkoura MG, Neelakantan N, Knekt P, Lajous M, Li Y, Magliano DJ, Malekzadeh R, Le Marchand L, Marques-Vidal P, Martinez-Gonzalez MA, Maskarinec G, Mishra GD, Mohammadifard N, O'Donoghue G, O'Gorman D, Popkin B, Poustchi H, Sarrafzadegan N, Sawada N, Schmidt MI, Shaw JE, Soedamah-Muthu S, Stern D, Tong L, van Dam RM, Völzke H, Willett WC, Wolk A, Yu C; EPIC-InterAct Consortium; Forouhi NG, Wareham NJ. Meat consumption and incident type 2 diabetes: an individual-participant federated meta-analysis of 1·97 million adults with 100 000 incident cases from 31 cohorts in 20 countries. Lancet Diabetes Endocrinol. 2024 Sep;12(9):619-630. doi: 10.1016/S2213-8587(24)00179-7. Erratum in: Lancet Diabetes Endocrinol. 2025 Feb;13(2):e2. doi: 10.1016/S2213-8587(25)00002-6. PMID: 39174161.

52. Malik VS, Popkin BM, Bray GA, Després JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. Diabetes Care. 2010 Nov;33(11):2477-83. doi: 10.2337/dc10-1079. Epub 2010 Aug 6. PMID: 20693348; PMCID: PMC2963518.

53. Wang M, Yu M, Fang L, Hu RY. Association between sugar-sweetened beverages and type 2 diabetes: A meta-analysis. J Diabetes Investig. 2015 May;6(3):360-6. doi: 10.1111/jdi.12309. Epub 2014 Dec 11. PMID: 25969723; PMCID: PMC4420570.

54. Jizhao Niu, Bai Li, Qing Zhang, Ge Chen, Angeliki Papadaki, Exploring the traditional Chinese diet and its association with health status—a systematic review, *Nutrition Reviews*, Volume 83, Issue 2, February 2025, Pages e237–e256, https://doi.org/10.1093/nutrit/nuae013

55. Emmett PM, Jones LR. Diet, growth, and obesity development throughout childhood in the Avon Longitudinal Study of Parents and Children. Nutr Rev. 2015 Oct;73 Suppl 3(Suppl 3):175-206. doi: 10.1093/nutrit/nuv054. PMID: 26395342; PMCID: PMC4586450.

56. Solsona, E.M., Johnson, L., Northstone, K. *et al.* Prospective association between an obesogenic dietary pattern in early adolescence and metabolomics derived and traditional cardiometabolic risk scores in adolescents and young adults from the ALSPAC cohort. *Nutr Metab (Lond)* **20**, 41 (2023). https://doi.org/10.1186/s12986-023-00754-z

57. Ambrosini GL, Emmett PM, Northstone K, Jebb SA. Tracking a dietary pattern associated with increased adiposity in childhood and adolescence. Obesity (Silver Spring). 2014 Feb;22(2):458-65. doi: 10.1002/oby.20542. Epub 2013 Sep 17. PMID: 23804590; PMCID: PMC3846445.

58. Foraita, R., Witte, J., Börnhorst, C. *et al.* A longitudinal causal graph analysis investigating modifiable risk factors and obesity in a European cohort of children and adolescents. *Sci Rep* **14**, 6822 (2024). https://doi.org/10.1038/s41598-024-56721-y

59. Laska MN, Murray DM, Lytle LA, Harnack LJ. Longitudinal associations between key dietary behaviors and weight gain over time: transitions through the adolescent years. Obesity (Silver Spring). 2012 Jan;20(1):118-25. doi: 10.1038/oby.2011.179. Epub 2011 Jun 23. PMID: 21701567; PMCID: PMC3402912.

60. Boulangé, C.L., Neves, A.L., Chilloux, J. *et al.* Impact of the gut microbiota on inflammation, obesity, and metabolic disease. *Genome Med* **8**, 42 (2016). https://doi.org/10.1186/s13073-016-0303-2

61. Jin Q, Black A, Kales SN, Vattem D, Ruiz-Canela M, Sotos-Prieto M. Metabolomics and Microbiomes as Potential Tools to Evaluate the Effects of the Mediterranean Diet. Nutrients. 2019 Jan 21;11(1):207. doi: 10.3390/nu11010207. PMID: 30669673; PMCID: PMC6356665.

62. Li D, Li Y, Yang S, Lu J, Jin X, Wu M. Diet-gut microbiota-epigenetics in metabolic diseases: From mechanisms to therapeutics. Biomed Pharmacother. 2022 Sep;153:113290. doi: 10.1016/j.biopha.2022.113290. Epub 2022 Jun 17. PMID: 35724509.

63. Putti R, Sica R, Migliaccio V, Lionetti L. Diet impact on mitochondrial bioenergetics and dynamics. Front Physiol. 2015 Apr 8;6:109. doi: 10.3389/fphys.2015.00109. PMID: 25904870; PMCID: PMC4389347.

64. Putti R, Sica R, Migliaccio V, Lionetti L. Diet impact on mitochondrial bioenergetics and dynamics. Front Physiol. 2015 Apr 8;6:109. doi: 10.3389/fphys.2015.00109. PMID: 25904870; PMCID: PMC4389347.

65. Niforou A, Konstantinidou V, Naska A. Genetic Variants Shaping Inter-individual Differences in Response to Dietary Intakes-A Narrative Review of the Case of Vitamins. Front Nutr. 2020 Dec 1;7:558598. doi: 10.3389/fnut.2020.558598. PMID: 33335908; PMCID: PMC7736113.

66. Sun H, Lin M, Russell EM, Minster RL, Chan TF, Dinh BL, Naseri T, Reupena MS, Lum-Jones A; Samoan Obesity, Lifestyle, and Genetic Adaptations (OLaGA) Study Group; Cheng I, Wilkens LR, Le Marchand L, Haiman CA, Chiang CWK. The impact of global and local Polynesian genetic ancestry on complex traits in Native Hawaiians. PLoS Genet. 2021 Feb 11;17(2):e1009273. doi: 10.1371/journal.pgen.1009273. PMID: 33571193; PMCID: PMC7877570.

67. Pacific islanders pay heavy price for abandoning traditional diet. Bull World Health Organ. 2010 Jul 1;88(7):484-5. doi: 10.2471/BLT.10.010710. PMID: 20616964; PMCID: PMC2897991.

68. Dela Cruz R, Wolfe E, Yonemori KM, Fialkowski MK, Wilkens LR, Coleman P, Lameko-Mua S, Johnson E, Gilmatam D, Sigrah C, Shomour M, Remengesau S, Alfred J, Acosta M, Ettienne R, Deenik J, Tanisha A, Salazar KA, Novotny R, Boushey CJ. Exploring Foods of the Pacific: Cultural Food Identity in the US Affiliated Pacific Region. Hawaii J Health Soc Welf. 2022 Sep;81(9):247-252. PMID: 36118154; PMCID: PMC9460762.

Figure S1. Overview for constructing and evaluating trans-ethnic polygenic score (PGS) and diet score prediction models. The estimations of trans-ethnic PGS for BMI and T2D were based on the GWAS from three ethnic groups including East Asian (EAS, representing largely EAS-like ancestry), European (EUR, representing largely EUR-like ancestry), and Native Hawaiians (NH, representing largely Polynesian, or PNS-like, ancestry) using Bayesian approaches (BridgePRS and PRS-CSx). In this case, MEC-NH individuals genotyped on MEGA array (N = 4,156-4,164) were used to train the PGS, while individuals genotyped on the GDA array (N = 1,218 − 1,226) were used as validation dataset. Diet score was trained and validated for each phenotype using three different machine learning-based algorithms, with all individuals with survey data only (N = 7026_{T2D}; 8372_{BMI}) used as training dataset, and individuals with survey and genetic data (N = 4632_{T2D} ; $5,222_{BMI}$) were used as validation dataset.

BMI			
Ancestry	**GWAS Dataset Composition**	**N**	**Source**
EAS-like	Taiwan Biobank + Biobank Japan	256,450	Chen et al, 2023
EUR-like	GIANT + UKBB	~700,000	Yengo et al, 2018
PNS-like	Native Hawaiians	4,156	Multi-Ethnic Cohort Study (MEC)
T2D			
Ancestry	**GWAS Dataset Composition**	**N**	**Source**
EAS-like	40 GWAS encompassing China, Hong Kong, Japan, Korea, Philippines, Singapore, Taiwan, USA	427,504	Suzuki et al, 2024
EUR-like	36 GWAS encompassing Denmark, Estonia, Finland, France, Germany, Greece, Iceland, Netherlands, Sweden, UK, USA, Various European countries	1,812,017	Suzuki et al, 2024
PNS-like	Native Hawaiians	4,164	Multi-Ethnic Cohort Study (MEC)

Figure S2. Summary of GWAS datasets that were used to train PGS models for BMI (top) and T2D (bottom) for Native Hawaiians.

BMI (n_{test} =1,226)				
Model	P-value	Beta ± SE	R^2 [95% CI]	Partial-R^2 [95% CI]
EAS + EUR	4.13×10^{-35}	0.39 ± 0.03	0.190 [0.160 to 0.240]	0.119 [0.084 to 0.155]
Meta (EAS+EUR)	3.40×10^{-20}	0.33 ± 0.03	0.141 [0.114 to 0.188]	0.067 [0.067 to 0.098]
Meta (EAS+EUR) + PNS	8.03×10^{-9}	0.29 ± 0.05	0.104 [0.081 to 0.149]	0.027 [0.027 to 0.049]
EAS + EUR + PNS	9.51×10^{-11}	0.38 ± 0.06	0.113 [0.089 to 0.158]	0.034 [0.016 to 0.057]

Figure S3. Comparison of trans-ethnic PGS models integrating GWAS information from two and three ancestral groups using PRS-CSx. We compared two additional approaches for constructing PGS prediction models for BMI: (1) Meta (EAS+EUR), which performs GWAS meta-analysis and then apply PRS-CS on the single GWAS summary statistics, and (2) Meta (EAS+EUR) + PNS, which we combined the meta-analyzed summary statistics from (1) with that from PNS-only for a trans-ethnic PGS model using PRS-CSx. This approach could in theory incorporate information from three separate populations.

		BMI (n_{test} =1226)			T2D (n_{test} =1218)		
Model	*GWAS*	*P-value*	*Beta ± SE*	*Partial-R^2 [95% CI]*	*P-value*	*AUC*	*Liability R^2 [95% CI]*
PRS-CS	EAS	6.00×10^{-22}	0.26 ± 0.03	0.074 [0.046 to 0.104]	2.54×10^{-8}	0.52	0.049 [0.023 to 0.086]
	EUR	1.26×10^{-27}	0.40 ± 0.04	0.093 [0.062 to 0.126]	1.16×10^{-11}	0.63	0.074 [0.040 to 0.116]
	NH	2.68×10^{-9}	0.33 ± 0.05	0.029 [0.014 to 0.051]	3.54×10^{-3}	0.56	0.013 [0.001 to 0.034]
PRS-CSx	EAS+NH	1.61×10^{-13}	0.39 ± 0.05	0.044 [0.024 to 0.069]	2.50×10^{-3}	0.53	0.014 [0.002 to 0.036]
	EUR+NH	2.12×10^{-11}	0.33 ± 0.05	0.036 [0.019 to 0.061]	4.20×10^{-8}	0.62	0.047 [0.020 to 0.082]
	EAS+EUR	4.13×10^{-35}	0.39 ± 0.03	0.119 [0.084 to 0.155]	9.18×10^{-14}	0.64	0.090 [0.055 to 0.133]
BridgePRS	EAS+NH	1.69×10^{-15}	0.22 ± 0.03	0.051 [0.030 to 0.077]	3.87×10^{-3}	0.56	0.013 [0.001 to 0.034]
	EUR+NH	2.31×10^{-19}	0.25 ± 0.03	0.065 [0.039 to 0.094]	5.41×10^{-4}	0.58	0.019 [0.003 to 0.043]
	EAS+EUR	7.16×10^{-29}	0.35 ± 0.03	0.098 [0.065 to 0.132]	4.68×10^{-14}	0.65	0.092 [0.054 to 0.138]

Table S1.   PGS accuracy on BMI and T2D across different PGS models. All BMI PGS were adjusted for the first 10 PCs. T2D PGS were adjusted for the age, sex, BMI, and first 10 PCs. PGS models were validated in held-out sample of Native Hawaiians that were not used in generating the GWAS dataset

Figure S4. Correlation among dietary variables available in the Multi-Ethnic Cohort. The correlations among (top) all 520 diet variables and (bottom) pruned (n=220) diet variables in Native Hawaiians (N = 13,595). The pruned dataset removes randomly one variable among all pairs of variables correlated with $R^2 > 0.5$.

	Distribution Pairs	Distance	95% CI	P-value
UMAP 1	Japanese vs NH:POLY >50%	0.23	0.20 to 0.26	0.0001
UMAP 2	Japanese vs NH:POLY >50%	0.17	0.15 to 0.19	0.0001
UMAP 1	Japanese vs NH:EAS >50%	0.08	0.06 to 0.10	0.0001
UMAP 2	Japanese vs NH:EAS >50%	0.12	0.09 to 0.14	0.0001
UMAP 1	NH:POLY >50% vs NH:EAS >50%	0.06	0.03 to 0.08	0.0001
UMAP 2	NH:POLY >50% vs NH:EAS >50%	0.01	-0.004 to 0.01	0.0162

	Distribution Pairs	Distance	95% CI	P-value
UMAP 1	White vs NH:POLY >50%	0.35	0.30 to 0.40	0.0001
UMAP 2	White vs NH:POLY >50%	0.52	0.47 to 0.57	0.0001
UMAP 1	White vs NH:EUR >50%	0.17	0.13 to 0.21	0.0001
UMAP 2	White vs NH:EUR >50%	0.23	0.19 to 0.27	0.0001
UMAP 1	NH:POLY >50% vs NH:EUR >50%	0.02	0.006 to 0.03	0.0141
UMAP 2	NH:POLY >50% vs NH:EUR >50%	0.08	0.04 to 0.10	0.0001

Figure S5. Distance between dietary pattern distributions between racial/ethnic groups. The Kullbak-Leibler (KL) divergence was applied to quantify the difference between dietary

distributions between racial/ethnic groups. Native Hawaiians were grouped based on their EAS, EUR, and Polynesian ancestry proportions. The KL distances were computed on the rank ordered UMAPs after bootstrapping (1000 times) and the p-values were estimated from the permutations (1000 times). The top figure shows the differential/similar dietary patterns between Japanese, Native Hawaiians with Polynesian ancestry > 50%, and Native Hawaiians with East Asian ancestry >50%. The bottom figure is the distance between dietary patterns between White participants and Native Hawaiians with European ancestry >50%, and Native Hawaiians with Polynesian ancestry >50%.

Partial R^2 [95% CI]			
	Elastic Net	Random Forest	Neural Network
Black	0.043 [0.038 to 0.047]	0.085 [0.079 to 0.091]	0.055 [0.050 to 0.061]
Japanese	0.041 [0.037 to 0.044]	0.089 [0.085 to 0.094]	0.056 [0.052 to 0.060]
Latino	0.027 [0.023 to 0.030]	0.056 [0.051 to 0.060]	0.035 [0.032 to 0.039]
White	0.071 [0.066 to 0.075]	0.117 [0.112 to 0.123]	0.078 [0.073 to 0.083]

Figure S6. Transferability of diet scores for BMI across other racial/ethnic groups in MEC participants. The partial R^2 for each diet score, based on a model adjusted for daily calorie intake is also shown in the table.

BMI				
	N	**P-value**	**Beta ± SE**	**Partial-R^2 [95% CI]**
Elastic Net	5222	$3.62\times10^{-\square1}$	0.27 ± 0.01	0.068 [0.054 to 0.081]
Random Forest	5222	$1.36\times10^{-1\square2}$	0.37 ± 0.01	0.117 [0.100 to 0.134]
Neural Network	5222	$2.17\times10^{-11\square}$	0.32 ± 0.01	0.0952 [0.0800 to 0.1110]
HEI total score	5224	$4.70\times10^{-\square}$	-0.05 ± 0.01	0.0032 [0.0009 to 0.0071]
AHEI total score	5224	0.0020	-0.04 ± 0.01	0.0018 [0.0002 to 0.0049]
DASH total score	5224	$1.90\times10^{-\square}$	-0.07 ± 0.01	0.0052 [0.0020 to 0.0101]
AMDS E total score	5224	$1.44\times10^{-\square}$	-0.07 ± 0.01	0.0053 [0.0022 to 0.0101]
AMDS total score	5224	$3.87\times10^{-\square}$	-0.07 ± 0.02	0.0041 [0.0015 to 0.0080]
Overall plant-based diet index	5224	7.50×10^{-12}	-0.09 ± 0.01	0.0090 [0.0044 to 0.0148]
Healthful plant-based diet index	5224	5.37×10^{-12}	-0.09 ± 0.01	0.0091 [0.0048 to 0.0148]
Unhealthful plant-based diet index	5224	0.5000	0.01 ± 0.01	$8.74\times10^{-\square}$ [$2.52\times10^{-\square}$ to 0.0015]
Empirical dietary index for hyperinsulinemia	5069	$1.12\times10^{-2\square}$	0.16 ± 0.01	0.0234 [0.0151 to 0.0331]
Empirical dietary inflammatory pattern	5069	$5.67\times10^{-3\square}$	0.17 ± 0.01	0.0305 [0.0210 to 0.0421]
Empirical dietary index for insulin resistance	5224	$4.26\times10^{-\square3}$	0.20 ± 0.01	0.0357 [0.0264 to 0.0471]
HEI-2015: Total vegetables	5224	0.7140	-0.0049 ± 0.01	$2.58\times10^{-\square}$ [$1.12\times10^{-\square}$ to 0.0012]
HEI-2015: Greens & beans	5224	0.0013	-0.04 ± 0.01	0.0020 [0.0004 to 0.0050]
HEI-2015: Total fruits	5224	0.0003	-0.05 ± 0.01	0.0025 [0.0005 to 0.0063]
HEI-2015: Whole fruits	5224	0.1760	-0.02 ± 0.01	0.0004 [$1.42\times10^{-\square}$ to 0.0023]
HEI-2015: Whole grains	5224	0.0021	-0.04 ± 0.01	0.0018 [0.0002 to 0.0048]
HEI-2015: Dairy	5224	0.4410	-0.01 ± 0.02	0.0001 [$4.79\times10^{-\square}$ to 0.0015]

HEI-2015: Total protein foods	5224	2.82×10⁻□	0.08 ± 0.01	0.0059 [0.0024 to 0.0103]
HEI-2015: Seafood & plant proteins	5224	0.8580	0.0026 ± 0.01	6.13×10⁻□ [1.13×10⁻□ to 0.0010]
HEI-2015: Fatty acids	5224	0.1730	0.02 ± 0.02	0.0004 [1.96×10⁻□ to 0.0019]
HEI-2015: Sodium	5224	4.75×10⁻¹²	-0.09 ± 0.01	0.0091 [0.0043 to 0.0145]
HEI-2015: Refined grains	5224	8.77×10⁻¹□	-0.10 ± 0.01	0.0106 [0.0056 to 0.0168]
HEI-2015: Saturated fats	5224	0.0103	-0.04 ± 0.02	0.0013 [7.69×10⁻□ to 0.0040]
HEI-2015: Added sugars	5224	0.0052	0.04 ± 0.01	0.0015 [9.94×10⁻□ to 0.0042]
Healthy Eating Iindex-2015: Total score	5224	2.64×10⁻□	-0.07 ± 0.01	0.0059 [0.0025 to 0.0111]
Energy-adjusted Dietary Inflammatory Index	5224	0.0087	0.03 ± 0.01	0.0013 [8.88×10⁻□ to 0.0041]
Dietary Inflammatory Index	5224	0.9130	-0.0020 ± 0.02	2.31×10⁻□ [1.57×10⁻□ to 0.0009]
EAT-Lancet Index: Vegetables	5222	0.4630	-0.01 ± 0.01	0.0001 [3.48×10⁻□ to 0.0015]
EAT-Lancet Index: Fruits	5222	0.0029	-0.04 ± 0.01	0.0017 [0.0002 to 0.0045]
EAT-Lancet Index: Unsaturated fats	5222	1.82×10⁻□	0.07 ± 0.01	0.0044 [0.0015 to 0.0086]
EAT-Lancet Index: Saturated fats	5222	3.00×10⁻□	-0.06 ± 0.01	0.0042 [0.0014 to 0.0084]
EAT-Lancet Index: Legumes (without soy)	5222	0.9460	0.0016 ± 0.02	8.74×10⁻□ [1.79×10⁻□ to 0.0010]
EAT-Lancet Index: Soy foods	5222	0.4050	0.01 ± 0.01	0.0001 [2.28×10⁻□ to 0.0015]
EAT-Lancet Index: Nuts	5222	0.0008	-0.05 ± 0.01	0.0021 [0.0003 to 0.0051]
EAT-Lancet Index: Whole grains	5222	1.27×10⁻□	-0.06 ± 0.01	0.0037 [0.0011 to 0.0079]
EAT-Lancet Index: Fish and shellfish	5222	0.0001	0.06 ± 0.02	0.0028 [0.0006 to 0.0065]
EAT-Lancet Index: Beef and lamb	5222	6.39×10⁻¹□	-0.13 ± 0.01	0.0125 [0.0075 to 0.0193]
EAT-Lancet Index: Pork	5222	1.86×10⁻¹³	-0.10 ± 0.01	0.0103 [0.0056 to 0.0166]
EAT-Lancet Index: Chicken and other poultry	5222	0.0120	-0.04 ± 0.01	0.0012 [8.02×10⁻□ to 0.0039]

EAT-Lancet Index: Eggs	5222	$1.33 \times 10^{-\square}$	-0.07 ± 0.01	0.0045 [0.0016 to 0.0091]
EAT-Lancet Index: Dairy foods	5222	0.1360	0.02 ± 0.02	0.0004 [$2.25 \times 10^{-\square}$ to 0.0023]
EAT-Lancet Index: Potatoes and Tubers	5222	0.6920	-0.01 ± 0.01	$3.02 \times 10^{-\square}$ [$1.38 \times 10^{-\square}$ to 0.0012]
EAT-Lancet Index: Added sugars	5222	$4.59 \times 10^{-\square}$	0.06 ± 0.01	0.0032 [0.0008 to 0.0067]
EAT-Lancet Index: Total score	5222	$2.74 \times 10^{-\square}$	-0.07 ± 0.01	0.0042 [0.0014 to 0.0083]

Table S2. Association of dietary scores and indices to BMI in Native Hawaiians. Evaluations were conducted only in held-out Native Hawaiians that were not used to develop the diet score in this study. The associations between dietary scores/indices and BMI were adjusted for calorie intake. We did not test the empirical lifestyle index for hyperinsulinemia and empirical lifestyle index for insulin resistance as these two indices used directly BMI as part of their model construction.

T2D				
Diet Indices/Scores	**N**	**OR [95% CI]**	**Liability R^2 [95% CI]**	**P-value**
Elastic Net (T2D)	4,632	1.13 [1.06 to 1.21]	6.04×10^{-3} [1.20×10^{-3} to 0.014]	1.65×10^{-4}
Random Forest (T2D)	4,632	1.21 [1.12 to 1.30]	0.012 [4.14×10^{-3} to 0.022]	1.96×10^{-7}
Neural Network (T2D)	4,632	1.16 [1.08 to 1.25]	8.14×10^{-3} [2.48×10^{-3} to 0.018]	1.47×10^{-5}
Elastic Net (BMI)	4,632	1.17 [1.08 to 1.26]	0.0072 [0.0017 to 0.0152]	4.03×10^{-5}
Random Forest (BMI)	4,632	1.22 [1.13 to 1.32]	0.0107 [0.0042 to 0.0207]	4.67×10^{-7}
Neural Network (BMI)	4,632	1.16 [1.08 to 1.24]	0.0072 [0.0019 to 0.0150]	3.46×10^{-5}
HEI total score	4,632	0.94 [0.88 to 1.00]	0.0015 [$1.77 \times 10^{-\square}$ to 0.0061]	0.06
AHEI total score	4,632	0.91 [0.85 to 0.97]	0.0031 [0.0003 to 0.0097]	0.01
DASH total score	4,632	0.91 [0.85 to 0.97]	0.0035 [0.0003 to 0.0097]	0.0040
AMDS E total score	4,632	0.97 [0.91 to 1.03]	0.0004 [$1.06 \times 10^{-\square}$ to 0.0037]	0.33
AMDS total score	4,632	0.98 [0.91 to 1.06]	0.0001 [$3.95 \times 10^{-\square}$ to 0.0025]	0.61
Overall plant-based diet index	4,632	1.01 [0.94 to 1.08]	$1.30 \times 10^{-\square}$ [$2.33 \times 10^{-\square}$ to 0.0025]	0.86
Healthful plant-based diet index	4,632	0.97 [0.91 to 1.03]	0.0004 [$1.19 \times 10^{-\square}$ to 0.0036]	0.31
Unhealthful plant-based diet index	4,632	1.04 [0.98 to 1.11]	0.0007 [$2.10 \times 10^{-\square}$ to 0.0042]	0.21
Empirical dietary index for hyperinsulinemia	4,632	1.09 [1.01 to 1.17]	0.0022 [$5.92 \times 10^{-\square}$ to 0.0076]	0.02
Empirical dietary inflammatory pattern	4,632	1.13 [1.05 to 1.22]	0.0052 [0.0008 to 0.0128]	0.0006
Empirical dietary index for insulin resistance	4,632	1.11 [1.02 to 1.23]	0.0035 [0.0002 to 0.0108]	0.01
Empirical lifestyle index for hyperinsulinemia	4,632	1.18 [1.03 to 1.35]	0.0025 [$8.63 \times 10^{-\square}$ to 0.0083]	0.02
Empirical lifestyle index for insulin resistance	4,632	1.26 [1.13 to 1.43]	0.0059 [0.0015 to 0.0130]	0.0003
HEI-2015: Total vegetables	4,632	1.00 [0.94 to 1.07]	$1.34 \times 10^{-\square}$ [$3.29 \times 10^{-\square}$ to 0.0021]	0.96

HEI-2015: Greens & beans	4,632	0.99 [0.93 to 1.06]	$3.97\times10^{-\square}$ [$5.13\times10^{-\square}$ to 0.0021]	0.76
HEI-2015: Total fruits	4,632	0.94 [0.89 to 1.00]	0.0016 [$1.41\times10^{-\square}$ to 0.0064]	0.05
HEI-2015: Whole fruits	4,632	0.97 [0.92 to 1.03]	0.0004 [$8.57\times10^{-\square}$ to 0.0036]	0.31
HEI-2015: Whole grains	4,632	0.95 [0.90 to 1.02]	0.0009 [$4.75\times10^{-\square}$ to 0.0044]	0.15
HEI-2015: Dairy	4,632	0.99 [0.92 to 1.06]	$5.67\times10^{-\square}$ [$7.43\times10^{-\square}$ to 0.0022]	0.71
HEI-2015: Total protein foods	4,632	0.97 [0.91 to 1.04]	0.0004 [$7.21\times10^{-\square}$ to 0.0033]	0.32
HEI-2015: Seafood & plant proteins	4,632	0.96 [0.90 to 1.02]	0.0007 [$2.66\times10^{-\square}$ to 0.0042]	0.21
HEI-2015: Fatty acids	4,632	1.03 [0.96 to 1.11]	0.0003 [$7.59\times10^{-\square}$ to 0.0031]	0.40
HEI-2015: Sodium	4,632	0.92 [0.86 to 0.98]	0.0028 [0.0002 to 0.0088]	0.01
HEI-2015: Refined grains	4,632	0.97 [0.91 to 1.04]	0.0003 [$7.11\times10^{-\square}$ to 0.0031]	0.37
HEI-2015: Saturated fats	4,632	1.03 [0.95 to 1.11]	0.0002 [$8.62\times10^{-\square}$ to 0.0029]	0.46
HEI-2015: Added sugars	4,632	0.95 [0.90 to 1.01]	0.0011 [$5.65\times10^{-\square}$ to 0.0054]	0.10
Healthy Eating lindex-2015: Total score	4,632	0.93 [0.87 to 0.99]	0.0020 [$3.69\times10^{-\square}$ to 0.0074]	0.03
Energy-adjusted Dietary Inflammatory Index	4,632	1.05 [0.99 to 1.11]	0.0009 [$2.03\times10^{-\square}$ to 0.0047]	0.14
Dietary Inflammatory Index	4,632	1.03 [0.95 to 1.12]	0.0002 [$8.72\times10^{-\square}$ to 0.0029]	0.54
EAT-Lancet Index: Vegetables	4,632	1.04 [0.97 to 1.11]	0.0005 [$1.43\times10^{-\square}$ to 0.0042]	0.26
EAT-Lancet Index: Fruits	4,632	0.97 [0.92 to 1.04]	0.0003 [$2.03\times10^{-\square}$ to 0.0034]	0.38
EAT-Lancet Index: Unsaturated fats	4,632	0.97 [0.91 to 1.03]	0.0004 [$1.01\times10^{-\square}$ to 0.0034]	0.31
EAT-Lancet Index: Saturated fats	4,632	1.04 [0.97 to 1.10]	0.0005 [$1.48\times10^{-\square}$ to 0.0037]	0.27
EAT-Lancet Index: Legumes (without soy)	4,632	1.04 [0.93 to 1.16]	0.0002 [$8.46\times10^{-\square}$ to 0.0029]	0.51
EAT-Lancet Index: Soy foods	4,632	1.05 [0.99 to 1.12]	0.0010 [$3.04\times10^{-\square}$ to 0.0049]	0.13
EAT-Lancet Index: Nuts	4,632	0.93 [0.86 to 0.99]	0.0018 [$2.30\times10^{-\square}$ to 0.0072]	0.04

EAT-Lancet Index: Whole grains	4,632	0.91 [0.84 to 0.98]	0.0029 [0.0002 to 0.0090]	0.01
EAT-Lancet Index: Fish and shellfish	4,632	0.99 [0.93 to 1.08]	$1.61\times10^{-\square}$ [$3.55\times10^{-\square}$ to 0.0020]	0.98
EAT-Lancet Index: Beef and lamb	4,632	1.01 [0.93 to 1.09]	$1.10\times10^{-\square}$ [$4.91\times10^{-\square}$ to 0.0023]	0.87
EAT-Lancet Index: Pork	4,632	0.96 [0.90 to 1.03]	0.0006 [$2.17\times10^{-\square}$ to 0.0041]	0.25
EAT-Lancet Index: Chicken and other poultry	4,632	1.03 [0.96 to 1.11]	0.0003 [$1.83\times10^{-\square}$ to 0.0035]	0.36
EAT-Lancet Index: Eggs	4,632	0.98 [0.91 to 1.05]	0.0001 [$7.10\times10^{-\square}$ to 0.0030]	0.58
EAT-Lancet Index: Dairy foods	4,632	1.02 [0.94 to 1.09]	$8.40\times10^{-\square}$ [$6.34\times10^{-\square}$ to 0.0025]	0.66
EAT-Lancet Index: Potatoes and Tubers	4,632	0.96 [0.90 to 1.03]	0.0005 [$2.52\times10^{-\square}$ to 0.0036]	0.29
EAT-Lancet Index: Added sugars	4,632	0.98 [0.91 to 1.04]	0.0002 [$6.71\times10^{-\square}$ to 0.0032]	0.45
EAT-Lancet Index: Total score	4,632	0.98 [0.92 to 1.05]	0.0001 [$8.09\times10^{-\square}$ to 0.0026]	0.59

Table S3. Association of dietary scores and indices to T2D in Native Hawaiians. Evaluations were conducted only in held-out Native Hawaiians that were not used to develop the diet score in this study. The associations between dietary scores/indices and T2D were adjusted for age, sex, BMI, and daily calorie intake.

Subjects by Geographical Location:　　　▲ Subjects in Hawaii　　　▲ Subjects in Los Angeles

Figure S7. Dietary patterns/behaviors across all MEC participants by geographical location and race/ethnicity. We applied uniform manifold approximation and projection (UMAP) to visualize the diet/nutrition variability using the first 10 principal components of dietary consumption data based on 220 diet variables. The UMAPs on the top left illustrates dietary patterns by recruitment area (Hawaii vs Los Angeles). The remaining UMAPs display dietary patterns by self-identified race/ethnicity (Black=B, Native Hawaiians= H, Japanese=J, Latino/Hispanic = L, White=W) in all MEC participants (top right), or those recruited in Los Angeles (bottom left), and Hawaii (bottom right).

Phenotype	P_value	N_Case	N_Control	OR (95% CI)
Alzheimer_dementia	1.26e-01	635	2858	1.09 [0.98 to 1.21]
Alzheimer_ever	2.85e-01	269	3197	1.09 [0.93 to 1.28]
Ami_ever	3.31e-02	249	3227	1.17 [1.00 to 1.36]
Anemia_ever	3.76e-03	1760	1800	1.12 [1.03 to 1.21]
Asthma_ever	3.46e-01	703	2798	1.05 [0.95 to 1.15]
Atrial_Fibrillation_ever	6.59e-01	714	2787	0.98 [0.89 to 1.08]
Cataract_ever	9.89e-01	2009	1529	1.00 [0.93 to 1.08]
Chronic_Kidney_dz	1.82e-02	289	2668	1.18 [1.03 to 1.36]
Chronic_Kidney	3.01e-06	1556	1989	1.20 [1.12 to 1.31]
COPD	6.28e-01	806	2696	1.02 [0.93 to 1.12]
Depression	5.94e-02	593	2891	1.10 [0.99 to 1.22]
Gout	1.88e-02	1061	2370	1.11 [1.02 to 1.21]
H.br	9.94e-01	317	2377	1.00 [0.88 to 1.13]
H_crc	3.40e-01	99	4180	0.90 [0.72 to 1.12]
H_lung	7.67e-01	153	4212	1.03 [0.85 to 1.22]
H_pc	2.63e-01	309	1835	1.08 [0.95 to 1.22]
Hyperl	7.83e-01	659	2821	1.02 [0.89 to 1.14]
Hyperp	7.83e-01	659	2821	1.02 [0.89 to 1.14]
Hypert	1.72e-02	2772	891	1.11 [1.01 to 1.22]
Hypoth	2.93e-01	588	2895	1.06 [0.95 to 1.18]
Ischemic_heart	1.33e-04	1627	1944	1.17 [1.08 to 1.27]
Osteoporosis	2.67e-01	521	2873	0.93 [0.83 to 1.06]
RA_OA	9.58e-01	1398	2106	1.00 [0.93 to 1.08]
Stroke_tia	7.14e-04	561	2921	1.20 [1.08 to 1.34]

Figure S8. Association of BMI diet score with other binary disease outcome in Native Hawaiians. We defined cases and controls for each disease based on self-reported disease status from the baseline questionnaire (question: "Has your doctor ever told you that you had disease?). The associations were assessed after adjusting for age, sex, BMI, and first 10 PCs.

Mediator: CRP					
Disease	N			Estimate [95% CI]	P-value
Anemia	1283		Total Effect	0.0132 [-0.0154 to 0.0489]	0.42
			ACME	0.0002 [-0.0012 to 0.0026]	0.82
			ADE	0.0130 [-0.0155 to 0.0496]	0.44
			Prop. Mediated	0.0136 [-1.1520 to 0.5578]	0.92
Chronic Kidney Disease	1273		Total Effect	0.0299 [0.0049 to 0.0664]	0.04
			ACME	0.0010 [-0.0007 to 0.0041]	0.28
			ADE	0.0289 [0.0033 to 0.0642]	0.04
			Prop. Mediated	0.0341 [-0.0363 to 0.2247]	0.32
Ischemic Heart Disease	1290		Total Effect	0.0096 [-0.0195 to 0.0387]	0.60
			ACME	0.0011 [-0.0005 to 0.0038]	0.24
			ADE	0.0085 [-0.0210 to 0.0379]	0.64
			Prop. Mediated	0.1162 [-1.3122 to 0.6877]	0.72
Stroke	1245		Total Effect	0.0373 [0.0089 to 0.0652]	<0.0001
			ACME	-0.0009 [-0.0026 to 0.0005]	0.22
			ADE	0.0382 [0.0092 to 0.0663]	<0.0001
			Prop. Mediated	-0.0247 [-0.1066 to 0.0147]	0.22
T2D	1879		Total Effect	0.0319 [0.0092 to 0.0604]	<0.0001
			ACME	0.0022 [5.9208×10^{-5} to 0.005]	0.06
			ADE	0.0297 [0.0059 to 0.0593]	0.02
			Prop. Mediated	0.0068 [0.0282 to 0.3688]	0.06
Mediator: LDL					
Disease	N			Estimate [95% CI]	P-value
Anemia	1277		Total Effect	0.0225 [-0.0119 to 0.0518]	0.12
			ACME	-1.5875×10^{-5} [-7.7907×10^{-4} to 0.0010]	0.86
			ADE	0.0225 [-0.0123 to 0.0518]	0.12
			Prop. Mediated	-7.0475×10^{-4} [-0.0289 to 0.0643]	0.94
Chronic Kidney Disease	1268		Total Effect	0.0444 [0.0144 to 0.0728]	<0.0001
			ACME	-0.0001 [-0.0024 to 0.0015]	0.92

			Estimate [95% CI]	P-value
		ADE	0.0445 [0.0148 to 0.0719]	<0.0001
		Prop. Mediated	-0.0033 [-0.0871 to 0.0283]	0.92
Ischemic Heart Disease	1285	Total Effect	0.0212 [-0.0051 to 0.0477]	0.17
		ACME	1.5875×10^{-5} [-9.9328×10^{-4} to 0.0011]	0.98
		ADE	0.0212 [-0.0052 to 0.0481]	0.14
		Prop. Mediated	9.1195 [-0.1737 to 0.2254]	1.00
Stroke	1248	Total Effect	0.0401 [0.0116 to 0.0667]	<0.0001
		ACME	-2.3742×10^{-6} [-5.9467×10^{-4} to 9.1074×10^{-4}]	0.84
		ADE	0.0401 [0.0114 to 0.0667]	<0.0001
		Prop. Mediated	-5.9182×10^{-5} [-0.0119 to 0.0431]	0.84
T2D	1806	Total Effect	0.0318 [0.0179 to 0.0607]	0.02
		ACME	-4.6180×10^{-5} [-4.253×10^{-4} to 5.9714×10^{-4}]	0.96
		ADE	0.0319 [0.0184 to 0.0609]	0.02
		Prop. Mediated	-0.0015 [-0.0238 to 0.0166]	0.98
Mediator: HDL				
Disease	**N**		**Estimate [95% CI]**	**P-value**
Anemia	1282	Total Effect	0.0269 [-0.0087 to 0.0527]	0.18
		ACME	-0.0009 [-0.0049 to 0.0020]	0.56
		ADE	0.0278 [-0.0081 to 0.0535]	0.18
		Prop. Mediated	-0.0338 [-0.9396 to 0.3865]	0.66
Chronic Kidney Disease	1274	Total Effect	0.0467 [0.0206 to 0.0771]	<0.001
		ACME	0.0028 [0.0003 to 0.0062]	0.02
		ADE	0.0439 [0.0192 to 0.0748]	<0.001
		Prop. Mediated	0.0603 [0.0076 to 0.1533]	0.02
Ischemic Heart Disease	1291	Total Effect	0.0222 [-0.0128 to 0.0544]	0.24
		ACME	0.0014 [-0.0006 to 0.0047]	0.20
		ADE	0.0208 [-0.0151 to 0.0521]	0.24
		Prop. Mediated	0.0650 [-0.4302 to 0.7218]	0.32
Stroke	1253	Total Effect	0.0406 [0.0163 to 0.0649]	0.02

Disease	N		Estimate [95% CI]	P-value
		ACME	0.0020 [-5.8239×10⁻⁶ to 0.0054]	0.06
		ADE	0.0387 [0.0118 to 0.0637]	0.02
		Prop. Mediated	0.0482 [-0.0044 to 0.1770]	0.08
T2D	1816	Total Effect	0.0367 [0.0117 to 0.0652]	<0.0001
		ACME	0.0014 [-0.0011 to 0.0034]	0.36
		ADE	0.0353 [0.0100 to 0.0637]	<0.0001
		Prop. Mediated	0.0372 [-0.0262 to 0.1597]	0.36
Mediator: Triglyceride				
Disease	*N*		*Estimate [95% CI]*	*P-value*
Anemia	1282	Total Effect	0.0212 [-0.0060 to 0.0511]	0.20
		ACME	0.0009 [-0.0014 to 0.0044]	0.34
		ADE	0.0204 [-0.0079 to 0.0499]	0.20
		Prop. Mediated	0.0408 [-0.3188 to 0.5523]	0.46
Chronic Kidney Disease	1272	Total Effect	0.0447 [0.0198 to 0.0808]	<0.0001
		ACME	0.0016 [-0.0002 to 0.0040]	0.16
		ADE	0.0432 [0.0164 to 0.0792]	<0.0001
		Prop. Mediated	0.0352 [-0.0038 to 0.1335]	0.16
Ischemic Heart Disease	1289	Total Effect	0.0211 [-0.0098 to 0.0507]	0.16
		ACME	0.0016 [-0.0004 to 0.0041]	0.24
		ADE	0.0196 [-0.0111 to 0.0495]	0.16
		Prop. Mediated	0.0735 [-0.1750 to 0.3818]	0.36
Stroke	1252	Total Effect	0.0389 [0.0151 to 0.0675]	<0.0001
		ACME	0.0008 [-0.0007 to 0.0027]	0.32
		ADE	0.0381 [0.0141 to 0.0669]	<0.0001
		Prop. Mediated	0.0206 [-0.0244 to 0.0842]	0.32
T2D	1814	Total Effect	0.0336 [0.0124 to 0.0615]	0.02
		ACME	0.0042 [0.0004 to 0.0082]	0.02
		ADE	0.0294 [0.0093 to 0.0575]	0.06
		Prop. Mediated	0.1256 [0.0095 to 0.3515]	0.04

Table S4. Mediation analysis of BMI diet score and diseases in Native Hawaiians. Biomarkers that may reflect inflammatory (C-reactive protein, CRP) and lipid pathways (low-density lipoproteins, LDL, high-density lipoproteins, HDL, and triglycerides) were used to explore the potential mediated/indirect effects of BMI diet score on other phenotypes. For each disease and mediator, mediation analysis was conducted to estimate total effect, which sums up direct and indirect effect of diet score, average causal mediated/indirect effect (ACME) of biomarkers, average direct effect (ADE) of diet score, and proportion of mediated effect.

Hyperparameter name	Candidate values	Optimal value (BMI)	Optimal value (T2D)
Batch size	16, 32, 64,128	64	32
Optimizer	Adam, RMSProp, AdaMax, Adadelta, Adagrad, Ftrl	RMSProp	RMSProp
Loss function	None	MSE	Binary cross entropy
Activation function (hidden layers)	relu, gelu, swish, LeakyReLU, elu, selu, PReLU, tanh, softplus	relu	relu
Activation function (output layer)	None	linear	sigmoid
Number of neurons (hidden layers)	(16, 16, 16), (32, 32, 32), (64, 64, 64), (32, 32, 32, 32, 32, 32), (64, 64, 64, 64, 64, 64), (32, 32, 32, 32, 32, 32, 32, 32), (64, 64, 64, 64, 64, 64, 64, 64)	(32, 32, 32, 32, 32, 32)	(64, 64, 64, 64, 64, 64)
Number of neurons (output layer)	None	1	2

Table S5. The candidate and optimal values of the hyperparameters of the neural network models for predicting BMI and T2D. Column Optimal value (BMI) and Optimal value (T2D) show the values of the hyperparameters in the models with the lowest validation loss for predicting BMI and T2D, respectively. None in Column Candidate values indicates that the corresponding optimal hyperparameters were determined by following common practices. For example, for predicting binary-valued T2D, the activation function and number of neurons at the output layer are commonly set to Binary cross-entropy and 2, respectively.