Gestational Age as a Predictor of Oxygen Needs and ABG Patterns in Preterm Neonates with RDS: Evidence from a Resource-Limited NICU in Pakistan

Gestational Age and Oxygen Needs in Preterm.

Shoaib Mehmood[1], Ruhamah Yousaf, PhD[2]

[1]Department of Health Professional Technologies (DHPT), Faculty of Allied Health Sciences (FAHS), The University of Lahore, Pakistan.
[2]Senior Lecturer, Department of Health Professional Technologies (DHPT), Faculty of Allied Health Sciences (FAHS), The University of Lahore, Pakistan.

Corresponding Author:
Ruhamah Yousaf, PhD
Senior Lecturer
Department of Health Professional Technologies (DHPT), Faculty of Allied Health Sciences (FAHS), The University of Lahore, Pakistan.
Email: ruhamah2009@yahoo.com

ABSTRACT

Background:

Respiratory Distress Syndrome (RDS) remains a leading cause of morbidity and mortality among premature infants. Their immature lungs lack surfactant, making breathing difficult. These infants need close monitoring with arterial blood gas (ABG) checks to guide oxygen therapy. While gestational age is known to influence disease severity, locally relevant data on oxygen requirements and arterial blood gas (ABG) changes across gestational age groups remain scarce in Pakistan.

Objective:

To determine the association of gestational age with oxygen requirements, ABG parameters, and respiratory support in preterm neonates with RDS in a resource-limited NICU, and to explore their implications for clinical management.

Methodology:

This four-month cross-sectional study was conducted in the NICU of the University of Lahore Teaching Hospital, Pakistan. Sixty-five preterm neonates (28–36 weeks) with RDS were enrolled through consecutive sampling. Data on gestational age, oxygen requirement (FiO₂), mode of respiratory support, and ABG parameters were collected before and after stabilization on respiratory support. Data were analysed in SPSS 27 using ANOVA, Kruskal–Wallis, Chi-square, paired t-tests, and Spearman's correlation.

Results:

Lower gestational age was linked to higher oxygen needs (FiO_2 83% at 28–30 weeks vs. 55% at 34–36 weeks, p < 0.001) and more invasive ventilation (60% vs. 10%, p = 0.011). ABG parameters correlated significantly with gestational age: earlier gestations had lower pH, higher $PaCO_2$, and lower PaO_2 and SaO_2 (all p < 0.05), while HCO_3 showed no correlation (p = 0.316). Despite stabilization, SaO_2 values remained below the target 90–94% range in all groups.

Conclusion:

Gestational age is a key predictor of oxygen needs, ABG derangements, and intensity of respiratory support in preterm neonates with RDS. This first locally relevant evidence from Pakistan underscores the need for gestation-specific oxygen protocols to improve care in resource-limited NICUs and similar low-resource settings worldwide.

Key words:

Respiratory Distress Syndrome, Gestational Age, Preterm neonates, Oxygen therapy, Arterial blood gas (ABG), Neonatal intensive care

Key Messages

What is already known on this topic

Respiratory distress syndrome is a leading cause of illness and death in preterm babies. Oxygen therapy and arterial blood gas monitoring are central to care, but both under- and over-treatment can cause harm. Gestational age strongly influences oxygen needs and outcomes, yet data from local NICUs remain limited.

What this study adds

This study shows that babies born at lower gestational ages need higher FiO_2 and more invasive support. Blood gas values such as pH, PaO_2, $PaCO_2$, and SaO_2 vary significantly with gestational age, while HCO_3 does not. It also provides locally relevant evidence from Pakistan to guide neonatal care.

How this study might affect research, practice or policy

These findings highlight the need for gestation-specific oxygen therapy and monitoring strategies in NICUs. They support the development of standardized oxygen protocols in resource-limited settings. The results may also inform future research to improve outcomes and reduce preventable complications in preterm infants.

INTRODUCTION

Respiratory distress syndrome (RDS) is a major cause of morbidity and mortality among preterm neonates. It results from pulmonary immaturity and insufficient surfactant, leading to alveolar collapse, impaired oxygenation, and respiratory failure. Infants born before 36 weeks are at greatest risk, although antenatal corticosteroids, surfactant therapy, and modern respiratory support have improved survival.(1, 2)

The pathophysiology of RDS involves alveolar collapse due to surfactant deficiency, which increases the work of breathing and reduces oxygen delivery to tissues. Many affected infants require supplemental oxygen, continuous positive airway pressure (CPAP), or invasive ventilation in severe cases. However, oxygen therapy carries risks. Hypoxia can result in organ dysfunction, while hyperoxia contributes to complications such as bronchopulmonary dysplasia (BPD), retinopathy of prematurity (ROP), and impaired neurodevelopment. Therefore, careful titration of FiO_2 with close arterial blood gas (ABG) monitoring is essential to balance safety and efficacy.(3)

Gestational age strongly influences the severity of RDS and the level of support required. Extremely preterm infants often require higher FiO_2, prolonged respiratory support, and demonstrate more deranged ABG values compared with more mature preterm.(4) Non-invasive strategies, particularly CPAP and high-flow nasal cannula (HFNC), are now widely preferred to reduce ventilator-induced lung injury.(5) In addition, less invasive surfactant administration techniques, such as thin-catheter delivery, have shown promise in reducing the need for intubation and improving outcomes.(6) In resource-limited settings where surfactant is scarce, CPAP has proven to be an effective first-line therapy and improves survival when initiated early.(7)

Management guidelines now recommend targeting oxygen saturation ranges between 90–95% to avoid both hypoxia and hyperoxia. This requires continuous FiO_2 adjustment and ABG evaluation.(8) Evidence shows that very preterm infants (<28 weeks) not only require higher oxygen concentrations initially but also take longer to stabilize compared with later gestations.(9) ABG parameters, including pH, $PaCO_2$, PaO_2, HCO_3, and SaO_2, remain critical indicators of respiratory and metabolic status, guiding therapy and predicting outcomes.(10)

Globally, preterm birth affects nearly 15 million infants each year, with a large proportion at risk of RDS. The burden is higher in low-resource countries where advanced interventions remain limited.(11) Despite improvements, oxygen-related complications continue to be a major concern.(12) Antenatal corticosteroids remain a cornerstone for promoting lung maturation and reducing disease severity.(13)

In Pakistan and other low- and middle-income countries, standardized oxygen protocols and advanced monitoring are not always available. There is limited local data on how oxygen requirements, ABG parameters, and respiratory support vary by gestational age in neonates

with RDS. Addressing this evidence gap is essential to optimize oxygen therapy and reduce preventable complications.

Therefore, this study was conducted to compare oxygen requirements, ABG parameters, and respiratory support across gestational age groups in preterm neonates with RDS. By providing locally relevant data, it aims to support individualized, evidence-based neonatal care and improve outcomes in resource-constrained NICUs.

METHODOLOGY

Study Design and Setting:

This hospital-based, observational analytical study with a comparative cross-sectional design was conducted in the Neonatal Intensive Care Unit (NICU) of University of Lahore Teaching Hospital, a tertiary care centre for preterm neonates with respiratory distress syndrome (RDS).

Participants:

Preterm neonates (<37 weeks gestation) diagnosed with RDS, admitted within 24 hours of birth, and requiring oxygen therapy with available arterial blood gas (ABG) analysis were included. Neonates with major congenital anomalies, sepsis, or birth asphyxia were excluded.

Sample Size and Sampling:

The sample size was calculated using a correlation-based formula:(14)

$$N = (Z\alpha + Z\beta \ / \ C)^2 + 3, \ C = 0.5 \times \ln 1 - r/1 + r$$

where $Z_\alpha = 1.96(\alpha = 0.05)$, $Z_\beta = 0.84$(power = 80%), and r = 0.35 (expected correlation). Based on this, 62 neonates were required; 65 were enrolled using consecutive sampling to account for potential exclusions.(14, 15)

Data Collection:

Demographic details, gestational age, oxygen delivery device, FiO₂, and ABG parameters (pH, PaO₂, PaCO₂, HCO₃, SaO₂) were extracted from medical records using a structured questionnaire (attached). ABGs were collected just before starting respiratory support and after stabilization. Gestational age was the independent variable; FiO₂, oxygen delivery device, and ABG parameters were dependent variables.

Statistical Analysis:

Continuous variables were summarized as mean ± SD and categorical variables as frequencies (%). Comparisons across gestational age groups used ANOVA or Kruskal-Wallis tests. Paired t-tests compared pre- and post-support values. Correlations between gestational

age and oxygen/ABG parameters were assessed using Pearson or Spearman correlation. A p-value <0.05 was considered statistically significant.

Patient and Public Involvement

Patients and/or the public were not involved in the design, conduct, reporting, or dissemination plans of this research.

RESULTS

A total of 65 preterm neonates with respiratory distress syndrome (RDS) were enrolled in the study. Among them, the largest proportion belonged to the 31–33-week gestational age group (38.5%), followed by 28–30 weeks (30.8%) and 34–36 weeks (30.8%). Males constituted 56.9% of the cohort, while 43.1% were females. Most neonates had low birth weight (60.0%), and 30.8% were very low birth weight. Caesarean delivery was the predominant mode of birth (73.8%), and 63.1% of neonates were inborn. Antenatal corticosteroid administration was reported in 78.5% of cases, and 63.1% received surfactant therapy. The 5-minute Apgar score was ≥ 7 in 60.0% of neonates.

Analysis of oxygen requirements across gestational age groups revealed that neonates of lower gestational age required significantly higher oxygen concentrations. The mean maximum FiO_2 was $83.1 \pm 18.5\%$ in the 28–30-week group, $68.5 \pm 20.4\%$ in the 31–33-week group, and $55.6 \pm 16.5\%$ in the 34–36-week group ($p < 0.001$). Similarly, the mean minimum FiO_2 decreased with advancing gestational age ($57.1 \pm 9.8\%$, $45.2 \pm 13.7\%$, and $37.3 \pm 9.8\%$, respectively; $p < 0.001$).

Baseline (pre-support) arterial blood gas (ABG) analysis showed mild respiratory acidosis in all groups, with a mean pH of 7.22 ± 0.03 in the 28–30-week group and 7.24 ± 0.04 in the 34–36-week group. $PaCO_2$ and PaO_2 values also improved with increasing gestational age. After initiation of respiratory support, all groups showed marked improvement in ABG parameters. The mean pH increased to 7.28–7.31, $PaCO_2$ decreased, and both PaO_2 and SaO_2 showed significant increases ($p < 0.001$ for within-group changes). However, no significant differences were observed in the magnitude of change (ΔpH, $\Delta PaCO_2$, ΔPaO_2, ΔHCO_3, or ΔSaO_2) among the three gestational groups (ANOVA, $p > 0.05$).

Spearman correlation analysis revealed a strong negative correlation between gestational age and both maximum ($\rho = -0.558$, $p < 0.001$) and minimum ($\rho = -0.571$, $p < 0.001$) FiO_2 requirements, indicating that younger gestational age was associated with higher oxygen demand. Moderate positive correlations were observed between gestational age and average pH ($\rho = 0.377$, $p = 0.002$), and weak positive correlations with PaO_2 ($\rho = 0.251$, $p = 0.044$) and SaO_2 ($\rho = 0.274$, $p = 0.027$). A moderate negative correlation existed between gestational age and $PaCO_2$ ($\rho = -0.464$, $p < 0.001$), while no significant correlation was found with HCO_3.

Regarding modes of respiratory support, 40% of neonates were managed with CPAP, 35.4% required mechanical ventilation, and 24.6% received oxygen via nasal cannula. The distribution of support types differed significantly across gestational age groups (p = 0.011), with the youngest infants (28–30 weeks) more frequently requiring mechanical ventilation, whereas those of higher gestational age were more commonly managed with CPAP or nasal cannula.

Overall, oxygenation status improved after respiratory support in all gestational categories, as reflected by increases in PaO_2 and SaO_2 levels. Neonates of lower gestational age required higher initial FiO_2 and demonstrated greater variability in oxygen requirements, while more mature preterm neonates achieved target saturations with lower oxygen concentrations.

DISCUSSION

This study demonstrated that gestational age strongly influences respiratory management in preterm infants with respiratory distress syndrome (RDS). Neonates born at 28–30 weeks required substantially higher FiO_2 and were more frequently managed with mechanical ventilation, whereas those at 34–36 weeks responded better to non-invasive support. These findings are consistent with earlier reports from high-resource countries, which have shown that lung immaturity and surfactant deficiency in very preterm infants increase oxygen needs and the likelihood of invasive ventilation.(3, 4, 7) However, unlike studies from high-income settings where surfactant use is routine and non-invasive support is increasingly successful, our cohort demonstrated a higher proportion of very preterm neonates requiring intubation and mechanical ventilation, reflecting differences in treatment resources and protocols in Pakistan.

Arterial blood gas parameters improved significantly after initiation of support, with increases in pH, PaO_2, and SaO_2 and a decline in $PaCO_2$, although the magnitude of change did not differ by gestational age. This suggests that oxygen therapy is beneficial across maturity levels when titrated appropriately, in line with earlier reports.(1, 8) yet the persistence of suboptimal SaO_2 values (<90%) across all gestational age groups contrasts with findings from centers equipped with advanced monitoring and oxygen blending devices. This highlights the practical challenges of achieving recommended saturation targets (90–94%) in resource-limited NICUs.

Our results therefore extend existing knowledge by providing the first evidence from Pakistan that gestational age not only predicts oxygen and ventilation needs but also influences the success of stabilization under conditions where surfactant, high-flow nasal cannula, and advanced monitoring are not routinely available. These insights are particularly relevant for low- and middle-income countries, where neonatal care guidelines must adapt to resource constraints.

This study has several limitations. It was conducted in a single centre with a relatively small cohort, which may limit generalizability. Only three modes of respiratory support (CPAP, mechanical ventilation, and nasal cannula) were assessed, while high-flow nasal cannula was

not included. Although variables such as birth weight, antenatal steroid use, and surfactant therapy were recorded, they were not analysed in detail as potential confounders. Other important factors, such as neonatal sepsis, were also not included. In addition, long-term outcomes like bronchopulmonary dysplasia or neurodevelopment were not evaluated. As with all observational studies, residual confounding cannot be excluded. Future multicentre studies with larger samples, inclusion of additional clinical variables, and follow-up of long-term outcomes are needed to strengthen these findings.

CONCLUSION

Gestational age is a key determinant of oxygen requirements and respiratory support in preterm neonates with RDS. Infants at earlier gestations required higher FiO_2 and more invasive ventilation, whereas those closer to term were more often stabilized with non-invasive support. Most ABG parameters (pH, $PaCO_2$, PaO_2, SaO_2) varied with gestational age, while HCO_3 did not, indicating that metabolic compensation was relatively uniform across groups. This study provides the first locally relevant evidence from Pakistan, demonstrating how gestational age can be used to anticipate oxygen needs and guide respiratory management in resource-limited NICUs. These findings highlight the importance of developing gestation-specific oxygen protocols to reduce the risks of both hypoxemia and hyperoxia. Future multicentre studies with larger cohorts are warranted to validate these results and inform national and regional neonatal care guidelines.

Acknowledgment:

The authors would like to thank the Neonatal Intensive Care Unit (NICU) staff of the University of Lahore Teaching Hospital, Lahore, Pakistan, for their cooperation during data collection and clinical support. The encouragement of colleagues and family members during manuscript preparation is also gratefully acknowledged.

Funding:

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Waiver / Discount Request (APC)

I respectfully request consideration for an APC waiver/discount as I am a student researcher from Pakistan, which is a lower-middle income country according to World Bank classification, and I do not have institutional or external funding to cover publication costs

Competing interests

The authors declare no competing interests.

Ethics approval

This study was approved by the Departmental Review Board of The University of Lahore, Pakistan approval letter no. *Centre/Admin FAHS/195/25*.

Patient and public involvement statement

Patients or the public were not involved in the design, conduct, reporting, or dissemination plans of this research.

Author contributions
SM: Conceptualization, data collection, data analysis, drafting of the manuscript.
RY: Supervision, methodology guidance, critical revision of the manuscript.
Both authors approved the final version and are accountable for the integrity of the work.

Data availability statement
The data that support the findings of this study are available from the corresponding author upon reasonable request. Due to patient confidentiality and institutional restrictions, the data are not publicly available.

Tables

Table 1: Distribution of Study Participants by Gestational Age Group

Gestational Age Group (weeks)	Frequency (n)	Percent (%)
28-30	20	30.8
31-33	25	38.5
34-36	20	30.8
Total	**65**	**100.0**

Values are presented as frequency (n) and percentage (%). GA = Gestational Age.

Table 2: Baseline Characteristics of Study Participants

Variable	Category	Frequency (n)	Percent (%)
Sex	Female	28	43.1
	Male	37	56.9
	Total	**65**	**100**
Birth Weight	VLBW (<1500 g)	20	30.8
	LBW (1500-2499 g)	39	60.0
	Normal (≥2500 g)	6	9.2
	Total	**65**	**100**
Delivery Mode	Caesarean	48	73.8
	Vaginal	17	26.2
	Total	**65**	**100.0**
Place of birth	Inborn	41	63.1
	Outborn	24	36.9
	Total	**65**	**100.0**
Antenatal Corticosteroids	No	14	21.5
	Yes	51	78.5
	Total	**65**	**100.0**
Surfactant	No	24	36.9
	Yes	41	63.1
	Total	**65**	**100.0**

5min APGAR	<7	26	40.0
	≥7	39	60.0
	Total	**65**	**100.0**

Values are expressed as frequency (n) and percentage (%). VLBW = Very low birth weight (<1500 g); LBW = Low birth weight (1500–2499 g). Apgar scores are reported at 5 minutes after birth.

Table 3: Comparison of Maximum and Minimum FiO₂ Requirements Across Gestational Age Groups (Kruskal–Wallis Test)

Gestational Age (weeks)	Frequency (n)	Maximum FiO₂ (%) Mean ± SD	*p*-value (Kruskal-Wallis)	Minimum FiO₂ (%) Mean ± SD	*p*-value (Kruskal-Wallis)
28-30	20	83.10 ± 18.49		57.05 ± 9.81	
31-33	25	68.52 ± 20.39	**< 0.001**	45.24 ± 13.73	**<0.001**
34-36	20	55.60 ± 16.54		37.25 ± 9.84	
Total	65	68.17 ± 21.41		46.42 ± 13.79	

Values are mean ± SD. FiO₂ = Fraction of inspired oxygen. Differences between groups tested with Kruskal–Wallis test; p < 0.05 considered significant

Table 4: Baseline (pre-support) and stabilized (post-support) arterial blood gas (ABG) parameters by gestational age group

Baseline (pre-support) ABGs by Gestational age (Group)						
Gestational Age (weeks)	Frequency (n)	PH pre mean ± SD	PaCO$_2$ pre (mmHg) mean ± SD	PaO$_2$ pre (mmHg) mean ± SD	HCO$_3^-$ pre (mmol/L) mean ± SD	SaO2 pre (%) mean ± SD
28-30	20	7.22 ± 0.03	61.85 ± 8.31	37.35 ± 6.47	19.20 ± 2.67	68.72 ± 7.60
31-33	25	7.22 ± 0.05	55.52 ± 7.35	43.16 ± 7.11	19.08 ± 1.847	75.66 ± 7.98
34-36	20	7.24 ± 0.04	52.00 ± 6.473	43.50 ± 10.18	19.70 ± 1.89	75.29 ± 10.53
Stabilized (post-support) ABGs by Gestational age (Group)						
Gestational Age (weeks)	Frequency (n)	PH post mean ± SD	PaCO$_2$ post (mmHg) mean ± SD	PaO$_2$ post (mmHg) mean ± SD	HCO$_3$ post (mmol/L) mean ± SD	SaO$_2$ post (%) mean ± SD
28-30	20	7.28 ± 0.03	53.30 ± 8.58	51.85 ± 6.75	20.35 ± 2.54	84.53 ± 4.06
31-33	25	7.30 ± 0.03	48.68 ± 6.90	55.96 ± 8.00	20.12 ± 1.79	87.45 ± 4.46
34-36	20	7.31 ± 0.03	44.95 ± 6.21	56.50 ± 9.25	21.10 ± 2.17	87.60 ± 4.91

Values are mean ± SD. PaO□ = Partial pressure of oxygen; PaCO□ = Partial pressure of carbon dioxide; HCO□□ = Bicarbonate; SaO□ = Arterial oxygen saturation.

Table 5. Within-group changes (Δ) in arterial blood gas (ABG) parameters and between-group comparison by gestational age

Parameter	28–30 wks (mean ± SD)	31–33 wks (mean ± SD)	34–36 wks (mean ± SD)	Paired p-value (all groups)	ANOVA F	ANOVA p-value
ΔPH	0.06 ± 0.04	0.07 ± 0.07	0.07 ± 0.04	< 0.001	0.595	0.555 (NS)
ΔPaCO☐	-8.55 ± 2.67	-6.84 ± 3.28	-7.05 ± 3.30	< 0.001	1.897	0.159 (NS)
ΔPaO☐	14.50 ± 5.36	12.8 ± 5.01	13.00 ± 3.67	< 0.001	0.804	0.452 (NS)
ΔHCO☐☐	1.15 ± 0.93	1.04 ± 0.84	1.40 ± 0.82	< 0.001	0.987	0.378 (NS)
ΔSaO$_2$	15.81 ± 5.76	11.79 ± 5.28	12.30 ± 6.63	< 0.001	2.93	0.061 (NS)

Values are mean ± SD. Δ = change from pre- to post-support. Within-group changes assessed by paired t-test (p-values shown); between-group comparisons by one-way ANOVA. PaO☐ = Partial pressure of oxygen; PaCO☐ = Partial pressure of carbon dioxide; HCO☐☐ = Bicarbonate; SaO☐ = Arterial oxygen saturation; NS = not significant.

All gestational age groups showed significant within-group improvements in ABG parameters after respiratory support ($p < 0.001$ for all). However, between-group comparisons revealed no statistically significant differences in ΔpH, ΔPaCO☐, ΔPaO☐, ΔHCO☐☐, or ΔSaO☐ (ANOVA, all $p > 0.05$).

Table 6. Correlation of gestational age with oxygen requirements and average arterial blood gas (ABG) parameters

Variable	Correlation Test	Correlation Coefficient	Sig. (2-tailed)	Interpretation
Min. FiO$_2$ vs. GA	Spearman's rho	-0.571**	< 0.001	↑ GA → ↓ FiO$_2$ (Strong negative)
Max. FiO$_2$ vs. GA	Spearman's rho	-0.558**	< 0.001	↑ GA → ↓ FiO$_2$ (Strong negative)
Min. FiO$_2$ vs. Max. FiO$_2$	Spearman's rho	+0.901**	< 0.001	very strong positive
pH_avg. vs. GA	Spearman's rho	0.377**	0.002	↑ GA → ↑pH (Moderate positive)
PaCO$_2$ avg. vs. GA	Spearman's rho	-0.464**	< 0.001	↑ GA → ↓ PaCO☐ (Moderate negative)
PaO$_2$ avg. vs GA	Spearman's rho	0.251**	0.044	↑ GA → ↑ PaO☐ (weak positive)
HCO$_3^-$ avg. vs. GA	Spearman's rho	0.126**	0.316	No meaningful correlation
SaO$_2$ avg. vs. GA	Spearman's rho	0.274**	0.027	↑ GA → ↑ SaO☐ (weak positive)

*Values are Spearman's correlation coefficients (ρ) with corresponding 2-tailed significance. "Average" values were calculated as the mean of pre- and post-support measurements. **p < 0.05 significant; *p < 0.01 highly significant. GA = Gestational Age; FiO☐ = Fraction of inspired oxygen; PaCO☐ = Partial pressure of carbon dioxide; PaO☐ = Partial pressure of oxygen; HCO☐☐ = Bicarbonate; SaO☐ = Arterial oxygen saturation*

Table 7: Type of respiratory support by gestational age group

Gestational Age Group (weeks)	Frequency (n)	CPAP (n, %)	MV (n, %)	NC (n, %)	*p*-value (Pearson Chi-Square)
28-30	20	7, 35%	12, 60%	1, 5%	
31-33	25	9, 36%	9, 36%	7, 28%	0.011
34-36	20	10, 50%	2, 10%	8, 40%	
Total	65	26, 40%	23, 35.4%	16, 24.6%	

Values are presented as number (percentage). CPAP = Continuous positive airway pressure; MV = Mechanical ventilation; NC = Nasal cannula. Differences between groups tested with Pearson's Chi-square test (p = 0.011).

Table 8: Oxygen requirement and oxygenation status of preterm neonates with RDS

Gestational Age (weeks)	Start FiO2 (%) Mean ± SD	FiO□ Band (IQR range: P25–P75)	PaO_2 Pre (mmHg) Mean ± SD	PaO2 Post (mmHg) Mean ± SD	SaO_2 Pre (%) Mean ± SD	SaO_2 Post (%) Mean ± SD	Target SaO2 (%)
28-30	57.05 ± 9.81	50-99	37.35 ± 6.47	51.85 ± 6.75	68.73 ± 7.60	84.53 ± 4.06	90-94
31-33	45.24 ± 13.73	32-92	43.16 ± 7.11	55.96 ± 7.99	75.66 ± 7.98	87.45 ± 4.46	90-94
34-36	37.25 ± 9.84	28-65	43.50 ± 10.19	55.50 ± 9.25	75.29 ± 10.53	87.60 ± 4.92	90-94

stratified by gestational age

Oxygen requirement and arterial blood gas parameters in preterm neonates with respiratory distress syndrome by gestational age. Data are presented as mean ± SD unless otherwise stated; FiO□ band shown as interquartile range (P25–P75). Pre = values before initiation of respiratory support; Post = values after stabilization.

REFERENCES

1.   Ali S, Mohammed N, Qureshi N, Gupta S. Oxygen therapy in preterm infants: recommendations for practice. Paediatrics and Child Health. 2020;31; https://doi.org/10.1016/j.paed.2020.10.001.
2.   Bahadue FL, Soll R. Early versus delayed selective surfactant treatment for neonatal respiratory distress syndrome. Cochrane Database of Systematic Reviews. 2012(11); 10.1002/14651858.CD001456.pub2.
3.   Mathias M, Chang J, Perez M, Saugstad O. Supplemental Oxygen in the Newborn: Historical Perspective and Current Trends. . Antioxidants. 2021;10(12):1879; https://doi.org/10.3390/antiox10121879.
4.   Norman M, Jonsson B, Söderling J, Björklund LJ, Håkansson S. Patterns of respiratory support by gestational age in very preterm infants. Neonatology. 2023;120(1):142–52; https://doi.org/10.1159/000527641.
5.   Manley BJ, Arnolda GRB, Wright IMR, Owen LS, Foster JP, Huang L, et al. Nasal High-Flow Therapy for Newborn Infants in Special Care Nurseries. N Engl J Med. 2019;380(21):2031-40; 10.1056/NEJMoa1812077.
6.   Abdel-Latif ME, Davis PG, Wheeler KI, De Paoli AG, Dargaville PA. Surfactant therapy via thin catheter in preterm infants with or at risk of respiratory distress syndrome. Cochrane Database Syst Rev. 2021;5(5):Cd011672; 10.1002/14651858.CD011672.pub2.
7.   Abdallah Y, Mkony M, Noorani M, Moshiro R, Bakari M, Manji K. CPAP failure in the management of preterm neonates with respiratory distress syndrome where surfactant is scarce. A prospective observational study. BMC Pediatrics. 2023;23(1):211; https://doi.org/10.1186/s12887-023-04038-6.
8.   Obst S, Herz J, Alejandre Alcazar MA, Endesfelder S, Möbius MA, Rüdiger M, et al. Perinatal Hyperoxia and Developmental Consequences on the Lung-Brain Axis. Oxidative Medicine and Cellular Longevity. 2022;2022(1):5784146; https://doi.org/10.1155/2022/5784146.
9.   Sweet DG CV, Greisen G, et al. European Consensus Guidelines on the management of RDS – 2022 update. Neonatology. 2023;119;
10.   Subramaniam P, Ho JJ, Davis PG. Prophylactic or very early initiation of continuous positive airway pressure (CPAP) for preterm infants. Cochrane Database Syst Rev. 2021;10(10):Cd001243; https://doi.org/10.1002/14651858.CD001243.pub4.
11.   Ohuma EO, Moller A-B, Bradley E, Chakwera S, Hussain-Alkhateeb L, Lewin A, et al. National, regional, and global estimates of preterm birth in 2020, with trends from 2010: a systematic analysis. The Lancet. 2023;402(10409):1261-71; https://doi.org/10.1016/S0140-6736(23)00878-4.
12.   Thébaud B, Goss KN, Laughon M, Whitsett JA, Abman SH, Steinhorn RH, et al. Bronchopulmonary dysplasia. Nature Reviews Disease Primers. 2019;5(1):78; https://doi.org/10.1038/s41572-019-0127-7.
13.   McGoldrick E, Stewart F, Parker R, Dalziel SR. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. Cochrane Database of Systematic Reviews. 2020;2020(12); https://doi.org/10.1002/14651858.CD004454.pub4.

14.   Hulley SB, Cummings SR, Browner WS, Grady D, Newman TB. Designing Clinical Research. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2013.
15.   Kohn MA. Sample Size Calculators 2022 [Available from: https://sample-size.net.

FIGURES

Figure 1. Mean FiO□ (%) across Gestational Age groups

Bars represent mean values. FiO□ = Fraction of inspired oxygen; GA = Gestational Age. Group differences tested with Kruskal–Wallis test.

Figure 2: Relationship Between Gestational Age and Average pH

Scatter plot depicting the association between gestational age and mean pH values, with a fitted linear regression line ($R^2 = 0.144$). Gestational age groups were coded as: 1 = 28–30 weeks, 2 = 31–33 weeks, and 3 = 34–36 weeks.

Figure 3: Relationship Between Gestational Age and Average P_aCO_2

Scatter plot depicting the association between gestational age and mean PaCO2 values, with a fitted linear regression line (R² = 0.205). Gestational age groups were coded as: 1 = 28–30 weeks, 2 = 31–33 weeks, and 3 = 34–36 weeks.

Figure 4: Relationship Between Gestational Age and Average P_aO_2

Scatter plot depicting the association between gestational age and mean PaO$_2$ values, with a fitted linear regression line (R² = 0.072). Gestational age groups were coded as: 1 = 28–30 weeks, 2 = 31–33 weeks, and 3 = 34–36 weeks.

Figure 5: Relationship Between Gestational Age and Average HCO_3^-

Scatter plot depicting the association between gestational age and mean HCO$_3^-$ values, with a fitted linear regression line (R² = 0.014). Gestational age groups were coded as: 1 = 28–30 weeks, 2 = 31–33 weeks, and 3 = 34–36 weeks.

Figure 6: Relationship Between Gestational Age and Average S_aO_2

Scatter plot depicting the association between gestational age and mean SaO$_2$ values, with a fitted linear regression line (R² = 0.078). Gestational age groups were coded as: 1 = 28–30 weeks, 2 = 31–33 weeks, and 3 = 34–36 weeks.

Figure 7: Types of Respiratory Support Across Different Gestational Ages

Distribution of respiratory support modalities among neonates by gestational age groups (28–30 weeks, 31–33 weeks, and 34–36 weeks). Respiratory support includes continuous positive airway pressure (CPAP), mechanical ventilation, and nasal cannula. The y-axis represents the number of neonates in each category.

FiO2 Requirements by Gestational Age

R^2 Linear = 0.205

$y=61.77-4.55*x$