Spread of the plague in Venice, 1630–1631: epidemic entropy in a "natural experiment"

Jonathan L. Hwang[1], Ariktha Srivathsan[1,2], Michael S. Deiner[1,3], Seth Blumberg[1,4], Travis
C. Porco[1,2,3], and Thomas M. Lietman[1,2,3,5,*]

[1] Francis. I. Proctor Foundation, University of California, San Francisco, California
[2] Department of Epidemiology and Biostatistics, University of California, San Francisco,
California
[3] Department of Ophthalmology, University of California, San Francisco, California
[4] Department of Medicine, University of California, San Francisco, California
[5] Institute for Global Health Sciences, University of California, San Francisco, California
*Tom.Lietman@ucsf.edu

ABSTRACT

Precise modeling of epidemic spread is difficult. One explanation is that disease spread is
inherently stochastic. This would suggest that the distribution of cases across geographic regions
would progress towards that more favored by chance. If the epidemic proceeds long enough, the
allocation of cases could approach that most expected, maximizing Boltzmann–Gibbs–Shannon
entropy. Here, we tested these hypotheses on mortality data from the Venetian 1630–1631
plague epidemic. Entropy per case (intensive) of the quantile function (distribution of parishes
ranked by case rates) increased from an effective number of 7.32 parishes (95% CI 3.32–12.55
parishes) to 47.9 parishes (47.5–48.9 parishes) out of 50 total, indicating that the quantile
function approached a uniform maximum entropy distribution. Intensive entropy of the
probability density function (parishes categorized by cumulative case rate) increased from 0.63
nats (0.32–0.93 nats) to 1.75 nats (1.53–1.87 nats). The PDF approached a Gaussian distribution.
The Kullback–Leibler divergence decreased from 0.84 nats (0.71–1.42 nats) to 0.12 nats (0.083–
0.35 nats). These findings quantify how disease spreads and demonstrate that observed
heterogeneity in infections between regions may in some circumstances be explained by chance
alone.

Introduction

Forecasting the time course and geographical extent of epidemics often fails.[1–6] However,
predicting the distribution of cases across communities may be achievable.[7–12] If disease
transmission were in large part driven by stochasticity, the accuracy of small-scale predictions
would be limited, while large-scale behavior would be predictable. In particular, observed case
distributions (macrostates) with more possible combinatorial arrangements (microstates) would
be favored over time. Therefore, the distribution of infections across communities would have
increasing Boltzmann–Gibbs–Shannon entropy over time. Furthermore, if the epidemic persisted
long enough, it is plausible that this distribution would approach the distribution with the greatest
number of such arrangements, a maximum entropy distribution. Specific maximum entropy
distributions, in particular, the uniform, exponential, or Gaussian distributions, can be
characterized by a small number of constraints (0, 1, and 2 sufficient parameters, respectively).

Beyond these constraints, maximum entropy distributions make minimal assumptions to model
the distribution of infections across geographic regions during an epidemic.
The 1630–1631 plague outbreak in Venice offers an opportunity to test these hypotheses.
Advantages of this for hypothesis testing include the absence of an effective intervention
(vaccines, antibiotics), restricted movement, detailed daily death records unrestricted by privacy
laws, and a complete, uninterrupted epidemic cycle rarely seen in modern outbreaks.[13–15] These
features, combined with clear geographical boundaries between parishes, make the Venice
outbreak an ideal "natural experiment" for testing basic generalizable principles of epidemic
spread.[16] Here, we test the hypotheses that the distribution of mortality across parishes both had
increasing entropy over the course of the 1630–1631 Venetian plague, and approached a
maximum entropy distribution.
52
Results
Descriptive Statistics
A total of 113,111 individuals across 50 parishes were considered at risk for infection on
January 1, 1629. Among the 1629 population estimates of the 50 parishes, the median parish
population was 1,734 (SD 1,802). A total of 42,538 deaths were recorded over the specified time
span. The mean cumulative mortality of a parish from January 1, 1629 to December 31, 1631
was 37% (SD 10%). The progression in the geographic distribution of cumulative mortality over
time is depicted in Figure 1 and Supplementary Video S1 online.
61

62
Fig. 1. (**a**) Choropleth maps, (**b**) quantile function bar charts (ranked parishes by their normalized
death rates), and (**c**) smooth probability density function (PDF) histograms (Epanechnikov

kernel, $N = 512$) depicting the geographic distribution of cumulative mortality at the beginning
of the epidemic (January 7, 1629), during the primary wave (October 31, 1630), and at the end of
the epidemic (December 31, 1631). In (**a**), darker shades of orange correspond to higher
cumulative mortalities, represented as the ratio of the total deaths in a parish to its population.
The maximum entropy distribution is overlaid in red on the quantile functions in (**b**) and the
PDFs in (**c**). The coefficient of variation (CV) was 1.32 (95% CI 1.00–1.65) on January 7, 1629,
0.34 (0.26–0.41) on October 31, 1630, and 0.28 (0.22–0.33) on December 31, 1631.
72
Cumulative Mortality
When we analyzed the temporal progression of mortality distribution across Venice's
parishes throughout the epidemic, the quantile function of cumulative mortality (ranked parishes
by their death rates) demonstrated a nearly monotonic increase in entropy, indicating a
progressive shift from concentrated to dispersed mortality patterns. For each entropy value, the
number of parishes with uniformly distributed mortality that would result in the same entropy
value (effective number of parishes) was also evaluated. At the outbreak's onset (January 1,
1629), the distribution showed relatively low entropy of 1.99 nats (95% confidence interval (CI)
1.20–2.53 nats), equivalent to an effective number of 7.32 parishes (3.32–12.55 parishes)—
equivalent to the same entropy as an even distribution across only 7-8 parishes. By the
epidemic's later stages (December 29, 1631), entropy had increased to 3.87 nats (3.86–3.89
nats), equivalent to an effective uniform distribution across 47.9 parishes (47.5–48.9 parishes),
approaching the maximum attainable 50 parishes. This transition toward uniformity was further
quantified using Kullback-Leibler (KL) divergence from a discrete uniform distribution
(negentropy from a uniform), a maximum entropy distribution, which decreased from 1.92 nats
(1.38–2.71 nats) at the outbreak's start to 0.038 nats (0.023–0.054 nats) by December 1631,
indicating the distribution had become nearly uniform. The time-series of entropy and
negentropy of the quantile function are displayed in Figure 2.

Quantile Function of Cumulative Mortality

Fig. 2. Time-series graphs for every day between January 1, 1629 and December 31, 1631 of (**a**) entropy (left axis) and effective number of parishes (right axis) of the distribution of cumulative mortality across parishes and (**b**) Kullback–Leibler divergence of the distribution of cumulative mortality across parishes from a uniform distribution. The red lines depict the maximum possible entropy and minimum possible negentropy, respectively, corresponding to an even distribution across all 50 parishes. Bootstrap 95% CIs ($N = 999$) for each day are depicted in gray.

The probability density function (PDF) of cumulative mortality (parish density in death rate bins) decreased in entropy prior to the largest wave in fall of 1630 and increased thereafter, indicating a shift in the distribution of variation in cumulative mortality toward a normal distribution after the height of the epidemic. At the outbreak's onset, the distribution showed relatively low entropy of 0.63 nats (0.32–0.93 nats). A peak of 1.82 nats (1.55–1.89 nats) was reached on June 17, 1631. By the end of the epidemic, entropy had slightly decreased to 1.75 nats (1.53–1.87 nats). The PDF transitioned toward a Gaussian distribution, another maximum entropy distribution, over the course of the epidemic. This was quantified with Kullback-Leibler (KL) divergence from a binned Gaussian distribution with identical mean and variance (negentropy from a Gaussian). Negentropy decreased from 0.84 nats (0.71–1.42 nats) at the outbreak's start to a minimum of 0.071 nats (0.050–0.31 nats) on May 5, 1629 before slightly increasing to 0.12 nats (0.083–0.35 nats) by its end, indicating the distribution had become relatively normal. Thus, the distribution of variation in cumulative mortality between parishes was consistent with that of a stochastic process. The time-series of entropy and negentropy of the PDF are displayed in Figure 3.

PDF of Cumulative Mortality

114
Fig. 3. Time-series graphs for every day between January 1, 1629 and December 31, 1631 of (**a**)
entropy of the binned distribution of parish density across cumulative mortalities and (**b**)
Kullback–Leibler divergence of the binned distribution of parish density across cumulative
mortalities from a binned Gaussian distribution with the same mean and variance as the data. The
red lines depict the maximum possible entropy and minimum possible negentropy, respectively.
Bootstrap 95% CIs (*N* = 999) for each day are depicted in gray.
121

Instantaneous Mortality Rate (Daily)

The distribution of instantaneous mortality rate across Venice's parishes (quantile
function of instantaneous mortality) approached maximal entropy during the largest wave in the
fall of 1630 and minimal entropy at the beginning and end of the epidemic, indicating that the
distribution of daily deaths was only relatively uniform at the height of the epidemic. At the
outbreak's onset, the distribution showed relatively low entropy of 1.99 nats (1.23–2.52 nats). In
the fall of 1630 (November 29), entropy had increased to a peak of 3.75 nats (3.68–3.81 nats).
However, by the end of the epidemic, the distribution once again showed relatively low entropy
of 1.96 nats (1.05–2.52 nats). The uniformity of the distribution of daily deaths was further
quantified using negentropy from a uniform, which decreased from 1.92 nats (1.40–2.68 nats) at
the outbreak's start to a minimum of 0.16 nats (0.10–0.23 nats) in the fall of 1630 (November
29) before increasing to 1.95 nats (1.39–2.86 nats) by its end, indicating that the distribution was
only relatively uniform at the height of the epidemic. The lack of uniformity of the distribution
before and after the peak in cases does not indicate that the likelihood of infection was
heterogeneous across parishes, but rather that new cases did not occur frequently enough for the

uniform pattern to be observed over a single day. The time-series of entropy and negentropy of
the instantaneous quantile function are displayed in Supplementary Fig. S2 online.
The distribution of parish density across instantaneous mortality rates (PDF of
instantaneous mortality) approached maximal entropy during the largest wave in the fall of 1630
and minimal entropy at the beginning and end of the epidemic, indicating that the distribution of
variation in new deaths was only relatively normal at the height of the epidemic. At the
outbreak's onset, the distribution showed relatively low entropy of 0.63 nats (0.32–0.95 nats). In
the fall of 1630 (November 7), entropy had increased to a peak of 1.90 nats (1.72–1.92 nats).
However, by the end of the epidemic, the distribution once again showed relatively low entropy
of 0.65 nats (0.27–0.93 nats). The PDF transitioned toward normality as quantified by
negentropy from a Gaussian, which decreased from 0.84 nats (0.71–1.43 nats) at the outbreak's
start to a minimum of 0.088 nats (0.075–0.27 nats) in the fall of 1630 (November 23) before
increasing to 0.92 nats (0.79–1.56 nats) by its end, indicating that the distribution of variation in
new deaths was only relatively normal at the height of the epidemic. As with the daily quantile
function, this lack of normality may reflect an insufficient number of cases to observe the normal
pattern that would be expected from a simple stochastic process on a daily basis. The time-series
of entropy and negentropy of the instantaneous PDF are displayed in Supplementary Fig. S3
online.
155
Sestieri
Venice comprises 6 districts known as sestieri, each of which contains an average of 8.33
parishes (SD 3.20 parishes). If cumulative mortality had been clustered geographically, the mean
cumulative mortality of the parishes in each sestiere would vary across sestieri. However, such
variation was not observed: across the 6 sestieri, the mean parish cumulative mortality rate was
38% with a standard deviation of only 1.5%. Analyzing entropy on the sestiere level provides
insight on the behavior of entropy in areas with fewer subregions. The quantile function of
cumulative mortality had increasing entropy and closely approximated a maximum entropy
distribution within all 6 sestieri (Figure 4).

Fig. 4. Entropy (left axis) and effective number of parishes (right axis) of the distribution of
cumulative mortality across parishes for each sestiere, indicated with colors. The thick dotted
lines indicate the maximum possible entropy of each sestiere, or the log of the number of
parishes in that sestiere, the thin dotted lines indicate bootstrap 95% CIs ($N = 999$), and the solid
lines indicate observed entropy.

Discussion

The entropy of the geographic distribution of mortality during the 1630–1631 Venetian
plague was analyzed to assess the degree to which stochastic factors influence disease spread.
After the primary wave in the fall of 1630, both the quantile function (Fig. 1b) and PDF (Fig. 1c)
of cumulative mortality had increasing entropy and approached a maximum entropy distribution,
indicating that the spread of plague could in large part be explained by stochastic factors. This
finding was robust to the choice of geographic scale, as the same behavior was observed in the
subsets of parishes within each of Venice's 6 sestieri. Additionally, at the height of the epidemic,
the geographic distribution of instantaneous mortality approached a maximum entropy
distribution. These findings suggest that heterogeneity in infections between regions during an
epidemic may arise from random fluctuations rather than predictable factors.

The distribution of infections across communities has had increasing entropy and
approached a maximum entropy distribution in outbreaks of other diseases, including SARS-

CoV-2 and Ebola virus disease.[17–19] During the COVID-19 pandemic, many models attributed
variation in the cumulative prevalences in US states to heterogeneous public health
practices.[6,20,21] If this were the case, regions with high initial prevalences would have had
consistently higher prevalences over the course of the epidemic. However, the distribution of
cumulative prevalence across US states eventually approached a uniform distribution, indicating
that this initial variation was primarily caused by stochastic factors (e.g. the timing of
introduction).[19] The distribution of infection over geographical regions in endemic diseases has
also been monitored. The stochastic SIS model predicts that the PDF will approach an
approximate geometric distribution when the reproduction number (R_0) is less than one and a
Gaussian distribution when R_0 is greater than one.[22] This has been verified in subcritical diseases
such as trachoma and onchocerciasis at the village level, and trachoma and leprosy at the district
level.[7,23–25] The parameters of the maximum entropy distribution approached by the PDF of an
epidemic may suffice to convey much of the epidemic's distributional information. Additionally,
for a given transmission model, it is possible to formulate a fundamental equation and equations
of state relating the epidemiological interpretation of entropy to the thermodynamic
definition.[11,26] For example, for an SIS model with $R_0 > 1$, the mean approaches $1-1/R$ and the
variance approaches N/R.[27] Because the entropy of the Gaussian distribution is known, it can be
written in a fundamental equation as a function of the other state variables, such as I at
equilibrium and N.[28] This framework allows epidemiological parameters such as R_0 to be
estimated.

Spatial entropy has previously been used to characterize other attributes of epidemics.[9–
11,26,29–31] For instance, while this study focused on the end stages of an epidemic, a previous study
used spatial entropy to predict the spatial source of epidemics.[31] In the study, the metapopulation
was represented with a directed graph and the MaxEnt method was used to reweight ensemble
trajectories. However, maximum entropy distributions themselves were not used as models for
the distribution of infections. The authors ran several epidemic simulations and found that their
model consistently included the true infectious source among the top 5 most probable
predictions. A second study used spatial entropy to identify when coronavirus infection was
maximally dispersed in various countries.[9] The authors considered non-cumulative incidence
rate, and therefore used entropy to determine the time of maximum diffusion of infections. They
found that all of the countries they considered reached maximal dispersion within a three-month
window, and that the same methodology worked on the city-level. However, the specific
maximum entropy distribution each region approached was not considered.

Limitations of this study include the consideration of deaths from all causes. While a
concurrent smallpox outbreak had the potential to create a substantial difference between the
observed distribution of deaths and the distribution of plague cases, particularly prior to the first
wave of plague (9/1/1630–12/31/1630).[13] However, if smallpox spread in such a way that
location within the city did not matter, as was the case with the plague, this disparity would be
minimal. The parish of San Marco was excluded from the dataset because it served as a
placeholder for deaths from drowning and those in prison, which resulted in an early spike in

cases not representative of parish transmission (see Supplementary Fig. S4–5 online).[14] As
deaths accumulated, instantaneous mortality was increasingly underestimated as parish
populations were assumed to be fixed. However, because cumulative mortality was uniform
across all parishes, this effect was symmetric. Finally, while entropy necessarily increases in the
early stages of an epidemic due to cases occurring in regions for the first time, entropy increased
at a faster rate than would be expected from this effect alone, indicating progression towards a
maximum entropy distribution.
The 'clean' experimental conditions of the Venetian outbreak (lacking modern
interventions, mass transit, or privacy restrictions) allowed us to observe fundamental patterns of
disease spread that may be obscured in modern datasets. Modern outbreaks are often controlled
before reaching distributional equilibrium, making historical datasets particularly valuable for
testing theories about long-term epidemic behavior.[32] Understanding these basic principles can
improve our ability to predict disease distribution patterns in any epidemic, regardless of the
specific pathogen or time period. For instance, the effect of interventions could be quantified
with Kullback–Leibler divergence of the observed distribution of infections from the maximum
entropy distribution being approached. Additionally, maximum entropy distributions could be
used to benchmark more complex models or guide early outbreak response under uncertainty due
to their low bias. This same entropy-based analysis could be conducted on geospatial data from
any epidemic, and we hypothesize that the entropy of other epidemics will similarly increase and
reach a maximum entropy distribution with sufficient time. Currently, the consistency of this
pattern across epidemics of varying diseases and populations is promising, as it suggests a
reliable method for predicting the distribution of cases across communities using maximum
entropy distributions. The findings of this study therefore indicate that before complex models
are developed to explain observed heterogeneity in the distribution of infections across areas, the
degree to which this heterogeneity could be explained by chance alone merits consideration.

Methods

Data collection and cleaning

Venetian census data of parish populations from both 1586 and 1696 was linearly
interpolated to obtain estimates of parish populations in 1629 (see Supplementary Fig. S6
online).[33] Death counts in each of 50 parishes recorded daily between January 1, 1629 and
December 31, 1631 in Venetian necrologies were normalized by parish populations to calculate
cumulative mortality (see Supplementary Fig. S7 online). Because all-cause deaths, rather than
infections, were recorded, all-cause deaths were used to approximate infections. However, all-
cause deaths closely match plague-related deaths because plague was the primary cause of death
during the time.[15] Cumulative mortality was used to avoid an upper bound on entropy values due
to insufficient cases. However, because cumulative mortality necessarily increases over the
course of an epidemic, we used intensive entropy, or the average entropy per case. Boltzmann-
Gibbs-Shannon entropy in nats (defined as $-\sum_{i=1}^{N} p_i \log(p_i)$, where p_i is the probability in bin i)
was used to relate our findings to the number of combinatorial arrangements of cases across

regions that would result in the observed distribution. The effective number of parishes, or the
number of evenly distributed parishes that would result in the same entropy value, was evaluated
as the exponential function of entropy.[34] Because parish names were inconsistent between
censuses and death records, manual matching was occasionally required. 20 parishes were
excluded from the initial dataset containing 70: one parish (S. Marcilian) was not recorded in the
1696 census data, 18 had no data in the primary outbreak period (9/1/1630–12/31/1630), and one
(S. Marco) was a placeholder for deaths from drowning, deaths in prison, and deaths with other
ambiguous locations due to its role as an administrative district.[14] For included parishes, missing
data points were replaced with zeroes.

Data analysis and modeling
The geographic distribution of mortality was represented with both a quantile function
and a probability density function (PDF). The quantile function was the bar chart of parishes
ordered by mortality while the PDF was approximated by a histogram of parish counts binned by
mortality under Sturges's Rule.[35] Kullback–Leibler (KL) divergence from maximum entropy
distributions, or negentropy, was used to measure the extent to which the distribution of
mortality approximated a maximum entropy distribution. KL divergence can be interpreted as
the expectation of the logarithmic difference between the observed probability distribution P and
the model distribution Q, defined as $\sum_{x \in \mathcal{X}} P(x) \log \left(\frac{P(x)}{Q(x)} \right)$. For the quantile function, KL
divergence from a discrete uniform distribution was calculated. For the PDF, KL divergence
from a binned Gaussian distribution with identical mean and variance to the data was also
calculated.[36]
Bootstrap 95% CIs for both entropy and KL divergence were calculated by resampling
parishes ($N = 999$) and evaluating the 2.5% and 97.5% intervals at each time point to test for bias
in parish selection. Entropy and KL divergence were also calculated and bootstrapped within
each sestiere to identify potential proximity-based heterogeneity in this pattern. The geographic
distribution of cumulative mortality for every month in the specified time period was visualized
with choropleth maps, the base code for which was sourced from Lazzari et al.[15] The entropy of
daily mortality and the entropy when the parish of S. Marco was included were calculated as
sensitivity analyses. All analyses were conducted using R (Version 4.2.0 for MacOS).[37]

Data Availability

The dataset analyzed during the current study is available in the Venice-plague-epidemic-paper
repository created by Lazzari et al., https://github.com/ggrrll/Venice-plague-epidemic-paper.

References

1. Ioannidis, J. P. A., Cripps, S. & Tanner, M. A. Forecasting for COVID-19 has failed. *Int. J.*

Forecast. **38**, 423–438 (2022).

2.   Wynants, L. *et al.* Prediction models for diagnosis and prognosis of covid-19: systematic

review and critical appraisal. *BMJ* **369**, m1328 (2020).

3.   Roda, W. C., Varughese, M. B., Han, D. & Li, M. Y. Why is it difficult to accurately predict

the COVID-19 epidemic? *Infect. Dis. Model.* **5**, 271–281 (2020).

4.   Chharia, A. *et al.* Accuracy of US CDC COVID-19 forecasting models. *Front. Public Health*

12, (2024).

5.   Scarpino, S. V. & Petri, G. On the predictability of infectious disease outbreaks. *Nat.*

Commun. **10**, 898 (2019).

6.   Rahimi, I., Chen, F. & Gandomi, A. H. A review on COVID-19 forecasting models. *Neural*

Comput. Appl. 1–11 (2021) doi:10.1007/s00521-020-05626-8.

7.   Lietman, T. M. *et al.* The distribution of the prevalence of ocular chlamydial infection in

communities where trachoma is disappearing. *Epidemics* **11**, 85–91 (2015).

8.   Salathé, M. & Jones, J. H. Dynamics and Control of Diseases in Networks with Community

Structure. *PLOS Comput. Biol.* **6**, e1000736 (2010).

9.   Lucia, U., Deisboeck, T. S. & Grisolia, G. Entropy-Based Pandemics Forecasting. *Front.*

Phys. **8**, (2020).

10. Wang, Z., Broccardo, M., Mignan, A. & Sornette, D. The dynamics of entropy in the

COVID-19 outbreaks. *Nonlinear Dyn.* **101**, 1847–1869 (2020).

11. Harding, N., Nigmatullin, R. & Prokopenko, M. Thermodynamic efficiency of contagions: a

statistical mechanical analysis of the SIS epidemic model. *Interface Focus* **8**, 20180036

(2018).

12. Srivathsan, A. *et al.* District-Level Forecast of Achieving Trachoma Elimination as a Public

Health Problem By 2030: An Ensemble Modelling Approach. *Clin. Infect. Dis.* **78**, S101–

S107 (2024).

13. Ell, S. R. Three Days in October of 1630: Detailed Examination of Mortality During an

Early Modern Plague Epidemic in Venice. *Rev. Infect. Dis.* **11**, 128–139 (1989).

14. Bamji, A. Medical Care in Early Modern Venice. *J. Soc. Hist.* **49**, 483–509 (2016).

15. Lazzari, G. *et al.* A digital reconstruction of the 1630–1631 large plague outbreak in Venice.

Sci. Rep. **10**, 17849 (2020).

16. Palmer, R. J. The Control of Plague in Venice and Northern Italy 1348-1600. (University of

Kent, 1978). doi:10.22024/UniKent/01.02.86158.

17. Almetwally, E. M., Dey, S. & Nadarajah, S. An Overview of Discrete Distributions in

Modelling COVID-19 Data Sets. *Sankhya A* **85**, 1403–1430 (2023).

18. Pigott, D. M. *et al.* Mapping the zoonotic niche of Ebola virus disease in Africa. *eLife* **3**,

e04395 (2014).

19. CDC. COVID Data Tracker. *Centers for Disease Control and Prevention*

https://covid.cdc.gov/covid-data-tracker (2020).

20. Unwin, H. J. T. *et al.* State-level tracking of COVID-19 in the United States. *Nat. Commun.*

11, 6189 (2020).

21. Dominik Güss, C. *et al.* The politics of COVID-19: Differences between U.S. red and blue

states in COVID-19 regulations and deaths. *Health Policy OPEN* **5**, 100107 (2023).

22. Nåsell, I. The Quasi-Stationary Distribution of the Closed Endemic SIS Model. *Adv. Appl.*

Probab. **28**, 895–932 (1996).

23. Surakat, O. A. *et al.* Geospatial distribution and predictive modeling of onchocerciasis in

Ogun State, Nigeria. *PLOS ONE* **18**, e0281624 (2023).

24. Wangara, F. *et al.* The spatial epidemiology of leprosy in Kenya: A retrospective study.

PLoS Negl. Trop. Dis. **13**, e0007329 (2019).

25. Lietman, T. M. *et al.* Models of Trachoma Transmission and Their Policy Implications:

From Control to Elimination. *Clin. Infect. Dis.* **66**, S275–S280 (2018).

26. Wang, W., Wu, Z., Wang, C. & Hu, R. Modelling the spreading rate of controlled

communicable epidemics through an entropy-based thermodynamic model. *Sci. China Phys.*

Mech. Astron. **56**, 2143–2150 (2013).

27. Nåsell, I. *Extinction and Quasi-Stationarity in the Stochastic Logistic SIS Model.* vol. 2022

(Springer, Berlin, Heidelberg, 2011).

28. Rumack, A., Tibshirani, R. J. & Rosenfeld, R. Recalibrating probabilistic forecasts of

epidemics. *PLOS Comput. Biol.* **18**, e1010771 (2022).

29. Erten, E. Y., Lizier, J. T., Piraveenan, M. & Prokopenko, M. Criticality and Information

Dynamics in Epidemiological Models. *Entropy* **19**, 194 (2017).

30. Momani, S., Ibrahim, R. W. & Hadid, S. B. Susceptible-Infected-Susceptible Epidemic

Discrete Dynamic System Based on Tsallis Entropy. *Entropy* **22**, 769 (2020).

31. Ansari, M., Soriano-Paños, D., Ghoshal, G. & White, A. D. Inferring spatial source of

disease outbreaks using maximum entropy. *Phys. Rev. E* **106**, 014306 (2022).

32. Phelps, M. D. *et al.* The importance of thinking beyond the water-supply in cholera

epidemics: A historical urban case-study. *PLoS Negl. Trop. Dis.* **11**, e0006103 (2017).

33. Vianello, F. M., Giovanni, F., Maria, M., Pierpaolo, S. & Francesca, T. Le anime dei

demografi. Fondi per la rilevazione della popolazione di Venezia nei secoli XVI e XVII.

(1991).

34. Jost, L. Entropy and diversity. *Oikos* **113**, 363–375 (2006).

35. Sturges, H. A. The Choice of a Class Interval. *J. Am. Stat. Assoc.* **21**, 65–66 (1926).

36. Marden, J. msos: Data Sets and Functions Used in Multivariate Statistics: Old School by

John Marden. 1.2.0 https://doi.org/10.32614/CRAN.package.msos (2014).

37. R Core Team (2022). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Figure Legends

Fig. 1. (**a**) Choropleth maps, (**b**) quantile function bar charts (ranked parishes by their normalized
death rates), and (**c**) smooth probability density function (PDF) histograms (Epanechnikov
kernel, $N = 512$) depicting the geographic distribution of cumulative mortality at the beginning
of the epidemic (January 7, 1629), during the primary wave (October 31, 1630), and at the end of
the epidemic (December 31, 1631). In (**a**), darker shades of orange correspond to higher
cumulative mortalities, represented as the ratio of the total deaths in a parish to its population.
The maximum entropy distribution is overlaid in red on the quantile functions in (**b**) and the
PDFs in (**c**). The coefficient of variation (CV) was 1.32 (95% CI 1.00–1.65) on January 7, 1629,
0.34 (0.26–0.41) on October 31, 1630, and 0.28 (0.22–0.33) on December 31, 1631.
382
Fig. 2. Time-series graphs for every day between January 1, 1629 and December 31, 1631 of (**a**)
entropy (left axis) and effective number of parishes (right axis) of the distribution of cumulative
mortality across parishes and (**b**) Kullback–Leibler divergence of the distribution of cumulative
mortality across parishes from a uniform distribution. The red lines depict the maximum possible
entropy and minimum possible negentropy, respectively, corresponding to an even distribution
across all 50 parishes. Bootstrap 95% CIs ($N = 999$) for each day are depicted in gray.
389
Fig. 3. Time-series graphs for every day between January 1, 1629 and December 31, 1631 of (**a**)
entropy of the binned distribution of parish density across cumulative mortalities and (**b**)
Kullback–Leibler divergence of the binned distribution of parish density across cumulative
mortalities from a binned Gaussian distribution with the same mean and variance as the data. The
red lines depict the maximum possible entropy and minimum possible negentropy, respectively.
Bootstrap 95% CIs ($N = 999$) for each day are depicted in gray.
396
397   Fig. 4. Entropy (left axis) and effective number of parishes (right axis) of the distribution of
398   cumulative mortality across parishes for each sestiere, indicated with colors. The thick dotted
399   lines indicate the maximum possible entropy of each sestiere, or the log of the number of
400   parishes in that sestiere, the thin dotted lines indicate bootstrap 95% CIs ($N = 999$), and the solid
401   lines indicate observed entropy.

Acknowledgements

We would like to thank Dr. Gianrocco Lazzari, Dr. Giovanni Colavizza, and their collaborators for compiling and digitizing the database in a previous study. The work presented in this article is funded by the National Eye Institute (NEI) at the National Institutes of Health (NIH) (grant number R01EY025350). This work was supported in part by a grant from an unrestricted grant from Research to Prevent Blindness (PI: JLD).

Author contributions

All authors participated in the design of the experiment. J.L.H. performed the analysis. J.L.H. and T.M.L. wrote the initial draft. All authors reviewed and approved the manuscript.

Competing Interests

The authors declare no competing interests.