1

Addressing the causes of 'missingness' in healthcare: a co-designed suite of interventions

Calum Lindsay[1]*, David Baruffati[1], Mhairi Mackenzie[2],David A. Ellis[3], Michelle Major[4], Catherine A O'Donnell[1], Sharon A Simpson[5], Andrea E Williamson[1], Geoff Wong[6]

*Corresponding Author: calum.lindsay@glasgow.ac.uk

[1] Department of General Practice and Primary Care, School of Health and Wellbeing, University of Glasgow, Glasgow, United Kingdom
[2] Department of Urban Studies, School of Social & Political Sciences, University of Glasgow, Glasgow, United Kingdom
[3] Centre for Healthcare Innovation and Improvement Information, Decisions and Operations, Centre for Business Organisations and Society (CBOS), University of Bath, United Kingdom
[4] Homeless Network Scotland, Glasgow, United Kingdom
[5] MRC/CSO Social & Public Health Sciences Unit, School of Health and Wellbeing, University of Glasgow, United Kingdom
[6] Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, United Kingdom

Abstract

Background: Missingness refers to a repeated tendency not to take up offers of care that has a negative impact on the person and their life chances, visible in patterns of missed health appointments. Epidemiological work has shown that patients experiencing 'missingness' are more likely to have multiple physical and mental health conditions, to live in adverse or precarious circumstances, and experience a range of negative health outcomes. Yet existing approaches designed to address missed appointments rarely focus on these patients; when they do, it is often through a punitive lens rather than one that engages meaningfully with the causes of missingness. As a result, existing interventions are often ineffective for these patients and may instead worsen access inequalities. This study addresses this gap by outlining a co-produced 'suite' of interventions aimed at addressing the specific, complex causes of missingness. *Methods and Findings:* The study synthesised findings from three co-occurring workstreams: an extensive realist review of 253 documents including peer-reviewed and grey literature; interviews with 61 'key informants' whose personal and professional experiences provided insight into causes and possible solutions; and a series of co-design workshops with a Stakeholder Advisory Group of 16 professionals and experts-by-experience aimed at designing and refining an intervention. The intervention consists of activities across several key domains: embedding a change of perspective around missed appointments; identification; relationships and communication; missingness coordinators; transport and logistics; flexibility; and contact around appointments. Unifying these activities, and crucial to their success, is a paradigm shift towards missed appointments that we term a 'missingness lens.' *Conclusions:* This intervention presented here is an evidence-informed, realistic and meaningful set of actions with the potential to address the complex causes of missingness and, by extension, access inequalities and wider health inequalities. Future interventional research into missed appointments should actively focus on or include 'missing' patients in design, implementation and the measurement of outcomes.

Introduction and background

Missed appointments have become central to policy discourse around healthcare globally (1). In multiple country contexts, health services generally – and primary care in particular – are impacted by overlapping crises of growing demand and limited capacity (2). Demand comes through aging populations with increasingly complex health needs, societal mental health crises, and a rise in poverty and inequality alongside the erosion of social safety nets (3, 4). Diminished capacity is caused by chronic underfunding, shortages of appropriate staffing and infrastructure; unsustainable workloads and staff burnout; long waiting lists, and thus services that often appear unresponsive and inaccessible to patients (2, 5, 6). In this context, policymakers often frame missed appointments as a growing source of wasted money and lost capacity , and addressing them becomes a financial and moral – even existential - imperative (7, 8). Much existing research is framed in these terms, as it seeks cost-effective and efficient reductions in missed appointments to benefit services. As outlined previously (9), the issue is often framed as a matter of 'fixing' problematic patient behaviours, with theoretical engagements limited to cognitive or behavioural domains, seeking to influence or 'nudge' behaviour: reminders; posters or letters seeking to induce commitment guilt or moral responsibility, and the threat sanctions including de-registrations, removal from waiting lists, or fines (10-14).

The missingness study

Where patients miss *multiple* appointments, the individual behavioural lens often results in overt or covert stigmatising and Othering narratives, with patients labelled as "repeat offenders" (15, 16), "chaotic" or irresponsible (14, 17). The complex health, personal and social circumstances of 'missing' patients' lives, evidenced in preceding epidemiological work (18-21), may also attract stigmatising narratives around poverty, substance use, mental health and long-term health conditions – areas that are also frequently ascribed to failures of individual responsibility (22-24).

Yet as our work has shown, the experience of 'missingness' – defined as "the repeated

tendency not to take up offers of care that has a negative impact on the person and their life

chances" (25) – has a range of complex, overlapping and mutually reinforcing causes that

occur at several points throughout patients' journeys to and through health services (19).

These include: the sense that healthcare is not 'for me' – not necessary, beneficial,

appropriate, or safe; past negative experiences including feelings of stigma, discrimination,

neglect and abuses of power, as well as broader issues of trauma and relational adversity; a

greater exposure to competing demands and crises, with fewer resources to reduce, mitigate

or manage those demands; issues managing gatekeeping and access systems, or the

unwritten rules of engagement; issues of travel, transport and the ability to move safely to

and through healthcare spaces; and an overall sense of mistrust and distrust (19). Societal

and institutional structures play a significant role in sustaining the conditions in which

missingness occurs. Stigma influences whether people possess the resources needed to

access or negotiate care and patterns their encounters with healthcare and other social

institutions **(**Baruffati et al forthcoming). Care itself is unequally distributed, withheld, and

often a poor fit with people's resources, their experiences of poor health, of their adverse

circumstances(26).

These experiences are rarely reflected in the design of interventions because they are

absent from the existing evidence base. Focused on service benefits, much of the evidence

is not stratified according to who is impacted and in what ways – and who may be missing or

absent (27, 28). In some studies, people are excluded by recruitment criteria – for example,

those with poor mental health, cognitive impairment, learning disabilities, with insufficient

English language skills, without phones, with incomplete practice records, or who are

otherwise deemed unsuitable (29-32)**.** Few studies reflect on self-selection bias and those

patient groups who may be less likely to accept or use the intervention being tested (33).

These absences are structurally patterned, with those excluded often at greater risk of

access issues or negative outcomes. Lacking a phone may be a sign of particular adversity

(34); the absence of practice data may reflect prior access issues, exclusion, or mis/distrust

(27, 35). The limited evidence that does consider patients most likely to be 'missing' shows

that they are either least likely to benefit from or are actively disadvantaged by interventions

designed to 'work' for all patients (27, 36, 37). Some consider cancellation as a positive

outcome as it reduces the unexpected non-attendance rate for the service (38-40) but this

still represents a missed opportunity for care and few studies explore *who* is most likely to

cancel, nor whether reductions represent a different form of exclusion as patients are

deterred from approaching health services in the first place (27). Thus, throughout the

evidence base, patients at risk of missingness also experience "structural missingness" (41),

with their situated and experiential knowledge absent from conversations about access to

healthcare. The aim of this paper is to address this substantial gap by reporting on a study

that synthesised extensive primary and secondary data to design a suite of co-produced,

evidence-informed interventions to address missingness in primary care.

Methods

The intervention development approach was guided by realist principles and by the Six

Steps in Quality Intervention Development (6SQuID) approach (42). Here we focus on the

first four steps: defining and understanding the problem; identifying malleable factors which

have scope for change; identifying how to bring about change (the active ingredient or

change mechanism); and exploring how to deliver the change mechanisms. The final two

stages involve implementing and evaluating the intervention and are not addressed here.

Interventions designed through 6SQuID involve "changing relationships, displacing existing

activities and redistributing and transforming resources" (42). They use a systems lens to

recentre the role of systemic and contextual causal factors, which are often absent from

predominantly behavioural approaches (43).

The stages of the 6SQuID process were addressed through the iterative synthesis of data

from the three workstreams of the "developing interventions to reduce 'missingness' in health

care" project. The study design was approved by the University of Glasgow College of

Medical, Veterinary and Life Sciences Ethics Committee in 2022 (Reference 00220187).

Workstream 1 was a realist review of literature, identified through database searching and

citation-tracking (see (9) for full methods) complemented by further citation tracking and

'umbrella' reviews of intervention ideas identified in other workstreams. A full PRISMA

diagram and list of included documents is available in sections 1 and 2 of the supporting

information. Workstream 2 consisted of realist interviews with 61 key informants whose

personal and professional experiences brought a broad range of clinical, social and inclusion

health perspectives on missingness. Participant recruitment began on 28[th] February 2023

and concluded on 11th July 2024. All participants provided informed consent in writing or,

where this was not possible, consent was provided verbally and audio-recorded prior to

interview. Interviews focused on participants' theories about the causes of missingness,

actions that might address these causes, and factors contributing to or inhibiting their

success (see (26) and **Baruffati et al forthcoming** for full methods). Interviews provided

further insight into missingness and addressed many of the gaps and shortcomings in the

research literature. A breakdown of interview participants is available in section 3 of the

supporting information.

The third workstream involved a series of Stakeholder Advisory Group (StAG) workshops.

StAG membership was largely drawn from interview participants, with 16 members (8

experts-by-experience and 8 professionals) who reflected a broad range of patient and

professional perspectives in the key areas of inclusion health; mental health; homelessness;

asylum and migration; substance use; and "severe and multiple disadvantage" (44). The

group met regularly throughout the project, acting as guides and sense-checkers for the

other workstreams and as co-designers of an overall interventional approach (45). The

contribution of each method to the 6SQuID approach is outlined in table 1, and the materials

used to guide stakeholder discussions are included in section 6 of the supporting information.

Participants in the StAG process provided informed consent in writing to participate in this

process.

TABLE 1: CONTRIBUTION OF WORKSTREAMS TO THE 6SQUID PROCESS

1. Defining and understanding the problem.
Review: Despite limitations, there is a body of critical literature providing an alternative to the dominant 'problematisation' of missingness (46) and a set of 'evidential fragments' (47) to build on. These provided empirical, conceptual and theoretical foundations for our theory of missingness.
Interviews: Helped address the gaps in the existing evidence base, brought insights from groups seldom heard in prior research, and provided nuanced understanding of the complexity of missingness including a contextual and structural understanding of the problem absent from the literature.
StAG: Provided guidance and a 'sense-check' for our understanding of the problem, helping refine, amend and address gaps. Also guided for recruitment strategies for interviews, and our approach to analysis and synthesis of the evidence.
2. Identifying malleable factors which have scope for change.
Review: Outlined a refined programme theory of the key causes of missingness and those factors considered amenable to change, organised into key domains (19).
Interviews: Provided further depth and breadth of insight into the refined programme theory, particularly relating to wider social structures and contextual conditions (26).
StAG: Validated the refined programme theory, then engaged in discussion of initial intervention ideas to address different causes at different 'levels' of the social-ecological model (48).
3. Identify how to bring about change (active ingredient/change mechanism).
Review and Interviews: Provided a set of intervention for discussion at the StAG.

StAG: Explored intervention activities and grouped them into domains linked to the causes they address**.** Using hypothetical patient case studies, the group considered which interventions might address the constellation of causes evident for each patient.

4. Exploring how to deliver the change mechanisms.

Review: Using priority domains from the StAG, an umbrella review looked in detail at several intervention domains not explored fully in the initial review and additional literature added further detail to other domains.

Interviews: Provided crucial contextual information from real-world examples of interventions: promising activities, key elements for successful implementation, and possible blockers or barriers.

StAG: The group outlined a set of core principles crucial to the success of intervention activities across different contexts (a 'missingness lens'). Through practice-level case studies, they explored resource requirements, challenges, drawbacks, risks, and unintended outcomes.

Moving back and forth between workstreams allowed each to address the limitations of the

others. For example, interventional literature is characterised by significant heterogeneity of

settings or approaches to the same activity, *and* by a lack of contextual description, making

assessment of implementation processes barriers, risks or unintended outcomes difficult

(49-51). These were then explored in interviews and stakeholder workshops. Similarly, the

alternative version of the problem evident in interviews and StAGs helped identify additional

causes at interpersonal, institutional, community and structural levels, and thus alternative

intervention activities for exploration in the iterative literature review process.

The process of analysis was similarly recursive. Literature was analysed in Nvivo, with data

relevant to the causal working and contextual dynamics of interventions extracted and

synthesised into an explanatory narrative in potential intervention domains. Interviews were

analysed separately in Nvivo, coded abductively according to a theoretical framework

combining fundamental causation theory and the candidacy framework (see **Baruffati et al**

(forthcoming) for further details). Findings from the literature were refined further by

findings from interviews, and both used to create materials presented to and refined further

by StAG members. Audio recordings and visual materials from StAG meetings were

summarised and synthesised with discussions from previous meetings into a final

intervention. The final suite of interventions detailed here was synthesised by the research

team using summary documents from all three workstreams and was validated by

participants at the final StAG. Given the quantity of the data collected, and the nature of the

synthesis process above, results are presented as a narrative without primary or secondary

data specifically referenced. The contribution of each work package to each intervention

domain is described instead in section 7 of the supporting information. Primary data are

available in the final project report (forthcoming) and are available on the Open Science

Framework (52), congruent with the principles of open research**.**

Results
177

178 ## Key principles: a missingness lens
179

The intervention presented here is more accurately a 'suite' of interventions, or a "recyclable

core set of processes" that can be applied according to the requirements of different

contexts (53). Their successful implementation depends on creating "congruence" (53)

between each part by rooting them within a set of principles termed 'the missingness lens'

(see figure 1). These principles were defined during a StAG meeting, and they represent a

fundamental shift in how the 'problem' of missingness is understood and how solutions to it

should be designed and implemented.

The 'situational' model		A missingness lens.
Patient 'responsibilisation.'	→	Services committed, resourced, incentivised to identify and address barriers.
Shallow, monocausal perspective.	→	Complex causality for individuals, in contexts

Firstly, instead of considering missingness as a problem for services caused by patient

behaviour, a missingness lens means services commit to directing resources to addressing

missingness as a problem for patients and their health. Rather than designing, implementing

and measuring interventions across an unstratified patient population, a missingness lens

involves both "proportionate universalism" (54)- shifting resources towards those in greatest

need - and "positive selectivism" (55) by targeting specific patients and evaluating outcomes

for them. It means moving from shallow, simple explanations towards deep and complex

causes at multiple levels: structural influences on patients' circumstances and resources;

institutional dynamics of service provision; and interpersonal and intrapersonal processes. It

also means moving beyond logistical, technical or practical solutions by recentring affective,

psychological and relational elements, particularly around *safety* - psychological safety

through trauma-informed practice and psychologically-informed environments (56); and

cultural and structural safety by identifying and addressing sources of stigma, discrimination

and power imbalance (57). This includes a central role for patient-centred, collaborative care.

Finally, a missingness lens moves beyond biomedical and clinical approaches to understand

health and healthcare in wider context through condition competency – "clinical

understanding of the lived-experience of specific conditions" (26) – and structural

competency, the ability to recognise and to influence the social and structural determinants

of health (58).

A 'suite' of interventions

The specific suite of actions outlined below represent an attempt to disrupt the systemic

causal dynamics sustaining missingness, either directly or by pursuing intermediate

outcomes in support of this goal (43, 59). They are limited to those actions which health

services have the power to change, while cognisant of the need for broader policy changes

to address social determinants of missingness and of illness. The actions are split into

several interconnected domains but are likely to be most impactful when synthesised rather

than as a set of discrete elements. The domains are represented in figure 2 below, followed

by detailed discussion of how each can by implemented in a way that is congruent with the

missingness lens.

FIGURE 2: A 'SUITE' OF INTERVENTIONS

Embedding a missingness lens

The first step in addressing missingness is in embedding the elements of this perspective

shift at all levels of health service organisation and delivery. Participants often described the

need for a "culture change" towards missed appointments at multiple levels - clinical,

administrative, policy, planning, and funding. Resources including money, time, staffing, data

infrastructure, knowledge and skills, need to be mobilised and prioritised to ensure this

culture change is matched with concrete changes in practices and systems. At the broadest

level, this means services having the money to ensure that missingness-focused work –

likely to be complex, time- and labour-intensive – is supported, encouraged and even

incentivised. Many participants felt that current NHS funding structures like the Quality and

Outcomes Framework actively disincentivise missingness-related work, with "exception-

reporting" used to exclude patients from monitoring and reporting, rather than incentivising

work to get patients into practices. Without specific funding, even the best intentions would

remain a "fantasy" or a "dream." Funding should permit localised innovation and plans for

making and monitoring change through multi-component quality improvement methods:

setting clear and transparent goals, planning and carrying out key actions, and regularly

reviewing or amending them through inclusive, proactive systems for accountability. Bringing

together staff and patient perspectives collaboratively is a crucial part of this process.

Targeted action, built on solid principles and localised planning, matched with resources

(staffing, knowledge, financial), acts as the foundation upon which the activities below can

be built. Monitoring and accountability mean going beyond service-side metrics of missed

appointment rates to measure impacts on patients qualitatively and quantitatively. They

should also occur above individual care delivery, at the strategic and policy level.

Several other actions are suggested to support the embedding process. The first involves

training for staff, using knowledge and skills-development to "displace" (42) existing

perspectives – particularly stigmatising attitudes or opinions among staff towards

missingness or 'missing' patients and their health-related or help-seeking behaviours.

Training should be provided to all staff at all levels – from administrative to clinical to

commissioning – to ensure congruence and consistency. In theory, raising awareness of the

negative health outcomes around missingness, of its main causes, and of actionable

solutions, may lead to increased staff motivation and empathy, to buy-in for meaningful

action, and for patients the sense that services are supportive and safe. All three

workstreams suggested other, complementary areas for professional development, including

poverty and inclusion health, stigma and discrimination, mental health, psychological trauma,

neurodivergence, and attachment. Training should be designed to help staff reflect on

practice systems and on relationships and communication dynamics: how and why patients

might present differently; how to manage challenging encounters; and how to enact

principles of relational care and to create safe environments. Expertise-by-experience is

important in formulating and delivering training activities to drive awareness, empathy and

change.

However, focusing exclusively on staff attitudes to embed the missingness lens risks simply

reversing the direction of blame without acknowledging staff who are motivated to change

practice but are inhibited by limited resources and capacity and the burnout they cause.

Knowledge, skills and motivation need to be matched by concrete resources to turn

motivation into meaningful action. Staff need the protected time and capacity to engage with

training and development, and providers the capacity to allow staff to attend as well as the

access to training resources and expertise. To allow the missingness lens to displace

existing practices, training needs to be complemented by supervision and reflective space,

positive reinforcement and feedback, and ongoing support for staff. There is also a role for

utilising existing staff knowledge and experience of missingness. Staffing changes may be

appropriate, including actively recruiting new clinical and non-clinical staff from communities

identified as being at risk of missingness, who may bring knowledge of barriers to care and

may act as a bridge to those communities.

Identification
273

To build a localised plan for missingness, and to monitor its success, practices and services

require IT infrastructure and data recording systems to monitor 'missingness' trends and

outcomes. Services need systems to identify who is missing through retrospective analysis

of appointment trends, and to identify and flag those who become missing in real-time. Past

work has demonstrated some difficulties in using UK primary care systems in this way (18,

21, 60). Practices should prioritise the creation of robust coding of patient records,

particularly of non-attendance, and should seek to address missing data which is often a site

of inequality (41). Services can complement this with other ways to identify those who are

missing, for example notifications of 'missingness' from referrals to secondary or tertiary care,

or staff knowledge of the patient population. While recently 'machine learning' approaches

have been designed to play this role, caution is needed due to concerns about inaccuracy,

bias and the absence of a critical approach to causes in these systems (**Ellis et al,**

forthcoming).

In carrying out identification, services must decide on their thresholds for 'missingness',

particularly considering current resource environments. The initial epidemiological work on

missingness selected those with 2 or more missed appointments over a three-year period,

based on a pilot study and stakeholder group validation (18, 21, 61). Missingness impacted

19% of the total patient population – a cohort of patients with complex health and social

needs (18, 20, 21). Uniformly applying this intervention to all these patients simultaneously

risks diluting available resources and limiting impact. To resolve this, services may elect to

focus only on those who miss the greatest number of appointments, or whose patient

profiles suggested particular vulnerabilities. Crucially, this work is dynamic and recursive and

can change as projects develop, and as ideas are tested and approaches refined over time.

Services should complement pattern-identification with identification of the causes of

missingness, carried out by contacting those experiencing it and having non-judgemental,

respectful and invitational conversations about their experiences. These conversations may

have instrumental benefits in identifying individual and collective causal patterns, and may

also represent the first step in a change in relationships (42). Staff become more aware and

understanding of patients' circumstances; patients feel a sense of care and effort on the part

of services that may previously have seemed uncaring or exclusionary. More broadly,

connections with key community stakeholders, community groups or settings may also

improve identification of access barriers among specific groups. Within a QI approach,

regular monitoring and feedback inputs from staff, individual patients, and across the

practice population on areas relevant to missingness acts as a point of ongoing identification.

The findings from identification can be used in two ways. Firstly, they can feed into broader

practice changes aimed at improving access to care. Secondly, they can underpin an

approach to positive selectivism and proportionate universalism, where particular patients or

groups are targeted for tailored responses and additional resources (54, 62). This is closer to

individual *needs* identification*,* covering the specific and proximate causes of missingness in

a collaborative and person-centred way, actively seeking and building on patients' priorities

and perspectives, and resulting in a concrete and tailored set of actions from services. A

Patient Individual Needs (PIN) document could be created, detailing a patient's key

information and priorities; their access needs and communication preferences; and the steps

to be taken by services to support them. Services may also seek insight from significant

people or services in patients' lives (with their consent). The findings of these assessments

become the foundations of consistency, continuity and accountability, reducing patients'

needs to advocate for access or additional support, or to repeatedly disclose sensitive

information to staff in their pursuit of care. Stakeholders felt that this document should be

NHS-wide, ensuring consistency across the system. In line with a person-centred approach

these assessments and the support following from them should not be limited to clinical or

access issues. Possessing few resources to manage competing demands in multiple

domains is a central cause of missingness; if not identified and addressed, these causes of

missingness and of ill health are likely to endure (9).

Relationships and Communication

327
328

All workstreams explored the importance of relationships in addressing missingness, and the

centrality of trust, consistency, continuity and safety for patients whose service interactions

have often been patterned by stigma, discrimination, exclusion, neglect, even hostility. A

missingness lens involves recentring the relational causes of missed appointments and

including relational components throughout the intervention – for example, the role of

training in relational care and trauma-informed, non-stigmatising practice in embedding a

missingness lens, or the relational work of reaching out to patients to identify their needs.

The outcome of these actions, and the core of a relational approach, is in creating

consistency and continuity at every stage of the patient journey, from registration to

consultation and follow-up. Patients' anxieties or fears will likely be reduced if they can trust

in a response that is consistently welcoming, invitational, non-judgemental and supportive,

not reprimanding, punitive, coercive or manipulative. For those with profoundly negative

relational histories, this is even more important. This is not to say that these relationships are

unboundaried – instead, clear, agreed boundaries and reciprocal expectations are central to

consistency.

Several concrete actions are suggested here. Extending efforts at continuity of care to

'missing' patients is central. Being able to see a familiar clinician may reduce anxieties about

attendance and increase feelings of safety and security and may also create a sense of

reciprocity or loyalty and a relationship to be protected by attending. Being familiar to a

clinician or key staff member may encourage "condition competency" (26) and "structural

competency" (63) and care plans built upon patients' needs, circumstances and preferences.

Continuity does not have to mean seeing the same person at every interaction. A dedicated

small team of clinical and non-clinical within practices focused on missing patients may

provide sufficient continuity and retain some flexibility for practices. If patients can contact

practices with the knowledge of who they might speak to and how they might be received, by

staff aware of their circumstances and needs, anxieties around gatekeeping and initial

access may be reduced. Other concrete actions include identifying and catering to patients'

communication needs and preferences in all interactions (e.g language barriers, literacy

issues, cognitive challenges, communication preferences).

Relational work takes time. Given the levels of mistrust and past trauma often evident in

missingness, people may not instantly respond to relationship-building work. Identification-

work will often require a relationship to be established over time, so that the individual feels

safe and able to disclose important information about themselves. Persistence is central, but

there is also a need to respect patients' right not be involved and to avoid being overly

persistent, coercive or manipulative, or perpetuating problematic power dynamics. A patient-

centred and collaborative approach to the intervention and to clinical practice, coupled with a

non-judgemental, supportive approach, may contribute to a sense that care is 'for me' both

instrumentally and relationally.

Missingness Coordination
367
368

The additional work required to address missingness brings the question of *who* is best

positioned to do it. Addressing missingness effectively requires a 'missingness coordinator',

a role akin to other non-clinical roles (e.g navigators, care coordinators, peer workers,

mentors, community outreach workers) but oriented specifically around missingness.

Missingness coordination involves several key tasks. Firstly, coordinators carry out

identification-work and build relationships with patients, seeking to develop trust through

active, empathetic, non-judgemental and collaborative support. Coordination is a core part of

relational continuity and consistency, a reliable point of contact for the patient and an

advocate for them and for good relational practice. In this, missingness coordinators engage

in bridging, brokering or mediating activities aimed at creating "safe passage" (64) to and

through services. Bridging activities may include working with patients to build their personal

resources – confidence, motivation, self-esteem, even expectations for health – as well as

supporting changes in service provision through individual, institutional and systemic

advocacy, such as the advocacy against austerity carried out by Deep End GPs (65). Care

needs to be taken to avoid these workers being made solely responsible for missingness, or

their work being marginalised or even co-opted by existing problematic practices. This

requires that coordinators be skilled and well-supported, and that they be embedded in

services in ways that allow them to change practice in line with the missingness lens.

Using a holistic, person-centred approach to identification, coordination extends beyond

primary care settings. It may involve coordinating healthcare across multiple service settings

to reduce treatment burden, or brokering access to wider resources aimed at addressing

social determinants of illness, social needs or the precariousness that impedes good health

and healthcare access. The shape of the work will vary from patient to patient, and thus

coordination requires flexibility, open-endedness and a tailored and person-centred

approach.

Additional considerations include who is best placed to act as a coordinator, and how they

might be funded or positioned to maximise their impact. Expertise-by-experience and 'peer-

ness' may provide a route to credibility, relationship-building and cultural and structural

safety, although stakeholders and interviewees felt that coordinators can come from a range

of personal and professional backgrounds as long as they are sufficiently trained, skilled and

supported to engage with complex and difficult circumstances. This, plus the need to 'embed'

workers as equal partners, suggested that paid staff were more appropriate than volunteers.

While NHS funding may provide a route to embeddedness, third-sector independence may

be a virtue for those mistrustful of the healthcare system.

Flexibility
404

Given inflexibility or 'impermeability' of services are a central cause of missingness (19),

addressing it means changing practice and service systems to be more flexible and suitable

for different patient circumstances. There is strong support across all strands of work for

prioritising 'missing' patients for different forms of tailored flexibility. This includes flexibility in

how appointments are made – in person, online, outside the usual appointment-making

windows, bypassing reception gatekeeping or telephone triage, booking via coordinators or

directly with 'missingness'-focused staff who can facilitate access. Offering greater choice in

when appointments occur may maximise convenience and minimise interference from other

demands, as may offering appointments outside of regular hours to those experiencing

barriers at other times. Prioritising missing patients for rapid access, drop-in access or open

appointment slots may allow providers to minimise delays or take advantage of the "window

of opportunity" (27) that exists at the point where care is sought. Offering longer

appointments, underpinned by person-centred consultation styles, may have multiple

benefits – more time and space to build rapport, for patients to describe needs,

circumstances, concerns or preferences, and to discuss and negotiate proposed courses of

action. These also become an opportunity to maximise the impact of each single

appointment, and thus the perceived benefits of attending in future. Similarly coordinated

care planning, or having the flexibility to combine appointments with other health or non-

health service providers, may increase the value of attendance and minimise the burdens of

managing multiple engagements.

Alongside temporal flexibility, services should offer flexibility in *who* patients see (see

relationships and communication above) and in *where* patients are seen. This may mean a

role for home visits, or for outreach/in-reach into key settings where identification work

suggests 'missing' patients are likely to be found (e.g. homelessness services, substance

use services, community organisations). In the early stages of missingness work, patients

may feel safer and more secure in a familiar environment and their transport and logistical

issues may be resolved. Outreach or in-reach may build the initial relational links that lead to

more regular engagement or contribute to word-of-mouth and the engagement of other

'missing' patients. Remote care through telephone or video appointments is also an option:

patients may feel safer or more secure in their home environments; travel costs, time or

difficulties may be eliminated, as may competing demands like work or childcare.

Flexibility also means making allowances or accommodations for lateness and avoiding

punitive or exclusionary responses to non-standard presentation or continued non-

attendance. The shape of flexibility and its limits can be discussed and agreed with patients

during initial identification, to ensure both they and service providers have a set of clear

expectations and boundaries, agreed and recorded in the PIN. Crucially, it should be offered

actively and gladly, neither guarded nor gatekept, as this may demonstrate a level of care

and commitment on the part of the service that patients respond to - but only if carried out

through a missingness lens without which interventions risk being ineffective or exclusionary.

Consider telephone or video appointments, where many of the causes of missingness

endure and new causes emerge: a lack of access to technology, limited tech literacy,

changing contact details, the costs of data or internet access, complex or unreliable systems,

mistrust and concerns about privacy or data security, the lack of safe space, the loss of

relational immediacy and limitations on rapport and communication. Carried out with an

embedded missingness lens, and designing systems with and for 'missing' patients, mean

these systems are likely to 'fit' patients' preferences or circumstances. This may include

using easily accessible systems, ensuring communication preferences are addressed, and

providing support through coordinators to access or use devices or data, connect to systems

and facilitate communication. Finally, there is a role for coordinators facilitating flexibility

beyond primary care, where 'missing' patients might be prioritised to overcome access

barriers in secondary and tertiary care including long delays, requirements for re-referral

after missed appointments, and exclusionary eligibility criteria.

Transport and Logistics

As well as being an aspect of flexibility, changing the site or the mode of an appointment is a

logistical intervention aimed at reducing barriers related to travel – costs and time, reliance

on support networks or unreliable transport systems, limited mobilities, poor mental health,

feelings of unsafety (19). There is a spectrum of possible actions here, their suitability

dependent on the circumstances of patients and the resources available to practices. At the

least intensive end, practices may consider reimbursing travel costs, although this depends

on patients having sufficient money to cover costs initially. Services may pre-emptively

provide tickets, tokens, vouchers or parking permits instead. Facilitated access to NHS or

voluntary transport services may also be beneficial. At the more intensive end, taxi transport

and home visits may be most appropriate, as may accompaniment by a missingness

coordinator to support safe "wayfinding" (66). Accompaniment can also mean supporting

patients within consultations – supporting the exchange of information, providing

reassurance or advocacy, and reducing anxieties or worries about attending. Systems for

arranging transport need to be accessible, inclusive, and actively offered, as often they are

described as inaccessible, complex, inflexible and either actively gatekept or implicitly

patterned by eligibility criteria or narratives of deservingness that deter access. Changing

service spaces using trauma-informed principles, or setting aside quiet or private waiting

rooms, are also crucial to ensuring people feel safe and secure.

Contact around appointments

Rather than a domain of 'reminders', our evidence suggests an expansive approach to

patient contact around appointments. Reminding has a role, but reminders alone will not

address deeper, broader and more enduring barriers than the simple forgetfulness that most

systems aim to address. Basic SMS, automated or email reminders are a minimum standard

– the medium determined by patients' contact preferences, with messages including details

of the appointment (time, location, staff member, even purpose). "Stepped" reminders (27)

may be beneficial – multiple messages, perhaps sent using different methods, with one

reminder far enough in advance to allow patients to plan and other closer to the time to

ensure patients do remember. Patients' preferences should again guide this action.

Concerns about data security, privacy and confidentiality should be discussed and

addressed where possible – and where patients express a preference to avoid reminders,

this should be respected. Practices should prioritise systems for patients to respond to

reminders with minimal barriers in order to discuss, cancel or rearrange their appointments.

They should work to obtain patients' contact details and preferences on practice systems,

and to maintain these systems to account for changes, otherwise these systems risk

perpetuating inequalities. Patients should have the option to identify significant others to

contact if they are unavailable.

Yet simply increasing the intensity or quantity of reminders, or tweaking wordings or formats,

are unlikely to address deeper causes. There is no support here for behavioural 'nudges' in

contact around appointments, which were seen to be stigmatising, coercive and alienating.

Instead, services should explore personalised pre-appointment contact with identified

patients. Personalised contact, carried out by coordinators, may help the relationship-

building process with an invitational, welcoming and supportive approach. It may help

support orientation as staff and patients discuss what to expect, explore concerns and

potential benefits, and identify barriers that are then addressed by actions in other

intervention domains.

Contact around appointments also includes contact after appointments, particularly if they

are missed, as an opportunity to express care and concern, maintain connection and to

check in on the patient's wellbeing and any health needs or unaddressed access barriers.

This should be framed as an invitational offer of support, rather than as a reprimand. The

approach to contact should be outlined during identification and agreed in the PIN, so

patients know they will be contacted and who will be contacting them, by what method, and

with what purpose. Crucially, contact around appointments needs to be two-way. Patients

must have the means to respond to written reminders to query or amend details or discuss

any concerns. There also needs to be room for patient-initiated contact between or around

appointments, whether via clinical teams, dedicated contact lines, or through coordinators,

otherwise contact systems risk perpetuating asymmetries where services can freely contact

patients but the patient cannot contact them in return.

Other approaches
518

Several possible actions are not included here. Some are excluded because stakeholders

did not discuss them or because they saw limited value or active drawbacks to using them.

Incentivisation of attendance (financial, in kind, or contingency management) did not feature

strongly, perhaps reflecting concerns about making healthcare transactional rather than

relational, and perpetuating power dynamics and trust. There was no support for fines,

punishments or sanctions, or for overbooking approaches that are essentially punitive and

were described by one StAG member as "sinister." Other approaches are promising but fall

beyond the purview of individual services – such as 'hub' models and integrated services,

bringing multiple support services into spaces and thus increasing coordination and reducing

the burdens of accessing multiple services. Other aspects are simply beyond the power of

health services to change – poverty, austerity, poor housing, the migration system,

stigmatising policy discourse, and other determinants of illness – although these continue to

act as major impediments to care and to good health. The health service does retain an

influencing and advocacy role here, however, as the site where many of the ill effects of

broader policy are often most visible.

534   Discussion and conclusions

This paper outlines the main components of a major paradigm and practice shift about

multiple missed appointments in primary care, based on the application of a missingness

lens and of key resources targeted at addressing the complex causes of missingness. The

suite of interventions outlined here represent a core set of activities to be applied flexibly

according to local contextual factors. A comprehensive approach to missingness would

include all the proposed actions, but we recognise the constraints of current primary care

systems and the need to prioritise some patients, and some actions, until resourcing permits

a more expansive approach.

This study has several strengths. Faced with a literature base built without key perspectives,

causal insights or theories of change, our stakeholders and interviewees brought alternative

(and seldom-heard) understandings and theories from their direct, contextualised

experiences (67). Using knowledge from within systems to build an intervention increases its

relevance, practicality and feasibility, and thus its potential for success (42, 68). While there

was a high degree of congruence between workstreams, areas of disagreement between

data sources could often be accounted for by the flaws of the literature base, or simply by

difference of experience, context, preference or positionality between participants. What

'works' for one person may not work for another. These differences illuminate, rather than

obscure, important causal and contextual dynamics, and highlight the importance of a

tailored approach. A further strength relates to the relevance of this work beyond primary

care, as both the primary and secondary data include evidence from across the healthcare

system.

Although this study is evidence-informed and based on thorough theorising, it does not test

or evaluate the proposed intervention in practice. However, it does provide some

implications for future research and evaluation of interventions into missingness and missed

appointments more generally. Some we have detailed previously – the need to change the

framing of missed appointments, to include data and analysis on missingness, and to use

inclusive methods of recruitment and data collection (19). Ensuring data is equally available

for 'missing' patients, acknowledging where and why it is not, and stratifying analysis of

intervention effects according to how they impact those experiencing missingness, are

crucial ways to counter "structural missingness" (41). Interventional research should also

carefully consider what outcomes are measured, with a stronger focus on patient health and

wellbeing and on patients' *chosen* outcomes in a person-centred approach. It should not

only measure numerical reductions in missed appointments, or in missingness, but should

also explore intermediate outcomes related to causes: do patients feel cared for, supported

and understood? Is their experience of care or its quality improved? Does their health or

quality of life improve? Crucially, how are these improvements unequally distributed? Finally,

greater reporting of the context and process of interventions is crucial to improving an

evidence base where these insights are limited.

A further limitation of this study relates to its 'level' of focus. During intervention design,

contributors considered interactions within and between multiple levels of focus – structural,

community, institutional, individual – before focusing on changes within the power of health

services. Consequently, many causal factors fall beyond our remit, particularly structural

factors: growing poverty and inequality, austerity politics, crumbling transport infrastructure,

the decimation of the third sector safety net; hostile immigration policies, and the systematic

underfunding of health and social services that erode their capacity and erode public trust

within them. If these issues continue to worsen there is a risk that institutional and practice

changes become "fantasy paradigms" unable to mitigate deeper and more powerful forces

(69).

The suite of interventions outlined here are highly congruent with several recent papers on addressing access and health inequalities in the UK (62, 70-72). Beyond specific Inclusion Health settings such as homelessness health care there are already encouraging mainstream care examples, such as the Inclusion Health Action in General Practice (HAGP) approach in Scotland. 'Deep End' practices were resourced to address inequalities, with missingness chosen as a point for action, and were able to evidence improved access and improved health, better relational environments, and improved staff morale and support (60). The unique contribution of this work is that missingness provides a tangible and measurable focal point for work to address both access inequalities and broader health inequalities. These problems are often framed as being particularly complex and intractable, overwhelming and demoralising to those seeking to address them. Addressing missingness provides an anchor point, a framework and a site for meaningful change, and with the application of resources to the problem evidence suggests that profound transformation can be achieved (60, 64, 73-75).

Acknowledgements

The research team would like to acknowledge the invaluable contribution of interview participants and members of our Stakeholder Advisory Group to this work. In addition Elspeth Rae, University of Glasgow, provided administrative support; Claire Duddy, formerly university of Oxford, was the information specialised who devised and conducted initial literature searches; and Jack Brougham, illustrator, provided illustrations used in the paper and throughout the project.

References

1.   Alturbag M. Factors and Reasons Associated With Appointment Non-attendance in Hospitals: A Narrative Review. Cureus. 2024;16(4):e58594.

2. Organization WH. World health statistics 2024: monitoring health for the
SDGs, sustainable development goals: World Health Organization; 2024.

3. Darzi A. Independent investigation of the NHS in England. London: HMSO;
2024.

4. Greener I, Powell M. Crisis in the UK National Health Service: What does it
mean, and what are the consequences? Social Policy & Administration.
2024;58(2):313-28.

5. Beech J, Fraser C, Gardner T, Buzelli L, Williamson S, Alderwick H. Stressed
and overworked. Health Foundation; 2023.

6. McPake B, Dayal P, Zimmermann J, Williams GA. How can countries respond
to the health and care workforce crisis? Insights from international evidence. The
International Journal of Health Planning and Management. 2024;39(3):879-87.

7. NHS drive to reduce 'no shows' to help tackle long waits for care [press
release]. NHS England, 28/01/2023 2023.

8. Missed GP appointments costing NHS millions [press release]. NHS England,
2/01/2019 2019.

9. Lindsay C, Baruffati D, Mackenzie M, Ellis DA, Major M, O'Donnell CA, et al.
Understanding the causes of missingness in primary care: a realist review. BMC
Med. 2024;22(1):235.

10. Aggarwal A, Davies J, Sullivan R. "Nudge" and the epidemic of missed
appointments Can behavioural policies provide a solution for missed appointments in
the health service? J Health Organ Manag. 2016;30(4):558-64.

11. Teo AR, Niederhausen M, Handley R, Metcalf EE, Call AA, Jacob RL, et al.
Using Nudges to Reduce Missed Appointments in Primary Care and Mental Health:
a Pragmatic Trial. Journal of General Internal Medicine. 2023;38(SUPPL 3):894-904.

12. Ruggeri K, Folke T, Benzerga A, Verra S, Buttner C, Steinbeck V, et al.
Nudging New York: adaptive models and the limits of behavioral interventions to
reduce no-shows and health inequalities. BMC Health Services Research.
2020;20(1).

13. Martin SJ, Bassi S, Dunbar-Rees R. Commitments, norms and custard
creams - a social influence approach to reducing did not attends (DNAs). J R Soc
Med. 2012;105(3):101-4.

14.   Dathan M. Patients could soon be charged for missing NHS appointments as
Health Secretary Jeremy Hunt backs the idea 'in principle'. The Independent. 2015
03/07/2015.

15.   BBC. Darlington sees 'shocking' increase in missed GP appointments. BBC.
2024 04/01/2024.

16.   Hussain-Gambles M, Neal RD, Dempsey O, Lawlor DA, Hodgson J. Missed
appointments in primary care: questionnaire and focus group study of health
professionals. British Journal of General Practice. 2004;54(499):108-13.

17.   Department for Levelling Up HaC. Frontline support models for people
experiencing multiple disadvantage: A Rapid Evidence Assessment. London: HMSO,
Department for Levelling Up HaC; 2023 April 2023.

18.   Ellis DA, McQueenie R, McConnachie A, Wilson P, Williamson AE.
Demographic and practice factors predicting repeated non-attendance in primary
care: a national retrospective cohort analysis. Lancet Public Health.
2017;2(12):E551-E9.

19.   Lindsay C, Baruffati D, Mackenzie M, Ellis DA, Major M, O'Donnell CA, et al.
Understanding the causes of missingness in primary care: a realist review. BMC
medicine. 2024;22(1):235.

20.   McQueenie R, Ellis DA, Fleming M, Wilson P, Williamson AE. Educational
associations with missed GP appointments for patients under 35 years old:
administrative data linkage study. BMC medicine. 2021;19(1):219.

21.   McQueenie R, Ellis DA, McConnachie A, Wilson P, Williamson AE. Morbidity,
mortality and missed appointments in healthcare: a national retrospective data
linkage study. BMC Medicine. 2019;17:9.

22.   Spratt TJR. Reconceptualising Judith Butler's theory of 'grievability'in relation
to the UK's 'war on obesity': Personal responsibility, biopolitics and disposability. The
Sociological Review. 2022;70(3):474-88.

23.   Roy A, Buchanan J. The paradoxes of recovery policy: Exploring the impact of
austerity and responsibilisation for the citizenship claims of people with drug
problems. Social Policy & Administration. 2016;50(3):398-413.

24.   Brown RC, Maslen H, Savulescu J. Against moral responsibilisation of health:
Prudential responsibility and health promotion. Public Health Ethics. 2019;12(2):114-
29.

25.   Lindsay C, Baruffati D, Mackenzie M, Ellis DA, Major M, O'Donnell K, et al. A
realist review of the causes of, and current interventions to address 'missingness' in
health care. NIHR Open Research. 2023;3.

26.   Mackenzie M, Baruffati D, Lindsay C, O'Donnell K, Ellis D, Simpson S, et al.
Fundamental causation and candidacy: Harnessing explanatory frames to better
understand how structural determinants of health inequalities shape disengagement
from primary healthcare. Social Science & Medicine. 2025;374:118043.

27.   McLean S, Gee M, Booth A, Salway S, Nancarrow S, Cobb M, et al. Targeting
the use of reminders and notifications for uptake by populations (TURNUP): a
systematic review and evidence synthesis. Health Services and Delivery Research.
2014;2(34).

28.   Fien S, Dowsett C, Hunter CL, Myooran J, Sahay A, Menzel K, et al.
Feasibility, satisfaction, acceptability and safety of telehealth for First Nations and
culturally and linguistically diverse people: a scoping review. Public health.
2022;207:119-26.

29.   McLean SM, Booth A, Gee M, Salway S, Cobb M, Bhanbhro S, et al.
Appointment reminder systems are effective but not optimal: results of a systematic
review and evidence synthesis employing realist principles. Patient Prefer
Adherence. 2016;10:479-99.

30.   Fairhurst K, Sheikh A. Texting appointment reminders to repeated non-
attenders in primary care: randomised controlled study. Quality & Safety in Health
Care. 2008;17(5):373-6.

31.   Shao CC, Katta MH, Smith BP, Jones BA, Gleason LT, Abbas A, et al.
Reducing no-show visits and disparities in access: the impact of telemedicine.
Journal of Telemedicine and Telecare. 2024:1357633X241241357.

32.   Carrillo de Albornoz S, Sia KL, Harris A. The effectiveness of
teleconsultations in primary care: systematic review. Fam Pract. 2022;39(1):168-82.

33.   Kannenberg B, Stadter G. Analysis and observations of telehealth in primary
care follow up appointments for vulnerable populations. WMJ. 2022;121(2):116-20.

34.   Anyaegbu CT. SMS reminders: reducing DNA at a community mental health
depot clinic. Journal of Community Nursing. 2021;35(1).

35.   Perron NJ, Dao MD, Kossovsky MP, Miserez V, Chuard C, Calmy A, et al.
Reduction of missed appointments at an urban primary care clinic: a randomised
controlled study. BMC Fam Pract. 2010;11.

36. Shah SJ, Cronin P, Hong CS, Hwang AS, Ashburner JM, Bearnot BI, et al.
Targeted Reminder Phone Calls to Patients at High Risk of No-Show for Primary
Care Appointment: A Randomized Trial. Journal of General Internal Medicine.
2016;31(12):1460-6.

37. Rowett M, Reda S, Makhoul S. Prompts to Encourage Appointment
Attendance for People With Serious Mental Illness. Schizophrenia Bulletin.
2010;36(5):910-1.

38. Robotham D, Satkunanathan S, Reynolds J, Stahl D, Wykes T. Using digital
notifications to improve attendance in clinic: systematic review and meta-analysis.
BMJ Open. 2016;6(10):14.

39. Bean AG, Talaga J. Appointment breaking: causes and solutions. Journal of
Health Care Marketing. 1992;12(4):14-25.

40. Mehra A, Hoogendoorn CJ, Haggerty G, Engelthaler J, Gooden S, Joseph M,
et al. Reducing patient no-shows: an initiative at an integrated care teaching health
center. Journal of Osteopathic Medicine. 2018;118(2):77-84.

41. Hopkins Walsh J, Dillard‐Wright J. The case for "structural missingness:"
A critical discourse of missed care. Nursing Philosophy. 2020;21(1):e12279.

42. Wight D, Wimbush E, Jepson R, Doi L. Six steps in quality intervention
development (6SQuID). J Epidemiol Community Health. 2016;70(5):520-5.

43. Moore GF, Evans RE, Hawkins J, Littlecott H, Melendez-Torres G, Bonell C,
et al. From complex social interventions to interventions in complex social systems:
future directions and unresolved questions for intervention development and
evaluation. Evaluation. 2019;25(1):23-45.

44. Bramley G, Fitzpatrick S, Sosenko F. Mapping the "hard edges" of
disadvantage in England: Adults involved in homelessness, substance misuse, and
offending. The Geographical Journal. 2020;186(4):390-402.

45. Malengreaux S, Martens M, Castellano Pleguezuelo V, Olujuwon I, Van Belle
S, Renmans D, et al. Stakeholder involvement in realist evaluation: A scoping review
and best fit framework synthesis. Evaluation. 2025;31(1):22-48.

46. Bacchi C. Why study problematizations? Making politics visible. Open journal
of political science. 2012;2(1):1-8.

47. Pawson R. Digging for Nuggets: How 'Bad' Research Can Yield 'Good'
Evidence. International Journal of Social Research Methodology. 2006;9(2):127-42.

48.   Bronfenbrenner U. The ecology of human development: Experiments by
nature and design: Harvard university press; 1979.

49.   Ali-Faisal SF, Colella TJ, Medina-Jaudes N, Scott LB. The effectiveness of
patient navigation to improve healthcare utilization outcomes: A meta-analysis of
randomized controlled trials. Patient Education and Counseling. 2017;100(3):436-48.

50.   Crable EL, Biancarelli DL, Aurora M, Drainoni ML, Walkey AJ. Interventions to
increase appointment attendance in safety net health centers: A systematic review
and meta‐analysis. Journal of Evaluation in Clinical Practice. 2021;27(4):965-75.

51.   Godoy L, Hodgkinson S, Robertson HA, Sham E, Druskin L, Wambach CG, et
al. Increasing mental health engagement from primary care: The potential role of
family navigation. Pediatrics. 2019;143(4).

52.   Developing interventions to reduce 'missingness' in health care [Internet].
2023.

53.   Pearson M, Brand S, Quinn C, Shaw J, Maguire M, Michie S, et al. Using
realist review to inform intervention development: methodological illustration and
conceptual platform for collaborative care in offender mental health. Implementation
Science. 2015;10:1-12.

54.   Marmot M. Fair Society, Healthy Lives: The Marmot Review: Strategic Review
of Health Inequalities in England Post-2010. . London: Marmot Review; 2010.

55.   Potter L, McDonagh L, Horwood J, Farr M, Feder G, Wong G. Improving
access to general practice for people with severe and multiple disadvantage: a
realist review protocol (The Connection Study). Wellcome Open Res. 2023;8:331.

56.   Machtinger EL, Cuca YP, Khanna N, Rose CD, Kimberg LS. From treatment
to healing: the promise of trauma-informed primary care. Women's Health Issues.
2015;25(3):193-7.

57.   Curtis E, Jones R, Tipene-Leach D, Walker C, Loring B, Paine S-J, et al. Why
cultural safety rather than cultural competency is required to achieve health equity: a
literature review and recommended definition. International journal for equity in
health. 2019;18(1):1-17.

58.   Mackenzie M, Gannon M, Stanley N, Cosgrove K, Feder G. 'You certainly
don't go back to the doctor once you've been told,"I'll never understand women like
you."'Seeking candidacy and structural competency in the dynamics of domestic
abuse disclosure. Sociology of health & illness. 2019;41(6):1159-74.

59. Hawe P, Shiell A, Riley T. Theorising interventions as events in systems.
American journal of community psychology. 2009;43:267-76.

60. Government S. Inclusion Health Action in General Practice: Early Evaluation
Report. 2024.

61. Williamson AE, Ellis DA, Wilson P, McQueenie R, McConnachie A.
Understanding repeated non-attendance in health services: a pilot analysis of
administrative data and full study protocol for a national retrospective cohort. BMJ
Open. 2017;7(2):11.

62. Potter LC, Stone T, Swede J, Connell F, Cramer H, McGeown H, et al.
Improving access to general practice for and with people with severe and multiple
disadvantage: a qualitative study. Br J Gen Pract. 2024;74(742):e330-e8.

63. Metzl JM, Hansen H. Structural competency: theorizing a new medical
engagement with stigma and inequality. Social science & medicine. 2014;103:126-33.

64. Mistry SK, Harris E, Harris M. Community Health Workers as Healthcare
Navigators in Primary Care Chronic Disease Management: a Systematic Review.
Journal of General Internal Medicine. 2021;36(9):2755-71.

65. Babbel B, Mackenzie M, Hastings A, Watt G. How do general practitioners
understand health inequalities and do their professional roles offer scope for
mitigation? Constructions derived from the deep end of primary care. Critical Public
Health. 2019;29(2):168-80.

66. Tonnesen M, Hedeager Momsen A-M. Bridging gaps in health? A qualitative
study about bridge-building and social inequity in Danish healthcare. International
Journal of Qualitative Studies on Health and Well-being. 2023;18(1):2241235.

67. Craig P, Di Ruggiero E, Frolich KL, Mykhalovskiy E, White M, Campbell R, et
al. Taking account of context in population health intervention research: guidance for
producers, users and funders of research. Southampton: National Institute for Health
Research: Canadian Institutes of Health Research (CIHR)–National Institute for
Health Research (NIHR) Context Guidance Authors Group; 2018.

68. Skivington K, Matthews L, Simpson SA, Craig P, Baird J, Blazeby JM, et al. A
new framework for developing and evaluating complex interventions: update of
Medical Research Council guidance. bmj. 2021;374.

69. Mackenzie M, Skivington K, Fergie G. "The state They're in": Unpicking
fantasy paradigms of health improvement interventions as tools for addressing
health inequalities. Social Science & Medicine. 2020;256:113047.

70.   Mercer SW, Patterson J, Robson JP, Smith SM, Walton E, Watt G. The
inverse care law and the potential of primary care in deprived areas. The Lancet.
2021;397(10276):775-6.

71.   Bidmead E, Hayes L, Mazzoli-Smith L, Wildman J, Rankin J, Leggott E, et al.
Poverty proofing healthcare: A qualitative study of barriers to accessing healthcare
for low-income families with children in northern England. PLoS One.
2024;19(4):e0292983.

72.   Gkiouleka A, Wong G, Sowden S, Kuhn I, Moseley A, Manji S, et al. Reducing
health inequalities through general practice: a realist review and action framework.
Health and Social Care Delivery Research. 2024;12(7):1-104.

73.   Brown S. Qualitative evaluation of Focused Care. 2019.

74.   Revolving Doors A. Navigating complexity: learning from Navigators across
Birmingham. 2020.

75.   Corrigan PW, Pickett S, Schmidt A, Stellon E, Hantke E, Kraus D, et al. Peer
navigators to promote engagement of homeless African Americans with serious
mental illness in primary care. Psychiatry Research. 2017;255:101-3.

33