Ask your big brothers and sisters! A study protocol on multidisciplinary exploratory research into a preventive community youth empowerment program using biopsychosocial measures

Marleen Verhagen[1], Carin Rots[2,#a], Linda M.A. van Loon[2], Marjolein Stoopendaal[2], Lieke G.M. Raaijmakers[2], Anja C. Huizink[3], Sjoerd A.A. van den Berg[4], Jolanda J.P. Mathijssen[5]

[1]Department of Child and Youth Healthcare, Municipal Health Service West-Brabant, Breda, The Netherlands

[2]Department of Research, Municipal Health Service West-Brabant, Breda, The Netherlands

[3]Department of Clinical, Neuro- and Developmental Psychology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[4]Department of Clinical Chemistry, Erasmus MC, Rotterdam, The Netherlands

[5]Tranzo, Scientific Center for Care and Wellbeing, Tilburg School of Social and Behavioral Sciences, Tilburg University, Tilburg, The Netherlands

[#a] Current Address: Stichting Delen geeft meer, Arnhem, The Netherlands

[*]Corresponding author

E-mail: m.verhagen@ggdwestbrabant.nl

Abstract

Introduction:

Children growing up in adversity often experience setbacks and toxic chronic stress. Throughout the particularly vulnerable developmental phase of adolescence, chronic stress can lead to significant mental and physical repercussions. However, these adverse outcomes can also be prevented or alleviated.

There are indications that preventive programs with an integrated approach can contribute to reducing chronic stress. Big Brother, Big Sister (BBBS) is a preventive community Youth Empowerment Program implemented in low socio-economic status neighborhoods. Developed for and by young people, it empowers participants to reconnect with education and employment through professionally facilitated peer group activities. The program is embedded in local networks.

Main objectives:

To assess the applicability of the following biopsychosocial outcome measures—stress(ful life events), subjective psychological well-being, and social participation—within a diverse group of young people in vulnerable positions engaged in the BBBS program. Additionally, to provide insight into how these outcomes change over time.

Measures:

This exploratory quantitative study has an observational design with three measurements (at the onset of the program, after 6 months of participation and at outflow > 9 months). Approximately 80 young people aged 14 to 27 will participate in this study.

Various methods will be utilized to assess outcomes; questionnaires will be completed (i.e. the Perceived Stress Scale; Basic Psychological Need Satisfaction and Frustration Scale,

Stressful Life Events Scale, social participation and general background (hair) characteristics questionnaires), and hair samples will be collected to conduct cortisol and cortisone analyses, as proxy for prolonged physiological stress.

Discussion:

The multidisciplinary collaboration of partners within the research group combined with the biopsychosocial approach of this study is innovative and provides opportunities for cross-domain learning and collaboration to pursue (research) opportunities for evidence-based (early life) stress prevention and improvement of psychological well-being during the critical developmental phase of adolescence.

Keywords: youth, (early life) stress, hair cortisol/cortisone, well-being, participation

Introduction

Early Life Stress (ELS) and resilience in adolescence

Children and young people growing up in adversity often experience setbacks and toxic chronic stress [1] often referred to as Early Life Stress [2,3]. Toxic chronic stress occurs when the stress system is repeatedly reactivated without sufficient opportunity for recovery. Throughout the particularly vulnerable developmental phase of adolescence (10-24 years) [4], persistently elevated levels of stress hormones can significantly affect (future) mental and physical health, learning, and behaviour through gene-environment interactions and can result in neurobiological changes [1,2,3,5,6,7]. Chronic stress also modifies the stress response; individuals react more quickly and violently to a stressor later in life, and the stress response persists longer [1,7]. As the brain undergoes rapid growth and development in adolescence, the adverse outcomes resulting from chronic stress can also be prevented or alleviated during this period and transformed into more favorable development by enhancing well-being and fostering resilience [5,8,9]. This way, the plastic brain can be reprogrammed [10]. Since individuals who have experienced stress in early life often struggle to cope with stress induced by their own children [1,3], it is crucial for the (mental) health of the next generation that adverse outcomes resulting from toxic chronic stress in early life are prevented, alleviated, or transformed during adolescence. Moreover, epigenetic changes (alterations in gene expression) during their lifetime may also contribute to whether or not early life stress is transmitted across generations [11].

Hair cortisol, hair cortisone and chronic stress

As a human response to acute and chronic stress, the hypothalamic-pituitary-adrenal (HPA) axis is activated. When repeatedly activated under chronic stress, cortisol is released to an increased extent [12,13]. Nonetheless, personal characteristics such as resilience are associated with a reduced activation of the HPA axis [14] and "dysregulated" HPA axis functioning can result in either increased or decreased cortisol levels [15].

Measurements of both cortisol and cortisone (inactivated cortisol) in human hair are increasingly utilized as biomarkers to gain insight into chronic stress exposure. Hair steroid measurements provide retrospective insights because the long-term cumulative release of cortisol and cortisone can be measured up to several months before [16].

Hair cortisol concentration (HCC) has been associated with adult mental health in previous research; for example, increased cortisol levels have been found in adults experiencing ongoing chronic stress, while reduced hair cortisol levels have been found in adults with anxiety disorders such as PTSD [15,17]. Although this research in children and young people, especially for those in vulnerable positions, is still in its early stages, a recent review and meta-analysis by Li et al. [18] demonstrated that chronic stress was positively correlated with HCC. Hence, hair cortisol could serve as a biomarker for chronic stress among children and young people (aged 5-21 years). Recently, reference ranges for hair cortisol in children (aged 0-18 years) have been identified [19]. This may provide an opportunity to compare the results of this study with these reference ranges.

In addition to cortisol, the hormone cortisone might be a useful biomarker for stress research. Very few studies in young people include measurements of both hair cortisol and cortisone [20]. Results from the study by Ullmann et al. [21] showed that the hair cortisol

and cortisone concentrations were significantly correlated with mental and physical stress, as well as with stress perception in students (aged 18-35 years).

Research on the association between socio-economic status and hair cortisol and cortisone levels showed that a low neighborhood-level socio-economic status was significantly associated with higher hair cortisol and cortisone levels in children and adolescents (aged 4-18 years) [22].

This study will provide insight into experienced (early life) stress, subjective psychological well-being, social participation, and the hair cortisol and cortisone levels of a group of young people of mixed descent who primarily grew up in low SES neighborhoods, are facing difficulties in connecting with society, and participated in the Big Brother, Big Sister program.

Community Youth Empowerment Program "Big Brother, Big Sister" Breda

'Big Brother, Big Sister' (BBBS) is an integrated community Youth Empowerment Program (YEP) that empowers participants to reconnect with society and make progress in fostering positive social connections and developing (psychosocial) skills to pursue meaningful goals [23]. The program is peer-developed by and for young people. Current participants, known as Big Brothers and Sisters, are individuals aged between 14 and 27, encompassing both boys and girls of mixed descent. Most participants have attained a practical level of vocational education. Building a trusting relationship between pedagogical employees ("workers") and participants is a cornerstone of the program, providing young people with low-threshold access to a safe space where they can reconnect with education or employment and develop personally and socially with support from peers. Big Brothers and Sisters aspire to serve as role models and enhance the quality of life within the

neighborhoods where they are active and where they have primarily grown up themselves.

BBBS currently offers placements for approximately 150 young people per year across three

locations in vulnerable neighborhoods in Breda. Figure 1 provides further details about the

program.

The operational core of the BBBS program comprises:

Street Work: Big Brothers and Sisters are actively engaged in the community on a daily basis, making connections with residents. They visit hangouts, problem areas, school playgrounds, shopping centers, and assess the needs and opportunities in the neighborhood.

Activities: Big Brothers and Sisters organize various daily activities in collaboration with each other and local partners. These activities include sports, creative endeavors (such as utilizing their own music studio), workshops, outings with the elderly, and neighborhood gatherings. A significant focus of these activities is on little children. By planning and executing these activities, young people develop social, practical, and creative skills. Additionally, Big Brothers and Sisters provide mentoring to the younger children, fostering a sense of responsibility and serving as role models themselves.

One-on-One Conversations: Participants receive guidance in their personal development through individual discussions with experienced pedagogical staff. These conversations cover topics such as psychological well-being, personal circumstances, self-reflection, learning objectives, and future prospects.

Competency Levels: Personal development is encouraged through competency levels (referred to as Bronze, Silver, Silver+, Gold). Each level entails specific tasks and responsibilities. Participants who develop more skills and competencies are entrusted with greater tasks and responsibilities, such as mentoring other youths or participating in municipal networks. Big Brothers and Sisters with Silver+ and Gold competency levels support the pedagogical staff.

Local networks: BBBS is integrated in local networks and collaborates with a diverse array of partners, including educational institutions, companies, municipalities, volunteers, law enforcement and youth (health) care organizations. Over the years, BBBS has forged close working relationships with these partners and has established a distinctive presence within these networks

---

Figure 1. The operational core of the BBBS program

In- and outflow at BBBS program

Young people can enter the program through various channels, including direct contact with

Big Brothers and Sisters, referrals from community partners such as welfare organizations,

housing associations, and youth health care services, as well as through schools.

Additionally, young people can register themselves for the program. An interview is conducted with each young person to assess their eligibility for participation in which factors such as their willingness to develop, learn, and serve as a role model (as a 'Big Brother' or 'Big Sister') are considered essential for inclusion in the program. When participants achieve their goals, find employment or make progress in school, they typically exit the BBBS program. Participants remain engaged in BBBS for a period ranging from six months to two years.

Previous research and effective elements/ theoretical underpinning

The BBBS program has been implemented for 15 years. Throughout its development, BBBS has undergone initial internal and external evaluations, as documented by Loef et al. [24] and Parren [25]. These evaluations have been used to refine the program.

In 2019, the research department of the Regional Public Health Service West-Brabant conducted a qualitative study on Big Brothers' self-perceived effects of the program [23]. This study revealed that participants feel comfortable and respected within BBBS. Moreover, they reported making progress in fostering positive social connections and developing (psychosocial) skills to pursue meaningful goals. Examples of these achievements include exhibiting reduced aggression, improved cooperation, enhanced sense of responsibility, and a sense of doing something meaningful for others and for themselves.

The way BBBS is designed aligns well with Self-Determination Theory. Self-Determination Theory posits that psychological well-being is determined by three basic human needs: autonomy, positive connection with others, and competence to achieve meaningful goals [26]. There are indications that preventive programs with an integrated approach, meeting these basic human needs, can increase well-being and contribute to reducing chronic stress

[3,10]. The following effective elements appointed by Pijpers et al. [3] are integral to the preventive Youth Empowerment Program BBBS:

1. Eliminating harmful stressors (e.g., aggressive encounters in the street, exposure to criminal pressures); 2. Enhancing (psychosocial) skills, autonomy and competencies, thereby boosting personal resilience and enhancing future opportunities; 3. Providing a social environment characterized by positive social interactions and connections with the community. Within a secure pedagogical climate, young people can confront challenges, fostering confidence in themselves and others.

Main objectives

Overall, 'Big Brother, Big Sister' serves as a preventive community Youth Empowerment Program with the potential to mitigate early-life stress while enhancing psychological well-being and social engagement among young people. Therefore, BBBS emerges as a promising environment to assess the applicability of the following biopsychosocial outcome measures—stress(ful life events), subjective psychological well-being, and social participation—within a diverse target group of young people in vulnerable positions engaged in the BBBS program and to provide insight in how these outcomes change over time.

To our knowledge, there has been hardly any multidisciplinary research on preventive youth programs in low socioeconomic status (SES) neighborhoods, such as BBBS, using biopsychosocial outcome measures that include (early life) stress, psychological well-being and social participation of youth both at the beginning of a program and over time. Recent literature underscores the particular necessity for biopsychosocial research among young people in vulnerable positions and of mixed descent. Such research is crucial for elucidating stress mechanisms and for pursuing evidence-based strategies aimed at preventing early-life

stress and enhancing psychological well-being during the sensitive developmental period of adolescence [1,3,7,19,27,28].

Methods and materials

Design

The quantitative study is exploratory in nature, employing an observational design with three measurement points: at the onset of the program, after 6 months of participation, and at outflow. Data collection started in September 2021 and is scheduled to conclude in December 2024.

Research population - involvement and recruitment

All young people who enrolled in the program between September 2021 and December 2023 were invited to participate in the study. Through preparatory discussions at BBBS, it was anticipated that approximately 80 young individuals aged 14-27 would take part. Participation in the study requires written informed consent from all participants, with additional parental consent for those aged 14-16 years. Following the acquisition of informed consent, the researcher will extend invitations to the young participants for measurement sessions at their respective BBBS locations, situated across three neighborhoods in Breda. These sessions were conducted in private settings and scheduled at mutually convenient times. To express appreciation for their involvement, participants received compensation in the form of shopping vouchers. The compensation, determined in consultation with BBBS, amounted to €10 for the first two measurement sessions and €20 for the third.

Inclusion criteria

The study will include all young people meeting the following criteria:

- Enrolled in BBBS between the end of September 2021 and December 2023.

- Aged between ≥ 14 years and ≤ 27 years.

Exclusion criteria

The study will exclude all young people who have participated in BBBS < 1 years ago
(determined at baseline).

Procedures

Participants will be invited for measurement within four weeks of the start of the program,
after 6 months of participation, and at outflow or when they achieve their original starting
goals (sometimes they continue in the program in another position). The measurement
process takes approximately 20 minutes and involves completing several self-reported
questionnaires covering perceived stress, stressful life events, and psychological well-being.
To maintain focus, questionnaires regarding social participation and general background and
hair characteristics, are then verbally administered by the researcher. With permission, a
small strand of hair is cut using nail scissors (approximately 100 hairs) in a discreet manner
(from the scalp, at the back of the head's posterior vertex, from a deep layer of hair) to
ensure minimal visibility. Cortisol and cortisone levels are then measured in the hair samples
[16]. Table 1 provides further details about the timing of the various measurement
moments.

Ethical considerations and approval

Throughout this study, data will be kept confidential, and all data will be pseudonymised. This means that no names or personal data are linked to the answers and hair measurements. Participants can choose not to complete certain questions or to stop participating in the study at any time, without providing a reason and without facing any consequences. A pedagogical supervisor is always present on-site if the participant requires support after the measurement. This study was approved by the Medical Ethics Review Committee Brabant on 22-10-2021 and was found not to be subject to the Medical Research Involving Human Subjects Act (WMO) (METC nr: NW2021-91).

Measures

Perceived stress

The ten-item Perceived Stress Scale (PSS-adolescent) [29] will be utilized to measure the extent to which respondents experience stress due to unpredictability, lack of control, and overload (e.g., How often have you felt stressed and nervous? Rated on a five point Likert scale). A Dutch version for adolescents is available. Previous studies have demonstrated a Cronbach's alpha > .70 [30,31].

Subjective psychological well-being

The twenty-four-item Basic Psychological Need Satisfaction and Frustration Scale (BPNSFS-child) [32], utilizing a five point Likert scale, will be employed to subjectively measure the satisfaction and frustration of basic psychological needs in one's life (e.g., I often doubt whether I am good at things). This scale is grounded in the Self-Determination Theory [26] which states that "satisfaction of the basic psychological needs for autonomy, competence,

and relatedness both fosters immediate well-being and strengthens inner resources contributing to subsequent resilience, whereas need frustration evokes ill-being and increased vulnerabilities for defensiveness and psychopathology" [33]. The Dutch version of BPNSFS-child is comprehensible for a diverse range of young individuals, as confirmed by a pretest involving approximately fifteen Big Brothers and Sisters (28-06-2021). In a sample of 359 Belgian adolescents (mean age 17.87), Cronbach's alphas for autonomy, relatedness, and competence satisfaction were, respectively, 0.69, 0.77, and 0.81 [32].

Stressful Life Events

A questionnaire based on the Stressful Life Events Scale (SLES) [30,34] will be utilized, comprising twenty-six events ranked by the degree to which the event aggravates the lives of children (e.g., divorce of parents or being a victim of violence). Participants will indicate whether or not they have experienced each event during their lifetime.

Social participation

Social participation is operationalized as active participation in education, employment, volunteer work, membership in (sports) clubs, level of physical activity, gaming, and engagement in conflicts or fights (yes or no and frequency in the past 3 months). A questionnaire has been collaboratively developed with BBBS to measure social participation. It incorporates pertinent questions on social participation drawn from existing youth monitors conducted by the Municipal Health Service GGD West-Brabant [35,36]. Additionally, the questionnaire assesses competence levels within BBBS (see Figure 1).

Background characteristics

The background characteristics questionnaire includes items such as gender, age, type of education, grade, completed education, place of residence, neighborhood where one grew

up, ethnicity, country of birth (of father and mother), living situation, financial situation, and a number of questions about participation in BBBS, which will be administered verbally.

Background hair characteristics

The background questionnaire on hair characteristics, based on the 2019 version for children from Erasmus MC [37], will include questions about hair color, dyeing/blonding/permanent treatments, frequency of washing, degree of excessive sweating on the head, use of hair products, use of corticosteroids, use of contraception, drug use, alcohol use, and general registration of notable life events in the past 3 months ("Have you experienced any stressful life events in the past 3 months?"). Additionally, BMI will be inquired. This questionnaire will be administered verbally and is a required part of the application for hair cortisol/cortisone analysis [37]. Questions regarding these background characteristics will only be asked when the hair analysis is conducted, except for the general registration of stressful life events in the past 3 months, which will be asked of all participants regardless of their participation in the hair analysis.

Hair sample

Measurements of cortisol and cortisone in hair of the participants are objective and non-invasive [38]. The hair analysis will be conducted using an improved LC-MS/MS-based method [16]. Hair grows at an average rate of 1 cm per month. Therefore, the 3 cm hair sample closest to the scalp measures the cumulative cortisol (chronic stress level) over the past three months prior to each measurement [39, 40].

An overview of the measuring instruments per measurement moment can be found in Table 1. At T0, all instruments will be administered. At T1 and T2, the registration of general background characteristics and Stressful Life Events will not be measured again.

Table 1. Measuring instruments per measuring moment

Measuring instruments	At influx T0	6 months after influx T1	At outflow T2
Perceived Stress Scale (PSS-adolescent)	x	x	x
Basic Psychological Need Satisfaction and Frustration Scale (BPNSFS-child)	x	x	x
Stressful Life Events Scale (SLES)	x		
Social participation questionnaire	x	x	x
Background characteristics questionnaire	x		
Background hair characteristics questionnaire version for children 2019 by Erasmus MC [37] including registration of life events (< 3 months ago) and BMI	x	x	x
Hair sample (cortisol/cortisone)	x	x	x

Data analysis

First, background characteristics, experienced (early life) stress, subjective psychological well-being, social participation, and the hair cortisol and cortisone levels of the respondents at T0 will be described and, if possible, compared to average Dutch young people. Chi-squared analyses, independent sample t-tests, and ANOVA will be used accordingly to assess differences in outcome variables for background variables (such as ethnicity, age, gender) and specific hair characteristics. Multivariable regression analyses will be performed to assess associations between background characteristics and outcome measures at T0. Linear mixed models are then used to measure changes over time in experienced (early life) stress, subjective psychological well-being, social participation, and the hair cortisol and cortisone levels of the respondents individually and relative to each other and to determine whether these changes are related to background characteristics.

Data management

The personal data of the participant will be encrypted and stored under a randomly assigned

respondent number. Completed questionnaires will then be pseudonymized. Only the

principal investigator will have access to the translation key to decrypt the questionnaire.

The questionnaires collected as part of the study will be digitized into the Enalyzer program

[41] and stored separately in sealed envelopes in a locked room. Hair samples will be kept in

sealed envelopes marked with a respondent number in a locked room and brought to the

laboratory by the researcher in batches.

The data will be used for scientific purposes only and will be destroyed 10 years after the

completion of the study (2034). Data management agreements have been made with

collaborating partners and the laboratory, which are documented in an overall consortium

collaboration agreement and a specific collaboration agreement with the laboratory

concerning the analyses of cortisol and cortisone. These collaboration agreements have

been reviewed by the privacy officer of the Municipal Health Service GGD West-Brabant.

Discussion

'Big Brother, Big Sister' is a preventive community Youth Empowerment Program that shows

promise in empowering young people of mixed descent who have primarily grown up in low

socio-economic status (SES) neighborhoods and are encountering challenges in integrating

with society. This group is typically underrepresented in research, particularly in studies

concerning adolescent mental health and the measurement of hair cortisol and cortisone

[19,42]. The program has proven its value in practice, with 15 years of practical experience

and accompanying research. So now is the time for the next step in scientific substantiation.

Conducting multidisciplinary research within the BBBS setting presents a valuable opportunity to deepen our understanding of stress mechanisms and the utilization of biopsychosocial outcome measures among a diverse group of young individuals in vulnerable circumstances. The chosen approach for this quantitative study is an observational design with an exploratory character, incorporating biopsychosocial measures. This research methodology aims to provide insights into the significance and viability of impact studies for future research. Such a phased approach is recommended in investigations into the outcomes of integrated, practice-driven interventions [43, 44].

Implementing research within this relatively small-scale practical program presents several limitations. Firstly, the examination will involve relatively small numbers of participants. Secondly, the assessment of Big Brothers and Sisters occurs within a multifactorial context; they participate in a preventive community youth empowerment program situated in the dynamic reality of adolescents, over which the researcher has no control. This reality includes factors such as diploma attainment, exam stress, financial independence, independent living, and social relationship maintenance. Additionally, the participants comprise a group of young people who struggle to connect with education, employment, or home, and they are not always readily accessible, which could pose challenges for conducting three follow-up measurements. Finally, the outcomes of the questionnaires do not fully parallel with the outcomes of the hair samples temporally; cortisol and cortisone levels reflect an average over the 3 months preceding the measurements, while most questionnaires assess the current status or events from the past 4 weeks. This temporal misalignment should be considered in the analyses.

Strengths of this research include the close collaboration with both professional workers and a diverse group of young individuals in vulnerable circumstances who possess valuable expertise derived from practical experiences. This practical knowledge is complemented by insights from a multidisciplinary scientific network of partners. The collaborative effort within the research group, coupled with the biopsychosocial approach, represents an innovative approach. It opens avenues for cross-domain learning and collaboration, enabling the pursuit of evidence-based strategies for preventing early-life stress and improving psychological well-being, particularly during the critical developmental phase of adolescence.

Acknowledgements: The authors would like to acknowledge the professional workers of 'Grote Broer, Grote Zus Breda' for their contribution to the study.

Supporting Information: not applicable

Authors declare no competing interests.

This project has received funding from "Stichting delen geeft meer" under project number 2021/11/01.

The data of this study are not publicly available due to privacy and ethical restrictions.

References

1. Shonkoff JP, Garner AS, Siegel BS, Dobbins MI, Earls MF, Garner AS, et al. The Lifelong Effects of Early Childhood Adversity and Toxic Stress. Pediatrics. 2012 January; 129: e232–e246.

2. Smith KE, Pollak SD. Early life stress and development: potential mechanisms for adverse outcomes. Journal of Neurodevelopmental Disorders. 2020 December; 12.

3. Pijpers F, Vanneste Y, Feron F. Stress bij kinderen: hoe houden we het gezond? Stress bezien vanuit de Jeugdgezondheidszorg. 2019. Nederlands Centrum Jeugdgezondheid (NCJ).

4. Sawyer SM, Azzopardi PS, Wickremarathne D, Patton GC. The age of adolescence. The Lancet Child & Adolescent Health. 2018 March; 2: 223–228.

5. Soleimanpour S, Geierstanger S, Brindis CD. Adverse Childhood Experiences and Resilience: Addressing the Unique Needs of Adolescents. Academic Pediatrics. 2017 September; 17: S108–S114.

6. Takeuchi H, Kawashima R. Neural Mechanisms and Children's Intellectual Development: Multiple Impacts of Environmental Factors. The Neuroscientist. 2016 July; 22: 618–631.

7. De Bellis MD, Zisk A. The Biological Effects of Childhood Trauma. Child and Adolescent Psychiatric Clinics of North America. 2014 April; 23: 185–222. doi: 10.1016/j.chc.2014.01.002

8. Arslan G. Psychological maltreatment, emotional and behavioral problems in adolescents: The mediating role of resilience and self-esteem. Child Abuse & Neglect. 2016 February; 52: 200–209.

9. Bethell CD, Newacheck P, Hawes E, Halfon N. Adverse Childhood Experiences: Assessing The Impact On Health And School Engagement And The Mitigating Role Of Resilience. Health Affairs. 2014 December; 33: 2106–2115.

10. McEwen BS, Gianaros PJ. Stress- and Allostasis-Induced Brain Plasticity. Annual Review of Medicine. 2011 February; 62: 431–445.

11. Roth TL, Lubin FD, Funk AJ, Sweatt JD. Lasting Epigenetic Influence of Early-Life Adversity on the BDNF Gene. Biological Psychiatry. 2009 May; 65: 760–769.

12. O'Connor DB, Thayer JF, Vedhara K. Stress and Health: A Review of Psychobiological Processes. Annual Review of Psychology. 2021 January; 72: 663–688.

13. Miller GE, Chen E, Zhou ES. If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. Psychological Bulletin. 2007; 133: 25–45.

14. Sun X, Dai X, Yang T, Song H, Yang J, Bai J, et al. Effects of mental resilience on neuroendocrine hormones level changes induced by sleep deprivation in servicemen. Endocrine. 2014 March; 47: 884–888.

15. Stalder T, Steudte-Schmiedgen S, Alexander N, Klucken T, Vater A, Wichmann S, et al. Stress-related and basic determinants of hair cortisol in humans: A meta-analysis. Psychoneuroendocrinology. 2017 March; 77: 261–274.

16. Mirzaian M, van Zundert SKM, Schilleman WF, Mohseni M, Kuckuck S, van Rossum EFC, van Schaik RHN, van den Berg SAA. Determination of cortisone and cortisol in human scalp hair using an improved LC-MS/MS-based method. Clinical Chemistry and Laboratory Medicine 2023 July;25;62(1):118-127.

17. Staufenbiel SM, Penninx BWJH, Spijker AT, Elzinga BM, van Rossum EFC. Hair cortisol, stress exposure, and mental health in humans: A systematic review. Psychoneuroendocrinology. 2013 August; 38: 1220–1235.

18. Li Y, Jia W, Yan N, Hua Y, Han T, Yang J, et al. Associations between chronic stress and hair cortisol in children: A systematic review and meta-analysis. Journal of Affective Disorders. 2023 May; 329: 438–447.

19. de Kruijff I, Noppe G, Kieviet N, Choenni V, Lambregtse-van den Berg MP, Begijn DGA, et al. LC-MS/MS-based reference intervals for hair cortisol in healthy children. Psychoneuroendocrinology. 2020 February; 112: 104539.

20. Chen Y, Petzold M, Rüetschi U, Dahlstrand J, Löfstedt P, Corell M, et al. Hair glucocorticoid concentration, self-perceived stress and their associations with cardiometabolic risk markers in Swedish adolescents. Psychoneuroendocrinology. 2022 December; 146: 105908.

21. Ullmann E, Barthel A, Petrowski K, Stalder T, Kirschbaum C, Bornstein SR. Pilot study of adrenal steroid hormones in hair as an indicator of chronic mental and physical stress. Scientific Reports. 2016 May; 6.

22. Vliegenthart J, Noppe G, van Rossum EFC, Koper JW, Raat H, van den Akker ELT. Socioeconomic status in children is associated with hair cortisol levels as a biological measure of chronic stress. Psychoneuroendocrinology. 2016 March; 65: 9–14.

23. Verhagen M, van Loon L, Rots-de Vries C. Vraag het je Grote Broers: Een verkennend kwalitatief onderzoek naar de ervaren effecten van een wijkgericht youth empowerment programma voor en door jongeren. Tijdschrift voor gezondheidswetenschappen. 2023 November; 101: 149-156.

24. Loef L, Fransen N, Dijk B. De evaluatie van KICK, Grote Broer en het Buurtinterventieteam. 2009. Amsterdam: DSP-groep. Available from: https://www.dsp-groep.eu/wp-content/uploads/15bdkick_xxx_evaluatie_KICK_Grote_Broer_Buurtinterventieteams.pdf

25. Parren J. Het Grote Broerproject Toekomstperspectief voor jongeren. M.Sc. Thesis, Avans Hogeschool Breda. 2012 May.

26. Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. American Psychologist. 2000; 55: 68–78.

27. van Dammen L, de Rooij SR, Behnsen PM, Huizink AC. Sex-specific associations between person and environment-related childhood adverse events and levels of cortisol and DHEA in adolescence. PLOS ONE. 2020 June; 15: e0233718.

28. Gray NA, Dhana A, Van Der Vyver L, Van Wyk J, Khumalo NP, Stein DJ. Determinants of hair cortisol concentration in children: A systematic review. Psychoneuroendocrinology. 2018 January; 87: 204–214.

29. Cohen, S. Perceived Stress Scale. 1994. Available from: https://www.mindgarden.com/documents/PerceivedStressScale.pdf [accessed 13-10-2020]

30. Van der Ploeg, J. Stress bij kinderen. 1st ed. Houten: Bohn Stafleu van Loghum; 2013.

31. Lee EH. Review of the Psychometric Evidence of the Perceived Stress Scale. Asian Nursing Research. 2012 December; 6: 121–127.

32. Chen B, Vansteenkiste M, Beyers W, Boone L, Deci EL, Van der Kaap-Deeder J, et al. Basic psychological need satisfaction, need frustration, and need strength across four cultures. Motivation and Emotion. 2014 November; 39: 216–236.

33. Vansteenkiste M, Ryan RM. On psychological growth and vulnerability: Basic psychological need satisfaction and need frustration as a unifying principle. Journal of Psychotherapy Integration. 2013 September; 23: 263–280.

34. Sandler IN, Block M. Life stress and maladaptation of children. American Journal of Community Psychology. 1979 August; 7: 425–440.

35. GGD West-Brabant. Corona Gezondheidsmonitor Jeugd 2021. Breda: GGD West-Brabant; 2022. Available from https://brabantscan.nl/

36. GGD West-Brabant. Gezondheidsmonitor Jeugd 2019. Breda: GGD West-Brabant; 2019. Available from https://brabantscan.nl/

37. Erasmus MC. Questionnaire belonging to application hair cortisol, version for children 2019.

38. Wosu AC, Valdimarsdóttir U, Shields AE, Williams DR, Williams MA. Correlates of cortisol in human hair: implications for epidemiologic studies on health effects of chronic stress. Annals of Epidemiology. 2013 December; 23: 797–811.e2.

39. Dettenborn L, Muhtz C, Skoluda N, Stalder T, Steudte S, Hinkelmann K, et al. Introducing a novel method to assess cumulative steroid concentrations: Increased hair cortisol concentrations over 6 months in medicated patients with depression. Stress. 2011 November; 15: 348–353.

40. Wennig R. Potential problems with the interpretation of hair analysis results. Forensic Science International. 2000 January; 107: 5–12.

41. Enalyzer. Enayzer Survey tool [internet]. Available from:

    https://www.enalyzer.com/nl/?s=survey#sidemenu=null

42. Lawrence RS, Gootman JA, Sim LJ. Adolescent Health Services, Missing Opportunities.

    Washington, DC: National Research Counsil and Institute of Medicine, The National

    Academies Press; 2009.

43. May A, Mathijssen J. Alternatieven voor RCT bij de evaluatie van effectiviteit van

    interventies!? Eindrapportage. 2015. Available from:

    https://www.researchgate.net/publication/299538473_Alternatieven_voor_RCT_bij_

    de_evaluatie_van_effectiviteit_van_interventies_Eindrapportage

44. Rots-de Vries MC. Rich evidence for poor families: Exploring the potential of practice-

    driven intervention research in Preventive Child Healthcare. Ph.D. dissertation,

    Tilburg University. 2010. Available from:

    https://research.tilburguniversity.edu/en/publications/rich-evidence-for-poor-

    families-exploring-the-potential-of-practi