TITLE: Circulating Cell-Free DNA Methylation Profiles Enable Disease-Specific

Detection of Alzheimer's, Parkinson's, and ALS from Blood

Chad Pollard[1,2], Erin Saito[2], Ryan Miller[2,3], Isaac Sirland[1,2], Mykle Keni[1,2], Andrew

Jenkins[1,2], Jonathon T. Hill[1,2], and Tim Jenkins[1,2]

[1] Department of Cell Biology & Physiology, Brigham Young University, Provo, UT 84602

[2] Resonant, LLC

[3] Department of Biology, Brigham Young University, Provo, UT 84602

Correspondence

Jonathon Hill, Department of Cell Biology & Physiology, 4005 LSB, Brigham Young

University, Provo, UT 84602, USA

E-mail: jhill@byu.edu

ABSTRACT

Noninvasive biomarkers for neurodegenerative diseases are urgently needed for earlier

detection, monitoring, and intervention. To address this, we developed a blood-based

cfDNA methylation platform. A whole-genome nanopore methylation atlas of six primary

human neural cell types—cortical, dopaminergic, and spinal motor neurons, astrocytes,

Schwann cells, microglia—was used to train classifiers that assign cfDNA to its

neuronal or glial origin. Classifiers were validated *in silico* and applied to 219 plasma

samples from patients with Alzheimer's disease (AD), Parkinson's disease (PD),

amyotrophic lateral sclerosis (ALS), and controls. Cortical cfDNA was elevated in AD,

dopaminergic cfDNA in PD, and spinal motor neuron cfDNA in ALS, with predictive

modeling achieving AUCs >0.98. In mild cognitive impairment, cortical cfDNA elevations

identified individuals who later progressed to AD, supporting predictive utility. A

multivariate model integrating multiple neurons improved accuracy, suggesting cfDNA

methylation profiling as a scalable framework for noninvasive neurodegeneration

detection and monitoring.

KEYWORDS: cell-free DNA, cfDNA methylation, neurodegeneration, Alzheimer's

disease, Parkinson's disease, amyotrophic lateral sclerosis, liquid biopsy, cortical

neurons, dopaminergic neurons, spinal motor neurons, epigenetics, ONT sequencing,

biomarker, cell-of-origin, disease diagnosis, machine learning

1. INTRODUCTION

Neurodegenerative diseases constitute one of the fastest-growing public health challenges worldwide. Alzheimer's disease (AD) alone currently affects ~7 million individuals over age 65 in the U.S., with prevalence projected to double by 2060[1]. Parkinson's disease (PD) impacts millions globally[2,3], and approximately 30,000 Americans live with amyotrophic lateral sclerosis (ALS)[4,5]. These diseases cause progressive and irreversible loss of neurons, leading to cognitive impairment, motor dysfunction, and premature mortality.

Despite decades of research, current diagnostics remain fundamentally reactive, applied in advanced disease stages after substantial neuron loss[6-8], when therapeutic interventions are limited and prognosis is poor[9]. Clinical symptoms, cerebrospinal fluid (CSF) analysis, and neuroimaging (MRI, PET) can confirm advanced disease but are invasive, expensive, and inaccessible for early screening or routine, longitudinal use[10-12].

Blood-based protein biomarkers such as NfL, GFAP, amyloid-β, tau, and α-synuclein are improving accessibility but remain insufficient for early screening or disease monitoring. General markers of neurodegeneration signal neuronal injury without resolving which cells or brain regions are affected[13,14]. Diagnostic protein biomarkers (e.g., amyloid in AD, α-synuclein in PD) are narrow in scope, limited in their ability to generalize across conditions, and often fail to correlate with disease progression[13,15]. Recent updates to AD diagnostic criteria underscore this limitation. While amyloid and tau biomarkers can confirm AD neuropathology, they are not

recommended for presymptomatic use due to limited sensitivity and the current lack of

effective presymptomatic disease-modifying therapies[15].

The result is a diagnostic catch-22: biomarkers confirm disease, but only in

advanced disease stages. Development of early detection tools has been sidelined

because they cannot yet demonstrate meaningful clinical impact. As a result, clinicians

still lack scalable tools to detect and track active neurodegeneration in real time, when

intervention would matter most.

Selective neuronal cell death is a fundamental hallmark of neurodegenerative

diseases[16], yet current protein biomarkers provide only indirect or nonspecific readouts

of this pathology. Circulating cell-free DNA (cfDNA) offers a fundamentally different

class of biomarkers that may address these challenges. Fragments of DNA released

into blood during neuronal apoptosis carry stable[17,18], "cell-of-origin" methylation

signatures of the dying cell. Thus cfDNA presents a biologically anchored, real-time

measure of active cell loss due to cfDNA's short circulating half-life[19]. Importantly, this

approach aims to monitor a defining feature of neurodegenerative disease—the loss of

neurons—providing unique insights that complement methods aimed at more indirect

indicators such as plaque formation or genetic risk factors.

Nucleic acid-based blood liquid biopsies have been suggested as an attractive

alternative due to their minimal invasiveness, scalability, and ability to capture cell and

disease-specific molecular features[20-26]. Applications in oncology are currently being

explored[27-30]. We have hypothesized that cfDNA methylation signatures, in particular,

could bridge the gap between broad but nonspecific staging markers (e.g., NfL, GFAP)

and narrow, disease-specific proteins (e.g., tau, α-synuclein), offering the specificity,

sensitivity, and scalability needed for earlier detection and intervention[13].

Impactful studies have demonstrated that methylation signatures can be

leveraged to identify cellular origin across tissues[31]. Whole genome methylation profiles

can be used to computationally deconvolute cfDNA fragments to their source tissues

and have been applied to prenatal testing, oncology, and organ transplants[32].

Subsequent studies have extended this concept with a more comprehensive human

cell-type methylation atlas, revealing that circulating cfDNA encodes distinct cell-of-

origin signatures in both healthy and diseased states[33]. For neurodegenerative disease,

specifically, we, and other groups, have shown that elevated circulating levels of cfDNA

from specific neurons and glia correlate with diseases like Alzheimer's disease[34],

traumatic brain injury, and MS[35].

However, these prior cfDNA methylation studies have relied on methods that

include bisulfite conversion, PCR amplification, and short-read sequencing. While

impactful in advancing the field, the approach introduces major limitations. Bisulfite- and

PCR-based methods degrade DNA[36] and introduce amplification bias[37], limiting

resolution and translational reliability.

In our pilot study, we detected elevated cortical neuron-derived cfDNA in the

blood of AD and MCI patients compared to controls, using a targeted bisulfite/PCR

assay at a single CpG site[34]. While this provided early proof of concept, it also

underscored the limitations of locus-restricted, bisulfite- and PCR-based methods. The

present study builds directly on that work, overcoming these constraints with native

methylation profiling via nanopore sequencing. To enhance cortical neuron specificity,

we expanded the methylation reference atlas to include additional neuronal and glial

populations, which also revealed potential applications across other neurodegenerative

diseases. Here, we establish proof of concept that cell type-resolved cfDNA

deconvolution provides a scalable and clinically feasible framework for detecting and

tracking neurodegeneration and differentiating neurodegenerative diseases.

2. METHODS

2.1 Sample Acquisition

2.1.1 Purified Neural Cell Types

Purified human neurons and glial cells were obtained from commercial vendors (Accegen and Creative BioArray; Supplemental Table 1). Cell types included both primary and induced pluripotent stem cell (iPSC)-derived cortical and dopaminergic neurons, as well as primary astrocytes, Schwann cells, and microglia.

Postmortem human spinal cord tissue was procured through BioIVT under institutional protocols for deidentified human biospecimen research (BioIVT ASTERAND® Normal Human Tissue). Upon receipt, tissue was rapidly processed to preserve DNA integrity. Spinal motor neurons were isolated using a Percoll gradient method adapted from previous methods[38], and optimized for mature human spinal cord tissue. Briefly, tissue was mechanically dissociated and layered onto dual isosmotic Percoll gradients (35% and 12.5%). Samples were centrifuged at 800 xg for 30 minutes, after which the 12.5% layer was collected for downstream processing. This fraction was enriched for spinal motor neurons and used for subsequent DNA extraction and methylation profiling. Optimization steps included adjusting gradient concentrations, centrifugation parameters, and handling conditions to account for the increased density and fragility of mature human neurons.

Genomic DNA extracted from these purified populations were used to generate whole genome nanopore methylation reference profiles for classifier training, evaluate cell type specificity, and validate the performance of methylation-based models for cfDNA deconvolution.

2.1.1 Human Patient Blood Samples

A total of 219 human plasma samples (1 mL each) were obtained from BioIVT

(Westbury, NY, USA), representing six clinically confirmed diagnostic categories:

Alzheimer's disease (AD; n = 45), mild cognitive impairment (MCI; n = 5), Parkinson's

disease (PD; n = 64), sporadic amyotrophic lateral sclerosis (SALS; n = 65), familial

amyotrophic lateral sclerosis (FALS, n = 10), and cognitively normal, age-matched

controls (HC, n = 30).

All samples were accompanied by case report forms (CRFs), informed consent

documentation, and detailed clinical metadata. Demographic variables included age,

sex, and ethnicity. Available clinical measures included Mini-Mental State Examination

(MMSE) scores at multiple timepoints and cerebrospinal fluid (CSF) concentrations of

disease-associated protein biomarkers. These included $A\beta1$-42, $A\beta1$-40, $A\beta1$-38, total

tau, α-synuclein, and PARK7, quantified using validated commercial assays (Meso

Scale Discovery [MSD], INNOTEST, and Covance platforms).

All BioIVT specimens were de-identified prior to receipt and used in accordance

with institutional review protocols and ethical guidelines governing secondary use of

human biological materials.

2.2 Sample Processing

All biospecimens were processed by Wasatch BioLabs (WBL), a high-throughput

sequencing provider specializing in clinical and research applications of Oxford

Nanopore Technologies sequencing. Sample processing followed standardized internal

quality control protocols and was managed through a laboratory information

management system (LIMS) to ensure traceability and data integrity.

2.2.1 Genomic DNA

Genomic DNA (gDNA) was extracted from purified cell types using the DNeasy

Blood & Tissue Kit (Cat # 69504) and libraries were prepared according to Wasatch

Biolabs' (WBL's) Direct Whole Methylome Sequencing (dWMS) protocol, a modified

whole genome ONT sequencing service optimized for methylation analysis. Extracted

DNA was fluorometrically quantified using the Qubit dsDNA High Sensitivity Assay

(Thermo Fisher Scientific) and assessed for purity and integrity using the NanoDrop

spectrophotometer (Thermo Fisher Scientific) and Agilent TapeStation 4200 (Agilent

Technologies), respectively.

Following gDNA dWMS sequencing, cell types included in model training

achieved an average of 15x global coverage, ensuring robust methylation signal

detection and accurate cell-type classification.

2.2.2 Cell-Free DNA

cfDNA was extracted from human plasma samples using the MAGicBead™

cfDNA Isolation Kit (Zymo, Cat # D4086). cfDNA library preparation was performed

according to WBL's Direct Methylation Sequencing (dMS) for cfDNA service, an ONT-

based cfDNA sequencing service for unbiased, native cfDNA sequencing. cfDNA

quantity was measured using Qubit fluorometric quantitation (Thermo Fisher Scientific),

and sample purity and integrity were assessed using NanoDrop spectrophotometry and

the Agilent TapeStation 4200, respectively, as described above for gDNA.

2.3 Data Generation

DNA methylation array data for purified cortical neurons, dopaminergic neurons,

and iPSC-derived spinal motor neurons were generated at the University of Utah using

the Illumina Infinium MethylationEPIC v2 array platform in alignment with manufacturer

protocols.

For nanopore sequencing, gDNA and cfDNA libraries were sequenced on the

ONT PromethION platform. Raw signal data (POD5 files) were basecalled and aligned

to the GRCh38 human reference genome using Dorado

(https://github.com/nanoporetech/dorado), an open-source ONT basecaller. Methylation

calls were extracted using ONT's modkit software

(https://github.com/nanoporetech/modkit), to produce per-read methylation summary

files.

All methylation data from gDNA and cfDNA sequencing were converted to CH3,

a proprietary file format developed by Wasatch BioLabs for downstream methylation

analysis, as detailed below.

2.4 Differential Methylation Analysis

All array samples underwent standard preprocessing and normalization. Raw

IDAT files were processed using the minfi R package. Beta values representing the

fraction of methylation at each CpG site were obtained following SWAN (Subset-

quantile Within Array Normalization). To assess data quality, beta value density plots

were examined to confirm the expected bimodal distribution, characterized by peaks in

the unmethylated (0.0–0.2) and methylated (0.8–1.0) ranges, and a flat valley in the

intermediate methylation range. Samples failing to meet these distribution criteria were

excluded from further analysis, and the remaining samples were re-normalized using

the same procedures.

For all nanopore sequencing data, CH3 methylation files were created for each

sample and processed with MethylSeqR, WBL's open-source R package that that

converts CH3 files into a queryable database, enables region/window summarization,

differential methylation testing, and built-in QC plots (https://github.com/Wasatch-

Biolabs-Bfx/MethylSeqR, v0.8.0). Methylation databases were indexed by individual

CpG positions with the make_pos_db() function. Using the summarize_windows()

function, these data were summarized into 200bp windows with a minimum of 5 CpGs.

Differential methylation analyses were performed between each cell type and pooled

human blood plasma using the calc_mod_diff() function. Differentially methylated

regions (DMRs) were identified by collapsing windows using the collapse_windows()

function, retaining regions with a minimum delta of at least 0.8 between cell type and

blood plasma.

DMRs between each cell type and blood plasma were further filtered to remove

overlapping DMRs between cell types. Resulting cell-specific DMRs were used in

downstream classification to evaluate cell-type-specific cfDNA contributions in plasma

from patients with neurodegenerative diseases.

2.5 Theoretical Model Validation

To evaluate the analytical performance of the cell type-specific classifiers, we

generated *in silico* mixtures of cell type-specific cfDNA in a background of plasma-

derived cfDNA.

Genomic DNA (gDNA) from each purified cell type was computationally

fragmented to more closely mimic the size distribution of circulating cfDNA (see

Supplemental Figure 1)[39]. This step ensured that cell-derived gDNA reads would be

more comparable in size and structure to those found in patient-derived cfDNA

samples.

Using these computationally fragmented reads, a series of *in silico* dilution

datasets was constructed. Each dataset contained approximately 1,000,000 total reads

and consisted of varying proportions of cell-specific reads spiked into a background of

healthy donor plasma cfDNA. The dilution series included nine fractional compositions:

0%, 0.1%, 1%, 10%, 15%, 25%, 50%, 75%, and 100% cell-type-specific reads, with the

remaining balance composed of plasma reads. This range was designed to reflect

physiologically plausible concentrations of cell-free DNA originating from neuronal and

glial sources. For each dilution point, cell-type-specific DMR classifiers were applied to

the synthetic mixtures. Classifier scores were calculated for individual reads based on

methylation concordance with the reference DMR set corresponding to the target cell

type. Predicted cell fractions were derived by aggregating classified reads and were

then compared to the known spiked-in proportions.

Classifier performance was assessed by calculating the coefficient of

determination (R^2) between predicted and actual cell fractions. Linearity and sensitivity

were evaluated across the dilution range, and classifier specificity was tested by

applying each model to unrelated cell types. Additional comparisons were performed

using DMRs derived from iPSC-based models to assess the impact of training source

on classification fidelity.

This theoretical validation enabled benchmarking of model accuracy and

demonstrated the potential of the classifier framework to detect low-frequency cfDNA

signals in a complex plasma background.

2.6 Clinical Analyses

As an initial exploratory step, the distribution of cell-type specific cfDNA

classifications was visualized across diagnostic categories. For each cell type of interest

(e.g., cortical, dopaminergic, or spinal motor neurons), boxplots displaying the

percentage of reads classified to that cell type across different disease states were

generated. These plots were overlaid with jittered individual points to highlight within-

group variability and potential diagnostic separation. This visualization provided early

evidence of cell-type specific cfDNA differences between disease and control groups,

motivating subsequent classification modeling.

To quantitatively assess the diagnostic value of these cell-type specific cfDNA

signals, a logistic regression classification framework was developed using a stratified

train-test split. Only samples from the disease group(s) of interest and comparator

group(s) were included. Binary labels were assigned accordingly. The dataset was then

partitioned into a training set (70%) and a testing set (30%) using stratified sampling to

maintain class balance.

The training set was used to fit a logistic regression model with one or more

specified predictors (e.g., % spinal-classified reads) corresponding to different neuronal

or glial populations. Importantly, the training set was also used to identify the optimal

decision threshold for classification, selected based on Youden's J statistic derived from

the receiver operator characteristic (ROC) curve. This threshold was fixed and carried

forward to the test set without modification, and is reported here as "optimal threshold".

All performance metrics, including sensitivity, specificity, accuracy, and area

under the ROC curve (AUC) with 95% confidence intervals were evaluated exclusively

on the test dataset to avoid overfitting and ensure generalizability. ROC curves with

confidence intervals and confusion matrix heatmaps were generated to visualize model

performance. All statistical analysis and figure generation were performed in R using the

caret, pROC, dplyr, tibble, and ggplot2 packages.

2.7 Classifier Performance and Statistical Analysis

All analyses were performed using R (v4.3.0) and Python (v3.10), supported by

in-house developed scripts. Statistical analyses were performed to evaluate the

quantitative accuracy and diagnostic performance of cfDNA methylation classifiers

across both computational and clinical datasets. For analytical validation using *in silico*

dilution series, predicted cfDNA fractions were compared to known input values, and

model performance was quantified using linear regression and R^2. These analyses

assessed the linearity and concordance of predicted cell fractions across a wide

dynamic range of cell-type-specific input levels.

To assess differences in cell classification signals between disease groups, we

employed the Wilcoxon rank-sum test (Mann–Whitney U test). This non-parametric

statistical test was selected due to its robustness against violations of normality and its

suitability for comparing distributions of continuous variables between two independent

groups. Specifically, we compared cell-type-specific classification scores (e.g., cortical

or dopaminergic cfDNA percentages) between disease states (e.g., Parkinson's disease

vs. age-matched controls, or ALS vs. Alzheimer's disease). All p-values obtained from

pairwise Wilcoxon tests were subsequently adjusted for multiple comparisons using the

Benjamini-Hochberg procedure to control the false discovery rate (FDR). Adjusted p-

values < 0.05 were considered statistically significant. These analyses were conducted

in R (v4.5.1) using the wilcox.test() function from base R and p.adjust() for FDR

correction.

The same statistical framework was applied to ancillary analyses of protein-

based biomarkers, including amyloid-β isoforms, total tau, and Mini-Mental State

Examination (MMSE) scores. These data were plotted as boxplots based on raw

concentration values or scores provided by the sample supplier (BioIVT). No

normalization, transformation, or additional statistical testing was performed, as we did

not have access to the proprietary diagnostic pipelines used by clinical providers.

Accordingly, these results are presented solely for qualitative context and visual

comparison with cfDNA-derived signals (Supplemental Figure 11).

For classification tasks in the clinical cohorts, diagnostic performance was

assessed through ROC curve analysis. Classifier scores for each sample were

generated based on read-level concordance with cell type-specific DMRs, and binary

classifications were evaluated for their ability to distinguish between disease and control

groups, as well as between other disease groups. The following performance metrics

were calculated: AUC, 95% confidence intervals for AUC (lower and upper bounds),

sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV),

overall accuracy, balanced accuracy, and the optimal classification threshold, which

was determined using Youden's J statistic. These metrics were computed for each

classifier individually, including cortical neuron-, dopaminergic-, and spinal motor

neuron-derived cfDNA, as well as for a combined model incorporating all cell types.

3. RESULTS

3.1 Whole-genome nanopore profiling identifies cell type-specific methylation signatures

To establish our methylation reference atlas for cfDNA classification, we

performed whole-genome methylation sequencing on genomic DNA from pooled human

blood plasma and eight primary neuron and glial cell types using Wasatch BioLabs'

(WBL) Direct Whole Methylome Sequencing (dWMS) platform. This Oxford Nanopore

Technologies (ONT)-based method captured native 5mC/5hmC methylation with single-

molecule and CpG resolution.

Differential methylation analyses between plasma and each neuronal and glial

cell type identified millions of differentially methylated CpGs for each comparison using

a threshold of $\Delta\beta \geq 0.8$ (Supplemental Table 2). Among the primary neurons, cortical

neurons exhibited the highest number of differentially methylated CpG sites (4.28

million), followed by dopaminergic (1.75 million) and spinal motor neurons (1.54 million).

Differential methylation analysis of glial cells—astrocytes, microglia, and Schwann

cells—compared to pooled blood plasma yielded between 1.24 and 1.43 million

differentially methylated CpG sites (Supplemental Table 2). Differentially methylated

regions (DMRs) identified with nanopore sequencing drastically outnumbered DMRs

identified via array (Supplemental Table 2).

To assess whether induced pluripotent stem cell (iPSC)-derived neurons could

serve as a more accessible alternative to primary cells for classifier development, we

included iPSC-derived cortical, dopaminergic, and spinal motor neurons in our

analyses. These cell types yielded a comparable number of differentially methylated

CpG sites relative to their primary counterparts (data not shown).

To assess cell type specificity, DMRs were compared across all neural

populations, including both primary and iPSC-derived cell types. Thousands of

overlapping and unique DMRs were identified per cell type. Figure 1A reports total,

overlapping, and unique DMRs, with directional visualizations of methylation change

(hypermethylation in red, hypomethylation in blue). Primary cortical neurons displayed

117,695 unique DMRs, primary dopaminergic neurons displayed 29,086 unique DMRs,

primary spinal motor neurons displayed 89,648 unique DMRs (Figure 1A). Unique, non-

redundant, cell type-specific DMRs for these cell types were used for downstream

classifier development.

Comparing DMRs between iPSC-derived neurons and blood plasma to their

primary counterparts revealed substantial divergence. Only 28.3% of DMRs in iPSC-

derived cortical neurons overlapped with primary cortical neurons, 38.1% with

dopaminergic neurons, and 6.8% with spinal motor neurons (Figure 1B). Although the

total number of DMRs was comparable, their content and genomic distribution differed

widely.

3.2 *In silico* validation demonstrates high analytical accuracy and specificity of cfDNA

classifiers

To assess the analytical accuracy and specificity of our DMR-based classifiers,

we generated *in silico* cfDNA-like gDNA mixtures containing defined proportions of cell

type-specific reads in a plasma background and applied our classifiers. To better

approximate native cfDNA properties, we computationally fragmented genomic DNA

reads from purified cells, which were initially longer than plasma-derived cfDNA.

Resultant fragment distributions more closely resembled circulating cfDNA

(Supplemental Figure 1).

The cortical neuron classifier displayed the highest predictive accuracy, with a

near-perfect correlation between predicted and actual cell fractions across the full

dilution series (R^2 to identity = 0.998; Figure 2A). Similarly high performance was

observed for primary dopaminergic neurons (R^2 to identity = 0.993, Figure 2B),

astrocytes (R^2 = 0.993, Figure 2D), Schwann cells (R^2 to identity = 0.986, Figure 2E),

and microglia (R^2 to identity = 0.992, Figure 2F), each achieving R^2 to identity values

above 0.98.

Strong linearity was also observed across iPSC-derived *in silico* mixtures,

including iPSC-derived cortical neurons (R^2 to identity = 0.946, Figure 2G),

dopaminergic neurons (R^2 to identity = 0.994), and spinal motor neurons (R^2 to identity

= 0.982, Figure 2I). In contrast, primary spinal motor neuron-derived fractions and

models developed from iPSC-derived neurons exhibited substantially reduced

performance (R^2 to identity = 0.876).

To assess cell type specificity, we also applied each classifier to *in silico* datasets

from other cell types. In all cases, slopes for the other cell types was near zero,

indicating high model specificity for each cell type (Supplemental Figures 2-10).

3.3 Clinical proof of concept for cfDNA classifiers in patient plasma

3.3.1 Neuron-specific cfDNA elevations distinguish Alzheimer's, Parkinson's, and ALS

samples from controls

After establishing the predictive ability of the models, we assessed their clinical

potential by applying the primary neuron classifiers to 219 plasma samples from healthy

controls and patients with MCI, AD, PD, or ALS. This analysis revealed cfDNA

elevations reflecting targeted neuronal cell loss across diseases, underscoring the

diagnostic potential of neuron-specific cfDNA methylation signals (Figure 3A–F).

Cortical neuron-derived cfDNA was significantly elevated in AD and MCI

(22.19%, SD = 5.45%) when compared to controls alone (5.73%, SD = 1.53%; p =

1.54e-28), and when compared to controls, PD, and ALS combined (10.48%, SD =

4.40%; p = 1.08e-21; Figure 3A, D). Dopaminergic neuron-derived cfDNA was

significantly elevated in PD (42.60%, SD = 10.60%) when compared to controls alone

(15.11%, SD = 5.19%; p = 6.92e-12; Figure 3B, E), and when compared to AD, MCI,

ALS, and controls combined (29.63%, SD = 12.12%; p = 1.00e-12). Spinal motor

neuron-derived cfDNA was significantly elevated in familial and sporadic ALS (7.80%,

SD = 4.45%) when compared to controls alone (4.61%, SD = 2.11%; p = 7.05e-08) and

when compared to all other groups (4.61%, SD = 2.11; p = 7.05e-8; Figure 3C, F).

By contrast, classifiers for astrocytes, Schwann cells, microglia, and iPSC-

derived neurons showed no consistent disease-associated changes in patient plasma

(Supplemental Figure 12). Astrocyte and microglial cfDNA remained stable across all

groups, and while Schwann cell cfDNA showed a modest increase in ALS relative to

controls, this effect was not significant when other diseases were included. Likewise,

iPSC-derived neuron classifiers failed to distinguish relevant disease states from

controls.

Together, these results are consistent with our broader observation that iPSC-

derived neuron methylation profiles differ substantially from those of primary neurons.

Importantly, they provide further support that the cfDNA signals we detected reflect true

cell-of-origin specificity, as cfDNA derived from distinct neuronal subtypes aligned

closely with the selectively vulnerable populations in AD, PD, and ALS. These findings

reinforce the diagnostic potential of cell type-resolved cfDNA profiling while also

underscoring that iPSC-derived neurons, although accessible, cannot accurately

recapitulate primary *in vivo* methylomes required for cell of origin classifiers.

3.3.2 Integrating multiple neuronal cfDNA signals enhances disease prediction

To further improve disease prediction accuracy, we applied cortical,

dopaminergic, and spinal motor neuron classifiers as an integrated model combining all

three. The integrated model was tested in both binary (e.g., AD vs. controls, PD vs.

controls, ALS vs. controls) and multi-disease (e.g., AD vs. all other diseases and

controls) contexts via ROC analysis, which enabled side-by-side comparisons of

univariate and multivariate models.

The multivariate classifiers yielded comparable or higher AUC values than the

univariate counterparts, generally displaying improved sensitivity, specificity, and overall

accuracy. For the binary AD vs. control analysis, the individual cortical neuron classifier

produced an AUC = 1.0000 (95% CI: 1.0000–1.0000), with 100% sensitivity, 100%

specificity, corresponding positive and negative predictive values (PPV and NPV) of

1.0000, and balanced accuracy of 100% at the optimal classification threshold of 0.5

(Figure 4). The multi-cell model that integrated all 3 neurons maintained perfect

performance, with AUC = 1.0000 (95% CI: 1.0000–1.0000), with 100% sensitivity, 100%

specificity, corresponding positive and negative predictive values (PPV and NPV) of

1.0000, and balanced accuracy of 100% at the optimal classification threshold of 0.5.

To further demonstrate the added value of integrating multiple cell-type

signatures into one model, we compared the performance of the combined classifier to

the corresponding univariate models across PD and ALS cases. In the PD comparisons

(Figure 5), while both the dopaminergic and combined models achieved perfect

separation from controls (AUC = 1.0000), only the combined model maintained strong

performance when distinguishing PD from ALS and AD. In contrast, cortical and spinal

motor classifiers, which performed well in PD vs. control settings (AUC = 0.9979 and

0.9575, respectively), showed steep declines in the three-class setting (AUC = 0.5997

and 0.5446). The combined model effectively aggregated discriminative signals across

relevant and partially overlapping cell types, preserving high accuracy across a more

complex clinical landscape.

A similar pattern held for ALS (Figure 6). The spinal motor classifier alone yielded

strong binary classification (AUC = 0.9680), but its specificity dropped in multi-disease

comparisons (AUC = 0.6538). Other univariate classifiers, while showing high AUCs in

ALS vs. control settings (cortical: 0.9696; dopaminergic: 0.9872), again lacked

discriminative power across multiple diseases (AUCs = 0.5600 and 0.5497). By

contrast, the combined ALS model sustained high discrimination (AUC = .9905),

highlighting its ability to integrate weak but complementary signals from across cell

types to improve disease specificity.

These cell type-specific disease predictions were completely lost when iPSC-

derived models were applied, whether individually or in combined multi-cell formats, and

in both binary and multi-disease contexts (Supplemental Figures 13–15). Together,

these results demonstrate that while neuron-specific classifiers capture relevant disease

signals, integrating cfDNA methylation features across neuronal populations provides

greater accuracy and specificity for predicting disease in real-world contexts where all

disease diagnoses are possible.

4. DISCUSSION

This study establishes a scalable liquid biopsy framework for cell type-resolved

detection of neurodegeneration from cfDNA methylation profiling. To address the

limitations of our prior Alzheimer's work[34], we generated a whole-genome nanopore

methylation atlas of six primary neural cell types and developed classifiers that

sensitively and specifically detect cortical, dopaminergic, and spinal motor neuron DNA.

Applying these classifiers to blood plasma from patients with AD, PD, and ALS, we

found that cfDNA from vulnerable neuronal populations can be detected in circulation:

cortical cfDNA was elevated in AD, dopaminergic cfDNA in PD, and spinal motor neuron

cfDNA in ALS, achieving an AUC > 0.98 (Figures 4–6). These signals were disease-

associated and orthogonal, enabling clear separation of AD, PD, and ALS in multiclass

comparisons, and distinction of cell type-specific pathologies rather than generic

neurodegeneration.

Integrating methylation signatures from multiple neuronal populations into a

multivariate model improved sensitivity, specificity, and overall accuracy compared to

individual classifiers, highlighting the value of capturing the broader landscape of

neurodegeneration. Together, these findings suggest potential utility of cfDNA

methylation analysis in detecting and monitoring cell type-specific neurodegeneration.

Applied in concert with other biomarkers, cfDNA methylation could enhance the

molecular precision of disease prediction.

Our approach builds on established workflows for cfDNA cell-of-origin

deconvolution pioneered in oncology, organ transplantation, and prenatal testing. These

applications rely on the observation that DNA methylation signatures are highly cell

type-specific, reproducible, and stable. Landmark studies have shown that methylation

patterns are >99% stable across 39 sorted human cell types derived from 205 healthy

tissues, and that replicates of the same cell type are >99.5% identical[31]. These findings

emphasize that methylation-based identity programs are robust to environmental

perturbation and therefore provide a biologically reliable substrate for cfDNA

deconvolution[30,32,33], and demonstrates that cfDNA carries rich, real-time information

about tissue-of-origin injury and cellular turnover.

While neurological applications of cfDNA deconvolution have lagged behind

oncology[27,40], early studies offer compelling evidence of its utility. These studies

demonstrate that neuron-derived cfDNA can be detected in circulation and suggest

concentrations correlate with clinical phenotypes in traumatic brain injury and multiple

sclerosis[35]. More recently, our own pilot study demonstrated elevated cortical neuron-

derived cfDNA in patients with AD and MCI compared to healthy controls[34].

While these and other studies established early feasibility, they have also

exposed limitations. Most prior efforts have relied on bisulfite conversion-based

sequencing that targets limited CpG loci or on array-based bisulfite methods that

interrogate only a fraction of the methylome at predefined sites. These approaches

degrade DNA[36,41], and introduce PCR biases[37]. Collectively, these constraints have

historically reduced both the biological specificity and the translational utility of cfDNA

methylation profiling.

Building on earlier work, our study overcomes these barriers by applying whole-

genome, native nanopore sequencing. The human genome contains ~30 million CpG

sites[31,42], yet Illumina's EPIC array only surveys ~900,000 (~3%), primarily within CpG

islands and promoters[43]. While arrays have expedited methylation breakthroughs and

advanced the field through standardized workflows, they systematically exclude ~97%

of the methylome and disproportionately emphasize well-characterized regulatory

regions.

In our prior work, we identified a cortical neuron cfDNA marker using array

data[34], but could not establish whether it was uniquely cortical, pan-neuronal, or

coincident. Here, whole-genome nanopore sequencing of purified primary cortical

neurons and human blood plasma captured the entire CpG context, revealing millions of

differentially methylated sites compared to the thousands by EPIC array, a lower-than-

expected discovery rate given probe coverage (Supplemental Table 2). Indeed, the

array captured only ~0.215% of the DMRs identified between cortical neurons and

plasma cfDNA, compared to the expected 3%. This highlights the disproportionate

representation of regulatory hotspots in array-based studies and underscores the value

of native sequencing for unbiased discovery of cell type-specific signatures.

We expanded the biological scope of our reference atlas—encompassing

cortical, dopaminergic, and spinal motor neurons as well as glial populations—which

further improved disease specificity (Figures 4–6). This expansion enabled us to

eliminate overlapping DMRs and focus on regions that were uniquely cell type-specific

relative to the other populations. When applied to human clinical plasma samples,

multivariate models from the broader atlas resolved cases that were ambiguous or

misclassified by single-cell classifiers (Figures 4-6), underscoring the additive value of

combining complementary cell type-specific signals. This highlights both the strengths

of native-read sequencing and the value of expanding the reference atlas, which

improved biological resolution and enhanced the accuracy of disease classification.

We acknowledge that true cell type specificity requires progressively eliminating

overlapping methylation signals by incorporating additional reference cell types. In

principle, a fully comprehensive atlas would require coverage of the >200 histologically

defined and >6,000 transcriptomically defined human cell types[45,46]. Such exhaustive

representation is not currently feasible, but for the purposes of this study, we

determined that our selected set of neural populations was sufficient to generate robust,

cell type-specific classifiers, a conclusion supported by our results. Moving forward, we

view this limitation as an opportunity. Our data suggest that expanding the atlas to

include additional CNS and peripheral cell types will likely enhance analytical accuracy

and broaden the clinical utility of the cfDNA classifiers for detecting and differentiating

neurodegeneration in disease. However, the acquisition of primary neuronal and glial

cell types is often logistically difficult due to the limited availability of human donor

material.

We encountered this limitation when sourcing primary human spinal motor

neurons, which are rarely available from commercial vendors. To address this, we

directly isolated neurons from postmortem spinal cord tissue using an optimized Percoll

gradient approach. Although this method was likely less specific than FACS-based

sorting used in prior atlases, the selective elevation of spinal motor neuron cfDNA in

ALS suggests that the core disease-relevant signatures were captured (Figure 3C,

Figure 6). As an exploratory alternative, we evaluated iPSC-derived neurons as a more

accessible substitute. However, these cells failed to perform in clinical testing

(Supplemental Figures 8–10), consistent with our qualitative observation that iPSC-

derived methylomes differ substantially from their primary counterparts. Together, these

findings underscore that primary tissue-derived references are critical for developing

robust, biologically relevant classifiers.

Several other limitations of the present study should be noted. Single-time-point

blood plasma samples are inherently retrospective and cross-sectional, limiting the

ability to assess temporal cfDNA dynamics. Longitudinal studies will be essential to

determine whether cfDNA changes can detect neurodegeneration before clinical

symptoms emerge and to evaluate how reliably they track progression and therapeutic

response. Notably, in our cohort, all individuals with MCI who subsequently progressed

to AD exhibited elevated cortical cfDNA at the time of sampling. While this suggests that

neuron-derived cfDNA can capture disease-relevant signals before clinical disease

conversion, direct longitudinal AD studies will be required to confirm this observation. In

addition, sample sizes of rarer diagnostic subgroups, such as MCI and familial ALS,

were small and lacked demographic diversity, highlighting the need for larger, more

representative cohorts to validate population-level generalizability.

We acknowledge that other neuronal and glial cell types with clear diagnostic

relevance, such as oligodendrocytes (multiple sclerosis), remain absent from our atlas

due to sourcing challenges. Incorporating these populations, along with additional

peripheral cell types, will be important for refining background correction, improving

disease-specific resolution, and broadening clinical applicability. Nevertheless, the

robust detection of highly specific cfDNA signals across hundreds of patient plasma

samples underscores the strength of our current framework and highlights its potential

for translational impact even at this stage of development.

Unlike static biomarkers such as protein concentrations or imaging, which often

change slowly and variably, cfDNA has a short half-life of minutes to hours[19,49],

providing a near real-time molecular snapshot of neuronal injury. In our cohort, we also

explored the protein biomarker data that accompanied the patient plasma samples

(Supplemental Figure 11). However, because only raw concentrations were available

and we did not have access to the proprietary diagnostic pipelines used by providers,

these analyses were considered ancillary and not central to our conclusions. We,

instead, focused on cfDNA methylation signals as the primary endpoint in alignment

with clinically-verified diagnoses, while recognizing that protein biomarkers remain an

important complementary modality that will be explored more explicitly in the future.

This dynamic nature makes cfDNA a particularly attractive candidate for

monitoring disease progression and therapeutic responses, offering sensitivity to short-

term molecular changes that may not be captured by traditional approaches. In clinical

practice, scalable cfDNA assays could support earlier diagnosis, presymptomatic risk

stratification, and real-time monitoring of disease activity. In therapeutic development,

cfDNA classifiers could accelerate clinical trials by improving patient selection and

providing dynamic, objective endpoints, while preclinical applications may extend to

animal models to track cell type-specific degeneration *in vivo.*

Our findings demonstrate that cfDNA methylation profiling can capture selective

neuronal cell death with high specificity, establishing a biologically grounded and

technically scalable framework for noninvasive detection and differentiation of

neurodegenerative diseases. Future work expanding the reference atlas, incorporating

additional neuronal and peripheral cell types, and conducting longitudinal studies will be

critical to establish clinical validity. With these refinements, cfDNA classifiers could

improve diagnostic timelines and enable earlier intervention across a spectrum of

conditions, including AD, PD, and ALS. Beyond these diseases, the modular

architecture of independent but combinable classifiers provides a flexible platform that

could be extended to other neurological disorders with selective cellular vulnerabilities,

such as multiple sclerosis, frontotemporal dementia, and cerebellar ataxias[16,50-52]. More

broadly, this work supports cfDNA methylation profiling as a versatile biomarker

platform adaptable to both research and clinical contexts.

ACKNOWLEDGEMENTS/CONFLICTS/FUNDING SOURCES:

This research was supported by two primary sources: (1) a Sponsored Research

Agreement between Brigham Young University (BYU) and Renew Biotechnologies,

LLC, which wholly owns Resonant, LLC, and (2) Research Catalyst Program funding

from the Alzheimer's Disease and Dementia Research Center (ADRC) at Utah State

University (USU). All authors are affiliated with Renew Biotechnologies and its

subsidiary, Resonant, LLC, both for-profit entities, representing potential financial

conflicts of interest. All work was conducted in accordance with institutional policies on

conflict of interest disclosure and management.

5. FIGURES AND LEGENDS

Figure 1 - (A) UpsetR plot showing cell-specific DMRs and overlap between each cell

type. (B) UpsetR plot showing cell specific DMRs and overlap between ISPC's and

primary cells. Both panels include hypo and hypermethylation annotations to explore

epigenetic commonalities and differences among diseases.

Figure 2 - Correlation plots comparing predicted versus actual cell-type proportions in

synthetic dilution datasets for each cell type. (A) Predicting the proportion of cortical

derived cfDNA using cortical specific DMRs. (B) Predicting the proportion of

dopaminergic neuron derived cfDNA using dopaminergic neuron specific DMRs. (C)

Predicting the proportion of spinal motor neuron derived cfDNA using dopaminergic

neuron specific DMRs. (D) Predicting the proportion of astrocyte derived cfDNA using

astrocyte specific DMRs. (E) Predicting the proportion of schwann cell derived cfDNA

using schwann cell specific DMRs. (F) Predicting the proportion of microglial derived

cfDNA using microglial specific DMRs. (G) Predicting the proportion of ipsc cortical

neuron derived cfDNA using ipsc cortical neuron specific DMRs. (H) Predicting the

proportion of ipsc dopaminergic neuron derived cfDNA using ipsc dopaminergic neuron

specific DMRs. (I) Predicting the proportion of ipsc spinal motor neuron derived cfDNA

using ipsc spinal motor neuron specific DMRs.

Figure 3 - Visualization of neuron-specific cfDNA classifications across patient groups.

(a), (b), and (c) show the proportion of cortical, dopaminergic, and spinal cfDNA,

respectively, across AD, MCI, PD, ALS, and control samples. (d) and (e) plot cortical vs.

dopaminergic cfDNA percentages in a bivariate space to illustrate disease separation

with and without ALS patients. (f) plots cortical vs. dopaminergic vs. spinal in a 3d

model to illustrate disease separation.

ROC Curves Showing Ability to Accurately Predict Parkinson's Disease (Primary Cells)

PD vs Age Matched Controls	Cortical %	Dopaminergic %	Spinal %	Combined Cell %'s
AUC	0.99788	1.0000	0.9575	1.0000
AUC 95% CI Lower	0.9427	1.0000	0.9064	1.0000
AUC 95% CI Upper	1.0000	1.0000	1.0000	1.0000
Sensitivity	1.0000	1.0000	0.9688	1.0000
Specificity	0.9200	1.0000	0.8400	1.0000
Positive Predictive Value	0.9412	1.0000	0.8857	1.0000
Negative Predictive Value	1.0000	1.0000	0.9545	1.0000
Accuracy	0.9649	1.0000	0.9123	1.0000
Balanced Accuracy	0.9600	1.0000	0.9044	1.0000
Optimal Threshold	0.4910	0.5000	0.2661	0.5000
PD vs Controls + Other NDD	**Cortical %**	**Dopaminergic %**	**Spinal %**	**Combined Cell %'s**
AUC	0.5997	0.9430	0.5446	0.9851
AUC 95% CI Lower	0.4881	0.9015	0.4307	0.9688
AUC 95% CI Upper	0.7113	0.9845	0.6586	1.0000
Sensitivity	0.0000	1.0000	0.0312	0.9375
Specificity	0.5714	0.7619	0.6190	0.9365
Positive Predictive Value	0.0000	0.6809	0.0400	0.8824
Negative Predictive Value	0.5294	1.0000	0.5571	0.9672
Accuracy	0.3789	0.8421	0.4211	0.9368
Balanced Accuracy	0.2857	0.8810	0.3251	0.9370
Optimal Threshold	0.3425	0.1936	0.3785	0.3824

Figure 4 - Receiver Operating Characteristic (ROC) curves for predicting Alzheimer's

Disease using Primary Cell derived models for (a)(e) cortical, (b)(f) dopaminergic, and

(c)(g) spinal neuron-derived cfDNA, alone and in (d)(h) combination. (i) shows the

respective stats associated with each panel. Panels compare classification performance

against both controls and other neurodegenerative diseases, showcasing the additive

value of combining multiple cell type signals.

AD vs Age Matched Controls	Cortical %	Dopaminergic %	Spinal %	Combined Cell %'s
AUC	1.0000	0.9938	0.9815	1.0000
AUC 95% CI Lower	1.0000	0.9792	0.9499	1.0000
AUC 95% CI Upper	1.0000	1.0000	1.0000	1.0000
Sensitivity	1.0000	0.9231	0.9231	1.0000
Specificity	1.0000	1.0000	0.9600	1.0000
Positive Predictive Value	1.0000	1.0000	0.9231	1.0000
Negative Predictive Value	1.0000	0.9615	0.9600	1.0000
Accuracy	1.0000	0.9737	0.9474	1.0000
Balanced Accuracy	1.0000	0.9615	0.9415	1.0000
Optimal Threshold	0.5000	0.7368	0.4787	0.5000
AD vs Controls + Other NDD	Cortical %	Dopaminergic %	Spinal %	Combined Cell %'s
AUC	1.0000	0.5553	0.6538	1.0000
AUC 95% CI Lower	1.0000	0.4322	0.5394	1.0000
AUC 95% CI Upper	1.0000	0.6785	0.7682	1.0000
Sensitivity	1.0000	0.9231	0.9231	1.0000
Specificity	1.0000	0.3902	0.5000	1.0000
Positive Predictive Value	1.0000	0.1935	0.2264	1.0000
Negative Predictive Value	1.0000	0.9697	0.9762	1.0000
Accuracy	1.0000	0.4632	0.5579	1.0000
Balanced Accuracy	1.0000	0.6567	0.7115	1.0000
Optimal Threshold	0.5000	0.1280	0.1331	0.5000

Figure 5 - Receiver Operating Characteristic (ROC) curves for predicting Parkinson's

Disease using Primary Cell derived models for (a)(e) cortical, (b)(f) dopaminergic, and

(c)(g) spinal neuron-derived cfDNA, alone and in (d)(h) combination. (i) shows the

respective stats associated with each panel. Panels compare classification performance

against both controls and other neurodegenerative diseases, showcasing the additive

value of combining multiple cell type signals.

ALS vs Age Matched Controls	Cortical %	Dopaminergic %	Spinal %	Combined Cell %'s
AUC	0.9696	0.9872	0.9680	1.0000
AUC 95% CI Lower	0.9265	0.9662	0.9301	1.0000
AUC 95% CI Upper	1.0000	1.0000	1.0000	1.0000
Sensitivity	1.0000	1.0000	1.0000	1.0000
Specificity	0.8800	0.9200	0.8000	1.0000
Positive Predictive Value	0.8929	0.9259	0.8333	1.0000
Negative Predictive Value	1.0000	1.0000	1.0000	1.0000
Accuracy	0.9400	0.9600	0.9000	1.0000
Balanced Accuracy	0.9400	0.9600	0.9000	1.0000
Optimal Threshold	0.2546	0.2358	0.1511	0.5000
ALS vs Controls + Other NDD	Cortical %	Dopaminergic %	Spinal %	Combined Cell %'s
AUC	0.5600	0.5497	0.8206	0.9905
AUC 95% CI Lower	0.4474	0.4380	0.7126	0.9753
AUC 95% CI Upper	0.6726	0.6614	0.9285	1.0000
Sensitivity	0.0000	0.0800	0.5600	0.9200
Specificity	0.6857	0.5857	0.9857	1.0000
Positive Predictive Value	0.0000	0.0645	0.9333	1.0000
Negative Predictive Value	0.6575	0.6406	0.8625	0.9500
Accuracy	0.5053	0.4526	0.8737	0.9683
Balanced Accuracy	0.3429	0.3329	0.7729	0.9600
Optimal Threshold	0.2815	0.2679	0.5586	0.6725

Figure 6 - Receiver Operating Characteristic (ROC) curves for predicting Amyotrophic

Lateral Sclerosis using Primary Cell derived models for (a)(e) cortical, (b)(f)

dopaminergic, and (c)(g) spinal neuron-derived cfDNA, alone and in (d)(h) combination.

Panels compare classification performance against both controls and other

neurodegenerative diseases, showcasing the additive value of combining multiple cell

type signals.

6. REFERENCES

1.   2023 Alzheimer's disease facts and figures. *Alzheimers Dement*. Apr
2023;19(4):1598-1695. doi:10.1002/alz.13016
2.   Yang W, Hamilton JL, Kopil C, et al. Current and projected future economic
burden of Parkinson's disease in the U.S. *npj Parkinson's Disease*. 2020/07/09
2020;6(1):15. doi:10.1038/s41531-020-0117-1
3.   Tolosa E, Garrido A, Scholz SW, Poewe W. Challenges in the diagnosis of
Parkinson's disease. *The Lancet Neurology*. 2021;20(5):385-397.
4.   Mehta P, Raymond J, Punjani R, et al. Prevalence of amyotrophic lateral
sclerosis in the United States using established and novel methodologies, 2017.
Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration. 2023;24(1-2):108-
116.
5.   Mehta P, Kaye W, Raymond J, et al. Prevalence of Amyotrophic Lateral
Sclerosis - United States, 2015. *MMWR Morb Mortal Wkly Rep*. Nov 23
2018;67(46):1285-1289. doi:10.15585/mmwr.mm6746a1
6.   Zissimopoulos J, Crimmins E, St Clair P. The Value of Delaying Alzheimer's
Disease Onset. *Forum Health Econ Policy*. Nov 2014;18(1):25-39. doi:10.1515/fhep-
2014-0013
7.   Rossi SL, Subramanian P, Bovenkamp DE. The future is precision medicine-
guided diagnoses, preventions and treatments for neurodegenerative diseases. *Front
Aging Neurosci*. 2023;15:1128619. doi:10.3389/fnagi.2023.1128619
8.   Hampel H, Hu Y, Cummings J, et al. Blood-based biomarkers for Alzheimer's
disease: Current state and future use in a transformed global healthcare landscape.
Neuron. Sep 20 2023;111(18):2781-2799. doi:10.1016/j.neuron.2023.05.017
9.   DeKosky ST, Marek K. Looking backward to move forward: early detection of
neurodegenerative disorders. *Science*. 2003;302(5646):830-834.
10.   Henriques AD, Benedet AL, Camargos EF, Rosa-Neto P, Nóbrega OT. Fluid and
imaging biomarkers for Alzheimer's disease: Where we stand and where to head to.
Experimental gerontology. 2018;107:169-177.
11.   Resende EdPF, Guerra JJL, Miller BL. Health and socioeconomic inequities as
contributors to brain health. *JAMA neurology*. 2019;76(6):633-634.

12.   Gavrilova S. The Evolution of Diagnostic Boundaries of Alzheimer's Disease and
Novel Therapeutic Options: Эволюция диагностических границ болезни
Альцгеймера и новые терапевтические возможности. *Consort Psychiatr*.
2022;3(1):8-15. doi:10.17816/cp152
13.   Pollard CA, Saito ER, Burns JM, Hill JT, Jenkins TG. Considering Biomarkers of
Neurodegeneration in Alzheimer's Disease: The Potential of Circulating Cell-Free DNA
in Precision Neurology. *Journal of Personalized Medicine*. 2024;14(11).
doi:10.3390/jpm14111104
14.   Montoliu-Gaya L, Salvadó G, Therriault J, et al. Plasma tau biomarkers for
biological staging of Alzheimer's disease. *Nature Aging*. 2025/08/22
2025;doi:10.1038/s43587-025-00951-w
15.   Jack Jr. CR, Andrews JS, Beach TG, et al. Revised criteria for diagnosis and
staging of Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimer's &*
Dementia. 2024;n/a(n/a)doi:https://doi.org/10.1002/alz.13859
16.   Wilson DM, III, Cookson MR, Van Den Bosch L, Zetterberg H, Holtzman DM,
Dewachter I. Hallmarks of neurodegenerative diseases. *Cell*. 2023;186(4):693-714.
doi:10.1016/j.cell.2022.12.032
17.   Dor Y, Cedar H. Principles of DNA methylation and their implications for biology
and medicine. *The Lancet*. 2018/09/01/ 2018;392(10149):777-786.
doi:https://doi.org/10.1016/S0140-6736(18)31268-6
18.   Gosselt HR, Griffioen PH, van Zelst BD, Oosterom N, de Jonge R, Heil SG.
Global DNA (hydroxy)methylation is stable over time under several storage conditions
and temperatures. *Epigenetics*. Jan 2021;16(1):45-53.
doi:10.1080/15592294.2020.1786318
19.   Mouliere F. A hitchhiker's guide to cell-free DNA biology. *Neuro-Oncology*
Advances. 2022;4(Supplement_2):ii6-ii14. doi:10.1093/noajnl/vdac066
20.   Soelter TM, Whitlock JH, Williams AS, Hardigan AA, Lasseigne BN. Nucleic acid
liquid biopsies in Alzheimer's disease: current state, challenges, and opportunities.
Heliyon. 2022;8(4)
21.   Schwarzenbach H, Hoon DS, Pantel K. Cell-free nucleic acids as biomarkers in
cancer patients. *Nature Reviews Cancer*. 2011;11(6):426-437.
22.   Rahat B, Ali T, Sapehia D, Mahajan A, Kaur J. Circulating cell-free nucleic acids
as epigenetic biomarkers in precision medicine. *Frontiers in genetics*. 2020;11:844.
23.   Pai M-C, Kuo Y-M, Wang I-F, Chiang P-M, Tsai K-J. The role of methylated
circulating nucleic acids as a potential biomarker in Alzheimer's disease. *Molecular*
Neurobiology. 2019;56:2440-2449.
24.   Miller RH, Pollard CA, Brogaard KR, et al. Tissue-specific DNA methylation
variability and its potential clinical value. *Front Genet*. 2023;14:1125967.
doi:10.3389/fgene.2023.1125967
25.   Martins-Ferreira R, Leal BG, Costa PP. The Potential of Circulating Cell-Free
DNA Methylation as an Epilepsy Biomarker. *Frontiers in Cellular Neuroscience*.
2022;16:852151.
26.   Luo H, Wei W, Ye Z, Zheng J, Xu RH. Liquid Biopsy of Methylation Biomarkers in
Cell-Free DNA. *Trends Mol Med*. May 2021;27(5):482-500.
doi:10.1016/j.molmed.2020.12.011

27.   Li W, Li Q, Kang S, et al. CancerDetector: ultrasensitive and non-invasive cancer
detection at the resolution of individual reads using cell-free DNA methylation
sequencing data. *Nucleic Acids Res*. Sep 6 2018;46(15):e89. doi:10.1093/nar/gky423
28.   Liu MC, Oxnard GR, Klein EA, et al. Sensitive and specific multi-cancer detection
and localization using methylation signatures in cell-free DNA. *Annals of Oncology*.
2020/06/01/ 2020;31(6):745-759. doi:https://doi.org/10.1016/j.annonc.2020.02.011
29.   Luo H, Wei W, Ye Z, Zheng J, Xu R-h. Liquid biopsy of methylation biomarkers in
cell-free DNA. *Trends in molecular medicine*. 2021;27(5):482-500.
30.   Mattox AK, Douville C, Wang Y, et al. The Origin of Highly Elevated Cell-Free
DNA in Healthy Individuals and Patients with Pancreatic, Colorectal, Lung, or Ovarian
Cancer. *Cancer Discov*. Oct 5 2023;13(10):2166-2179. doi:10.1158/2159-8290.Cd-21-
1252
31.   Loyfer N, Magenheim J, Peretz A, et al. A DNA methylation atlas of normal
human cell types. *Nature*. 2023;613(7943):355-364.
32.   Sun K, Jiang P, Chan KA, et al. Plasma DNA tissue mapping by genome-wide
methylation sequencing for noninvasive prenatal, cancer, and transplantation
assessments. *Proceedings of the National Academy of Sciences*. 2015;112(40):E5503-
E5512.
33.   Moss J, Magenheim J, Neiman D, et al. Comprehensive human cell-type
methylation atlas reveals origins of circulating cell-free DNA in health and disease.
Nature Communications. 2018/11/29 2018;9(1):5068. doi:10.1038/s41467-018-07466-6
34.   Pollard C, Aston K, Emery BR, Hill J, Jenkins T. Detection of neuron-derived
cfDNA in blood plasma: a new diagnostic approach for neurodegenerative conditions.
Original Research. *Frontiers in Neurology*. 2023-October-31
2023;14doi:10.3389/fneur.2023.1272960
35.   Lehmann-Werman R, Neiman D, Zemmour H, et al. Identification of tissue-
specific cell death using methylation patterns of circulating DNA. *Proc Natl Acad Sci U S
A*. Mar 29 2016;113(13):E1826-34. doi:10.1073/pnas.1519286113
36.   Tanaka K, Okamoto A. Degradation of DNA by bisulfite treatment. *Bioorganic &
medicinal chemistry letters*. 2007;17(7):1912-1915.
37.   Kanagawa T. Bias and artifacts in multitemplate polymerase chain reactions
(PCR). *Journal of Bioscience and Bioengineering*. 2003/01/01/ 2003;96(4):317-323.
doi:https://doi.org/10.1016/S1389-1723(03)90130-7
38.   Strong MJ, Garruto RM. Isolation of fetal mouse motor neurons on discontinuous
Percoll density gradients. *In vitro cellular & developmental biology*. 1989;25(10):939-
945.
39.   Qi T, Pan M, Shi H, Wang L, Bai Y, Ge Q. Cell-Free DNA Fragmentomics: The
Novel Promising Biomarker. *Int J Mol Sci*. Jan 12 2023;24(2)doi:10.3390/ijms24021503
40.   Kang S, Li Q, Chen Q, et al. CancerLocator: non-invasive cancer diagnosis and
tissue-of-origin prediction using methylation profiles of cell-free DNA. *Genome Biology*.
2017/03/24 2017;18(1):53. doi:10.1186/s13059-017-1191-5
41.   Olova N, Krueger F, Andrews S, et al. Comparison of whole-genome bisulfite
sequencing library preparation strategies identifies sources of biases affecting DNA
methylation data. *Genome biology*. 2018;19(1):1-19.

42.   Bibikova M, Barnes B, Tsan C, et al. High density DNA methylation array with
single CpG site resolution. *Genomics*. 2011/10/01/ 2011;98(4):288-295.
doi:https://doi.org/10.1016/j.ygeno.2011.07.007
43.   Pidsley R, Zotenko E, Peters TJ, et al. Critical evaluation of the Illumina
MethylationEPIC BeadChip microarray for whole-genome DNA methylation profiling.
Genome biology. 2016;17:1-17.
44.   Oehler JB, Wright H, Stark Z, Mallett AJ, Schmitz U. The application of long-read
sequencing in clinical settings. *Human Genomics*. 2023/08/08 2023;17(1):73.
doi:10.1186/s40246-023-00522-3
45.   Quake SR. The cell as a bag of RNA. *Trends in Genetics*. 2021;37(12):1064-
1068. doi:10.1016/j.tig.2021.08.003
46.   Khan YS, Farhana A. Histology, cell. *StatPearls [Internet]*. StatPearls Publishing;
2023.
47.   Tom SE, Hubbard RA, Crane PK, et al. Characterization of dementia and
Alzheimer's disease in an older population: updated incidence and life expectancy with
and without dementia. *American journal of public health*. 2015;105(2):408-413.
48.   Wong W. Economic burden of Alzheimer disease and managed care
considerations. *The American journal of managed care*. 2020;26(8 Suppl):S177-S183.
49.   Kustanovich A, Schwartz R, Peretz T, Grinshpun A. Life and death of circulating
cell-free DNA. *Cancer Biol Ther*. 2019;20(8):1057-1067.
doi:10.1080/15384047.2019.1598759
50.   Fu H, Hardy J, Duff KE. Selective vulnerability in neurodegenerative diseases.
Nature neuroscience. 2018;21(10):1350-1358.
51.   Saxena S, Caroni P. Selective neuronal vulnerability in neurodegenerative
diseases: from stressor thresholds to degeneration. *Neuron*. 2011;71(1):35-48.
52.   Giguère N, Burke Nanni S, Trudeau L-E. On cell loss and selective vulnerability
of neuronal populations in Parkinson's disease. *Frontiers in neurology*. 2018;9:383041.
833