(Hydroxy-) Chloroquine, G6PD deficiency, and sex differences in mortality during the COVID-19 pandemic: An ecological study in Peru

Marius Krämer[1*] (https://orcid.org/0009-0002-4274-1838)

Abstract

Background: Peru implemented a nationwide chloroquine/hydroxychloroquine (CQ/HCQ) treatment policy during the early phase of the COVID-19 pandemic. Whether these drugs trigger hemolytic crises in individuals with glucose-6-phosphate dehydrogenase deficiency (G6PDd), a genetic condition primarily affecting men, with a relatively high prevalence in certain regions, is a matter of debate. This study investigated whether the CQ/HCQ policy contributed to sex-specific differences in mortality, potentially mediated by regional G6PDd prevalence.

Methods: I conducted an ecological analysis using both all-cause excess and COVID-19 mortality data at the state level for adults aged ≥45 years. Using sex differences in age standardized mortality rates (ASMR) as the outcome in two-way fixed effects (TWFE) models, interaction analyses were performed to examine the influence of regional G6PDd prevalence, CQ/HCQ availability, and elevated healthcare system strain during pandemic waves with the CQ/HCQ recommendation.

Results: Male all-cause excess and COVID-19 mortality consistently exceeded female mortality, with the largest sex differences coinciding with periods with CQ/HCQ treatment recommendations. The TWFE models revealed that states with higher G6PDd prevalence, greater CQ/HCQ availability, and elevated healthcare system strain presented increased male–female mortality gaps. These results are consistent

---

[1] Technical University of Munich, Germany; TUM School of Social Sciences and Technology
[*] Correspondence: marius.kraemer@tum.de

with the hypothesized mechanism in which oxidative stress from both COVID-19 infection and CQ/HCQ

exposure contributed to hemolytic crises among (undiagnosed) G6PD-deficient outpatients, potentially

misidentified as severe COVID-19 during periods of healthcare strain.

Conclusion: While causality cannot be established owing to the ecological design, these findings suggest

that pandemic treatment policies may have unintended, sex-specific consequences if population-specific

genetic vulnerabilities are overlooked. They underscore the broader public health implication that drugs

considered safe under normal conditions may have unanticipated risks during infectious crises that induce

additional oxidative stress, emphasizing the importance of integrating genetic risk awareness into future

emergency response frameworks.

Keywords: Peru, COVID-19, Hydroxychloroquine, Chloroquine, G6PD, Sex difference, Gender gap,

Hemolysis

Introduction

Mortality during the COVID-19 crisis has differed between males and females across countries and regions, reflecting both biological vulnerabilities and contextual factors [1, 2]. Some studies show higher mortality among men [3–5], whereas others reveal complex patterns, including periods or locations where women experienced equal or higher mortality [6, 7]. Given the biological plausibility of sex-specific drug risks, treatment policy may represent an overlooked driver of these patterns.

At the beginning of the pandemic, chloroquine (CQ) and hydroxychloroquine (HCQ) were seen as potential COVID-19 treatments. However, international trials led by the WHO were discontinued in July 2020 because of safety concerns and a lack of efficacy [8]. While most countries remained cautious, Peru adopted CQ/HCQ widely. Initial protocols restricted use to moderate or severe cases [9], but by mid-April 2020, HCQ was authorized for mild outpatient cases [10]. Peru retained HCQ in official guidelines until September 2020, months after international trials had halted its use [8, 11]. Given that Peru has one of the highest all-cause excess mortality rates globally [12], it provides a unique context for studying the impact of treatment policies.

The safety profile of CQ and especially HCQ in the context of COVID-19 has been debated. In a meta-analysis of 14 trials testing CQ/HCQ in hospitalized patients, it was associated with an 11% (95% CI 2–20%) increase in all-cause mortality [13]. An observational preprint study by the Peruvian Medical Research Institute (IETSI) found that compared with standard care, HCQ combined with azithromycin was associated with an 84% (95% CI 12–202%) increase in all-cause mortality in hospitalized COVID-19 patients [14]. Evidence from outpatient studies generally reported few safety concerns, but these trials consistently excluded individuals with contraindications to CQ/HCQ [15], often explicitly mentioning glucose-6-phosphate dehydrogenase deficiency (G6PDd) as one such exclusion [16–19].

Among the factors that may influence CQ/HCQ-related risk, G6PDd is particularly relevant: an X-linked

enzymatic disorder that renders hemizygous males especially vulnerable, whereas females are usually

carriers or only partially deficient; G6PDd can predispose individuals to acute hemolytic anemia when

exposed to oxidative stress, which may be triggered by infections or certain drugs [20]. While CQ/HCQ use

in malaria-endemic regions has not been associated with widespread hemolytic events, according to a

systematic review of the available evidence [21], case reports during the COVID-19 pandemic document

acute hemolytic anemia in G6PDd patients treated with HCQ [22–26].

Moreover, COVID-19 itself has been reported to be associated with acute autoimmune hemolytic anemia,

often presenting with delayed hemolytic symptoms that may be overlooked due to the predominance of

respiratory manifestations [27].

The combination of the oxidative stressors COVID-19 and CQ/HCQ may increase the risk of acute hemolytic

anemia among G6PDd patients. In particular, during outpatient CQ/HCQ use, hemolysis symptoms may

have been delayed or mistaken for COVID-19 symptoms (e.g., dyspnea, fever), complicating detection,

especially in an underdeveloped and heavily strained healthcare system. Furthermore, Peruvian treatment

protocols with CQ/HCQ did not include warnings about potential adverse reactions or contraindications

for G6PDd patients. Consequently, given that G6PDd disproportionately affects males, the nationwide

CQ/HCQ policy could plausibly have sex-specific consequences for mortality.

To my knowledge, no prior study has examined whether national COVID-19 treatment policies contributed

to sex-specific differences in all-cause excess or COVID-19 mortality. This ecological study addresses this

gap by testing two hypotheses:

1. If CQ/HCQ triggers acute hemolytic anemia in G6PDd COVID-19 patients and Peru's healthcare

system failed to manage these reactions, periods of widespread CQ/HCQ use should coincide with

increased male–female mortality differences. Within these periods, increased healthcare system

strain is expected to further amplify these differences.

2. If regional G6PDd prevalence modifies risk, areas with higher prevalence should show stronger

male–female mortality differences during periods of widespread CQ/HCQ use.

By integrating publicly available all-cause and COVID-19 mortality data with treatment policy information

and regional G6PDd prevalence estimates [28], this study explores the plausibility of the described

mechanism.

Data and Methods

83 For this analysis, I used Python 3.11, primarily Pandas (v2.2.3) for data handling and Statsmodels (v0.14.1)

84 along with Linearmodels (v5.4) for statistical analysis.

Data

86 All datasets are publicly available and were aggregated at the state and week levels. Mortality data from

87 2017–2019 were used to construct baseline mortality, while all other datasets cover 2020–2022. The

88 compiled dataset includes 24 states over 157 iso-weeks.

Mortality data

Estimating the true mortality burden of COVID-19 is challenging, particularly in countries with limited

healthcare system capacity and incomplete vital registration [29–31]. Consequently, some researchers

prefer all-cause excess mortality, defined as the difference between observed and expected deaths, as a

more reliable measure [6, 32]. In this study, I use both all-cause excess and COVID-19 mortality data.

All-cause mortality data were obtained from Peru's National Information System of Deaths (SINADEF) [33].

An assessment concluded that SINADEF represents a substantial improvement over earlier death registries

in Peru; although some incompleteness remains, no systematic difference in completeness by sex was

observed [34]. In addition, COVID-19 case fatality data were analyzed [35]. These figures are subject to

underreporting and have been subsequently adjusted by the authorities [30]. However, this subgroup was

more likely affected by the CQ/HCQ treatment policy.

Medicament data

101 I collected data on CQ/HCQ distributions from the database of the National Center for the Supply of

102 Strategic Health Resources (CENARES) [36]. Since official treatment protocols treated CQ and HCQ as

103 equivalents, I do not distinguish between them and summed their distributions after the conversion of CQ

104 to HCQ-equivalents via protocol dosage ratios (see Table I, Appendix for details).

Prevalence of G6PD deficiency

106 I used a map of Latin America provided by Monteiro et al. [28] showing model-based estimates of the

107 prevalence of glucose-6-phosphate dehydrogenase deficiency (G6PDd). I overlaid the Peruvian state

108 boundaries onto it (Figure V, Appendix) to assign prevalence values to each state (Table IV, Appendix).

COVID-19 test results and vaccination data

110 I obtained COVID-19 test results from Peruvian authorities, which included both molecular (i.e., PCR) [37]

111 and nonmolecular tests (i.e., rapid antibody tests, antigen tests, and chemiluminescence assays) [38]. Like

112 official treatment protocols, I treated all types of tests equivalently. Additionally, I obtained COVID-19

113 vaccination data from Peruvian authorities [39] treating all vaccine types equivalently.

Other Data

115 Data on the population were obtained from official statistics [40]. For reasons of feasibility, all rate

116 calculations are based on the 2021 mid-year population estimates provided by the Ministry of Health of

117 Peru [40], which readily include the necessary breakdowns by sex, age group, and region. Moreover, I

118 consulted official documents containing CQ/HCQ treatment protocols for COVID-19 patients [9, 10, 41,

119 42].

Methods

Data preparation

Deaths categorized as violent (accidents, homicides, or suicides) and rare observations with missing

information on sex, age, or location were excluded. All datasets were aggregated to the state level and to

a weekly timeframe. The state of Callao is completely embedded within Lima and was therefore assigned

to Lima instead of treating it separately. I stratified the data by the age groups 0–14, 15–44, 45–64, 65–

79, and 80+ and created the aggregated group 45+, which is the main focus of this analysis, as this group

is considered to be most affected by COVID-19 and related treatments.

Coding of COVID-19 treatment protocols (recommendation_HCQ)

Official COVID-19 treatment protocols varied in their stance on CQ/HCQ [9, 10, 41, 42]. I coded these

recommendations in an ordinal variable called *recommendation_HCQ* from 0 (no recommendation), to 1

(recommended only for hospitalized, i.e., medium and severe COVID-19 cases), 2 (additionally

recommended for confirmed outpatient, i.e., mild cases), and 3 (additionally recommended for suspected

outpatient, i.e., mild cases). Protocol changes were aligned with the weekly dataset by assigning each

protocol to all calendar weeks between its issuance and the release of the subsequent update.

Definition of the CQ/HCQ exposure period (period_HCQ)

In line with the hypotheses, I constructed the binary variable *period_HCQ*[2] to indicate weeks when outpatient treatment with CQ/HCQ likely occurred and acute hemolytic anemia may have been mismanaged due to a strained healthcare system.

The variable *period_HCQ* was set to 1 if (i) the official COVID-19 treatment protocol recommended CQ/HCQ for outpatients, i.e., mild cases (*recommendation_HCQ* = 2 or 3), and (ii) the proxy for healthcare system strain and pandemic waves, i.e., the z-score of all-cause mortality (all ages) exceeded the threshold of 3 for at least two consecutive weeks. To account for potential delays between treatment and death, I extended the period by two additional weeks after the z-score went below the threshold.

In at least 11 states, however, CQ/HCQ was soon displaced by ivermectin (IVM) for outpatient treatment. In Piura, for example, the state health minister stated in early June 2020 (cw23) that HCQ for mild cases has been replaced by IVM owing to its perceived efficacy and fewer side effects [43]. Moreover, ten states (Ayacucho, Cajamarca, Cusco, Huancavelica, Huánuco, Junín, Moquegua, Pasco, Puno, Tacna) implemented mass IVM distributions through the Ministry of Defense's Mega-Operación Tayta (MOT). During its first phase, MOT conducted household visits to identify vulnerable individuals; one week later, it began distributing IVM broadly, regardless of test status or symptoms, alongside other medications such as acetaminophen and azithromycin [44]. HCQ was not distributed as part of the MOT. Similarly, the regional government of Tumbes carried out a locally organized campaign with mass IVM distributions (Vanessa 2020a, 2020b). In these 11 states, I therefore shortened *period_HCQ* by ending it two weeks after the start of the mass IVM distribution to reflect a reasonable transition period.

---

[2] For simplicity, I use here 'HCQ' to refer to both chloroquine and hydroxychloroquine.

Since *period_HCQ* is based on assumptions and reasoning, I reconstructed it under alternative definitions

for robustness checks, using different z-score thresholds (2.5, 3, 3.5, 4) both with and without adjustments

for IVM campaigns.

State-level CQ/HCQ availability (HCQ_per_pop)

Reliable state-level time series data on CQ/HCQ consumption are unavailable. Instead, I used data on the

distribution of CQ/HCQ tablets to regional health authorities (i.e., stock data). For each state, I summed all

distributions that occurred before or during *period_HCQ* and created a time-invariant variable,

HCQ_per_pop[3]. To make the states comparable, I standardized the different types of drugs according to

treatment protocols and divided the number of these tablets by the state population, expressed per

100,000 people (for details, see Table I, Appendix).

In most states, the quantities of distributed CQ/HCQ doses were limited relative to the total number of

COVID-19 cases. On average, only 1.16 doses per positive test in the 45+ age group were delivered,

decreasing to 0.4 doses when positive tests across all ages were considered (Table II, Appendix).

Symptomatic but untested cases were also treated [42], indicating relative scarcity in most states and

supporting the correspondence between stock data and actual usage. Moreover, self-medication with

CQ/HCQ seems to have been uncommon in Peru [45–47], and web searches indicate limited public interest

[48]. Therefore, *HCQ_per_pop* is unlikely to be strongly confounded by privately obtained doses. Although

this measure does not capture actual intake or distinguish between inpatient and outpatient

administration, *HCQ_per_pop* is expected to reasonably reflect interstate differences in CQ/HCQ

availability.

---

[3] For simplicity, I use 'HCQ' to refer to both chloroquine and hydroxychloroquine.

Statistical analysis

Estimating all-cause excess mortality rates

All-cause excess mortality is defined as the difference between the expected mortality and the observed

mortality. I estimated weekly absolute numbers of expected deaths (baselines) for 2020–22 for each state,

sex, and age group on the basis of historical data from 2017–19 via a time series regression model by

Karlinsky and Kobak [12]. Absolute numbers of all-cause excess deaths were then calculated by subtracting

the baseline from observed deaths, and crude mortality rates for each group were derived by dividing

excess deaths by the corresponding population and scaling per 100,000 individuals. For feasibility and

consistency, all rate calculations for 2020-22 used the 2021 mid-year population estimates provided by

the Ministry of Health of Peru [40], which already include necessary breakdowns by sex, age group, and

locality. This approach may not yield the most precise estimates of all-cause excess mortality rates.

However, more sophisticated methods would likely provide little additional insight, as only three years of

training data (2017-19) are available.

Age-standardized mortality rates (ASMR)

The confounding effect of different age structures makes comparisons between sexes for aggregated age

groups misleading. Therefore, for the 45+ age group, I calculated the age-standardized mortality rates

(ASMR) for males and females in each state via the direct method with Peru's age structure as the standard

population. The sex difference was then calculated as the difference in the ASMR between males and

females. This was done separately for both all-cause excess and COVID-19 mortality rates.

To quantify the statistical uncertainty of the estimated all-cause excess ASMRs, I calculated 95%

confidence intervals (CIs) using the predictive standard errors from the baseline mortality regressions. For

each sex, the derived variances were summed across age groups using the squared standard-population

weights. The CIs were constructed as the point estimate ± 1.96 times the square root of this variance.

   TWFE model specification

The sex difference in all-cause excess, as well as in COVID-19 ASMR, for the 45+ age group was used as the

dependent variable in two-way fixed-effects (TWFE) models, specified as follows:

$ASMR\_male_{i,t} - ASMR\_female_{i,t}$

$\qquad\qquad = \gamma\, period\_HCQ_{i,t} + \theta\, (period\_HCQ_{i,t} \times G6PDD_i)$

$\qquad\qquad + \phi\, (period\_HCQ_{i,t} \times HCQ\_per\_pop_i) + \psi\, (period\_HCQ_{i,t} \times z - score_{i,t}) + \beta\, X_{i,t}$

$\qquad\qquad + \eta_i + \delta_t + \varepsilon_{i,t}$

where $period\_HCQ_{i,t}$ indicates the weeks with likely outpatient treatment with CQ/HCQ during a strained

healthcare system in state i at time t, $(period\_HCQ_{i,t} \times G6PDD_i)$, $(period\_HCQ_{i,t} \times HCQ\_per\_pop_i)$,

and $(period\_HCQ_{i,t} \times z - score_{i,t})$ are interaction terms capturing the conditional effects of G6PDd

prevalence, QC/HCQ availability, and healthcare system strain—as measured by the all-cause mortality z-

score for both sexes and all age groups—during *period_HCQ*, reflecting the hypotheses. $X_{i,t}$ is a vector of

control variables, η_i and δ_t are state- and week-specific fixed effects, respectively, and $\varepsilon_{i,t}$ is the error

term[4]. I used Driscoll-Kraay standard errors to account for cross-sectional dependence and serial

correlation [49] and used wild cluster bootstrap inference [50] as a robustness check, given the small

number of states.

The TWFE setup addresses several sources of confounding. First, time-invariant differences across states

not included in the interaction terms are absorbed by the state fixed effects. Second, week fixed effects

account for shocks common to all states at a given week. Third, using the sex difference in ASMR as the

dependent variable implicitly removes factors that influence men and women in a similar way.

---

[4] For simplicity, I use 'HCQ' in the variable names to refer to both chloroquine and hydroxychloroquine.

Control variables include the z-score, age-standardized sex differences in COVID-19 test positive rates, and

cumulative sex differences in first-dose vaccination rates for individuals aged 45+ years. Analogous to the

age standardization of mortality rates, test positivity rates were standardized via the national age

distribution of tests to account for differences in testing by age. The cumulative vaccination rate sex

differences were calculated as the sum over time of the male first-dose rates minus the female rates.

The z-score control variable estimates the 'baseline' relationship between overall excess mortality and sex

differences in ASMR independent of *period_HCQ.* It captures the general male-biased mortality pattern

observed during periods of increased all-cause mortality, whereas the z-score interaction term captures

any additional effect during *period_HCQ*, reflecting the hypothesis that greater healthcare system strain

could have further amplified sex differences in mortality by increasing the likelihood of adverse effects

remaining unrecognized.

Because testing capacity in Peru was limited, particularly during the first wave, reported COVID-19 test

results are an imperfect proxy for true infections. Consequently, the observed sex differences in test

positivity may not fully capture actual infection patterns. Nonetheless, several considerations suggest that

the resulting bias is limited. Nationwide lockdowns and contact restrictions were not gender-specific and

are absorbed by week fixed effects, leaving only state-level reopening policies as a possible source of

differential exposure. Moreover, household transmission was high—a study in Lima reported a secondary

attack rate of 53% [51]—which likely reduced sustained disparities between sexes. Finally, Peruvian

seroprevalence studies consistently show no significant sex differences in infection levels [52–54]. Overall,

while short-term sex-specific fluctuations in infections may be partially captured by the control variable,

persistent sex-specific differences in infection risk were small and unlikely to explain the observed sex

difference in all-cause excess mortality.

Results

Sex differences during pandemic waves

Except for Apurímac, all Peruvian states recorded higher male than female ASMR in the 45+ age group during pandemic waves (i.e., periods ≥ 2 consecutive weeks with z-scores ≥ 3), leading to predominantly positive average sex differences in all-cause excess ASMR, as well as COVID-19 ASMR (Table 1). In most states, these differences were greater during the first pandemic wave (when CQ/HCQ was recommended) than in subsequent waves, with several cases showing confidence intervals that (nearly) exclude zero or p-values ≤0.05. Moreover, the size of the observed differences varied considerably, ranging from negligible or negative (Apurímac, Pasco) to very large (e.g., Madre de Dios, Moquegua, Ucayali).

Table 1: Average sex differences in all-cause excess and COVID-19 ASMR during pandemic waves

State	Weeks Wave 1	Weeks Waves 2+	Avg. all-cause excess sex diff. Wave 1 [95% CI]	Avg. all-cause excess sex diff. Waves 2+ [95% CI]	Diff. all-cause excess (W1–W2+) [95% CI]	COVID-19 Avg. Sex diff. Wave 1	COVID-19 Avg. Sex diff. Waves 2+	Diff. COVID-19 (W1–W2+) (p-value)
Amazonas	8	10	41.9 [18.3, 65.4]	17.3 [-6.3, 40.8]	24.6 [-8.7, 57.9]	34.4	19.2	15.2 (0.040)
Ancash	17	21	28.1 [12.6, 43.6]	19.0 [3.5, 34.5]	9.1 [-12.8, 31.1]	24.2	18.4	5.8 (0.037)
Apurimac	4	31	-6.5 [-31.7, 18.8]	2.9 [-22.3, 28.2]	-9.4 [-45.1, 26.4]	-1.1	5.4	-6.6 (0.367)
Arequipa	14	22	49.8 [36.0, 63.7]	16.7 [2.8, 30.5]	33.2 [13.6, 52.7]	3.3	14.6	-11.3 (0.078)
Ayacucho	2	8	54.0 [28.4, 79.6]	3.8 [-21.8, 29.4]	50.2 [14.0, 86.4]	27.2	8.9	18.3 (0.043)
Cajamarca	10	20	24.8 [12.1, 37.5]	11.9 [-0.9, 24.6]	12.9 [-5.1, 30.9]	19.8	12.3	7.5 (0.058)
Cusco	6	18	33.4 [15.6, 51.2]	10.8 [-6.9, 28.6]	22.6 [-2.5, 47.7]	7.4	14.7	-7.4 (0.053)
Huancavelica	6	18	30.1 [-8.5, 68.8]	6.1 [-32.5, 44.8]	24.0 [-30.7, 78.6]	19.8	7.8	12.0 (0.030)
Huánuco	12	35	12.8 [-2.9, 28.6]	8.0 [-7.8, 23.7]	4.8 [-17.4, 27.1]	11.3	11.4	-0.1 (0.966)
Ica	23	30	36.6 [18.9, 54.4]	21.3 [3.6, 39.1]	15.3 [-9.8, 40.4]	23.5	19.4	4.1 (0.337)
Junín	11	23	32.4 [16.4, 48.4]	25.7 [9.6, 41.7]	6.7 [-15.9, 29.4]	17.6	24.6	-7.0 (0.152)
La Libertad	18	25	30.2 [18.4, 42.1]	9.9 [-1.9, 21.7]	20.3 [3.6, 37.0]	20.7	11.3	9.5 (0.010)
Lambayeque	18	33	32.0 [11.6, 52.3]	9.6 [-10.8, 29.9]	22.4 [-6.4, 51.2]	23.2	12.3	10.9 (0.004)
Lima + Callao	27	31	28.7 [22.3, 35.0]	12.9 [6.5, 19.2]	15.8 [6.8, 24.8]	20.7	16.5	4.2 (0.201)
Loreto	14	15	58.8 [43.2, 74.4]	18.2 [2.5, 33.8]	40.6 [18.5, 62.7]	63.0	15.0	48.0 (0.001)
Madre de Dios	10	7	196.4 [131.4, 261.4]	100.3 [35.3, 165.3]	96.1 [4.2, 188.0]	194.8	101.0	93.7 (0.012)
Moquegua	10	12	119.3 [88.0, 150.5]	19.4 [-11.9, 50.6]	99.9 [55.7, 144.1]	122.4	29.8	92.6 (0.000)
Pasco	9	22	15.1 [-12.3, 42.6]	19.9 [-7.6, 47.3]	-4.7 [-43.5, 34.1]	11.0	11.9	-0.9 (0.889)
Piura	12	10	52.4 [34.6, 70.2]	13.4 [-4.4, 31.3]	39.0 [13.8, 64.2]	45.8	20.6	25.2 (0.002)
Puno	8	28	53.3 [39.9, 66.7]	18.8 [5.4, 32.2]	34.4 [15.5, 53.4]	21.0	15.1	5.8 (0.084)
San Martin	12	16	41.8 [21.3, 62.2]	18.1 [-2.3, 38.6]	23.6 [-5.3, 52.5]	40.4	18.3	22.1 (0.002)
Tacna	10	25	71.0 [42.7, 99.4]	35.1 [6.8, 63.5]	35.9 [-4.2, 76.0]	57.2	27.9	29.2 (0.012)
Tumbes	11	17	58.1 [22.2, 94.1]	36.8 [0.8, 72.7]	21.4 [-29.4, 72.2]	54.7	42.0	12.7 (0.238)
Ucayali	6	9	110.0 [85.2, 134.7]	39.1 [14.3, 63.8]	70.9 [35.9, 105.9]	95.3	50.7	44.6 (0.017)
Median	10.5	20.5	41.8	17.8	23.6	23.4	15.8	10.2
Mean	11.6	20.3	50.2	21.3	28.9	39.2	21.8	17.5
Std. Dev.	5.6	8.1	39.9	18.5	26.4	42.1	19.0	26.5

 Note: Average ASMR sex differences (males minus females) for individuals aged 45+ in deaths per 100,000. Waves are defined as periods of ≥2 consecutive weeks with a z-score ≥3. The

 95% confidence intervals (95% CIs) are based on the predictive variance of the baseline models. P-values are from two-sided permutation tests.

Figure 1 shows the male and female all-cause excess ASMR for those aged 45+ over time, with markedly

higher values for males, especially during the 2020 wave, and a close correspondence with the all-age

mortality z-score. There is no period where female all-cause excess mortality significantly exceeds male

mortality (i.e., a negative sex difference) and regional heterogeneity is evident. Periods where *period_HCQ*

=1 (i.e., weeks of substantial healthcare system strain (z-score ≥3) combined with the CQ/HCQ

recommendation for COVID-19 outpatients) are also indicated. Comparable figures for all-cause sex

differences and for the COVID-19 ASMR are provided in the Appendix (Figure I, Figure II, Appendix).

Figure 1: Male and female all-cause excess ASMR for individuals aged 45+ over time by state

 Note: 95% CI stands for 95% confidence interval.

TWFE model results

Table 2: Results of the TWFE models

Model	1 (all-cause excess)	2 (all-cause excess)	3 (COVID-19)	4 (COVID-19)
R² (Within)	0.3863	0.4706	0.4606	0.6115
F-statistic	224.37 (p = 0.000)	241.69 (p = 0.000)	328.76 (p = 0.000)	492.17 (p = 0.000)
Constant	2.771*** (0.813)	4.219*** (0.492)	2.094*** (0.741)	3.324*** (0.408)
period_HCQ	22.049*** (5.124)	-41.808*** (5.077)	15.689*** (4.424)	-47.428*** (5.690)
period_HCQ x G6PDD_prevalence	-	8.209*** (1.845)	-	8.232*** (1.761)
period_HCQ x HCQ_per_pop	-	0.770*** (0.177)	-	0.823*** (0.167)
period_HCQ x z-score	-	3.050*** (0.671)	-	2.653*** (0.624)
z-score	2.611*** (0.529)	1.529*** (0.279)	2.486*** (0.441)	1.571*** (0.241)
Sex difference test positive rate (lag 4)	34.474*** (11.454)	31.550*** (10.166)	31.927*** (8.870)	28.668*** (8.717)
Sex difference vaccination rate (lag 3)	-27.394** (11.570)	-9.402 (7.082)	-34.252*** (12.506)	-15.584** (6.881)

Note: Dependent variable: all-cause excess or COVID-19 sex differences in ASMR (males minus females) for individuals aged 45+ years. The z-score is based on all-cause excess mortality in all models. Driscoll–Kraay standard errors in parentheses (see Table III, Appendix for p-values from wild cluster bootstrap inference). Fixed Effects: entity, time. Observations: 3454 (22 states, 157 weeks). The states Apurímac and Ayacucho were excluded because no clear period_HCQ > 4 weeks was identified.

I present two models for each of the two outcomes (all-cause excess, COVID-19) to separate the main and interaction effects. In models 2 and 4 (Table 2), the main effect of *period_HCQ* should be interpreted with caution, since it is not only identified at unrealistic values of the interactions (all zero), but is also inflated by multicollinearity with the interaction terms. Models 1 and 3 (Table 2) show a positive and significant coefficient for *period_HCQ* for both outcome variants (22.049, $p \leq 0.001$; 15.689, $p \leq 0.001$), indicating that the difference between male and female ASMR increased during the period when CQ/HCQ was officially

recommended for COVID-19 outpatients and the healthcare system was strained (z-score ≥ 3). However,

this allows no conclusions regarding the hypothesized mechanism because it does not disentangle the

contributions of G6PDd prevalence, regional HCQ availability, or intensity of healthcare system strain.

Models 2 and 4 introduce interaction terms with G6PDd prevalence, regional CQ/HCQ availability, and the

all-cause *z-score*. This specification allows the effect of *period_HCQ* to vary by regional risk factors and

healthcare system strain (proxied by the *z-score*), effectively isolating the hypothesized mechanism. All

three interactions are positive and highly significant for both outcome variants: *period_HCQ × G6PDd*

prevalence (8.209, p ≤0.001; 8.232, p ≤0.001), *period_HCQ × HCQ_per_pop* (0.770, p ≤0.001; 0.823, p

≤0.001), and *period_HCQ × z-score* (3.050, p ≤0.001; 2.653, p ≤0.001). For example, a one-category increase

in G6PDd prevalence – corresponding to a two-percentage-point increase (Table IV, Appendix) – alters the

effect of *period_HCQ* by an additional sex difference of 8.209 all-cause excess deaths or 8.232 COVID-19

deaths per 100,000. Marginal effects plots for G6PDd prevalence are provided in the appendix (Figures III

and IV). The coefficients' magnitudes of the interaction terms should be interpreted with caution, as they

vary slightly over the tested definitions of *period_HCQ*. However, significance and signs are robust and

consistent with the tested mechanism.

Taken together, the results of the interaction terms suggest that in regions with higher G6PDd prevalence,

greater CQ/HCQ availability, and stronger healthcare system strain, sex differences in all-cause excess and

COVID-19 ASMR increased disproportionately during the CQ/HCQ treatment policy for outpatients,

supporting the hypothesized mechanism of drug-induced hemolytic anemia under stressed healthcare

system conditions.

The control variables also behaved as expected: the *z-score* captures the general male-biased mortality

pattern observed during periods of increased all-cause mortality and is positive and significant. The

coefficient of the 4-week lagged sex difference in test positive rates formally reflects a 100% difference

(i.e., only men tested positive and no women) and is positive and significant in all models. It should be

interpreted as capturing short-term shifts in infection burdens between sexes, as discussed in the Methods

section. Although this variable is an imperfect proxy for true infection rates, its significance reflects that

observed mortality patterns overlapped with periods and regions in which one sex was more likely to be

registered as infected.

The 3-week lagged sex difference in first-dose vaccination rates has a negative coefficient, suggesting that

weeks with relatively higher rates of female vaccination are associated with smaller sex differences in

mortality. As with the test-positivity variable, the coefficient formally reflects a 100% difference and should

be interpreted as capturing relative shifts in vaccination coverage between the sexes during the

nationwide vaccination campaign starting in April 2021. The effect is somewhat unstable across

specifications and was observed only in 2021, so it should be interpreted cautiously. While it provides

some insight into sex-differential dynamics during the vaccination campaign, this variable does not

contribute to the identification of *period_HCQ* and serves as a supplementary result.

To further assess which sex drives the effect of increased sex differences in ASMR during *period_HCQ*,

separate models were estimated with male and female ASMR as outcomes. While these specifications are

subject to omitted variable bias and endogeneity, they serve as a diagnostic check and show a strong effect

for males but only negligible and mostly insignificant effects for females, suggesting that the observed sex

differences stem from higher male mortality rather than unusually low female mortality.

Discussion

The findings from Peru align with those of prior studies showing among men higher COVID-19 mortality

[4, 5] and all-cause excess mortality [3]. Significant periods with higher female than male all-cause excess

mortality, as observed in Thailand [6], were not present in Peru. This difference can be explained partly by

the exclusion of traffic accidents in this analysis, which were probably responsible for the observed pattern

in Thailand [6]. The correlation between general excess mortality (z-score) and sex differences, as found

in Europe [3], was also confirmed by this analysis.

Moreover, the results are consistent with those of previous studies linking CQ/HCQ to increased all-cause

mortality [13], especially in the Peruvian context [14].

Except for some case reports [22–26], the risk of CQ/HCQ-induced acute hemolytic anemia in G6PD-

deficient patients is generally considered low according to a systematic review of the available evidence

[21]. This study, however, suggests that this risk may have been underestimated in the specific context of

Peru during the COVID-19 crisis, which was characterized by panic over an unknown virus, nontransparent

treatment protocols [55], misuse of medications [56, 57], and an overburdened healthcare system. Under

such circumstances, uncontrolled outpatient use of CQ and HCQ may have exposed G6PD-deficient

individuals to increased hemolytic risk, potentially contributing to the observed male–female mortality

gap. This hypothesis warrants further investigation using individual-level data linking medication use,

hematologic indicators, and G6PD status.

The Peruvian prison system may be a suitable sample for future research. From an ecological perspective,

a temporal association between the CQ/HCQ distribution and male mortality was also observed there.

Until October 2020, 445 Peruvian prison inmates have died from COVID-19, 251 (56%) of whom died

between April and May [58], which correlates with the delivery of 10,800 tablets of 400 mg HCQ in mid-

April [36].

There are several limitations to this study that are worth mentioning. First, the model is ecological in

nature and cannot establish individual-level causality. Second, the state-level measure of G6PDd

prevalence [28] is modeled and relatively coarse, which may blunt interaction effects. Third, the measure

of CQ/HCQ availability is based on stock data per population and does not capture actual intake,

compliance, informal distribution, black-market flows, or tablets sold via the private health sector. Fourth,

the z-score of the all-cause ASMR serves only as an indirect indicator of healthcare system strain and does

not reflect ICU capacity, staffing shortages, etc. for which no data is available. Finally, the all-cause excess

ASMR were predicted by the model by Karlinsky and Kobak [12] and were used directly in the TWFE

analysis without formally accounting for estimation uncertainty, which may have resulted in

underestimated standard errors.

Conclusion

This study indicates that Peru's nationwide CQ/HCQ treatment policy during the early COVID-19 pandemic

may have contributed to sex-specific differences in mortality among adults aged 45+, particularly in

regions with higher glucose-6-phosphate dehydrogenase deficiency (G6PDd) prevalence and during

periods of elevated healthcare system strain. Across states, both male all-cause excess and COVID-19

mortality consistently exceeded female mortality, with the largest male–female differences coinciding

with periods of plausible widespread CQ/HCQ use among COVID-19 outpatients. Interaction analyses

indicate that regions with higher G6PDd prevalence, greater CQ/HCQ availability, and elevated all-age

excess mortality (proxying healthcare system strain) experienced amplified male-female mortality gaps.

These findings are consistent with the hypothesis that CQ/HCQ-induced oxidative stress may have

triggered hemolytic crises in (undiagnosed) G6PDd patients. In the context of COVID-19, where infection

itself causes oxidative stress and potentially hemolysis [27] and presents with overlapping symptoms (e.g.,

dyspnea, fever), such adverse reactions may have been overlooked or misclassified as severe COVID-19 in

overstretched healthcare systems.

While causality cannot be established owing to the ecological study design, these findings underscore the

importance of incorporating genetic and sex-specific risk factors into pandemic treatment guidelines,

particularly in resource-limited settings with high uncertainty. The oxidative stress induced by COVID-19

infection represents a distinct clinical context, in which drugs previously considered safe — such as

CQ/HCQ for widespread malaria treatment — may pose unforeseen risks for G6PD-deficient individuals.

Given the likelihood that hemolysis may have been unrecognized, ecological analyses remain crucial for

detecting population-level associations and generating hypotheses for future research and policy-making.

Declarations

Ethics approval and consent to participate

371 Not applicable

Consent for publication

373 Not applicable

Availability of data and materials

375 The datasets used and/or analysed during the current study are available from the corresponding author

376 on reasonable request.

Competing interests

378 The authors declare that they have no competing interests.

Funding

380 This research received no specific grant from any funding agency in the public, commercial, or not-for-

381 profit sectors.

Authors' contributions

MK conceived the study, collected and analyzed the data, interpreted the results, and wrote the manuscript. The author read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Peretz C, Rotem N, Keinan-Boker L, Furshpan A, Green M, Bitan M, Steinberg DM. Excess mortality in Israel associated with COVID-19 in 2020-2021 by age group and with estimates based on daily mortality patterns in 2000-2019. Int J Epidemiol. 2022;51:727–36. doi:10.1093/ije/dyac047.

2. Henry NJ, Elagali A, Nguyen M, Chipeta MG, Moore CE. Variation in excess all-cause mortality by age, sex, and province during the first wave of the COVID-19 pandemic in Italy. Sci Rep. 2022;12:1077. doi:10.1038/s41598-022-04993-7.

3. Nielsen J, Nørgaard SK, Lanzieri G, Vestergaard LS, Moelbak K. Sex-differences in COVID-19 associated excess mortality is not exceptional for the COVID-19 pandemic. Sci Rep. 2021;11:20815. doi:10.1038/s41598-021-00213-w.

4. Bambra C, Albani V, Franklin P. COVID-19 and the gender health paradox. Scand J Public Health. 2021;49:17–26. doi:10.1177/1403494820975604.

5. Peckham H, Gruijter NM de, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. Nat Commun. 2020;11:6317. doi:10.1038/s41467-020-19741-6.

6. Pothisiri W, Prasitsiriphon O, Apakupakul J, Ploddi K. Gender differences in estimated excess mortality during the COVID-19 pandemic in Thailand. BMC Public Health. 2023;23:1900. doi:10.1186/s12889-023-16828-9.

7. Akter S. The Gender Gap in COVID-19 Mortality in the United States. Feminist Economics. 2021;27:30–47. doi:10.1080/13545701.2020.1829673.

8. World Health Organization. WHO discontinues hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-19. 2020. https://www.who.int/news/item/04-07-2020-who-discontinues-hydroxychloroquine-and-lopinavir-ritonavir-treatment-arms-for-covid-19. Accessed 4 Aug 2025.

9. Ministerio de Salud (MINSA). Resolución Ministerial N.° 139-2020-MINSA; 2020.

10. Ministerio de Salud (MINSA). Resolución Ministerial N.° 193-2020-MINSA; 2020.

11. Gestión. Minsa anuncia que la hidroxicloroquina se retirará de la guía de medicamentos para el
COVID-19. 2020. https://gestion.pe/peru/coronavirus-peru-pilar-mazzetti-anuncia-que-la-
hidroxicloroquina-se-retirara-de-la-guia-de-medicamentos-para-tratamiento-del-covid-19-nndc-
noticia/. Accessed 4 Sep 2025.

12. Karlinsky A, Kobak D. Tracking excess mortality across countries during the COVID-19 pandemic with
the World Mortality Dataset. Elife 2021. doi:10.7554/eLife.69336.

13. Axfors C, Schmitt AM, Janiaud P, Van't Hooft J, Abd-Elsalam S, Abdo EF, et al. Mortality outcomes
with hydroxychloroquine and chloroquine in COVID-19 from an international collaborative meta-
analysis of randomized trials. Nat Commun. 2021;12:2349. doi:10.1038/s41467-021-22446-z.

14. Soto-Becerra P, Culquichicón C, Hurtado-Roca Y, Araujo-Castillo RV. Real-world effectiveness of
hydroxychloroquine, azithromycin, and ivermectin among hospitalized COVID-19 patients: results of
a target trial emulation using observational data from a nationwide healthcare system in Peru; 2020.

15. Mokhtari M, Mohraz M, Gouya MM, Namdari Tabar H, Tabrizi J-S, Tayeri K, et al. Clinical outcomes
of patients with mild COVID-19 following treatment with hydroxychloroquine in an outpatient
setting. Int Immunopharmacol. 2021;96:107636. doi:10.1016/j.intimp.2021.107636.

16. Lofgren SM, Nicol MR, Bangdiwala AS, Pastick KA, Okafor EC, Skipper CP, et al. Safety of
Hydroxychloroquine Among Outpatient Clinical Trial Participants for COVID-19. Open Forum Infect
Dis. 2020;7:ofaa500. doi:10.1093/ofid/ofaa500.

17. Lagier J-C, Million M, Gautret P, Colson P, Cortaredona S, Giraud-Gatineau A, et al. Outcomes of
3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in
Marseille, France: A retrospective analysis. Travel Med Infect Dis. 2020;36:101791.
doi:10.1016/j.tmaid.2020.101791.

18. Million M, Lagier J-C, Gautret P, Colson P, Fournier P-E, Amrane S, et al. Early treatment of COVID-19
patients with hydroxychloroquine and azithromycin: A retrospective analysis of 1061 cases in
Marseille, France. Travel Med Infect Dis. 2020;35:101738. doi:10.1016/j.tmaid.2020.101738.

19. Mohana A, Sulaiman T, Mahmoud N, Hassanein M, Alfaifi A, Alenazi E, et al. Hydroxychloroquine
Safety Outcome within Approved Therapeutic Protocol for COVID-19 Outpatients in Saudi Arabia. Int
J Infect Dis. 2021;102:110–4. doi:10.1016/j.ijid.2020.10.031.

20. Luzzatto L, Nannelli C, Notaro R. Glucose-6-Phosphate Dehydrogenase Deficiency. Hematol Oncol
Clin North Am. 2016;30:373–93. doi:10.1016/j.hoc.2015.11.006.

21. Kane M. Hydroxychloroquine Therapy and G6PD Genotype. 2023.
https://www.ncbi.nlm.nih.gov/books/NBK591356/. Accessed 4 Sep 2025.

22. Aguilar J, Averbukh Y. Hemolytic Anemia in a Glucose-6-Phosphate Dehydrogenase-Deficient Patient
Receiving Hydroxychloroquine for COVID-19: A Case Report. Perm J 2020. doi:10.7812/TPP/20.158.

23. Sgherza N, Dalfino L, Palma A, Vitucci A, Campanale D, Grasso S, Musto P. "Hemolysis, or not
Hemolysis, that is the Question". Use of Hydroxychloroquine in a Patient with COVID-19 Infection

and G6PD Deficiency. Mediterr J Hematol Infect Dis. 2020;12:e2020076.
doi:10.4084/MJHID.2020.076.

24. Chaney S, Basirat A, McDermott R, Keenan N, Moloney E. COVID-19 and hydroxychloroquine side-
effects: glucose 6-phosphate dehydrogenase deficiency (G6PD) and acute haemolytic anaemia. QJM.
2020;113:890–1. doi:10.1093/qjmed/hcaa267.

25. Mastroianni F, Colombie V, Claes G, Gilles A, Vandergheynst F, Place S. Hydroxychloroquine in a
G6PD-Deficient Patient with COVID-19 Complicated by Haemolytic Anaemia: Culprit or Innocent
Bystander? Eur J Case Rep Intern Med. 2020;7:1875. doi:10.12890/2020_001875.

26. Maillart E, Leemans S, van Noten H, Vandergraesen T, Mahadeb B, Salaouatchi MT, et al. A case
report of serious haemolysis in a glucose-6-phosphate dehydrogenase-deficient COVID-19 patient
receiving hydroxychloroquine. Infect Dis (Lond). 2020;52:659–61.
doi:10.1080/23744235.2020.1774644.

27. Ayyad M, Abu Alya W, Barabrah AM, Darawish SM, AlHabil Y, MohammedAli M, et al. Autoimmune
hemolytic anemia in COVID-19 patients: A systematic review of 105 cases on clinical characteristics
and outcomes. Clin Immunol. 2025;277:110512. doi:10.1016/j.clim.2025.110512.

28. Monteiro WM, Val FFA, Siqueira AM, Franca GP, Sampaio VS, Melo GC, et al. G6PD deficiency in
Latin America: systematic review on prevalence and variants. Mem Inst Oswaldo Cruz.
2014;109:553–68. doi:10.1590/0074-0276140123.

29. Whittaker C, Walker PGT, Alhaffar M, Hamlet A, Djaafara BA, Ghani A, et al. Under-reporting of
deaths limits our understanding of true burden of covid-19. BMJ. 2021;375:n2239.
doi:10.1136/bmj.n2239.

30. Andina Peruvian News Agency. Covid-19: segundo informe para actualizar cifra de fallecidos se
conocerá esta semana. 26.07.2020. https://andina.pe/agencia/noticia-covid19-segundo-informe-
para-actualizar-cifra-fallecidos-se-conocera-esta-semana-807333.aspx. Accessed 4 Sep 2025.

31. Sempé L, Lloyd-Sherlock P, Martínez R, Ebrahim S, McKee M, Acosta E. Estimation of all-cause excess
mortality by age-specific mortality patterns for countries with incomplete vital statistics: a
population-based study of the case of Peru during the first wave of the COVID-19 pandemic. Lancet
Reg Health Am. 2021;2:None. doi:10.1016/j.lana.2021.100039.

32. Kupek E. How many more? Under-reporting of the COVID-19 deaths in Brazil in 2020. Trop Med Int
Health. 2021;26:1019–28. doi:10.1111/tmi.13628.

33. Ministerio de Salud (MINSA). SINADEF - Defunciones Registradas. 2025.
https://www.minsa.gob.pe/reunis/index.asp?op=1&niv=1&tbl=1#. Accessed 14 Aug 2025.

34. Castanheira HC, Da Monteiro Silva JHC. Examining sex differences in the completeness of Peruvian
CRVS data and adult mortality estimates. Genus. 2022;78:3. doi:10.1186/s41118-021-00151-5.

35. Ministerio de Salud (MINSA). Fallecidos por COVID-19. 2025.
https://www.datosabiertos.gob.pe/dataset/fallecidos-por-covid-19-ministerio-de-salud-minsa.
Accessed 14 Aug 2025.

36. Ministerio de Salud (MINSA). Centro Nacional de Abastecimiento de Recursos Estratégicos en Salud
(CENARES) - Distribucion por Producto. 2024.
https://intranet.cenares.gob.pe/cenares/abastecimiento/DISTRIBUCION/wf_distribucion_por_produ
cto.aspx. Accessed 5 Dec 2024.

37. Instituto Nacional de Salud (INS). Molecular Tests COVID-19: NETLAB. 2025.
https://www.datosabiertos.gob.pe/dataset/dataset-de-pruebas-moleculares-del-instituto-nacional-
de-salud-para-covid-19-ins. Accessed 11 Mar 2025.

38. Ministerio de Salud (MINSA). Non-Molecular Tests COVID-19: SISCOVID F100. 2025.
https://www.datosabiertos.gob.pe/dataset/siscovid-f100-pruebas/resource/7f897963-f4c0-4a68-
ae2f-10d49369edf9. Accessed 7 Feb 2025.

39. Ministerio de Salud (MINSA). Vacunación COVID19. 2025.
https://www.datosabiertos.gob.pe/dataset/vacunacion/resource/c2296574-f1ca-46a4-9d1d-
1aee320678a7. Accessed 20 Jun 2025.

40. Ministerio de Salud (MINSA). Población INEI 2021. 2025.
https://datosabiertos.gob.pe/dataset/poblaci%C3%B3n-peru/resource/6e31247b-8057-450c-873d-
d1ba28cf136a. Accessed 3 Aug 2025.

41. Ministerio de Salud (MINSA). Resolución Ministerial N.° 270-2020-MINSA; 2020.

42. Ministerio de Salud (MINSA). Resolución Ministerial N.° 375-2020-MINSA; 2020.

43. RCR Peru. CORONAVIRUS EN PIURA: EL 100% DE PACIENTES CON SÍNTOMAS LEVES SE HAN
RECUPERADO CON IVERMECTINA. 2020. https://www.rcrperu.com/coronavirus-en-piura-el-100-de-
pacientes-con-sintomas-leves-se-han-recuperado-con-ivermectina/. Accessed 4 Aug 2025.

44. Chamie JJ, Hibberd JA, Scheim DE. COVID-19 Excess Deaths in Peru's 25 States in 2020: Nationwide
Trends, Confounding Factors, and Correlations With the Extent of Ivermectin Treatment by State.
Cureus. 2023;15:e43168. doi:10.7759/cureus.43168.

45. Soriano-Moreno DR, Fernandez-Guzman D, Sangster-Carrasco L, Quispe-Vicuña C, Grados-Espinoza
P, Ccami-Bernal F, et al. Factors Associated With Drug Consumption Without Scientific Evidence in
Patients With Mild COVID-19 in Peru. J Patient Saf. 2022;18:e1189-e1195.
doi:10.1097/PTS.0000000000001053.

46. Quispe-Cañari JF, Fidel-Rosales E, Manrique D, Mascaró-Zan J, Huamán-Castillón KM, Chamorro-
Espinoza SE, et al. Self-medication practices during the COVID-19 pandemic among the adult
population in Peru: A cross-sectional survey. Saudi Pharm J. 2021;29:1–11.
doi:10.1016/j.jsps.2020.12.001.

47. Castro NE, García DR, Rivera MT, Rondán-Guerrero P, García-Rojas F, Taype-Rondan A. Tendencias
en el uso de fármacos para la COVID-19 durante la primera ola de la pandemia en un hospital de
Lima, Perú. [Trends in drugs usage for COVID-19 during the first wave of the pandemic in a hospital
in Lima, Peru]. Rev Peru Med Exp Salud Publica. 2021;38:608–14.
doi:10.17843/rpmesp.2021.384.8820.

48. Casas Bueno, Guerrero Martínez. Medicamentos buscados en la web para la COVID-19 en Perú:
Estudio infodemiológico [Tesis (Médico Cirujano)]. Lima, Perú: Universidad Peruana Cayetano
Heredia; 2021.

49. Hoechle D. Robust Standard Errors for Panel Regressions with Cross-Sectional Dependence. The
Stata Journal: Promoting communications on statistics and Stata. 2007;7:281–312.
doi:10.1177/1536867X0700700301.

50. Fischer A, Michuda A. wildboottest (Python module); 2022.

51. Angulo-Bazan Y, Solis G, Cardenas F, Jorge A, Acosta J, Cabezas C. Household transmission in people
infected with SARS-CoV-2 (COVID-19) in Metropolitan Lima; 2020.

52. Huamaní C, Velásquez L, Montes S, Mayanga-Herrera A, Bernabé-Ortiz A. SARS-CoV-2
seroprevalence in a high-altitude setting in Peru: adult population-based cross-sectional study.
PeerJ. 2021;9:e12149. doi:10.7717/peerj.12149.

53. Moreira-Soto A, Pachamora Diaz JM, González-Auza L, Merino Merino XJ, Schwalb A, Drosten C, et
al. High SARS-CoV-2 seroprevalence in rural Peru, 2021; a cross-sectional population-based study;
2021.

54. Moyano LM, Toledo AK, Chirinos J, Vilchez Barreto PMQ, Cavalcanti S, Gamboa R, et al. SARS-CoV-2
seroprevalence on the north coast of Peru: A cross-sectional study after the first wave. PLoS Negl
Trop Dis. 2023;17:e0010794. doi:10.1371/journal.pntd.0010794.

55. Taype-Rondan A, Herrera-Añazco P, Málaga G. Sobre la escasa transparencia en los documentos
técnicos para el tratamiento de pacientes con COVID-19 en Perú. Acta Med Peru 2020.
doi:10.35663/amp.2020.372.982.

56. Caira-Chuquineyra B, Fernandez-Guzman D, Alvarez-Arias PM, Zarate-Curi ÁA, Herrera-Añazco P,
Benites-Zapata VA. Association between prehospital medication and fatal outcomes in a cohort of
hospitalized patients due to coronavirus disease-2019 in a referral hospital in Peru. Travel Med
Infect Dis. 2022;50:102472. doi:10.1016/j.tmaid.2022.102472.

57. Martinez-Rivera RN, Taype-Rondan A. Overmedication in COVID-19 Context: A Report From Peru. J
Clin Pharmacol. 2020;60:1155–6. doi:10.1002/jcph.1704.

58. Sindeev A, Martínez-Álvarez BM. Clinical and epidemiological characteristics of prisoners infected
and deceased by COVID-19, National Penitentiary Institute of Peru, 2020. Rev Esp Sanid Penit.
2022;24:15–22. doi:10.18176/resp.00045.

Appendix

Figure I: Sex differences in all-cause excess ASMR (males-females) for individuals aged 45+ over time by state

556   *Note: The variance of the sex difference was obtained as the sum of male and female ASMR variances (see the* Statistical Analysis

557   section*), and the 95% confidence interval (95% CI) was constructed as the point estimate ± 1.96 times the square root of this*

558   *variance, assuming the independence of errors (Cov(m, f) = 0).*

559 *Figure II: Sex differences in COVID-19 ASMR (males-females) for individuals aged 45+ over time by state*

    Table I: Sum of delivered HCQ tablets per state population

State	HCQ_per_pop			
	period_HCQ z-score = 4	period_HCQ z-score = 3.5	period_HCQ z-score = 3	period_HCQ z-score = 2.5
Amazonas	39.62	39.62	39.62	39.62
Ancash	14.74	14.74	14.74	14.74
Apurimac	-	-	20.86	20.86
Arequipa	19.47	19.47	19.47	19.47
Ayacucho	-	-	27.35	27.35
Cajamarca	8.33	8.33	8.33	8.33
Cusco	10.30	10.30	10.30	10.30
Huancavelica	11.65	11.65	11.65	11.65
Huánuco	26.92	26.92	26.92	26.92
Ica	36.48	36.48	36.48	36.48
Junín	15.32	15.32	15.32	15.32
La Libertad	17.16	17.16	17.16	17.16
Lambayeque	16.36	21.55	21.55	21.55
Lima + Callao	18.69	18.69	18.69	18.69
Loreto	18.64	18.64	20.61	21.81
Madre de Dios	178.36	178.36	178.36	178.36
Moquegua	35.60	35.60	35.60	35.60
Pasco	25.17	25.17	25.17	25.17
Piura	5.51	5.51	5.51	5.51
Puno	7.65	7.65	7.65	7.65
San Martin	36.37	36.37	36.37	36.37

Tacna	19.75	19.75	19.75	19.75
Tumbes	12.29	12.29	12.29	12.29
Ucayali	20.57	20.57	20.57	39.68
Median	18.66	19.08	20.16	20.31
Mean	26.68	26.92	26.82	27.64
Std. Dev.	33.68	33.62	32.25	32.30

Note: The different CQ/HCQ drugs used were 'CLOROQUINA 150 MG TAB', 'CLOROQUINA FOSFATO 250 MG (EQUIV. 155 MG

CLOROQUINA BASE) TAB', 'HIDROXICLOROQUINA [SULFATO] 400 MG TAB', 'HIDROXICLOROQUINA SULFATO 200 MG (EQUIV. 155

MG HIDROXICLOROQUINA) TAB' [36]. According to the treatment protocol, 1 CQ is considered equivalent to 1/6 HCQ [9]. All drugs

were standardized using 1 tablet '155 MG HIDROXICLOROQUINA' as the base. All of these standardized tablets that were

distributed before or during *period_HCQ* were summed and divided by the state population, with *period_HCQ* constructed with

different definitions (z-score thresholds) as described in the Data Preparation section. Missing values stem from the lack of

identification of *period_HCQ* with the given z-score threshold.

Table II: Scarcity CQ/HCQ: Doses per positive test

State	CQ/HCQ-Doses per positive tests in age group 45+				Total tests per population during Period_HCQ (z-score = 2.5)
	period_HCQ z-score = 4	period_HCQ z-score = 3.5	period_HCQ z-score = 3	period_HCQ z-score = 2.5	
Amazonas	1.36	1.21	1.21	1.11	0.19
Ancash	0.46	0.46	0.42	0.41	0.21
Apurimac	NaN	NaN	4.48	2.12	0.13
Arequipa	0.23	0.22	0.22	0.22	0.50
Ayacucho	NaN	NaN	3.30	2.11	0.06
Cajamarca	0.59	0.59	0.57	0.57	0.10

Cusco	0.41	0.41	0.41	0.41	0.14
Huancavelica	1.36	1.09	1.09	1.09	0.09
Huánuco	1.23	1.14	1.14	1.14	0.14
Ica	0.54	0.54	0.54	0.54	0.39
Junín	1.12	1.03	1.03	1.03	0.10
La Libertad	0.51	0.51	0.48	0.48	0.19
Lambayeque	0.94	0.80	0.70	0.59	0.17
Lima + Callao	0.27	0.27	0.27	0.27	0.40
Loreto	1.38	1.38	1.09	0.94	0.08
Madre de Dios	3.66	3.35	3.35	3.05	0.26
Moquegua	0.40	0.40	0.39	0.39	0.48
Pasco	1.80	1.68	1.68	1.68	0.14
Piura	0.64	0.64	0.64	0.64	0.04
Puno	0.56	0.56	0.56	0.56	0.08
San Martin	1.36	1.36	1.29	1.29	0.11
Tacna	0.51	0.51	0.51	0.49	0.26
Tumbes	1.01	1.01	1.01	1.01	0.05
Ucayali	1.80	1.80	1.80	2.17	0.07
Median	0.79	0.72	0.85	0.79	0.14
Mean	1.00	0.94	**1.16**	1.00	0.18
Std. Dev.	0.73	0.67	1.04	0.70	0.13

571   Note: CQ/HCQ deliveries [36] that occurred before or during *period_HCQ* were considered, and 1 CQ/HCQ dose corresponds to

572   18 tablets of HIDROXICLOROQUINA SULFATO 200 MG [41]. All positive COVID-19 tests in the 45+ age group that occurred during

573 *period_HCQ* were counted, with *period_HCQ* constructed with different specifications (z-score thresholds) as described in the

574 *Data Preparation section.*

576 *Table III: Inference with Wild Cluster Bootstrap [50] for Models 1, 2, 3, and 4 (Table 2, main text)*

	p-value (Wild Cluster Bootstrap)			
Model	1 (all-cause excess)	2 (all-cause excess)	3 (COVID-19)	4 (COVID-19)
period_HCQ	0.000	0.001	0.002	0.000
period_HCQ x G6PDD prevalence	-	0.075	-	0.023
period_HCQ x HCQ_per_pop	-	0.046	-	0.003
period_HCQ x z-score	-	0.000	-	0.004
z-score	0.000	0.000	0.000	0.000
Sex difference test positive rate (lag 4)	0.028	0.029	0.003	0.003
Sex difference vaccination rate (lag 3)	0.002	0.051	0.000	0.007

Figure III: Marginal effect of period_HCQ conditional on G6PDd prevalence. Dependent variable: sex difference in all-cause excess

ASMR (males-females) at age 45+

Note: The variable *HCQ_per_pop* (time-invariant) is held at its mean. *The variable z-score is held at its mean during period_HCQ.*

Figure IV: Marginal effect of period_HCQ conditional on G6PDd prevalence. Dependent variable: sex difference in Covid-19 ASMR

(males-females) ages 45+

Note: The variable *HCQ_per_pop* (time-invariant) is held at its mean. *The variable z-score is held at its mean during period_HCQ.*

Figure V: States of Peru and modeled G6PDd prevalence [28]

Table IV: G6PDd prevalence by state, derived from [28]

State	G6PDd prevalence [28]
Amazonas	4.5 (6-10%)
Ancash	2.0 (2-4%)
Apurimac	2.0 (2-4%)
Arequipa	2.0 (2-4%)
Ayacucho	2.0 (2-4%)
Cajamarca	4.0 (6-8%)
Cusco	2.0 (2-4%)
Huancavelica	1.5 (0-4%)
Huanuco	2.0 (2-4%)

Ica	1.5 (0-4%)
Junin	1.5 (0-4%)
La Libertad	3.0 (4-6%)
Lambayeque	4.0 (6-8%)
Lima + Callao	1.0 (0-2%)
Loreto	4.5 (6-10%)
Madre de Dios	3.0 (4-6%)
Moquegua	2.0 (2-4%)
Pasco	1.5 (0-4%)
Piura	5.0 (8-10%)
Puno	3.0 (4-6%)
San Martin	3.5 (4-8%)
Tacna	2.0 (2-4%)
Tumbes	5.0 (8-10%)
Ucayali	3.0 (4-6%)

591 Note: G6PDd prevalence was coded on the basis of Monteiro et al. (2014); values in parentheses indicate the original modeled

592 prevalence [28].