(Hydroxy-) Chloroquine, G6PD-deficiency, and sex differences in mortality during the COVID-19 pandemic: An ecological study in Peru

Marius Krämer[1*]

Abstract

Background: Peru implemented a nationwide chloroquine/hydroxychloroquine (CQ/HCQ) treatment policy during the early phase of the COVID-19 pandemic. It is subject to debate if these drugs trigger hemolytic crises in individuals with glucose-6-phosphate dehydrogenase (G6PD) deficiency, a genetic condition primarily affecting men, with higher prevalence in certain regions. This study investigates whether the CQ/HCQ policy contributed to sex-specific differences in mortality, potentially mediated by regional G6PDD prevalence.

Methods: I conducted an ecological analysis using both all-cause excess and COVID-19 mortality data on the state-level for adults aged ≥45 years. Using sex differences in age-standardized mortality rates (ASMR) as the outcome in a two-way fixed effects (TWFE) model, interaction analyses examined the influence of regional G6PDD prevalence, CQ/HCQ availability, and elevated healthcare system strain during pandemic waves with CQ/HCQ recommendation.

Results: Male all-cause excess and COVID-19 mortality consistently exceeded female mortality, with the largest sex differences coinciding with periods with CQ/HCQ treatment recommendations. TWFE models showed that states with higher G6PDD prevalence, greater CQ/HCQ availability, and elevated healthcare system strain exhibited amplified male-female mortality gaps. These patterns are consistent with CQ/HCQ-induced oxidative stress triggering hemolytic crises in (undiagnosed) G6PD-deficient COVID-19 outpatients, which may have been misclassified as severe COVID-19 during periods of a strained healthcare system.

Conclusion: While causality cannot be established due to the ecological design, these findings suggest that pandemic treatment policies may have unintended, sex-specific consequences if population-specific genetic vulnerabilities are overlooked, particularly in resource-limited settings.

Keywords: Peru, COVID-19, Hydroxychloroquine, Chloroquine, G6PD, Sex difference

---

[1] Technical University of Munich, Germany; TUM School of Social Sciences and Technology

[*] Correspondence: marius.kraemer@tum.de

Introduction

Mortality during the COVID-19 pandemic has exhibited differences between males and females across countries and regions, reflecting both biological vulnerabilities and contextual factors [1, 2]. Some studies show higher mortality among men [3–5], others reveal complex patterns, including periods or locations where women experienced equal or higher mortality [6, 7]. Given the biological plausibility of sex-specific drug risks, treatment policy may represent an overlooked driver of these patterns.

At the start of the pandemic, chloroquine (CQ) and hydroxychloroquine (HCQ) were seen as potential COVID-19 treatments. International trials led by the WHO were discontinued in July 2020 due to safety concerns and lack of efficacy [8]. While most countries remained cautious, Peru adopted CQ/HCQ widely. Initial protocols restricted use to moderate or severe cases [9], but by mid-April 2020, HCQ was authorized for mild outpatient cases [10]. Peru retained HCQ in official guidelines until September 2020, months after international trials had halted its use [8, 11]. Combined with one of the highest all-cause excess mortality rates globally [12], Peru provides a unique context to study the impact of treatment policies.

The safety profile of CQ and especially HCQ in the context of COVID-19 has been debated. In a meta-analysis of 14 trials testing CQ/HCQ in hospitalized patients, it was associated with an 11% (95%CI 2–20%) increase in all-cause mortality [13]. An observational preprint study by the Peruvian medical research institute (IETSI) found that HCQ combined with azithromycin was even associated with an 84% (95%CI 12–202%) increase in all-cause mortality in hospitalized COVID-19 patients compared to standard care [14]. Evidence from outpatient studies generally reported few safety concerns, but these trials consistently excluded individuals with contraindications to CQ/HCQ [15], often explicitly mentioning glucose-6-phosphate dehydrogenase deficiency (G6PDD) as one such exclusion [16–19].

Among factors that may influence CQ/HCQ-related risk, G6PDD is particularly relevant: an X-linked enzymatic disorder that renders hemizygous males especially vulnerable, while females are usually carriers or only partially deficient; G6PDD can predispose individuals to acute hemolytic anemia when exposed to oxidative stress, which may be triggered by infections or certain drugs [20]. While CQ/HCQ use in malaria-endemic regions has not been associated with widespread hemolytic events, according to a systematic review of the available evidence [21], case reports during the COVID-19 pandemic document acute hemolytic anemia in G6PDD patients treated with HCQ [22–26].

Moreover, COVID-19 itself has been reported to be associated with acute autoimmune hemolytic anemia, often presenting with delayed hemolytic symptoms that may be overlooked due to the predominance of respiratory manifestations [27].

The combination of the oxidative stressors COVID-19 and CQ/HCQ may increase the risk of acute hemolytic anemia among G6PDD patients. Particularly during outpatient CQ/HCQ use, hemolysis symptoms may have been delayed or mistaken for COVID-19 symptoms (e.g., dyspnea, fever), complicating detection, especially in an underdeveloped and heavily strained healthcare system. Peruvian treatment protocols with CQ/HCQ did furthermore not include warnings about potential adverse reactions or contraindications for G6PDD patients. Consequently, given that G6PDD disproportionately affects males, the nationwide CQ/HCQ policy could plausibly have sex-specific consequences for mortality.

To my knowledge, no prior study has examined whether national COVID-19 treatment policies contributed to sex-specific differences in all-cause excess or COVID-19 mortality. This ecological study addresses this gap by testing two hypotheses:

1. If CQ/HCQ triggers acute hemolytic anemia in G6PDD COVID-19 patients and Peru's healthcare system failed to manage these reactions, periods of widespread CQ/HCQ use should coincide with increased male-female mortality differences. Within these periods, higher healthcare system strain is expected to further amplify these differences.
2. If regional G6PDD prevalence modifies risk, areas with higher prevalence should show stronger male-female mortality differences during periods of widespread CQ/HCQ use.

By integrating publicly available all-cause and COVID-19 mortality data with treatment policy information and regional G6PDD prevalence estimates [28], this study explores the plausibility of the described mechanism.

Data and Methods

For this analysis, I used Python 3.11, primarily Pandas (v2.2.3) for data handling and Statsmodels (v0.14.1) for statistical analysis.

Data

All datasets are publicly available and were aggregated at the state and week level. Mortality data from 2017–2019 were used to construct baseline mortality, while all other datasets cover 2020–2022. The compiled dataset includes 24 states over 157 iso-weeks.

Mortality data

Estimating the true mortality burden of COVID-19 is challenging, particularly in countries with limited healthcare system capacity and incomplete vital registration [29][29–31]. Consequently, some researchers prefer all-cause excess mortality, defined as the difference between observed and expected deaths, as a more reliable measure of the pandemic's impact [6, 32]. In this study, I use both, all-cause excess and COVID-19 mortality data.

All-cause mortality data were obtained from Peru's National Information System of Deaths (SINADEF) [33]. An assessment concluded that SINADEF represents a substantial improvement over earlier death registries in Peru; although some incompleteness remains, no systematic difference in completeness by sex was observed [34]. In addition, COVID-19 case fatality data were analyzed [35]. These figures are subject to underreporting and have been subsequently adjusted by the authorities [30]. However, this subgroup was more likely affected by the CQ/HCQ treatment policy.

Medicament data

I collected data on CQ/HCQ distributions from the database of the National Center for the Supply of Strategic Health Resources (CENARES) [36]. Since official treatment protocols treated CQ and HCQ as

equivalent, I did not distinguish between them and summed their distributions after conversion of CQ to HCQ-equivalents using protocol dosage ratios [37].

Prevalence of G6PD Deficiency

I used model-based estimates of the prevalence of glucose-6-phosphate dehydrogenase deficiency (G6PDD) form 28 [28], who provide a map of Latin America. I overlaid the Peruvian state boundaries onto their map (Figure V, Appendix) to assign prevalence values to each state (Table IV, Appendix).

COVID-19 test results and vaccination data

I obtained COVID-19 test results from Peruvian authorities, which included both molecular (i.e., PCR) [38] and non-molecular tests (i.e., rapid antibody tests, antigen tests, chemiluminescence assays) [39]. Like official treatment protocols, I treated all types of tests equivalently. Additionally, I obtained COVID-19 vaccination data from Peruvian authorities [40] treating all vaccine types equivalently.

Other Data

Data on population were obtained from official statistics [41]. For reasons of feasibility, all rate calculations are based on the 2021 mid-year population estimates provided by the Ministry of Health of Peru [41], which readily include the necessary breakdowns by sex, age group, and region. Moreover, I consulted official documents containing CQ/HCQ treatment protocols for COVID-19 patients [9, 10, 37, 42].

Methods

Data preparation

Deaths categorized as violent (accidents, homicides, suicides) and rare observations with missing information on sex, age, or location were excluded. All datasets were aggregated to the state-level and to a weekly timeframe. The state of Callao is completely embedded within Lima and was therefore assigned to Lima instead of treating it separately. I stratified the data by the age groups 0–14, 15–44, 45–64, 65–79, 80+, and created the aggregated group 45+ which is the main focus of this analysis, as this group is considered to be most affected by COVID-19 and related treatments.

Coding of COVID-19 treatment protocols

Official COVID-19 treatment protocols varied in their stance on CQ/HCQ [9, 10, 37, 42]. I coded these recommendations in an ordinal variable called *recommendation_HCQ* from 0 (no recommendation), to 1 (recommended only for hospitalized medium and severe cases), 2 (additionally recommended for confirmed mild outpatient cases), and 3 (additionally recommended for suspected mild outpatient cases). Protocol changes were aligned with the weekly dataset by assigning each protocol to all calendar weeks between its issuance and the release of the subsequent update.

Definition of the CQ/HCQ exposure period (period_HCQ)

In line with the hypotheses, I constructed the binary variable *period_HCQ*[2] to indicate weeks when outpatient treatment with CQ/HCQ likely occurred and acute hemolytic anemia may have been mismanaged due to a strained healthcare system.

period_HCQ was set to 1 if (i) the official COVID-19 treatment protocol recommended CQ/HCQ for outpatients, i.e., mild cases (*recommendation_HCQ* = 2 or 3), and (ii) the proxy for healthcare system strain and pandemic waves, i.e., the z-score of all-cause mortality (all ages) exceeded the threshold of 3 for at least two consecutive weeks. To account for delays between treatment and death, I extended the period by two additional weeks after the z-score went below the threshold.

In at least 11 states, however, CQ/HCQ was soon displaced by ivermectin (IVM) for outpatient treatment. In Piura, for instance, the state health minister stated in early June 2020 (cw23) that HCQ for mild cases has been replaced by IVM due to its perceived efficacy and fewer side effects [43]. Moreover, ten states (Ayacucho, Cajamarca, Cusco, Huancavelica, Huánuco, Junín, Moquegua, Pasco, Puno, Tacna) implemented mass IVM distributions through the Ministry of Defense's Mega-Operación Tayta (MOT). During its first phase, MOT conducted household visits to identify vulnerable individuals; one week later, it began distributing IVM broadly, regardless of test status or symptoms, alongside other medications such as acetaminophen and azithromycin [44]. HCQ was not distributed as part of MOT. Similarly, the regional government of Tumbes carried out a locally organized campaign with mass IVM distributions (Vanessa 2020a, 2020b). In these 11 states, I therefore shortened *period_HCQ* by ending it two weeks after the start of mass IVM distribution, to reflect a reasonable transition period.

Since *period_HCQ* is based on assumptions and reasoning, I reconstructed it under alternative definitions for robustness checks, using different z-score thresholds (2.5, 3, 3.5, 4) each both with and without adjustments for IVM campaigns.

State-level CQ/HCQ availability (HCQ_per_pop)

Reliable state-level time series data on CQ/HCQ consumption are unavailable. Instead, I used data on the distribution of CQ/HCQ tablets to regional health authorities (i.e., stock data). For each state, I summed all distributions that occurred before or during *period_HCQ* and created a time-invariant variable, *HCQ_per_pop*[3]. To make states comparable, I standardized the different types of drugs according to treatment protocols and divided the number of these tablets by the state population, expressed per 100,000 people (for details see Table I, Appendix).

In most states, the quantities of distributed CQ/HCQ doses were limited relative to the total number of COVID-19 cases. On average, only 1.16 doses per positive test in the 45+ age group were delivered (Table II, Appendix), decreasing to 0.4 doses when considering positive tests across all ages. Symptomatic but untested cases were also treated [42], indicating relative scarcity in most states, and supporting the correspondence between stock data and actual usage. Moreover, self-medication with CQ/HCQ seemed to be uncommon in Peru [45–47], and web searches indicate limited public interest [48]. Therefore,

---

[2] For simplicity, I use here 'HCQ' to refer to both chloroquine and hydroxychloroquine.
[3] For simplicity, I use 'HCQ' to refer to both chloroquine and hydroxychloroquine.

HCQ_per_pop is unlikely to be strongly confounded by privately obtained doses. Although this measure does not capture actual intake, nor does it distinguish between inpatient and outpatient administration, *HCQ_per_pop* is expected to reflect inter-state differences in CQ/HCQ availability.

Statistical Analysis

Estimating all-cause excess mortality rates

All-cause excess mortality is defined as the difference between the expected (baseline) mortality and the observed mortality. I estimated weekly absolute numbers of expected deaths (baselines) for 2020–22 for each state, sex, and age group based on historical data from 2017–19, using the time-series regression model by 12 [12]. Absolute numbers of all-cause excess deaths were then calculated by subtracting the baseline from observed deaths, and crude mortality rates for each group were derived by dividing excess deaths by the corresponding population and scaling per 100,000 individuals. For feasibility and consistency, all rate calculations for 2020-22 used the 2021 mid-year population estimates provided by the Ministry of Health of Peru [41], which already include necessary breakdowns by sex, age group, and locality. This approach may not yield the most precise estimates of all-cause excess mortality rates. However, more sophisticated methods would likely provide little additional insight, as only three years of training data (2017-19) are available.

Age-standardized mortality rates (ASMR)

The confounding effect of different age structures makes the comparison between sexes for aggregated age groups misleading. Therefore, for the age group 45+, I calculated the age standardized mortality rates (ASMR) for males and females in each state, using the direct method with Peru's age structure as the standard population. The sex difference was then calculated as the difference of ASMR for males and females. This was done separately for both, all-cause excess and COVID-19 mortality rates.

To quantify the statistical uncertainty of the estimated all-cause excess ASMRs, I calculated 95% CIs using predictive standard errors from the baseline mortality regressions. For each sex, the derived variances were aggregated across age groups using the squared standard-population weights. The confidence interval was constructed as the point estimate ± 1.96 times the square root of this variance.

Model specification

The sex difference in all-cause excess ASMR, as well as in COVID-19 ASMR, for the 45+ age group, is used as the dependent variable in two-way fixed-effects (TWFE) models, specified as follows:

$$
\begin{aligned}
ASMR\_male_{i,t} &- ASMR\_female_{i,t} \\
&= \gamma \, period\_HCQ_{i,t} + \theta \, (period\_HCQ_{i,t} \times G6PDD_i) \\
&+ \phi \, (period\_HCQ_{i,t} \times HCQ\_per\_pop_i) + \psi \, (period\_HCQ_{i,t} \times z - score_{i,t}) + \beta \, X_{i,t} \\
&+ \eta_i + \delta_t + \varepsilon_{i,t}
\end{aligned}
$$

where $period\_HCQ_{i,t}$ indicates the weeks with likely outpatient mass treatment with CQ/HCQ during a strained healthcare system in state i at time t, $(period\_HCQ_{i,t} \times G6PDD_i)$, $(period\_HCQ_{i,t} \times HCQ\_per\_pop_i)$, and $(period\_HCQ_{i,t} \times z - score_{i,t})$ are interaction terms capturing the conditional effects of G6PDD prevalence, QC/HCQ availability, and healthcare system strain – as measured by the z-

score based on all-cause mortality for both sexes and all age groups – during *period_HCQ*, reflecting the hypotheses. $X_{i,t}$ is a vector of control variables, η_i and δ_t are state- and week-specific fixed effects, respectively, and $\varepsilon_{i,t}$ is the error term[4]. I used Driscoll-Kraay standard errors to account for cross-sectional dependence and serial correlation [49] and used wild cluster bootstrap inference [50] as a robustness check, given the small number of states.

The TWFE setup addresses several sources of confounding. First, time-invariant differences across states not included in the interaction terms are absorbed by the state fixed effects. Second, week fixed effects account for shocks common to all states at a given week. Third, using the sex difference in ASMR as the dependent variable implicitly removes factors that influence men and women in a similar way.

Control variables include the z-score, age-standardized sex differences in COVID-19 test positive rates, and cumulative sex differences in first-dose vaccination rates for ages 45+. Analogous to age-standardization of mortality rates, test positivity rates were standardized using the national age distribution of tests to account for differences in testing by age. Cumulative vaccination rate sex differences were calculated as the sum over time of male first-dose rates minus female rates. The z-score control variable estimates the 'baseline' relationship between sex differences in all-cause excess mortality and overall excess mortality independent of *period_HCQ*, while the z-score interaction term captures any additional effect during *period_HCQ*, reflecting the hypothesis that higher healthcare system strain could have further amplified sex differences in mortality by increasing the likelihood of adverse effects remaining unrecognized.

Because testing capacity in Peru was limited, particularly during the first wave, reported COVID-19 test results are an imperfect proxy for true infections. Consequently, observed sex differences in test positivity may not fully capture actual infection patterns. Nonetheless, several considerations suggest that the resulting bias is limited. Nationwide lockdowns and contact restrictions were not gender-specific and are absorbed by week fixed effects, leaving only state-level reopening policies as a possible source of differential exposure. Moreover, household transmission was high – a study in Lima reported a secondary attack rate of 53% [51] – which likely reduced sustained disparities between sexes. Finally, Peruvian seroprevalence studies consistently show no significant sex differences in infection levels [52–54]. Overall, while short-term gender-specific fluctuations in infections may be partially captured by the control variable, persistent sex-specific differences in infection risk were small and unlikely to explain the observed sex difference in all-cause excess mortality.

Results

Sex Differences during pandemic waves

Except for Apurímac, all Peruvian states recorded higher male than female ASMR in the age group 45+ during pandemic waves (i.e., periods ≥ 2 consecutive weeks with z-score ≥ 3), leading to predominantly positive average sex differences in all-cause excess ASMR, as well as COVID-19 ASMR (Table 1). In most states, these differences were larger during the first pandemic wave (when CQ/HCQ was recommended) than in subsequent waves, with several cases showing confidence intervals that (nearly) exclude zero, or

---

[4] For simplicity, I use 'HCQ' in the variable names to refer to both chloroquine and hydroxychloroquine.

p-values < 0.05, respectively. At the same time, the size of the observed differences varied considerably, ranging from negligible or negative (Apurímac, Pasco) to very large (e.g., Madre de Dios, Moquegua, Ucayali).

Table 1: Average sex difference (male minus female) in all-cause excess and COVID-19 ASMR for ages 45+ during pandemic waves

State	Weeks Wave 1	Weeks Waves 2+	Avg. all-cause excess sex diff. Wave 1 [95% CI]	Avg. all-cause excess sex diff. Waves 2+ [95% CI]	Diff. all-cause excess (W1–W2+) [95% CI]	COVID-19 Avg. Sex diff. Wave 1	COVID-19 Avg. Sex diff. Waves 2+	Diff. COVID-19 (W1–W2+) (p-value)
Amazonas	8	10	41.9 [18.3, 65.4]	17.3 [-6.3, 40.8]	24.6 [-8.7, 57.9]	34.4	19.2	15.2 (0.040)
Ancash	17	21	28.1 [12.6, 43.6]	19.0 [3.5, 34.5]	9.1 [-12.8, 31.1]	24.2	18.4	5.8 (0.037)
Apurimac	4	31	-6.5 [-31.7, 18.8]	2.9 [-22.3, 28.2]	-9.4 [-45.1, 26.4]	-1.1	5.4	-6.6 (0.367)
Arequipa	14	22	49.8 [36.0, 63.7]	16.7 [2.8, 30.5]	33.2 [13.6, 52.7]	3.3	14.6	-11.3 (0.078)
Ayacucho	2	8	54.0 [28.4, 79.6]	3.8 [-21.8, 29.4]	50.2 [14.0, 86.4]	27.2	8.9	18.3 (0.043)
Cajamarca	10	20	24.8 [12.1, 37.5]	11.9 [-0.9, 24.6]	12.9 [-5.1, 30.9]	19.8	12.3	7.5 (0.058)
Cusco	6	18	33.4 [15.6, 51.2]	10.8 [-6.9, 28.6]	22.6 [-2.5, 47.7]	7.4	14.7	-7.4 (0.053)
Huancavelica	6	18	30.1 [-8.5, 68.8]	6.1 [-32.5, 44.8]	24.0 [-30.7, 78.6]	19.8	7.8	12.0 (0.030)
Huanuco	12	35	12.8 [-2.9, 28.6]	8.0 [-7.8, 23.7]	4.8 [-17.4, 27.1]	11.3	11.4	-0.1 (0.966)
Ica	23	30	36.6 [18.9, 54.4]	21.3 [3.6, 39.1]	15.3 [-9.8, 40.4]	23.5	19.4	4.1 (0.337)
Junin	11	23	32.4 [16.4, 48.4]	25.7 [9.6, 41.7]	6.7 [-15.9, 29.4]	17.6	24.6	-7.0 (0.152)
La Libertad	18	25	30.2 [18.4, 42.1]	9.9 [-1.9, 21.7]	20.3 [3.6, 37.0]	20.7	11.3	9.5 (0.010)
Lambayeque	18	33	32.0 [11.6, 52.3]	9.6 [-10.8, 29.9]	22.4 [-6.4, 51.2]	23.2	12.3	10.9 (0.004)
Lima + Callao	27	31	28.7 [22.3, 35.0]	12.9 [6.5, 19.2]	15.8 [6.8, 24.8]	20.7	16.5	4.2 (0.201)
Loreto	14	15	58.8 [43.2, 74.4]	18.2 [2.5, 33.8]	40.6 [18.5, 62.7]	63.0	15.0	48.0 (0.001)
Madre de Dios	10	7	196.4 [131.4, 261.4]	100.3 [35.3, 165.3]	96.1 [4.2, 188.0]	194.8	101.0	93.7 (0.012)
Moquegua	10	12	119.3 [88.0, 150.5]	19.4 [-11.9, 50.6]	99.9 [55.7, 144.1]	122.4	29.8	92.6 (0.000)
Pasco	9	22	15.1 [-12.3, 42.6]	19.9 [-7.6, 47.3]	-4.7 [-43.5, 34.1]	11.0	11.9	-0.9 (0.889)
Piura	12	10	52.4 [34.6, 70.2]	13.4 [-4.4, 31.3]	39.0 [13.8, 64.2]	45.8	20.6	25.2 (0.002)
Puno	8	28	53.3 [39.9, 66.7]	18.8 [5.4, 32.2]	34.4 [15.5, 53.4]	21.0	15.1	5.8 (0.084)
San Martin	12	16	41.8 [21.3, 62.2]	18.1 [-2.3, 38.6]	23.6 [-5.3, 52.5]	40.4	18.3	22.1 (0.002)
Tacna	10	25	71.0 [42.7, 99.4]	35.1 [6.8, 63.5]	35.9 [-4.2, 76.0]	57.2	27.9	29.2 (0.012)
Tumbes	11	17	58.1 [22.2, 94.1]	36.8 [0.8, 72.7]	21.4 [-29.4, 72.2]	54.7	42.0	12.7 (0.238)
Ucayali	6.0	9	110.0 [85.2, 134.7]	39.1 [14.3, 63.8]	70.9 [35.9, 105.9]	95.3	50.7	44.6 (0.017)
Median	11.0	18.6	41.8	17.8	23.6	39.9	22.0	17.8
Mean	11.6	18.6	50.2	21.3	28.9	23.5	16.5	10.9
Std. Dev.	5.6	6.3	39.9	18.5	26.4	42.1	19.0	26.5

Note: Average sex differences in deaths per 100,000. Waves are defined as periods of ≥2 consecutive weeks with a z-score ≥3. 95% confidence intervals (95% CI) are based on the predictive variance of baseline models. P-values are from two-sided permutation tests.

Figure 1 shows the male and female all-cause excess ASMR for ages 45+ over time, with markedly higher values for males, especially during the 2020 wave, and a close correspondence with the all-ages mortality z-score. There is no period where female all-cause excess mortality exceeded male mortality significantly, i.e., a negative sex difference and regional heterogeneity is evident. Periods where *period_HCQ* = 1 (i.e., weeks of substantial healthcare system strain (z-score ≥ 3) combined with CQ/HCQ recommendation for COVID-19 outpatients) are also indicated. Comparable figures for all-cause sex differences and for COVID-19 ASMR are provided in the Appendix (Figure I, Figure II, Appendix).

Figure 1: Male and female all-cause excess ASMR for ages 45+ over time by state

Note: 95%CI stands for 95% confidence interval.

TWFE model results

Table 2: Results TWFE models

Model	1 (all-cause excess)	2 (all-cause excess)	3 (COVID-19)	4 (COVID-19)
R² (Within)	0.3863	0.4706	0.4606	0.6115
F-statistic	224.37 (p = 0.000)	241.69 (p = 0.000)	328.76 (p = 0.000)	492.17 (p = 0.000)
Constant	2.771*** (0.813)	4.219*** (0.492)	2.094*** (0.741)	3.324*** (0.408)
period_HCQ	22.049*** (5.124)	-41.808*** (5.077)	15.689*** (4.424)	-47.428*** (5.690)
period_HCQ x G6PDD prevalence	-	8.209*** (1.845)	-	8.232*** (1.761)
period_HCQ x HCQ_per_pop	-	0.770*** (0.177)	-	0.823*** (0.167)
period_HCQ x z-score	-	3.050*** (0.671)	-	2.653*** (0.624)
z-score	2.611*** (0.529)	1.529*** (0.279)	2.486*** (0.441)	1.571*** (0.241)
Sex difference test positive rate (lag 4)	34.474*** (11.454)	31.550*** (10.166)	31.927*** (8.870)	28.668*** (8.717)
Sex difference vaccination rate (lag 3)	-27.394** (11.570)	-9.402 (7.082)	-34.252*** (12.506)	-15.584** (6.881)

Note: Dependent variable: all-cause excess or COVID-19 sex differences in ASMR (ASMR_m − ASMR_f) for ages 45+. z-score is based on all-cause excess mortality in all models. Observations: 3454 (22 states, 157 weeks). Driscoll-Kraay standard errors in parentheses (see Table III, Appendix for p-values from wild cluster bootstrap inference). Fixed Effects: entity, time. States Apurímac and Ayacucho were excluded because no clear period_HCQ > 4 weeks was identified.

I present two models for each of the two outcomes (all-cause excess, COVID-19) to separate main and interaction effects. In models 2 and 4 (Table 2), the main effect of *period_HCQ* should be interpreted with caution, since it is not only identified at the unrealistic baseline values of the interactions (all zero), but is also inflated by multicollinearity with the interaction terms. Models 1 and 3 (Table 2) show a positive and significant coefficient for *period_HCQ* for both outcome variants (22.049, $p \leq 0.001$; 15.689, $p \leq 0.001$), indicating that the difference between male and female ASMR increased during the period when CQ/HCQ was officially recommended for COVID-19 outpatients and the healthcare system was strained (z-score $\geq$ 3). However, this allows no conclusions regarding the hypothesized mechanism because it does not disentangle the contributions of G6PDD prevalence, regional HCQ availability, or intensity of healthcare system strain.

Models 2 and 4 introduce interaction terms with G6PDD prevalence, regional CQ/HCQ availability, and the all-cause *z-score*. This specification allows the effect of *period_HCQ* to vary by regional risk factors and healthcare system strain (proxied by *z-score*), effectively isolating the hypothesized mechanism. All three interactions are positive and highly significant for both outcome variants: *period_HCQ × G6PDD prevalence* (8.209, $p \leq 0.001$; 8.232, $p \leq 0.001$), *period_HCQ × HCQ_per_pop* (0.770, $p \leq 0.001$; 0.823, $p \leq 0.001$), and *period_HCQ × z-score* (3.050, $p \leq 0.001$; 2.653, $p \leq 0.001$). For example, a one-category increase in G6PDD prevalence – corresponding to a two-percentage-point increase (Table IV, Appendix) – alters the effect of *period_HCQ* by an additional sex difference of 8.209 all-cause excess deaths or 8.232 COVID-19 deaths per 100,000. Marginal effects plots for G6PDD prevalence are provided in the appendix (Figures III and IV). The coefficients' magnitudes should be interpreted with caution as these vary slightly over tested definitions of *period_HCQ*. However, significance and signs are robust and consistent with the tested mechanism.

The control variable *z-score* captures the baseline sex difference in ASMR, independent of CQ/HCQ treatment, consistently predicting a positive coefficient. By contrast, *the z-score* interaction indicates that during *period_HCQ*, additional healthcare system strain disproportionately amplified male–female ASMR differences, consistent with the idea that adverse effects of CQ/HCQ may have gone unrecognized in overburdened regions.

Taken together, the results suggest that in regions with higher G6PDD prevalence, greater CQ/HCQ availability, and stronger healthcare system strain, sex differences in all-cause excess and COVID-19 ASMR increased disproportionately, supporting the hypothesized mechanism of drug-induced hemolytic anemia under stressed healthcare system conditions.

Control variables also behaved as expected: the coefficient of the 4-week lagged sex difference in test positive rates formally reflects a 100% difference (i.e., only men tested positive and no women) and is positive and significant in all models. It should be interpreted as capturing short-term shifts in infection burdens between sexes as discussed in the Methods section. Although this variable is an imperfect proxy for true infection rates, its significance likely reflects that observed mortality patterns overlapped with periods and regions in which one sex was more likely to be registered as infected.

The 3-week lagged sex difference in first-dose vaccination rates has a negative coefficient, suggesting that weeks with relatively higher female vaccination are associated with smaller sex gaps in mortality. As with the test-positivity variable, the coefficient formally reflects a 100% difference and should be interpreted as capturing relative shifts in vaccination coverage between sexes during the nationwide vaccination campaign starting in April 2021. The effect is somewhat unstable across specifications and was only observed in 2021, so it should be interpreted cautiously. While it provides some insight into sex-differential dynamics during the vaccination campaign, this variable does not contribute to the identification of *period_HCQ* and serves as a supplementary result.

To further assess which sex drives the effect of increased sex differences in ASMR during *period_HCQ*, models were re-estimated with male and female ASMR as outcomes. While these specifications are subject to omitted variable bias and endogeneity, they serve as a diagnostic check. Results show a strong effect for males but only negligible effects for females, suggesting that the observed sex differences stem from higher male mortality rather than unusually low female mortality.

Discussion

The findings from Peru align with prior studies showing higher COVID-19 mortality [4, 5] and all-cause excess mortality [3] among men. Significant periods with higher female than male all-cause excess mortality as observed in Thailand [6], were not present in Peru. This difference can be explained partly by the exclusion of traffic accidents in this analysis, which were probably responsible for the observed pattern in Thailand. The correlation between general excess mortality (z-score) and sex differences as found in Europe [3] was also confirmed by this analysis.

Moreover, the results are consistent with previous studies linking CQ/HCQ to an increased all-cause mortality [13], especially in the Peruvian context [14].

Except for some case reports [22–26], the risk of CQ/HCQ-induced acute hemolytic anemia in G6PD-deficient patients is generally considered low according to a systematic review of the available evidence [21]. This study, however, suggests that this risk may have been underestimated in the specific context of Peru during the COVID-19 crisis, which was characterized by panic over an unknown virus, an underdeveloped and overburdened healthcare system, non-transparent treatment protocols [55], and misuse of medications [56, 57]. Under such circumstances, uncontrolled outpatient use of CQ and HCQ may have exposed G6PD-deficient individuals to increased hemolytic risk, potentially contributing to the observed male-female mortality gap. This hypothesis warrants further investigation using individual-level data linking medication use, hematologic indicators, and G6PD status.

The Peruvian prison system may be a suitable sample for future research. From an ecological perspective, a temporal association between CQ/HCQ distribution and male mortality was also observed there. Until October 2020, 445 Peruvian prison inmates died due to COVID-19, 251 (56%) of them between April and May [58] which correlates with a delivery of 10,800 tablets of 400mg HCQ in mid-April [36].

There are several limitations to this study that are worth mentioning. First, the model is ecological in nature and cannot establish individual-level causality. Second, the state-level measure of G6PDD prevalence [28] is modeled and relatively coarse, which may blunt interaction effects. Third, the measure of CQ/HCQ availability is based on stock data per population and does not capture actual intake, compliance, informal distribution, black-market flows, or tablets sold via the private health sector. Fourth, the z-score of all-cause ASMR serves only as an indirect indicator of healthcare system strain and does not reflect ICU capacity, staffing shortages, etc. Finally, the all-cause excess ASMR was predicted by the model by 12 [12] and are used directly in the TWFE analysis, without formally accounting for their estimation uncertainty, which may have resulted in underestimated standard errors.

Conclusion

This study indicates that Peru's nationwide CQ/HCQ treatment policy during the early COVID-19 pandemic may have contributed to sex-specific differences in mortality among adults aged 45+, particularly in regions with higher glucose-6-phosphate dehydrogenase deficiency (G6PDD) prevalence and during periods of elevated healthcare system strain. Across states, male all-cause and COVID-19 excess mortality consistently exceeded female mortality, with the largest male-female differences coinciding with periods of plausible widespread CQ/HCQ use among COVID-19 outpatients. Interaction analyses further indicate that regions with higher G6PDD prevalence, greater CQ/HCQ availability, and elevated all-ages excess mortality—i.e., healthcare system strain—experienced amplified male-female mortality gaps.

These findings are consistent with the hypothesis that CQ/HCQ-induced oxidative stress may have triggered hemolytic crises in (undiagnosed) G6PDD patients. In the context of COVID-19, where infection itself causes oxidative stress, hemolysis, and presents with overlapping symptoms, such adverse reactions may have been overlooked or misclassified as severe COVID-19 in overstretched healthcare systems.

While causality cannot be established due to the ecological study design, these findings underscore the importance of incorporating genetic and sex-specific risk factors into pandemic treatment guidelines, particularly in resource-limited settings with high uncertainty. Emergency treatment decisions should

consider not only efficacy data but also potential safety risks for vulnerable subpopulations. Given the likelihood that hemolysis may have been unrecognized, ecological analyses remain crucial for detecting population-level associations and generating hypotheses for future research and policy-making.

Appendix

Figure I: Sex differences in all-cause excess ASMR for ages 45+ over time by state

Note: The variance of the sex difference was obtained as the sum of male and female ASMR variances (see section Statistical Analysis), and the 95% confidence interval (95% CI) was constructed as the point estimate ± 1.96 times the square root of this variance, assuming independence of errors (Cov(m, f) = 0).

Figure II: Sex differences in COVID-19 ASMR for ages 45+ over time by state

Table I: Sum of delivered HCQ Tablets per state population

State	HCQ_per_pop			
	period_HCQ z-score = 4	period_HCQ z-score = 3.5	period_HCQ z-score = 3	period_HCQ z-score = 2.5
Amazonas	39.62	39.62	39.62	39.62
Ancash	14.74	14.74	14.74	14.74
Apurimac	-	-	20.86	20.86
Arequipa	19.47	19.47	19.47	19.47
Ayacucho	-	-	27.35	27.35
Cajamarca	8.33	8.33	8.33	8.33
Cusco	10.30	10.30	10.30	10.30
Huancavelica	11.65	11.65	11.65	11.65
Huanuco	26.92	26.92	26.92	26.92
Ica	36.48	36.48	36.48	36.48
Junin	15.32	15.32	15.32	15.32
La Libertad	17.16	17.16	17.16	17.16
Lambayeque	16.36	21.55	21.55	21.55
Lima + Callao	18.69	18.69	18.69	18.69
Loreto	18.64	18.64	20.61	21.81
Madre de Dios	178.36	178.36	178.36	178.36
Moquegua	35.60	35.60	35.60	35.60
Pasco	25.17	25.17	25.17	25.17
Piura	5.51	5.51	5.51	5.51
Puno	7.65	7.65	7.65	7.65
San Martin	36.37	36.37	36.37	36.37
Tacna	19.75	19.75	19.75	19.75
Tumbes	12.29	12.29	12.29	12.29
Ucayali	20.57	20.57	20.57	39.68
Median	18.66	19.08	20.16	20.31
Mean	26.68	26.92	26.82	27.64
Std. Dev.	33.68	33.62	32.25	32.30

Note: The different CQ/HCQ drugs used were: 'CLOROQUINA 150 MG TAB', 'CLOROQUINA FOSFATO 250 MG (EQUIV. 155 MG CLOROQUINA BASE) TAB', 'HIDROXICLOROQUINA [SULFATO] 400 MG TAB', 'HIDROXICLOROQUINA SULFATO 200 MG (EQUIV. 155 MG HIDROXICLOROQUINA) TAB' [36]. According to the treatment protocol, 1 CQ is considered as equivalent to 1/6 HCQ [9]. All drugs were standardized using 1 tablet '155 MG HIDROXICLOROQUINA' as base. All of these standardized tablets that were distributed before or during *period_HCQ* were summed up and divided by the state population, with *period_HCQ* constructed with different definitions (z-score thresholds) as described in section Data Preparation. Missing values stem from no identification of *period_HCQ* with given z-score threshold.

Table II: Scarcity CQ/HCQ: Doses per positive tests

State	CQ/HCQ-Doses per positive tests in age group 45+				Total tests per population during Period_HCQ (z-score = 2.5)
	period_HCQ z-score = 4	period_HCQ z-score = 3.5	period_HCQ z-score = 3	period_HCQ z-score = 2.5	
Amazonas	1.36	1.21	1.21	1.11	0.19
Ancash	0.46	0.46	0.42	0.41	0.21
Apurimac	NaN	NaN	4.48	2.12	0.13
Arequipa	0.23	0.22	0.22	0.22	0.50
Ayacucho	NaN	NaN	3.30	2.11	0.06
Cajamarca	0.59	0.59	0.57	0.57	0.10
Cusco	0.41	0.41	0.41	0.41	0.14
Huancavelica	1.36	1.09	1.09	1.09	0.09
Huanuco	1.23	1.14	1.14	1.14	0.14
Ica	0.54	0.54	0.54	0.54	0.39
Junin	1.12	1.03	1.03	1.03	0.10
La Libertad	0.51	0.51	0.48	0.48	0.19
Lambayeque	0.94	0.80	0.70	0.59	0.17
Lima + Callao	0.27	0.27	0.27	0.27	0.40
Loreto	1.38	1.38	1.09	0.94	0.08
Madre de Dios	3.66	3.35	3.35	3.05	0.26
Moquegua	0.40	0.40	0.39	0.39	0.48
Pasco	1.80	1.68	1.68	1.68	0.14
Piura	0.64	0.64	0.64	0.64	0.04
Puno	0.56	0.56	0.56	0.56	0.08
San Martin	1.36	1.36	1.29	1.29	0.11
Tacna	0.51	0.51	0.51	0.49	0.26
Tumbes	1.01	1.01	1.01	1.01	0.05
Ucayali	1.80	1.80	1.80	2.17	0.07
Median	0.79	0.72	0.85	0.79	0.14
Mean	1.00	0.94	**1.16**	1.00	0.18
Std. Dev.	0.73	0.67	1.04	0.70	0.13

Note: CQ/HCQ deliveries [36] that occurred before or during *period_HCQ* were considered, 1 CQ/HCQ dose corresponds to 18 tablets of HIDROXICLOROQUINA SULFATO 200 MG [37]. All positive COVID-19 tests in the age group 45+ that occurred during

period_HCQ were counted, with *period_HCQ* constructed with different specifications (z-score thresholds) as described in section *Data Preparation*.

Table III: Inference with Wild Cluster Bootstrap [50] for Models 1, 2, 3, 4 (Table 2, main text)

Model	p-value (Wild Cluster Bootstrap)			
	1 (all-cause excess)	2 (all-cause excess)	3 (COVID-19)	4 (COVID-19)
period_HCQ	0.000	0.001	0.002	0.000
period_HCQ x G6PDD prevalence	-	0.075	-	0.023
period_HCQ x HCQ_per_pop	-	0.046	-	0.003
period_HCQ x z-score	-	0.000	-	0.004
z-score	0.000	0.000	0.000	0.000
Sex difference test positive rate (lag 4)	0.028	0.029	0.003	0.003
Sex difference vaccination rate (lag 3)	0.002	0.051	0.000	0.007

Figure III: Marginal effect of period_HCQ conditional on G6PDD prevalence. Dependent variable: sex difference all-cause excess ASMR ages 45+

Note: Variable *HCQ_per_pop* (time-invariant) is held at its mean. Variable *z-score* is held at its mean during *period_HCQ.*

Figure IV: Marginal effect of period_HCQ conditional on G6PDD prevalence. Dependent variable: sex difference Covid-19 ASMR ages 45+

Note: Variable *HCQ_per_pop* (time-invariant) is held at its mean. Variable *z-score* is held at its mean during *period_HCQ*.

Figure V: States of Peru and modeled G6PDD prevalence [28]

Table IV: G6PDD prevalence by state, derived from [28]

State	G6PDD prevalence [28]
Amazonas	4.5 (6-10%)

Ancash	2.0 (2-4%)
Apurimac	2.0 (2-4%)
Arequipa	2.0 (2-4%)
Ayacucho	2.0 (2-4%)
Cajamarca	4.0 (6-8%)
Cusco	2.0 (2-4%)
Huancavelica	1.5 (0-4%)
Huanuco	2.0 (2-4%)
Ica	1.5 (0-4%)
Junin	1.5 (0-4%)
La Libertad	3.0 (4-6%)
Lambayeque	4.0 (6-8%)
Lima + Callao	1.0 (0-2%)
Loreto	4.5 (6-10%)
Madre de Dios	3.0 (4-6%)
Moquegua	2.0 (2-4%)
Pasco	1.5 (0-4%)
Piura	5.0 (8-10%)
Puno	3.0 (4-6%)
San Martin	3.5 (4-8%)
Tacna	2.0 (2-4%)
Tumbes	5.0 (8-10%)
Ucayali	3.0 (4-6%)

Note: G6PDD prevalence was coded based on Monteiro et al. (2014); values in parentheses indicate the original modeled prevalence from 28 [28].

References

1. Peretz C, Rotem N, Keinan-Boker L, Furshpan A, Green M, Bitan M, Steinberg DM. Excess mortality in Israel associated with COVID-19 in 2020-2021 by age group and with estimates based on daily mortality patterns in 2000-2019. Int J Epidemiol. 2022;51:727–36. doi:10.1093/ije/dyac047.

2. Henry NJ, Elagali A, Nguyen M, Chipeta MG, Moore CE. Variation in excess all-cause mortality by age, sex, and province during the first wave of the COVID-19 pandemic in Italy. Sci Rep. 2022;12:1077. doi:10.1038/s41598-022-04993-7.

3. Nielsen J, Nørgaard SK, Lanzieri G, Vestergaard LS, Moelbak K. Sex-differences in COVID-19 associated excess mortality is not exceptional for the COVID-19 pandemic. Sci Rep. 2021;11:20815. doi:10.1038/s41598-021-00213-w.

4. Bambra C, Albani V, Franklin P. COVID-19 and the gender health paradox. Scand J Public Health. 2021;49:17–26. doi:10.1177/1403494820975604.

5. Peckham H, Gruijter NM de, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ITU admission. Nat Commun. 2020;11:6317. doi:10.1038/s41467-020-19741-6.

6. Pothisiri W, Prasitsiriphon O, Apakupakul J, Ploddi K. Gender differences in estimated excess mortality during the COVID-19 pandemic in Thailand. BMC Public Health. 2023;23:1900. doi:10.1186/s12889-023-16828-9.

7. Akter S. The Gender Gap in COVID-19 Mortality in the United States. Feminist Economics. 2021;27:30–47. doi:10.1080/13545701.2020.1829673.

8. World Health Organization. WHO discontinues hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-19. 2020. https://www.who.int/news/item/04-07-2020-who-discontinues-hydroxychloroquine-and-lopinavir-ritonavir-treatment-arms-for-covid-19. Accessed 4 Aug 2025.

9. Ministerio de Salud (MINSA). Resolución Ministerial N.° 139-2020-MINSA; 2020.

10. Ministerio de Salud (MINSA). Resolución Ministerial N.° 193-2020-MINSA; 2020.

11. Gestión. Minsa anuncia que la hidroxicloroquina se retirará de la guía de medicamentos para el COVID-19. 2020. https://gestion.pe/peru/coronavirus-peru-pilar-mazzetti-anuncia-que-la-hidroxicloroquina-se-retirara-de-la-guia-de-medicamentos-para-tratamiento-del-covid-19-nndc-noticia/. Accessed 4 Sep 2025.

12. Karlinsky A, Kobak D. Tracking excess mortality across countries during the COVID-19 pandemic with the World Mortality Dataset. Elife 2021. doi:10.7554/eLife.69336.

13. Axfors C, Schmitt AM, Janiaud P, Van't Hooft J, Abd-Elsalam S, Abdo EF, et al. Mortality outcomes with hydroxychloroquine and chloroquine in COVID-19 from an international collaborative meta-analysis of randomized trials. Nat Commun. 2021;12:2349. doi:10.1038/s41467-021-22446-z.

14. Soto-Becerra P, Culquichicón C, Hurtado-Roca Y, Araujo-Castillo RV. Real-world effectiveness of hydroxychloroquine, azithromycin, and ivermectin among hospitalized COVID-19 patients: results of a target trial emulation using observational data from a nationwide healthcare system in Peru; 2020.

15. Mokhtari M, Mohraz M, Gouya MM, Namdari Tabar H, Tabrizi J-S, Tayeri K, et al. Clinical outcomes of patients with mild COVID-19 following treatment with hydroxychloroquine in an outpatient setting. Int Immunopharmacol. 2021;96:107636. doi:10.1016/j.intimp.2021.107636.

16. Lofgren SM, Nicol MR, Bangdiwala AS, Pastick KA, Okafor EC, Skipper CP, et al. Safety of Hydroxychloroquine Among Outpatient Clinical Trial Participants for COVID-19. Open Forum Infect Dis. 2020;7:ofaa500. doi:10.1093/ofid/ofaa500.

17. Lagier J-C, Million M, Gautret P, Colson P, Cortaredona S, Giraud-Gatineau A, et al. Outcomes of 3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in

Marseille, France: A retrospective analysis. Travel Med Infect Dis. 2020;36:101791. doi:10.1016/j.tmaid.2020.101791.

18. Million M, Lagier J-C, Gautret P, Colson P, Fournier P-E, Amrane S, et al. Early treatment of COVID-19 patients with hydroxychloroquine and azithromycin: A retrospective analysis of 1061 cases in Marseille, France. Travel Med Infect Dis. 2020;35:101738. doi:10.1016/j.tmaid.2020.101738.

19. Mohana A, Sulaiman T, Mahmoud N, Hassanein M, Alfaifi A, Alenazi E, et al. Hydroxychloroquine Safety Outcome within Approved Therapeutic Protocol for COVID-19 Outpatients in Saudi Arabia. Int J Infect Dis. 2021;102:110–4. doi:10.1016/j.ijid.2020.10.031.

20. Luzzatto L, Nannelli C, Notaro R. Glucose-6-Phosphate Dehydrogenase Deficiency. Hematol Oncol Clin North Am. 2016;30:373–93. doi:10.1016/j.hoc.2015.11.006.

21. Kane M. Hydroxychloroquine Therapy and G6PD Genotype. 2023. https://www.ncbi.nlm.nih.gov/books/NBK591356/. Accessed 4 Sep 2025.

22. Aguilar J, Averbukh Y. Hemolytic Anemia in a Glucose-6-Phosphate Dehydrogenase-Deficient Patient Receiving Hydroxychloroquine for COVID-19: A Case Report. Perm J 2020. doi:10.7812/TPP/20.158.

23. Sgherza N, Dalfino L, Palma A, Vitucci A, Campanale D, Grasso S, Musto P. "Hemolysis, or not Hemolysis, that is the Question". Use of Hydroxychloroquine in a Patient with COVID-19 Infection and G6PD Deficiency. Mediterr J Hematol Infect Dis. 2020;12:e2020076. doi:10.4084/MJHID.2020.076.

24. Chaney S, Basirat A, McDermott R, Keenan N, Moloney E. COVID-19 and hydroxychloroquine side-effects: glucose 6-phosphate dehydrogenase deficiency (G6PD) and acute haemolytic anaemia. QJM. 2020;113:890–1. doi:10.1093/qjmed/hcaa267.

25. Mastroianni F, Colombie V, Claes G, Gilles A, Vandergheynst F, Place S. Hydroxychloroquine in a G6PD-Deficient Patient with COVID-19 Complicated by Haemolytic Anaemia: Culprit or Innocent Bystander? Eur J Case Rep Intern Med. 2020;7:1875. doi:10.12890/2020_001875.

26. Maillart E, Leemans S, van Noten H, Vandergraesen T, Mahadeb B, Salaouatchi MT, et al. A case report of serious haemolysis in a glucose-6-phosphate dehydrogenase-deficient COVID-19 patient receiving hydroxychloroquine. Infect Dis (Lond). 2020;52:659–61. doi:10.1080/23744235.2020.1774644.

27. Ayyad M, Abu Alya W, Barabrah AM, Darawish SM, AlHabil Y, MohammedAli M, et al. Autoimmune hemolytic anemia in COVID-19 patients: A systematic review of 105 cases on clinical characteristics and outcomes. Clin Immunol. 2025;277:110512. doi:10.1016/j.clim.2025.110512.

28. Monteiro WM, Val FFA, Siqueira AM, Franca GP, Sampaio VS, Melo GC, et al. G6PD deficiency in Latin America: systematic review on prevalence and variants. Mem Inst Oswaldo Cruz. 2014;109:553–68. doi:10.1590/0074-0276140123.

29. Whittaker C, Walker PGT, Alhaffar M, Hamlet A, Djaafara BA, Ghani A, et al. Under-reporting of deaths limits our understanding of true burden of covid-19. BMJ. 2021;375:n2239. doi:10.1136/bmj.n2239.

30. Andina Peruvian News Agency. Covid-19: segundo informe para actualizar cifra de fallecidos se conocerá esta semana. 26.07.2020. https://andina.pe/agencia/noticia-covid19-segundo-informe-para-actualizar-cifra-fallecidos-se-conocera-esta-semana-807333.aspx. Accessed 4 Sep 2025.

31. Sempé L, Lloyd-Sherlock P, Martínez R, Ebrahim S, McKee M, Acosta E. Estimation of all-cause excess mortality by age-specific mortality patterns for countries with incomplete vital statistics: a population-based study of the case of Peru during the first wave of the COVID-19 pandemic. Lancet Reg Health Am. 2021;2:None. doi:10.1016/j.lana.2021.100039.

32. Kupek E. How many more? Under-reporting of the COVID-19 deaths in Brazil in 2020. Trop Med Int Health. 2021;26:1019–28. doi:10.1111/tmi.13628.

33. Ministerio de Salud (MINSA). SINADEF - Defunciones Registradas. 2025. https://www.minsa.gob.pe/reunis/index.asp?op=1&niv=1&tbl=1#. Accessed 14 Aug 2025.

34. Castanheira HC, Da Monteiro Silva JHC. Examining sex differences in the completeness of Peruvian CRVS data and adult mortality estimates. Genus. 2022;78:3. doi:10.1186/s41118-021-00151-5.

35. Ministerio de Salud (MINSA). Fallecidos por COVID-19. 2025. https://www.datosabiertos.gob.pe/dataset/fallecidos-por-covid-19-ministerio-de-salud-minsa. Accessed 14 Aug 2025.

36. Ministerio de Salud (MINSA). Centro Nacional de Abastecimiento de Recursos Estratégicos en Salud (CENARES) - Distribucion por Producto. 2024. https://intranet.cenares.gob.pe/cenares/abastecimiento/DISTRIBUCION/wf_distribucion_por_produc to.aspx. Accessed 5 Dec 2024.

37. Ministerio de Salud (MINSA). Resolución Ministerial N.° 270-2020-MINSA; 2020.

38. Instituto Nacional de Salud (INS). Molecular Tests COVID-19: NETLAB. 2025. https://www.datosabiertos.gob.pe/dataset/dataset-de-pruebas-moleculares-del-instituto-nacional-de-salud-para-covid-19-ins. Accessed 11 Mar 2025.

39. Ministerio de Salud (MINSA). Non-Molecular Tests COVID-19: SISCOVID F100. 2025. https://www.datosabiertos.gob.pe/dataset/siscovid-f100-pruebas/resource/7f897963-f4c0-4a68-ae2f-10d49369edf9. Accessed 7 Feb 2025.

40. Ministerio de Salud (MINSA). Vacunación COVID19. 2025. https://www.datosabiertos.gob.pe/dataset/vacunacion/resource/c2296574-f1ca-46a4-9d1d-1aee320678a7. Accessed 20 Jun 2025.

41. Ministerio de Salud (MINSA). Población INEI 2021. 2025. https://datosabiertos.gob.pe/dataset/poblaci%C3%B3n-peru/resource/6e31247b-8057-450c-873d-d1ba28cf136a. Accessed 3 Aug 2025.

42. Ministerio de Salud (MINSA). Resolución Ministerial N.° 375-2020-MINSA; 2020.

43. RCR Peru. CORONAVIRUS EN PIURA: EL 100% DE PACIENTES CON SÍNTOMAS LEVES SE HAN RECUPERADO CON IVERMECTINA. 2020. https://www.rcrperu.com/coronavirus-en-piura-el-100-de-pacientes-con-sintomas-leves-se-han-recuperado-con-ivermectina/. Accessed 4 Aug 2025.

44. Chamie JJ, Hibberd JA, Scheim DE. COVID-19 Excess Deaths in Peru's 25 States in 2020: Nationwide Trends, Confounding Factors, and Correlations With the Extent of Ivermectin Treatment by State. Cureus. 2023;15:e43168. doi:10.7759/cureus.43168.

45. Soriano-Moreno DR, Fernandez-Guzman D, Sangster-Carrasco L, Quispe-Vicuña C, Grados-Espinoza P, Ccami-Bernal F, et al. Factors Associated With Drug Consumption Without Scientific Evidence in Patients With Mild COVID-19 in Peru. J Patient Saf. 2022;18:e1189-e1195. doi:10.1097/PTS.0000000000001053.

46. Quispe-Cañari JF, Fidel-Rosales E, Manrique D, Mascaró-Zan J, Huamán-Castillón KM, Chamorro-Espinoza SE, et al. Self-medication practices during the COVID-19 pandemic among the adult population in Peru: A cross-sectional survey. Saudi Pharm J. 2021;29:1–11. doi:10.1016/j.jsps.2020.12.001.

47. Castro NE, García DR, Rivera MT, Rondán-Guerrero P, García-Rojas F, Taype-Rondan A. Tendencias en el uso de fármacos para la COVID-19 durante la primera ola de la pandemia en un hospital de Lima, Perú. [Trends in drugs usage for COVID-19 during the first wave of the pandemic in a hospital in Lima, Peru]. Rev Peru Med Exp Salud Publica. 2021;38:608–14. doi:10.17843/rpmesp.2021.384.8820.

48. Casas Bueno, Guerrero Martínez. Medicamentos buscados en la web para la COVID-19 en Perú: Estudio infodemiológico [Tesis (Médico Cirujano)]. Lima, Perú: Universidad Peruana Cayetano Heredia; 2021.

49. Hoechle D. Robust Standard Errors for Panel Regressions with Cross-Sectional Dependence. The Stata Journal: Promoting communications on statistics and Stata. 2007;7:281–312. doi:10.1177/1536867X0700700301.

50. Fischer A, Michuda A. wildboottest (Python module); 2022.

51. Angulo-Bazan Y, Solis G, Cardenas F, Jorge A, Acosta J, Cabezas C. Household transmission in people infected with SARS-CoV-2 (COVID-19) in Metropolitan Lima; 2020.

52. Huamaní C, Velásquez L, Montes S, Mayanga-Herrera A, Bernabé-Ortiz A. SARS-CoV-2 seroprevalence in a high-altitude setting in Peru: adult population-based cross-sectional study. PeerJ. 2021;9:e12149. doi:10.7717/peerj.12149.

53. Moreira-Soto A, Pachamora Diaz JM, González-Auza L, Merino Merino XJ, Schwalb A, Drosten C, et al. High SARS-CoV-2 seroprevalence in rural Peru, 2021; a cross-sectional population-based study; 2021.

54. Moyano LM, Toledo AK, Chirinos J, Vilchez Barreto PMQ, Cavalcanti S, Gamboa R, et al. SARS-CoV-2 seroprevalence on the north coast of Peru: A cross-sectional study after the first wave. PLoS Negl Trop Dis. 2023;17:e0010794. doi:10.1371/journal.pntd.0010794.

55. Taype-Rondan A, Herrera-Añazco P, Málaga G. Sobre la escasa transparencia en los documentos técnicos para el tratamiento de pacientes con COVID-19 en Perú. Acta Med Peru 2020. doi:10.35663/amp.2020.372.982.

56. Caira-Chuquineyra B, Fernandez-Guzman D, Alvarez-Arias PM, Zarate-Curi ÁA, Herrera-Añazco P, Benites-Zapata VA. Association between prehospital medication and fatal outcomes in a cohort of hospitalized patients due to coronavirus disease-2019 in a referral hospital in Peru. Travel Med Infect Dis. 2022;50:102472. doi:10.1016/j.tmaid.2022.102472.

57. Martinez-Rivera RN, Taype-Rondan A. Overmedication in COVID-19 Context: A Report From Peru. J Clin Pharmacol. 2020;60:1155–6. doi:10.1002/jcph.1704.

58. Sindeev A, Martínez-Álvarez BM. Clinical and epidemiological characteristics of prisoners infected and deceased by COVID-19, National Penitentiary Institute of Peru, 2020. Rev Esp Sanid Penit. 2022;24:15–22. doi:10.18176/resp.00045.