Lower pre-treatment TMS-evoked cortical reactivity and alpha-band oscillatory dynamics predict efficacy of primary motor cortex neuromodulation for chronic pain

Enrico De Martino[1], Margit Midtgaard Bach[1*], Anne Jakobsen[1*], Bruno Nascimento Couto[1], Nahian S. Chowdhury[2,3], Stian Ingemann-Molden[1], Adenauer Casali[4], Thomas Graven-Nielsen[1], Daniel Ciampi de Andrade[1]

[1] Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark
[2] University of New South Wales, Sydney, New South Wales, Australia,
[3] Pain IMPACT, Neuroscience Research Australia, Sydney, Australia
[4] Institute of Science and Technology, Federal University of São Paulo, São Paulo, Brazil
[*] Contributed equally to this work and share second authorship

Abstract

Repetitive transcranial magnetic stimulation (rTMS) targeting the primary motor cortex (M1) provides significant pain relief in approximately 45% of patients with chronic pain. Identifying markers that predict rTMS treatment responders to M1 before initiating treatment is crucial for informing decision-making and improving patient outcomes in clinical practice. In this secondary analysis of a clinical trial assessing the effects of rTMS on different clinical targets (NCT06395649), we employed a combination of transcranial magnetic stimulation and electroencephalography (TMS-EEG) to investigate cortical reactivity and oscillatory dynamics in 43 patients with chronic pain prior to 12 sessions of therapeutic 10 Hz rTMS applied to M1 over an 8-week period. Responders were defined as those reporting a ≥30% reduction in pain intensity on a visual analogue scale at the end of treatment. TMS-evoked cortical reactivity was assessed using global mean field power (GMFP), calculated across all electrodes, and local mean field power (LMFP), computed in 4 electrodes at the TMS stimulation site. TMS-evoked oscillatory dynamics were evaluated with event-related spectral perturbation (ERSP) and intertrial coherence (ITC) across three frequency bands: alpha (8–12 Hz), beta-1 (13–20 Hz), and beta-2 (21–30 Hz) at the stimulation site. 20 patients (47%) were responders at the end of the rTMS treatment. Compared to non-responders, responders exhibited lower GMFP and lower LMFP at the stimulation region (both $P<0.05$). They also showed lower alpha-band ERSP and ITC at the stimulation region (both $P<0.05$). These lower neurophysiological features were associated with greater reductions in pain intensity (all $P<0.05$). Exploratory supervised machine learning using three baseline TMS-EEG features (GMFP, alpha-band ERSP, and ITC) predicted responder status with acceptable accuracy (ROC-AUC = 0.70, PR-AUC = 0.76). Together, these results suggest that lower pre-treatment TMS-evoked cortical reactivity and alpha-band oscillatory dynamics activity predict better clinical response to rTMS in patients with chronic pain (≈ 3 in 4 Responders correctly identified). Future prospectively designed clinical studies should implement TMS-EEG assessment of the target to be used for rTMS before treatment is started, in order to prospectively test the utility of these personalized neuromodulatory interventions.

Introduction

High-frequency (10 Hz) repetitive transcranial magnetic stimulation (rTMS) targeting the primary motor cortex (M1) is a non-invasive neuromodulation technique that induces analgesic effects in patients with chronic pain [1] by repetitively depolarizing myelinated axons [2], consistent with activity-dependent (LTP-like) plasticity in corticothalamic networks [3]. International guidelines currently recommend TMS for the management of different chronic pain conditions, such as chronic neuropathic pain [4], migraine [5], and fibromyalgia [6]. However, clinical trials have demonstrated high inter-individual response variability in chronic pain patients receiving M1 rTMS treatment, with approximately 45% showing a pain reduction of more than 30% [7,8]. Determining the likelihood of treatment response at the individual level is highly relevant in clinical practice, as it may help direct therapy to those who are more likely to benefit from it [9]. A recent feasibility pilot study combining transcranial magnetic stimulation with electroencephalography (TMS-EEG), a neurophysiological approach for assessing cortical function [10], showed that pre-treatment global phase-based connectivity may help select rTMS strategies for chronic pain relief [11]. Although innovative, this feasibility pilot study investigated only a single, global phase-based connectivity measure, and a more comprehensive analysis could provide greater insights [12].

TMS-EEG enables the assessment of many aspects of cortical physiology, including TMS-evoked global and local reactivity and TMS-evoked oscillatory dynamics, through measures such as global mean field power (GMFP), local mean field power (LMFP), event-related spectral perturbation (ERSP, a measure of changes in power across frequencies in response to stimuli), and intertrial coherence (ITC, a measure of phase consistency across trials) [10]. Previous studies have shown that experimental acute pain reduced both local reactivity (LMFP measured peri-cortical target) and global reactivity (GMFP) compared to non-painful stimuli [13]. Additionally, reductions in amplitude-based (ERSP) and phase-based (ITC) cortical oscillatory dynamics were also shown to be reduced during acute pain in the alpha band following M1 stimulation and in the low beta band following left dorsolateral prefrontal cortex (DLPFC) stimulation [14], suggesting a less synchronized and more variable cortical response to TMS probing of both M1 and DLPFC during short-lasting experimental pain. Notably, decreases in TMS-evoked oscillatory dynamics did not occur uniformly across participants: approximately 50% of healthy individuals exhibited predominant reductions in local cortical function in either the DLPFC or M1, but rarely at both sites concurrently [13]. Also, individual trait sensitivity to pain was significantly related to the magnitude of M1-evoked oscillatory dynamics during experimental pain, with lower reductions in phase-based connectivity correlated to higher individual pain sensitivity [14]. These data consistently suggest that even in healthy individuals under experimental pain conditions, personal characteristics may influence how pain affects cortical function [13]. Although those findings are in experimental pain in healthy participants, they suggest trait-like differences in state-dependent M1 responsiveness that could generalize to patient populations. It would therefore be possible to hypothesize that individuals with different profiles of TMS M1 responses during pain respond differently to modulatory therapeutic interventions aiming at this brain area. Based on homeostatic plasticity principles, which propose that cortical activity is regulated via compensatory adjustments whenever it deviates from the optimal range [15], we hypothesized that chronic pain patients with low cortical reactivity and oscillatory dynamics upon probing M1 before treatment would be more likely to benefit from therapeutic M1 rTMS compared to those without these features. To test this hypothesis, we compared TMS-evoked cortical reactivity, as well as oscillatory dynamics, measured before therapy to clinical outcomes in responders and non-responders to M1 rTMS in people with chronic pain. We also provided single-patient classification for each TMS-EEG metric based on normative data from 125 healthy pain-free individuals.

Methods

Patients

This study is a secondary exploratory analysis from a larger trial that assessed the use of pre-treatment TMS-EEG to select the cortical target for rTMS in each patient. In this double blind, three-arm randomized clinical trial (clinical trial registration number NCT06395649) [11] M1 was used as a target in the active control comparator arm, representing the current best neuromodulatory treatment. Among 90 patients with chronic pain, 43 patients received rTMS treatment targeting M1 and were analyzed in the present study. Eligible participants were males and females aged 18 to 80 years with pain persisting for at least 6 months. All chronic pain mechanisms (neuropathic, nociplastic, and nociceptive) with at least moderate pain (pain intensity higher than 3/10 on a 0–10 numerical rating scale; NRS) were allowed in the trial. Exclusion criteria included diagnosed psychiatric conditions, substance abuse, unstable medical conditions, contraindications to rTMS, such as severe head trauma, prior neurosurgery, current or past epilepsy, intracranial hypertension, or implanted ferromagnetic devices. Additional reasons for exclusion included inability to complete study follow-up, current pregnancy, or breastfeeding. The study was approved by the local Ethics Committee (N-20230076) and conducted in accordance with the Helsinki Declaration. Written informed consent was obtained before the start of the study.

Study design

After recruitment and before rTMS, patients participated in a TMS-EEG session probing M1 with single-pulse TMS under EEG recording. rTMS sessions began the week after the TMS-EEG assessment and lasted 8 weeks. The protocol included five sessions over five consecutive weekdays (induction phase), followed by a maintenance phase with one session per week for seven weeks, totaling 12 rTMS sessions. At the end of the study, participants were divided into Responders or Non-responders based on their level of pain reduction and used for the current study. Responders were defined based on a predefined primary outcome by reduction of more than 30% in mean pain intensity, calculated by comparing scores on the last day of the maintenance phase (week 8) with the average pain intensity during the week preceding the TMS-EEG assessment. Pain intensity was assessed using a visual analogue scale (VAS), which allows participants to rate their pain on a continuous scale ranging from 0 to 100, where '0' indicates 'no pain' and '100' indicates 'the worst pain imaginable.'

Clinical trial outcome

The primary outcome measure was the reduction in pain intensity, evaluated after each rTMS session using the VAS [16], referring to the patient's pain at that moment in time. Furthermore, at the beginning and the end of the study, additional questionnaires were completed by the patients, including average pain over the last 7 days, pain in the last 24 hours, a body chart, the short-form brief pain inventory (BPI), the hospital anxiety and depression scale (HADS), and quality of life (EQ-5D), the number and type of medications, and patient global impression of change (PGIC). From the BPI, two indices were extracted: pain severity, calculated as the average of four pain intensity items (worst, least, average, and current pain), and pain interference, calculated as the average of seven items assessing interference with general activity, mood, walking ability, normal work, relations with others, sleep, and enjoyment of life. The number of painful body regions was calculated based on participants' reported body maps to quantify the spatial distribution of pain [17]. The HADS is a 14-item self-report questionnaire designed to assess symptoms of anxiety and depression, with higher scores indicating greater symptom severity [18]. The EQ-5D is a measure of health-related quality of life, assessing five dimensions: mobility, self-care, usual activities, pain/discomfort, and anxiety/depression. Each domain is rated by the participant based on perceived problems using

a VAS [19]. The PGIC is a 7-point Likert-type scale used to assess a participant's subjective
evaluation of overall improvement following treatment. Responses range from "Very much
worse" to "Very much improved", capturing the individual's perceived change in health status
compared to baseline [20].

TMS-EEG assessment
200 single-pulse TMS was delivered using a biphasic stimulator (MagPro R30, MagVenture
A/S, Farum, Denmark) and a figure-of-eight coil (Cool-B35, MagVenture A/S, Farum,
Denmark) [21]. TMS was applied to the hemisphere contralateral to the side of the most intense
pain. If patients did not report a predominant site, stimulation was delivered over the left
hemisphere.
To record TMS-evoked potentials (TEPs), a TMS-compatible passive electrode cap with 63
electrodes (10-20 system; g.tec-medical engineering GmbH, Schiedlberg, Austria) was used,
with the Cz electrode positioned at the vertex. The ground electrode was placed on the right
zygoma, and a reference electrode was located on the right mastoid [13]. Electrode impedance
was monitored and kept below 5 kΩ. Raw signals were sampled at 4800 Hz using a TMS-
compatible EEG amplifier (g.HIamp, g.tec-medical engineering GmbH, Schiedlberg, Austria).
To minimize auditory responses from the click generated by the TMS coil from interfering with
the TEPs, a TMS-click sound masking toolbox (TAAC, [22]) was used, and participants wore
noise-canceling in-ear headphones (ER3C, Etymotic, 50 Ohm, Illinois, USA). To reduce
somatosensory sensations caused by the TMS coil, two net caps (GVB-geliMED GmbH,
Ginsterweg Bad Segeberg, Germany) with a plastic wrap were applied to the EEG [13].
A navigated brain stimulation software (InVesalius Navigator, 3.1) was used to calibrate the
participant's head and the TMS coil position with the aid of an optical (infrared) motion capture
camera (Polaris Vicra, NDI, Ontario, Canada) [23]. The M1 target was identified by locating the
hand muscle representation in the left or right hemisphere at the largest observable hand
movement. The rMT was determined as the TMS intensity needed to evoke a visible hand
movement. An intensity of 90% of the rMT was used for the TEPs to prevent sensory feedback
contamination [24]. Furthermore, to ensure a detectable TEP, a real-time visualization tool (rt-
TEP) was used [25]. This interface enabled adjustments to the TMS coil orientation and intensity
across participants to minimize unwanted artifacts and ensure a minimum early peak-to-peak
amplitude of 6 μV, as measured in the average of 30 trials at the EEG electrode closest to the
M1 target. The InVesalius Navigator system and rt-TEP tool were used throughout the study to
monitor the TMS coil position (keeping it within 3 mm of the cortical targets), with the
interstimulus interval randomly jittered between 2000 and 2400 ms to prevent substantial
reorganization or plasticity processes from affecting longitudinal TMS-EEG measurements [26].
A total of 200 single-pulse TMS stimulations were administered to each patient. During TMS
stimulation, participants sat comfortably on an ergonomic chair with their eyes open, focused
on a fixation point on the wall.

TMS-EEG analysis
The pre-processing was performed using Python (Python Software Foundation). EEG data
were segmented into epochs of ±800 ms around the TMS trigger. The signal between 0 and 15
ms containing the TMS artifact was replaced by temporally mirrored data from the pre-stimulus
period [11]. Epochs and channels containing decay artifacts, 50 Hz line noise, eye blinks, eye
movements, or facial and neck muscle activity were visually inspected and manually rejected,
ensuring the highest possible signal quality while retaining a minimum of 150 trials. A single
independent component analysis (fast ICA) was used to remove residual artifacts [27]. Finally,
TEPs were band-pass filtered between 1 and 80 Hz using a third-order Butterworth filter, down-

sampled to 1200 Hz, EEG channels re-referenced to the average reference, and any bad
channels were interpolated using spherical splines [28].
Approximately half of the patients received stimulation over the left hemisphere (due to right-
sided pain or bilateral pain), and the other half received stimulation over the right hemisphere
(due to left-sided pain). To enable consistent group-level comparisons, data were reorganized
according to stimulation laterality. Specifically, results were analysed relative to the stimulated
hemisphere, regardless of whether stimulation was applied to the left or right hemisphere.
A predefined time window of 20–120 ms post-TMS was selected for extracting TMS-EEG
outcomes of TMS-evoked cortical reactivity and TMS-evoked oscillatory dynamics. This early
interval has been adopted in prior TMS-EEG studies because it captures the most specific
cortical responses to TMS [14,21], while minimizing contamination from remaining
somatosensory and auditory evoked potentials, which are most pronounced at 180-250 ms [29].
After preprocessing, we applied Group Task-Related Component Analysis (gTRCA) to extract
response components that maximized trial-to-trial reproducibility within each group
(Responders and Non-responders). Figure 1 shows the resulting group-level waveforms and
their corresponding topographical distributions (see Supplementary Figure 1 for individual
data). This approach allowed us to visualize the most consistent neural responses at the group
level [30].
210
TMS-EEG outcomes
To assess global cortical reactivity to TMS, GMFP was calculated as the root-mean-squared
value of the TEP across all electrodes following TMS stimulation [13]. This measure reflects the
overall strength of TEPs across the entire cortex, providing insight into widespread cortical
activity following stimulation [31]. Furthermore, to assess local cortical reactivity to TMS, LMFP
was calculated for the local cluster of electrodes, thereby focusing the analysis on the direct
effects of TMS [32]. Specifically, the central clusters consisted of C1, C3, CP3, and CP1 on the
left hemisphere, and C2, C4, CP2, and CP4 on the right hemisphere. To assess TMS-evoked
oscillatory dynamics in these clusters in the stimulated hemisphere, time-frequency maps were
computed in the 8–45 Hz range using Morlet wavelets with 3.5 cycles [14]. The analysis focused
on three frequency bands: alpha (8–13 Hz), low beta (14–20 Hz), and high beta (21–30 Hz)
based on previous study results [14,21]. Two main parameters were extracted: ERSP and ITC.
ERSP was used to quantify changes in power across time and frequency over trials. It was
computed as the average spectral power ratio of individual EEG trials relative to a pre-stimulus
baseline (-600 to -50 ms). This measure captures how TMS-evoked oscillatory power evolves
across specific frequency bands. Statistical significance was determined by comparing ERSP
values to baseline using bootstrapping (500 permutations, two-sided test, $p < 0.05$, with FDR
correction for multiple comparisons). Significant ERSP values were then averaged across time
points and electrodes to characterize frequency-specific power modulation evoked by TMS
[14,21]. ITC was extracted to evaluate phase-reset of cortical oscillatory activity across trials,
serving as a marker of phase synchronization. ITC was calculated by normalizing the complex-
valued time-frequency data from each trial, averaging across trials, and then computing the
absolute value. Higher ITC values (approaching 1) reflect stronger phase-locking to the TMS
pulse and indicate greater neural synchrony across trials. Significance was assessed using
bootstrapping against the baseline (500 permutations, one-sided test, $p < 0.05$, FDR-corrected),
and significant ITC values were averaged across electrodes, time windows, and frequency bins
[14,21].
238
Normative data
For the same outcomes, a control group of 125 healthy pain-free participants (age range: 20–
71 years; 67 females, 58 males) was used to establish the distribution of values and define the

normal ranges. These TMS-EEG data derive from previously published studies [11,14,21] as well
as ongoing work currently in preparation. This included identifying key statistical parameters,
such as the median value, as well as the 25th and 75th percentiles, which serve as reference
points for comparison with individuals without chronic pain.
246
rTMS Treatment
Magnetic stimulation was delivered using a transcranial magnetic stimulator (MagPro R30,
MagVenture A/S, Farum, Denmark) using a figure-of-eight coil (Cool-B35). Patients were
seated comfortably in a reclining chair and instructed to stay as relaxed as possible during each
session. Each rTMS session involved 30 trains of 100 TMS pulses given over 10 seconds (10
Hz), with 20-second pauses between trains, totaling 3,000 pulses over 15 minutes [7,11,21].
Stimulation intensity was set at 90% of the rMT. If patients reported any sensation in the hand
during treatment, the stimulation intensity was reduced until no sensation was no longer felt.
rMT was checked before each rTMS session to ensure proper stimulation. As for the TMS-
EEG assessment, the rTMS treatment was delivered to the hemisphere contralateral to the side
of the most intense pain. If no dominant pain side was reported, stimulation was delivered over
the left hemisphere. All adverse events and reasons for treatment discontinuation were recorded
throughout the study period. During the rTMS treatment, patients were permitted to continue
their concurrent medication and other treatments. Supplementary Table 1 presents the
pharmacological treatments at baseline and the end of the intervention.
262
Statistical analyses
All statistical analyses were performed using the Statistical Package for Social Sciences (SPSS,
version 25; IBM, Chicago, IL). All outcomes are reported as means and standard deviations,
unless otherwise specified. Statistical significance was set to maintain a two-sided 5% level
(using multiplicity adjustment where appropriate) for comparisons. The normality of data
distributions for each clinical and neurophysiological outcome was assessed in Responders and
Non-responders using the Shapiro–Wilk test and visual inspection of histograms for extreme
violations. Separate two-way mixed model analysis of variance (ANOVA) was used to compare
the clinical outcomes, including pain at the time of the visit (primary clinical outcome), in the
last 24 hours, over the last 7 days, quality of life, BPI (pain severity and pain interference),
HADS, and number of medications. The within-subject factor was Time, and the between-
subject factor was Group. An interaction effect between Group and Time was included in the
models. Bonferroni-corrected multiple pairwise comparisons, and corresponding adjusted 95%
confidence intervals (CIs) and P values were generated and reported in the results. The
Greenhouse–Geisser approach was used to correct for violations of sphericity, and effect sizes
(partial eta-squared $\eta^2_{partial}$) were calculated. For the neurophysiological outcomes, depending
on the results of the normality test, Student's t-tests (for normally distributed data) or the
Mann–Whitney U test (a non-parametric alternative) were used to compare the groups. In total,
8 comparisons were made: GMFP, LMFP, ERSP, and ITC in three frequency bands in the
central stimulated cluster. Levene's test of equality of variance was used, and Cohen's was used
to assess the effect size (Cohen's d). In addition to group comparisons, neurophysiological
values were compared to the normative data. For each measure, the 25th and 75th percentiles of
the normative distribution were calculated to define the interquartile range. The proportion of
participants in each group falling below the 25th percentile was computed and compared using
a Chi-square test of independence (2×2 contingency table) to determine whether Responders
were significantly more likely than Non-responders to fall outside the normative range.
Additionally, Pearson correlation analyses, or Spearman correlation analyses when
distributions were non-normal, were conducted to investigate associations between the
percentage of pain reduction and the four neurophysiological outcomes that showed

differences. Finally, as an exploratory analysis, we performed logistic regression with k-fold
cross-validation (CV) to test whether baseline TMS-EEG features predicted responder status
(Responder = 1, Non-responder = 0), following prior modelling work on TMS and EEG pain
biomarkers [33]. To keep dimensionality small relative to the number of training samples per fold
(given N = 43), we set three predictors: (i) GMFP, included as a broad measure of cortical
excitability; (ii) alpha ERSP in the stimulated cluster, chosen to capture complementary aspects
of alpha-band activity (supported by prior evidence for alpha oscillations in pain processing
[33,34]) and (iii) alpha ITC in the stimulated cluster. All preprocessing and modeling were
embedded in a leakage-safe scikit-learn pipeline (median imputation -> z-scaling -> penalized
logistic regression). We fit elastic-net logistic regression (saga; class_weight=balanced), with
hyperparameters (C, l1_ratio) tuned by an inner stratified CV. Because class counts allowed
balanced splits, we used 5-fold cross-validation. Performance was obtained via stratified nested
CV with an outer 5-fold split repeated 20 times (100 outer evaluations total). For each run, after
inner-CV tuning, we selected a probability cutoff to target 80% sensitivity (as we gave more
weight to correctly identify Responders as opposed to Non-responders), then evaluated this
fixed cutoff on the held-out outer fold. Primary metrics were ROC-AUC (discrimination) and
PR-AUC (positive-class performance with a baseline equal to the responder prevalence), and
sensitivity at the 0.80-operating point. Results were summarized across the 100 outer
evaluations as mean ± SD.
311

Results
Patient characteristics
Of the 43 patients included in the study, 20 were Responders (47%) to M1 rTMS treatment,
and 23 were Non-responders (53%). Responders and Non-responders were similar in age,
height, and sex distribution (demographic and clinical characteristics are summarized in Table
1). The proportion of patients with nociceptive, neuropathic, or nociplastic pain was also
similar, as well as the pain duration (Responders: 9 ± 7 years; Non-responders: 12 ± 10 years).
Among the Responders, 14 patients received left-hemisphere rTMS and 6 received right-
hemisphere rTMS, whereas among Non-responders, 18 received left-hemisphere stimulation
and 5 received right-hemisphere stimulation ($\chi^2 = 0.38$, $p = 0.536$).
322

Clinical trial outcomes
The ANOVA on VAS pain intensity, rated immediately after each rTMS session, revealed a
significant Time × Group interaction ($F_{1,41} = 9.923$; $P < 0.001$; $\eta^2_{partial} = 0.19$). Post-hoc pairwise
comparisons showed that pain VAS scores were lower in Responders at day 3 of the induction
phase ($P = 0.011$; 95% CI [-32.1 -9.9]), week 2 ($P = 0.029$; 95% CI [-26.8 -5.6]), week 3 ($P = 0.001$; 95%CI [-28.9 -10.8])
0.001; 95%CI [-28.9 -10.8]), and from week 5 through week 8 (all $P < 0.001$; week 5: 95% CI
[-35.6 -16.2]; week 6: 95% CI [-35.6 -15.3]; week 7: 95% CI [-36.9 -16.2]; week 8: 95% CI [-
47.7 -30.9]) of the maintenance phase (Figure 2).
For the other clinical outcomes assessed before and after the rTMS intervention, ANOVA
revealed significant Time × Group interactions for average pain VAS intensity in the last 7 days
($F_{1,41} = 20.464$; $P < 0.001$; $\eta^2_{partial} = 0.33$), average pain VAS intensity in the last 24 hours ($F_{1,41} = 8.663$; $P = 0.005$; $\eta^2_{partial} = 0.17$), pain severity ($F_{1,41} = 18.271$; $P < 0.001$; $\eta^2_{partial} = 0.31$), and
= 8.663; P = 0.005; $\eta^2_{partial}$ = 0.17), pain severity ($F_{1,41} = 18.271$; $P < 0.001$; $\eta^2_{partial} = 0.31$), and
quality of life ($F_{1,41} = 8.456$; $P = 0.006$; $\eta^2_{partial} = 0.17$). Post hoc pairwise comparisons indicated
pain reduction in Responders compared with Non-responders over the last 7 days ($P = 0.001$;
95% CI [-31.4 -10.2] – Figure 3A), the last 24 hours ($P = 0.001$; 95% CI [-27.3 -8.2] – Figure
3B), pain severity ($P = 0.001$; 95% CI [-2.7 -0.9] – Figure 3C), and an increase in the quality
of life ($P = 0.046$; 95% CI [0.3 3.0] - Figure 3D).
The PGIC scale revealed that, among Responders, 9 participants (39.1%) reported being "Much
improved," followed by 8 patients (34.8%) reporting "Minimally improved" at the end of

treatment. However, 3 Responders (13.0%) reported "No improvement". Non-responders most
commonly reported "No improvement" (12 participants, 52.2%), followed by "Minimally
improved" (7 participants, 30.4%). One Non-responder (4.3%) rated "Much improved", while
2 (8.7%) as "Minimally worse". For pain interference, the number of body areas with pain, and
HADS, no differences were found (Supplementary Figure 2).
347
Compliance and side effects
Out of the 12 total rTMS sessions, the average number of attendances to therapeutic rTMS
sessions was 11.6 ± 0.6 (97%) among Responders and 11.9 ± 0.2 (99%) among Non-
responders. The maximum number of missed sessions per patient was two. The main side effect
was headache in 14% of sessions, followed by "feel dizzy" in 2%. Majority of side effects (61
of all) were reported during the 5 days of the induction week, and the rest were reported during
the 7 weeks of maintenance.
The number of medications used by patients did not significantly change during treatment in
both groups (Time effect: $F_{1,41} = 4.001$, $P = 0.052$, $\eta^2_{partial} = 0.09$; Time × Group interaction:
$F_{1,41} = 0.621$, $P = 0.435$, $\eta^2_{partial} = 0.02$; Supplementary Figure 2). The list of medications is
provided in Supplementary Table 1.
359
TMS-EEG and rTMS intensities
The intensity of TMS applied to the M1 during EEG recording was $50 \pm 8.8\%$ for Responders
(rMT $55 \pm 10\%$) and $45.7 \pm 5.7\%$ for Non-responders ($50 \pm 6\%$), with no significant difference
between the groups ($t_{41} = 1.636$; $P = 0.110$; Cohen's $d = 0.500$). The average rTMS intensity
used for the treatments was $42.6 \pm 7.3\%$ for the Responders (starting from $42.4 \pm 7.1\%$ at D1 and
finishing at $42.8 \pm 7.4\%$ at W8) and $38.3 \pm 7.4\%$ for the Non-responders (starting from
$38.4 \pm 10.2\%$ at D1 and finishing at $40.8 \pm 6.8\%$ at W8).
367
TMS-EEG global and local mean field power
GMFP values were lower in Responders compared to Non-responders ($t_{41} = 2.571$; $P = 0.011$;
Cohen's $d = 0.786$; Figure 4A). A total of 65% of Responders and 44% of Non-responders had
values that fell below the 25th percentile of the normative dataset ($\chi^2 = 1.99$; $P = 0.158$).
Significantly lower LMFP was also identified in the Responder group in the stimulated cluster
($t_{41} = 2.858$, $P = 0.007$, Cohen's $d = 0.874$; Figure 4B). 55% of Responders and 26% of Non-
responders had values below the 25th percentile of the normative dataset ($\chi^2 = 3.74$; $P = 0.053$).
375
Time-frequency analysis
In Responders compared with Non-responders, lower alpha ERSP (Mann-Whitney U-
Test = 2.070, $P = 0.038$; Figure 4E) and alpha ITC ($t_{41} = 2.374$, $P = 0.022$, 95% CI [-0.24 -0.02],
Cohen's $d = 0.726$; Figure 4F) were found. For ERSP, 70% of Responders and 48% of Non-
responders had values below the 25th percentile of the normative dataset ($\chi^2 = 2.16$; $P = 0.142$).
For ITC, 55% of Responders and 22% of Non-responders fell below the 25th percentile ($\chi^2 =$
5.07; $P = 0.021$).
383
384   *Correlation analysis*
385   The percentage of pain reduction significantly correlated with GMFP (Pearson $r = 0.40$,
386    $P = 0.008$ - Figure 5A), LMFP (Pearson $r = 0.38$, $P = 0.012$ - Figure 5B), alpha ERSP (Pearson
387    $r = 0.31$, $P = 0.046$ - Figure 5C) and alpha ITC (Spearman $r = 0.35$, $P = 0.020$ - Figure 5D).
388
389   *Logistic regression for the prediction of response to rTMS*
390   Using three predictors: GMFP, alpha ERSP, and alpha ITC, the model achieved acceptable
391   performance across the 100 folds. On average, ROC-AUC was 0.70 ± 0.18 and PR-AUC was

0.76 ± 0.15 (Figure 6 shows the means and distributions). Table 2 summarizes the standardized coefficients of the predictors, along with their median values and interquartile ranges. At the cut-off of 80% chosen to maximize sensitivity, sensitivity was 0.74 ± 0.26 with the expected drop in specificity (0.48 ± 0.28). These results suggest, on average, the model ranked a randomly chosen responder above a non-responder 74% of the time. Moreover, with Responders comprising ~47% of the cohort, as the threshold for identifying Responders is varied, the model maintains ~76% precision on average, versus a 47% baseline from random guessing, indicating acceptable positive-class performance. Lastly, using a cutoff chosen on the training split to target 80% sensitivity, the realized sensitivity on held-out data averaged 0.74 (≈ 3 in 4 Responders correctly identified).

Discussion

The current study showed that patients responding to M1 rTMS treatment had, upon pre-therapy probing of the planned therapeutic target, lower global and local cortical reactivity. Furthermore, these patients showed reduced local TMS-evoked oscillatory dynamics in the alpha frequency compared to patients who did not respond to the rTMS treatment. These electrophysiological changes correlated with the percentage of pain reduction. Taken together, the findings suggest that lower pre-treatment global and local cortical reactivity and reduced alpha-band power and synchronization in the targeted region may reflect a neurophysiological state more prone to undergo adaptive neuroplastic changes, thereby increasing the likelihood of a favorable response to M1 rTMS.

Clinical outcomes

In the current study, 47% of patients (20 out of 43) were responders to M1 rTMS. This percentage aligns with the recent findings of Quesada et al. [8], who reported a 47% response rate (17 out of 36 patients) and Hodaj et al. (27 out of 57 patients) [35], both using a ≥30% pain reduction criterion. Attal et al. identified 44% of patients as responders (22 out of 49) using a higher threshold of ≥ 50% pain reduction. Applying a ≥ 50% pain reduction threshold to our data, the responder rate was 39.5% (17 out of 43). Consistent with all previous studies [8,35–37], analgesic effects of rTMS mainly appeared after several days of rTMS treatment, indicating a cumulative therapeutic effect. In addition to reductions in pain intensity reported during the visit, the present study also found decreased pain severity, less pain in the previous 24 hours and 1 week, and improved quality of life after rTMS. Conversely, no significant changes were observed in pain interference, the number of body areas affected by pain, depression, anxiety, or medication use. Furthermore, it is important to note that a few patients in the Non-responder group reduced their medication use despite reporting similar pain intensity levels, suggesting a possible analgesic effect of the treatment, something previously reported in multicenter randomized sham-controlled trials on rTMS for chronic pain [6]. An important finding of the current study is that the analgesic effects of rTMS were observed in responders across various types of chronic pain conditions, not limited to neuropathic pain [7] and fibromyalgia [36]. Therefore, while most previous studies have focused on rTMS efficacy in specific chronic pain conditions [7,8,35,37], our results suggest that similar therapeutic benefits may extend to multiple pain etiologies.

Global and local cortical reactivity

Consistent with the hypothesis that future responders to M1 stimulation had pre-treatment reductions in cortical reactivity, we found significantly reduced GMFP in responders compared with non-responders. These findings were confirmed by the presence of more GMFP results falling below normative ranges in the responder group. Cortical reactivity can be defined as the relationship between the strength of the stimulus and the magnitude of the evoked response,

with GMFP representing the global amplitude of TMS-evoked brain activity across the scalp [31,38]. The fact that responders also showed lower LMFP values in the stimulated hemisphere suggests that reduced local cortical reactivity may facilitate the therapeutic effects of rTMS. This may reflect a greater capacity for excitability enhancement through rTMS-induced plasticity, consistent with homeostatic mechanisms [15], which are more salient and likely to occur when baseline reactivity is low. Taking together, these findings suggest that reduced global and local M1 reactivity to single probing pulses of TMS may create a more favorable condition for upregulation through rTMS. In contrast, when global and local cortical reactivity is closer to normative levels, the capacity for further modulation may be limited, potentially resulting in weaker therapeutic effects.

Local oscillatory dynamics

In the current study, patients responding to the rTMS treatment showed reduced TMS-evoked alpha power and phase coherence over the stimulated cortical area compared to non-responders. TMS-evoked oscillatory dynamics refer to the rhythmic activity generated by neuronal populations within a specific cortical region in response to the TMS stimuli, characterized by frequency-specific changes in power and phase from the pre-stimulus baseline [39]. These TMS-evoked oscillatory dynamics provide insight into the functional state of the underlying cortical circuitry [40]. The synchronization of the rhythmic activity across multiple frequency bands enables communication between spatially distributed brain regions through interareal phase locking [41]. This process creates time windows during which information can be integrated concurrently in sparse neuronal clusters, allowing complexity to emerge [42]. The observed efficacy of rTMS in chronic pain patients with the lowest alpha-band oscillatory dynamics may be interpreted in several ways. One possibility is that this pattern reflects bigger disruptions in thalamocortical and corticocortical synchronization [43], potentially creating a neurophysiological environment more conducive to plastic reorganization by rTMS. Alpha-band oscillations are thought to propagate from higher-order to lower-order cortical regions and toward the thalamus via short-range supragranular feedback projections [44]. A reduction in alpha ERSP may indicate weakened local synchronization of neural circuits, while lower ITC suggests impaired phase alignment across repeated stimuli, indicative of reduced temporal precision and rhythmic coordination locally. Together, diminished alpha synchrony and temporal coordination may reflect a more "permissive" cortical state, facilitating greater susceptibility to modulation and plasticity through rTMS. Previous studies have shown that neuropathic pain and fibromyalgia are related to lower intracortical inhibition and facilitation [45], a sign of reduced GABA tonus within M1. rTMS to M1 has been shown to restore defective GABA-dependent intracortical inhibition coupled with pain improvement [36]. Alpha oscillations are implicated in large-scale cortical network integration, with phase locking and coherence modulating both local and long-range neural interactions [46,47].

An alternative explanation is that 10 Hz rTMS may be particularly effective in a specific subpopulation of patients who exhibit altered cortical oscillatory profiles near the alpha band frequency. Increasing evidence suggests that some individuals with chronic pain exhibit EEG frequency abnormalities, including reduced α- and low β-band activity (8–20 Hz), or increased high β-band activity (20–30 Hz) [48]. These frequency alterations are highly heterogeneous and highly variable across chronic pain patients [49]. A recent study reported that the closer proximity of the resting-state peak alpha frequency to the 10 Hz rTMS frequency predicted a greater analgesic benefit of 10 Hz rTMS during tonic experimental pain [50], suggesting that rTMS may be more effective when delivered at or near an individual's intrinsic M1 "natural" oscillatory frequency. In the current study, the observed reduction in alpha power and phase coherence among responders may reflect a cortical state closer to an optimal oscillatory dynamic, in which 10 Hz rTMS can more effectively induce plasticity and therapeutic effects. Conversely, in non-

responders, stimulation at a different frequency may be necessary to engage their intrinsic oscillatory dynamics and achieve comparable analgesic outcomes.

Another possibility concerns the target of rTMS. Individual differences in cortical pain processing may determine which cortical target, such as the M1, dorsolateral prefrontal cortex, or posterosuperior insula, is most responsive to neuromodulatory interventions. Experimental pain models in healthy participants have shown a polarization of pain-induced cortical suppression, with some individuals exhibiting reductions in excitability in one cortical area but not the other, reflecting marked inter-individual variability in pain-related cortical dynamics [13]. It is possible that, in the current study, the most appropriate cortical target for rTMS in non-responders was not M1, but stimulating an alternative cortical target may have yielded a therapeutic effect.

Potential clinical applications
We used a supervised machine learning approach to test whether three baseline TMS-EEG features (ITC and ERSP Alpha in the stimulated cluster, and GMFP) could predict responder status to rTMS. We used a leakage-safe, stratified nested cross-validation framework with reduced dimensionality and penalized logistic regression to mitigate overfitting in a small sample size. Across 100 outer evaluations, discrimination and positive class performance of the model were acceptable on average, with lower ITC, ERSP, and GMFP predicting a higher likelihood of responding to rTMS. Moreover, with a cutoff chosen on the training split to target 80% sensitivity, the realized held-out sensitivity was acceptable, with the expected trade-off in specificity. These findings suggest that baseline TMS-EEG features may contain predictive signals for identifying rTMS responders, with an accuracy of about 75% (≈3 in 4 correctly identified), compared to the ~45% response rate typically observed with the current use of M1 rTMS. In principle, an optimized model could support treatment selection by allowing clinicians to input patient-level values and estimate response probability, thereby informing individualized care. However, we do not claim to have such a model yet. Given the modest sample and exploratory design (reduced dimensionality, penalized logistic regression, and stratified nested cross-validation), results should be interpreted cautiously and viewed as proof-of-concept. Larger, prospectively designed studies in clinical populations are needed before clinical deployment.

Conclusion
This study provides the first evidence that baseline neurophysiological markers of global and local cortical reactivity and oscillatory dynamics recorded by probing the same target to be used for treatment is feasible and may predict the long-term response to rTMS in people with chronic pain. The presence of local decreased cortical responses may indicate a more maladaptive or dysfunctional network hub that is more likely to be positively influenced by neuromodulation. These findings support the potential utility of cortical excitability and synchronization measures as a predictive strategy for identifying individuals most likely to benefit from rTMS interventions. Prospective trials with larger sample sizes specifically designed for this purpose are needed to test this new perspective and research paradigm.

Declaration of conflict of interest

Acknowledgements

The current study was supported by an ERC Horizon Europe Consolidator grant (PersoNINpain 101087925) and a Novo Nordisk grant (Grant NNF21OC0072828). The Center

542 for Neuroplasticity and Pain is supported by the Danish National Research Foundation
543 (DNRF121). Relevant to this study, Thomas Graven-Nielsen has received funding from the
544 Lundbeck Foundation (R441-2023-232).
545

Author contribution

Enrico De Martino: Conceptualization, Methodology, Formal analysis, Investigation, Data
curation, Visualization, Writing- Original draft preparation; Project administration; **Margit**
Midtgaard Bach: Investigation, Data curation, Writing - Review & Editing; **Anne Jakobsen**:
Investigation, Writing - Review & Editing; **Stian Ingemann-Molden**: Investigation, Writing -
Review & Editing; **Nahian S. Chowdhury**: Data curation, Writing - Review & Editing; **Bruno**
Andry Nascimento Couto: Software, Formal analysis, Data Curation, Visualization, Writing
- Review & Editing; **Adenauer Girardi Casali**: Software; Writing - Review & Editing;
Thomas Graven-Nielsen: Writing - Review & Editing; **Daniel Ciampi de Andrade**:
Conceptualization, Methodology, Visualization, Writing - Review & Editing; Project
administration, Funding acquisition.

References

1. Lefaucheur JP, Aleman A, Baeken C, et al. Evidence-based guidelines on the therapeutic use of repetitive transcranial magnetic stimulation (rTMS): An update (2014–2018). *Clinical Neurophysiology*. 2020;131(2):474-528. doi:10.1016/j.clinph.2019.11.002

2. Siebner HR, Funke K, Aberra AS, et al. Transcranial magnetic stimulation of the brain: What is stimulated? – A consensus and critical position paper. *Clinical Neurophysiology.Elsevier Ireland Ltd*. 2022;140:59-97. doi:10.1016/j.clinph.2022.04.022

3. Klomjai W, Katz R, Lackmy-Vallée A. Basic principles of transcranial magnetic stimulation (TMS) and repetitive TMS (rTMS). *Ann Phys Rehabil Med.Elsevier Masson SAS*. 2015;58(4):208-213. doi:10.1016/j.rehab.2015.05.005

4. Soliman N, Moisset X, Ferraro MC, et al. Pharmacotherapy and non-invasive neuromodulation for neuropathic pain: a systematic review and meta-analysis. *Lancet Neurol*. 2025;24(5):413-428. doi:10.1016/S1474-4422(25)00068-7

5. Moisset X, Pereira B, Ciampi de Andrade D, Fontaine D, Lantéri-Minet M, Mawet J. Neuromodulation techniques for acute and preventive migraine treatment: a systematic review and meta-analysis of randomized controlled trials. *Journal of Headache and Pain.BioMed Central Ltd*. 2020;21(1). doi:10.1186/s10194-020-01204-4

6. Silva VA, Baptista AF, Fonseca AS, et al. Motor cortex repetitive transcranial magnetic stimulation in fibromyalgia: a multicentre randomised controlled trial. *Br J Anaesth*. 2025;134(6):1756-1764. doi:10.1016/j.bja.2024.12.045

7. Attal N, Poindessous-Jazat F, De Chauvigny E, et al. Repetitive transcranial magnetic stimulation for neuropathic pain: a randomized multicentre sham-controlled trial. *Brain*. Published online 2021:3328-3339. doi:10.1093/brain/awab208

8. Quesada C, Pommier B, Fauchon C, et al. New procedure of high-frequency repetitive transcranial magnetic stimulation for central neuropathic pain: A placebo-controlled randomized crossover study. *Pain*. 2020;161(4):718-728. doi:10.1097/j.pain.0000000000001760

9. Ciampi de Andrade D, García-Larrea L. Beyond trial-and-error: Individualizing therapeutic transcranial neuromodulation for chronic pain. *European Journal of Pain (United Kingdom).John Wiley and Sons Inc*. 2023;27(9):1065-1083. doi:10.1002/ejp.2164

10. Tremblay S, Rogasch NC, Premoli I, et al. Clinical utility and prospective of TMS – EEG. *Clinical Neurophysiology*. 2019;130:802-844. doi:10.1016/j.clinph.2019.01.001

11. De Martino E, Nascimento Couto BA, Jakobsen A, et al. Pretreatment TMS-EEG connectivity assessment as a potential predictor of rTMS effectiveness in chronic pain: A feasibility pilot study. *Clinical Neurophysiology*. Published online 2025. doi:10.1016/j.clinph.2025.02.268

12. Russo S, Rosanova M. Stimulate to probe, stimulate to modulate: Single-pulse brain stimulation to titrate neuromodulation. *Clinical Neurophysiology.Elsevier Ireland Ltd*. Preprint posted online August 1, 2025. doi:10.1016/j.clinph.2025.2110737

13. De Martino E, Casali A, Casarotto S, et al. Acute pain drives different effects on
local and global cortical excitability in motor and prefrontal areas: insights into
interregional and interpersonal differences in pain processing. *Cerebral Cortex*.
2023;33(18):9986-9996. doi:10.1093/cercor/bhad259

14. De Martino E, Casali A, Casarotto S, et al. Evoked oscillatory cortical activity
during acute pain: Probing brain in pain by transcranial magnetic stimulation
combined with electroencephalogram. *Hum Brain Mapp*. 2024;45(6).
doi:10.1002/hbm.26679

15. Bienenstock EL, Cooper LN, Munro PW. Theory for the development of neuron
selectivity: orientation specificity and binocular interaction in visual cortex. *The
Journal of Neurosci ence*. 1982;2(1):32-48.

16. British Pain Society. *Outcome Measures: Recommendations for Use in Adult Pain
Management Services*.; 2019. Accessed July 16, 2025.
https://www.britishpainsociety.org/static/uploads/resources/files/Outcome_Mea
sures_January_2019.pdf

17. Tan G, Jensen MP, Thornby JI, Shanti BF. Validation of the brief pain inventory for
chronic nonmalignant pain. *Journal of Pain*. 2004;5(2):133-137.
doi:10.1016/j.jpain.2003.12.005

18. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta
Psychiatr Scand*. 1983;67(6):361-370. http://www.hqlo.com/content/1/1/29

19. Group E. *EuroQol*-a New Facility for the Measurement of Health-Related Quality
of Life*.; 1990.

20. Hurst H, Bolton J. Assessing the clinical significance of change scores recorded
on subjective outcome measures. *J Manipulative Physiol Ther*. 2004;27(1):26-35.
doi:10.1016/j.jmpt.2003.11.003

21. De Martino E, Casali AG, Nascimento Couto BA, Graven-Nielsen T, Ciampi de
Andrade D. Increase in beta frequency phase synchronization and power after a
session of high frequency repetitive transcranial magnetic stimulation to the
primary motor cortex. *Neurotherapeutics*. Published online November
2024:e00497. doi:10.1016/j.neurot.2024.e00497

22. Russo S, Sarasso S, Puglisi GE, et al. TAAC - TMS Adaptable Auditory Control: A
universal tool to mask TMS clicks. *J Neurosci Methods*. 2022;370:109491.
doi:10.1016/j.jneumeth.2022.109491

23. Souza VH, Matsuda RH, Peres ASC, et al. Development and characterization of
the InVesalius Navigator software for navigated transcranial magnetic
stimulation. *J Neurosci Methods*. 2018;309:109-120.
doi:10.1016/j.jneumeth.2018.08.023

24. Fecchio M, Pigorini A, Comanducci A, et al. The spectral features of EEG
responses to transcranial magnetic stimulation of the primary motor cortex
depend on the amplitude of the motor evoked potentials. *PLoS One*.
2017;12(9):1-15. doi:10.1371/journal.pone.0184910

25. Casarotto S, Fecchio M, Rosanova M, et al. The rt-TEP tool: real-time visualization
of TMS-Evoked Potentials to maximize cortical activation and minimize artifacts. *J
Neurosci Methods*. 2022;370:109486. doi:10.1016/j.jneumeth.2022.109486

26. Casarotto S, Lauro LJR, Bellina V, et al. EEG responses to TMS are sensitive to
changes in the perturbation parameters and repeatable over time. *PLoS One*.
2010;5(4). doi:10.1371/journal.pone.0010281

27.   Makeig S, Westerfield M, Jung T, et al. Dynamic Brain Sources of Visual Evoked
651   Responses. *Science (1979)*. 2002;295(January):690-694.

652   28.   Delorme A, Makeig S. EEGLAB : an open source toolbox for analysis of single-trial
653   EEG dynamics including independent component analysis. *J Neurosci Methods*.
654   2004;134:9-21. doi:10.1016/j.jneumeth.2003.10.009

655   29.   Rocchi L, Di A, Brown K, et al. Brain Stimulation Disentangling EEG responses to
656   TMS due to cortical and peripheral activations. *Brain Stimul*. 2021;14:4-18.
657   doi:10.1016/j.brs.2020.10.011

658   30.   Couto BA, Fecchio M, Russo S, et al. Extracting Reproducible Components from
659   Electroencephalographic Responses to Transcranial Magnetic Stimulation with
660   Group Task-Related Component Analysis. *bioRxiv*. Published online June 6, 2025.
661   doi:https://doi.org/10.1101/2025.06.02.657489

662   31.   Kähkönen S, Wilenius J, Komssi S, Ilmoniemi RJ. Distinct differences in cortical
663   reactivity of motor and prefrontal cortices to magnetic stimulation. *Clinical
664   Neurophysiology*. 2004;115(3):583-588. doi:10.1016/j.clinph.2003.10.032

665   32.   Kähkönen S, Komssi S, Wilenius J, Ilmoniemi RJ. Prefrontal TMS produces smaller
666   EEG responses than motor-cortex TMS: Implications for rTMS treatment in
667   depression. *Psychopharmacology (Berl)*. 2005;181(1):16-20. doi:10.1007/s00213-
668   005-2197-3

669   33.   Chowdhury NS, Bi C, Furman AJ, et al. Predicting Individual Pain Sensitivity Using
670   a Novel Cortical Biomarker Signature. *JAMA Neurol*. 2025;82(3):237-246.
671   doi:10.1001/jamaneurol.2024.4857

672   34.   Furman AJ, Prokhorenko M, Keaser ML, et al. Sensorimotor Peak Alpha Frequency
673   Is a Reliable Biomarker of Prolonged Pain Sensitivity. *Cerebral Cortex*. Published
674   online 2020:1-14. doi:10.1093/cercor/bhaa124

675   35.   Hodaj H, Payen JF, Hodaj E, et al. Long-term treatment of chronic orofacial,
676   pudendal, and central neuropathic limb pain with repetitive transcranial magnetic
677   stimulation of the motor cortex. *Clinical Neurophysiology*. 2020;131(7):1423-
678   1432. doi:10.1016/j.clinph.2020.03.022

679   36.   Mhalla A, Baudic S, Ciampi de Andrade D, et al. Long-term maintenance of the
680   analgesic effects of transcranial magnetic stimulation in fibromyalgia. *Pain*.
681   2011;152(7):1478-1485. doi:10.1016/j.pain.2011.01.034

682   37.   Hosomi K, Sugiyama K, Nakamura Y, et al. A randomized controlled trial of 5 daily
683   sessions and continuous trial of 4 weekly sessions of repetitive transcranial
684   magnetic stimulation for neuropathic pain. *Pain*. 2020;161(2):351-360.
685   doi:10.1097/j.pain.0000000000001712

686   38.   Komssi S, Kähkönen S, Ilmoniemi RJ. The Effect of Stimulus Intensity on Brain
687   Responses Evoked by Transcranial Magnetic Stimulation. *Hum Brain Mapp*.
688   2004;21(3):154-164. doi:10.1002/hbm.10159

689   39.   Makeig S, Westerfield M, Jung TP, et al. Dynamic Brain Sources of Visual Evoked
690   Responses. *Science (1979)*. 2002;295(5555):690-694.

691   40.   Rosanova M, Casali A, Bellina V, Resta F, Mariotti M, Massimini M. Natural
692   frequencies of human corticothalamic circuits. *Journal of Neuroscience*.
693   2009;29(24):7679-7685. doi:10.1523/JNEUROSCI.0445-09.2009

694   41.   Buzsáki G, Draguhn A. Neuronal Oscillations in Cortical Networks. *Science
695   (1979)*. 2004;304(5679):1926-1929. doi:10.1126/science.1099745

696   42.   Fries P. A mechanism for cognitive dynamics: Neuronal communication through
697   neuronal coherence. *Trends Cogn Sci*. 2005;9(10):474-480.
698   doi:10.1016/j.tics.2005.08.011

699   43.   Sarnthein J, Stern J, Aufenberg C, Rousson V, Jeanmonod D. Increased EEG power
700   and slowed dominant frequency in patients with neurogenic pain. *Brain*.
701   2006;129(1):55-64. doi:10.1093/brain/awh631

702   44.   Halgren M, Ulbert I, Bastuji H, et al. The generation and propagation of the human
703   alpha rhythm. *Proceedings of the National Academy of Sciences (PNAS)*.
704   2019;116:23772-23782. doi:10.6084/m9.figshare.9927125.v2

705   45.   Mhalla A, de Andrade DC, Baudic S, Perrot S, Bouhassira D. Alteration of cortical
706   excitability in patients with fibromyalgia. *Pain*. 2010;149(3):495-500.
707   doi:10.1016/j.pain.2010.03.009

708   46.   Sadaghiani S, Scheeringa R, Lehongre K, et al. Alpha-band phase synchrony is
709   related to activity in the fronto-parietal adaptive control network. *Journal of*
710   *Neuroscience*. 2012;32(41):14305-14310. doi:10.1523/JNEUROSCI.1358-12.2012

711   47.   Chapeton JI, Haque R, Wittig JH, Inati SK, Zaghloul KA. Large-Scale
712   Communication in the Human Brain Is Rhythmically Modulated through Alpha
713   Coherence. *Current Biology*. 2019;29(17):2801-2811.e5.
714   doi:10.1016/j.cub.2019.07.014

715   48.   Mussigmann T, Bardel B, Lefaucheur JP. Resting-state electroencephalography
716   (EEG) biomarkers of chronic neuropathic pain. A systematic review.
717   *Neuroimage*.*Academic Press Inc*. 2022;258.
718   doi:10.1016/j.neuroimage.2022.119351

719   49.   Bott FS, Zebhauser PT, Hohn VD, et al. EEG-based analysis of intrinsic brain
720   network function in chronic pain: Insights from a comprehensive multi-data set
721   study. *bioRxiv*. Published online August 12, 2024. doi:10.1101/2024.08.12.607584

722   50.   Millard SK, Chiang AKI, Chowdhury N, et al. Peak alpha frequency is not
723   significantly altered by five days of experimental pain and repetitive transcranial
724   stimulation of the left dorsolateral prefrontal cortex. *bioRxiv*. Published online
725   June 14, 2024. doi:10.1101/2024.06.14.599003

726
727

Figure legend

Figure 1: A: Normalized eigenvalue spectra of gTRCA components for Responders (blue) and Non-responders (red), indicating the amount of reproducible task-related variance (Eigenvalue) captured by each component. **B:** Scalp topographies of the first gTRCA component show the spatial distribution of the most reproducible TMS-evoked responses in each group. **C:** Time series projections of the dominant component for Responders (blue) and Non-responders (red), with shaded areas representing the standard error of the mean. The vertical dashed line indicates TMS onset (0 s).

Figure 2: Pain intensity (Visual Analogue Scale, VAS: 0–100) across time points for Responders (blue) and Non-responders (red). Squares and error bars represent means ± standard deviations, with individual participant data shown as dots. The pre-treatment period includes assessments conducted 7 days and 24 hours before the TMS-EEG session, as well as 24 hours before the first rTMS session (shaded in light gray). The induction phase spans five consecutive treatment days (D1–D5, shaded in light yellow), followed by a 7-week maintenance phase (V2–V8, shaded in light green), during which pain was assessed immediately after each rTMS session. Asterisks indicate statistically significant differences between groups at specific time points after Bonferroni correction (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 3: Secondary clinical outcomes in Responders (blue) and Non-responders (red) before and after rTMS treatment. Bars represent means ± standard deviations, with individual participant data shown as dots. **A)** Pain intensity 7 days before and after treatment; **B)** Pain intensity 24 hours before and after treatment; **C)** Pain severity; **D)** Quality of life. Significant post-treatment between-group differences are indicated (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

Figure 4: Neurophysiological outcomes in Responders (blue) and Non-responders (red) before rTMS treatment. Boxplots represent median ± quartiles, with individual participant data shown as dots. **A)** Global mean field power; **B)** Local mean field power; **C)** Event-related spectral perturbation (ERSP) in the alpha band; **D)** Inter-trial coherence (ITC) in the alpha band. Significant post-treatment between-group differences are indicated (*$p < 0.05$, **$p < 0.01$).

Figure 5: Correlation between percentage change in pain intensity and neurophysiological outcomes. Responders are shown in blue; Non-responders in red. **A)** Global mean field power (μV); **B)** Local mean field power (μV); **C)** Event-related spectral perturbation (ERSP) in the alpha band (dB). **F)** Inter-trial coherence (ITC) in the alpha band (0–1).

Figure 6: A) Mean receiver operating characteristic (ROC) curve with area under the curve (AUC = 0.70) and 95% confidence interval (CI). **B)** Mean precision–recall (PR) curve with average precision (PR = 0.47) and 95% CI; the dashed line indicates the baseline precision. **C)** Distribution of ROC-AUC values across 100 outer folds, with the mean indicated by the dashed line. **D)** Distribution of PR-AUC values across 100 outer folds, with the mean indicated by the dashed line.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 6

Table
Table 1
Demographic and clinical characteristics of Responders (N = 20) and Non-responders (N = 23)

	Responders	Non-responders
Age (years)	53±12	52±13
Height (cm)	173±10	174±8
Weight (kg)	91±20	85±18
Male	8	9
Female	12	14
Nociceptive pain (n)	10	8
Neuropathic pain (n)	4	7
Nociplastic pain (n)	6	8
Pain duration (years)	9±7	12±10

Table 2

Standardized coefficients and descriptive statistics of predictors used in the logistic regression model.

Outcome	Standardized Coefficient	Cut-off at Other Variable Means (median)	IQR (25th–75th)
GMFP	-0.037	0.163	[0.152, 0.176]
Alpha ITC	-0.036	0.673	[0.571, 0.722]
Alpha ERSP	-0.020	0.800	[0.662, 0.975]

GMFP – global mean field power; ERSP – event-related spectral perturbation; ITC - inter-trial coherence