1

Relationship between psychosocial factors and cardiovascular autonomic function among junior doctors: A cross-sectional study

Short title: Psychosocial factors and cardiovascular autonomic function among junior doctors

Kofi Tekyi Asamoah[1,2,3*], Richard Dei-Asamoa[4], Eugene Baafi Ampofo[1,2], Felix Razak Awindaogo[2], Francis Agyekum[2,5], Alfred Doku[1,2,5], Sammy Ohene[6], Eugene Amable[5]

1- National Cardiothoracic Centre, Korle-Bu Teaching Hospital, Accra, Ghana

2- Department of Medicine and Therapeutics, Korle-Bu Teaching Hospital, Accra, Ghana

3- Division of Cardiology, Department of Medicine, Yonsei University College of Medicine, Seoul, South Korea

4- Department of Psychiatry, Korle-Bu Teaching Hospital, Accra, Ghana

5- Department of Medicine, University of Ghana School of Medicine and Dentistry, Accra, Ghana

6- Department of Psychiatry, University of Ghana School of Medicine and Dentistry, Accra, Ghana

Corresponding author: asamoahfiifi@gmail.com (KTA)

Abstract

Background: Psychosocial factors such as stress, anxiety, and depression, are important causes of CVD by contributing to autonomic dysfunction, predisposing to endothelial dysfunction, arrhythmogenesis, among others. Doctors are at a high risk of psychosocial distress due to factors such as heavy workload and taking end-of-life care decisions. We sought to determine the prevalence of psychosocial factors among junior doctors and explore their relationship with cardiovascular autonomic function.

Methods: A cross-sectional study was conducted among junior doctors at the Korle-bu Teaching Hospital from April to August 2023. Data were collected using an electronic questionnaire and a treadmill stress test. The Depression, Anxiety, and Stress Scale -21 items (DASS-21) was incorporated into the questionnaire to assess depressive, anxiety and stress symptoms. Cardiovascular autonomic function was determined by heart rate recovery and chronotropic index. The data collected were analysed with the Stata version 16.1, using descriptive and inferential statistics.

Results: 244 junior doctors participated in the study, with a mean age of 31.46. The prevalences of depressive, anxiety and stress symptoms were 29.51%, 48.77% and 48.36% respectively, correlating moderately with one another. Chronotropic index was abnormal for 9.43% of respondents (95% CI = 6.07% - 13.81%) and heart rate recovery-at-1-minute (HRR_1), 3.28% (1.43% - 6.36%). There were no differences in autonomic function profile based on tertiles of psychosocial factors using the Kruskal-Wallis test and one-way analysis of variance. Stress scores were associated with increased HRR_1 (β per SD 1.53 bpm, 95% CI 0.03 - 3.02). Higher depression scores were associated with lower odds of chronotropic incompetence (adjusted OR 0.44, 95% CI 0.22 - 0.88).

Conclusion: There is a high prevalence of depressive, anxiety and stress symptoms among junior doctors. This is however associated with augmented rather than impaired cardiovascular autonomic function.

Introduction

Cardiovascular disease (CVD) is a leading cause of morbidity and death in sub-Saharan Africa (SSA) [1], accounting for a significant disability-adjusted life year (DALY) burden [2]. CVD rates have increased in low- and middle-income countries (LMICs) by up to 50% over the last 3 decades [1,2].

Psychosocial stressors, which include stress, anxiety and depression, contribute to CVD symptoms acutely through sympathetic nervous system (SNS) activation [3,4], and chronically through various mechanisms [4] (Fig. 1). They also worsen pre-existing CVD like heart failure and myocardial infarction [5]. Acute stressors evoke a physiological response through the SNS [4,6], followed over time by a psychological response [7]. The physiological response to stress involves two major biochemical pathways; the SNS and the hypothalamic-pituitary-adrenal (HPA) axis [4,6–8]. If personal coping mechanisms are unable to resolve the psychosocial disturbance, the chronic SNS activation in response to stress will result in noradrenaline secretion, activation of the renin-angiotensin-aldosterone system (RAAS), and the elaboration of inflammatory cells and cytokines [8,9]. Conversely, the parasympathetic nervous system (PSNS) output diminishes [7], upsetting sympathovagal balance and increasing resting heart rate (HR) and blood pressure (BP) [4,10]. The resting HR may therefore be an index of one's cardiovascular autonomic function, though Lankhorst et al.[11]

and Bustamante-Sánchez et al.[12] identified heart rate recovery (HRR) as an earlier and

more sensitive sign of autonomic disturbance than resting HR (Fig 1).

Fig 1. Diagram showing the study's conceptual framework from psychosocial stress exposure

to development of cardiovascular disease

SNS-induced vasoconstriction alters laminar vascular blood flow [6] through reduced

endothelial nitric oxide synthase (eNOS) activity [13]. Cortisol, produced by the HPA axis,

also inhibits eNOS [13]. Nitric oxide depletion favours platelet aggregation, elaboration of E-

selectin and vascular cell adhesion molecule [14], creating a proinflammatory milieu that

leads to endothelial dysfunction and increased atherogenesis [4,14]. The INTERHEART

study reported that psychosocial factors confer an almost 3-fold increase in risk of acute

myocardial infarction [15]. The INTERSTROKE study also reported that permanent or

repeated episodes of stress increased the incidence of stroke, more with occupational stress

than other forms [16].

Sympathovagal imbalance may be assessed by various non-invasive means. These include

HRR and chronotropic index [17]. HRR is assessed using the treadmill stress test (TMT) and

is the difference between HR at peak exercise and HR at various stages of recovery [17].

Within thirty seconds of exercise cessation, PSNS output rises rapidly and is reflective of

cardiovascular autonomic function [17], peaking at approximately 2 minutes. The HR and BP

therefore drop significantly within the first 2 minutes and plateau. A suboptimal HR drop

after 2 minutes increases cardiovascular mortality [18]. This is expected to be faster among

younger people [19]. Reduced HRR has however also been reported among obese, otherwise

healthy females [20]. Chronotropic index is the percentage of one's heart rate reserve for age

utilised during exercise [21,22]. Chronotropic incompetence occurs when the chronotropic

index is less than 0.8, or 0.62 in patients taking beta blockers [23,24]. It gives an

individualised age-based assessment of cardiovascular autonomic function. Chronotropic

index alone better predicts autonomic dysfunction and cardiovascular risk than HRR, while

derangements in both parameters confer significantly higher cardiovascular risk [23].

Psychosocial factors are more prevalent in younger populations [25]. Among health workers,

factors contributing to psychosocial stress include shift systems (including night and weekend

shifts), emergency and/or critical care duties, taking end-of-life care decisions and relaying

such decisions to patients and relatives, and having regular inpatient and outpatient services

[26,27]. Few studies have reported the psychosocial factors experienced by junior doctors in

Ghana, and even less is known about their association with changes in cardiovascular

physiology. There is therefore a potential risk to this population that is uncharacterised.

Understanding the psychosocial profile of junior doctors is relevant for developing

interventions that address their overall wellbeing which will, in turn, improve their quality of

life and healthcare provision.

This study therefore sought to determine the burden of depressive, anxiety and stress

symptoms among junior doctors at the Korle-Bu Teaching Hospital (KBTH) and explore how

these relate with their cardiovascular autonomic function, measured primarily by post-

exercise heart rate recovery after one (1) minute (HRR_1) and chronotropic index, and

secondarily by heart rate recovery after two (2) minutes (HRR_2)

Materials and methods

The study was conducted at the Korle-Bu Teaching Hospital, the premier teaching hospital

and largest public referral facility in Ghana. It also serves as the largest facility for

postgraduate medical training across all specialties in Ghana.

A cross-sectional study was conducted from 17[th] of April to the 18[th] of August 2023. Junior

doctors from departments with predisposing factors to psychosocial stress described earlier

were considered for this study. Therefore, the selected departments were Internal Medicine,

Surgery, Child Health, Obstetrics and Gynaecology, Anaesthesia, Emergency Medicine, and

Family Medicine. These doctors had been at post for at least 6 months prior to the study.

Exclusion criteria, due to potential effects on cardiovascular autonomic function, were:

- Pregnancy

- Self-declared previous or current diagnosis of hypertension, diabetes mellitus,

hypothyroidism, hyperthyroidism, heart failure, chronic kidney disease or psychiatric

illness.

- Doctors on beta blockers, renin-angiotensin-aldosterone system (RAAS) blockers or

psychotropic medications

- Doctors with absolute contraindications for the TMT, e.g., inability to exercise, severe

aortic stenosis, left bundle branch block on baseline electrocardiogram, etc.

Sample size determination and sampling

The minimum sample size calculated using the Cochran formula at 95% confidence [28] and

adjusting for a finite population of 864 junior doctors in the departments selected for this

study was 241. With an anticipated 10% non-response rate, this was adjusted to 268. Though

there is a paucity of data on stress, anxiety and depression among doctors in the West African

sub-region, an Ethiopian study determined a 68.2% prevalence of stress among healthcare

workers [26]. This gave a larger sample size than that obtained when calculated using a

pooled prevalence of depression (29%) among doctors [29]. Participants were sampled using

proportionate stratified sampling techniques based on department, rank and gender.

Lists of junior doctors were obtained from each selected department's secretariats per rank in

random order. Names of doctors in each rank were extracted, categorised based on gender

and assigned serial numbers. The total number of doctors to be recruited per rank in each

department was then determined proportionately based on gender. Subsequently, a sampling

frame was applied based on the numbers of junior doctors in the departments per rank and

sex, with the starting point determined by ballot.

Fig 2. Flow chart showing the recruitment process for the study

Operational definitions

House officer – A doctor who has completed medical school and is enrolled in the mandatory

2-year internship in 4 selected core rotations prior to permanent registration with the Medical

and Dental Council of Ghana.

Junior doctor – Medical doctor who has not completed fellowship training in a

medical/surgical specialty under a recognised postgraduate medical college.

Junior resident – A doctor enrolled in a postgraduate clinical training programme leading to

the award of Membership of a postgraduate medical college.

Medical officer – A doctor who has completed the mandatory 2-year housemanship

programme, is permanently licensed with the Ghana Medical and Dental Council but not

enrolled in a residency training programme with a recognised postgraduate medical college.

Senior resident: A specialist who has enrolled in a Fellowship programme with a recognised

postgraduate medical college.

Specialist – A doctor who has completed a clinical Membership programme with a

recognised postgraduate medical college. They share the same professional responsibilities as

a senior resident.

184 Study Instruments

Data was obtained using a semi-structured questionnaire, anthropometric tools and the TMT.

The questionnaire collected data on demographics, professional history, and psychosocial

symptoms. The DASS-21 questionnaire was incorporated in the semi-structured

questionnaire for the study. This was validated by Coker, Coker and Sanni in a Nigerian

study [30]. The questionnaire was pre-tested on 10 junior doctors sampled from the various

ranks using the same procedure intended for the study.

Weight was measured with a standard scale, measuring in kilograms (kg) to the nearest

0.1kg. A locally manufactured wooden stadiometer was used to measure height and

subsequently converted from centimetres (cm) to metres (m), measured to the nearest 0.01m.

The TMTs were performed in line with recommendations on the conduct of TMTs by Smarz

et al. [31], with a setup for resuscitation available if required. These were done during

working hours with flexibility to accommodate varied work schedules (between 9am and 2pm) to ensure physiologic conditions were similar to what participants experience on a regular day.

Data collection procedure

Selected participants were called on phone a week before the stress ECG to introduce the study. Consequently, the electronic questionnaire with an attached consent form detailing the study and unique serial number were forwarded via WhatsApp or electronic mail (e-mail) based on the participant's preference. Regular reminders to complete the questionnaire were sent till the end of the following week. The TMT was scheduled once the questionnaire had been completed. The first question on the electronic questionnaire established consent to participate, while the second detailed the exclusion criteria. These were compulsory questions which required a "yes" and "no" response respectively to be allowed to participate. Any answers other than these rendered the participant disqualified. All those who met exclusion criteria were made to discontinue the study, and a replacement of the same rank, sex and department was selected randomly by ballot. Those who declined to fill the questionnaire were termed non-responders, as were those who had not filled the questionnaire by the end of the overall data collection period despite repeated reminders.

On arrival for the TMT, exclusion criteria were re-assessed to ensure qualification to participate. Anthropometric data were then obtained. Participants were ushered to a couch where they rested for 5 minutes, and vital signs measured. A portable electrocardiogram (Norav PC-ECG-1200 device) was fixed using the modified Mason-Likar technique and wirelessly connected to a console for continuous ECG monitoring. The baseline ECG was studied for ECG-based exclusion criteria. A symptom-limited Bruce protocol TMT was then

performed, with no post-exercise cool-down stage. Indications for terminating the test

included fatigue, limiting chest pain, light-headedness, leg cramping, ventricular tachycardia,

new-onset ST segment elevations more than 1mm above baseline ECG, systolic BP drop by

10mmHg during exercise, and participant's request to discontinue for other reasons.

Participants who were unfamiliar with the use of the treadmill had a 1-minute warm-up

session at speeds from 0.5km/h up to 1.5km/h, to allow them to adjust to its use prior to

commencing the Bruce protocol. On completion of the exercise stages, participants had a 3-

minute recovery stage.

Data management

Autonomic function assessed by HRR_1 and chronotropic index from the TMT was the

primary outcome. As a secondary outcome, HRR_2 was also assessed.

The maximal predicted heart rate (MPHR) was defined as "220-age in years" [31].

HRR_1 was determined by "heart rate at peak exercise minus heart rate after the 1^{st} minute of

recovery". HRR_1 values $\leq$ 18bpm were considered abnormal [18,32,33]. All heart rate

determinations were based on ECG tracings from the wireless electrocardiographic device.

The chronotropic index was calculated as (peak HR-resting HR) / (maximum predicted HR –

resting HR). Chronotropic incompetence was defined as an index < 0.80 [23].

HRR_2 was calculated as the difference between HR at peak exercise minus HR at 2 minutes

of recovery, with values of $\leq$42bpm considered abnormal [22].

Psychosocial assessment

Scores obtained from answers to questions for depressive, anxiety and stress symptom categories from the DASS-21 scale were summed and doubled. Consequently, they were used to determine the classification of the respective symptoms as shown in Table 1 below.

Table 1 Categorisation of depressive, anxiety and stress symptoms according to the DASS-21 scale

	Depression	Anxiety	Stress
Normal	0-9	0-7	0-14
Mild	10-13	8-9	15-18
Moderate	14-20	10-14	19-25
Severe	21-27	15-19	26-33
Extremely severe	28+	20+	34+

Statistical analysis

The completed questionnaire was exported from Google Forms to a Microsoft Excel spreadsheet. The dataset was then exported, cleaned, coded, and analysed using Stata statistical software version 16.1. Missing responses within the DASS-21 subscale were handled using person mean substitution, since at most one item within the respective subscale was missing. Continuous variables were summarized using mean and standard deviation (SD) for approximately normally distributed data, and median with interquartile range (IQR) for skewed distributions. Categorical variables were summarized as frequencies and percentages. Internal consistency of the DASS-21 subscales was assessed using Cronbach's alpha, while sampling adequacy was evaluated with the Kaiser-Meyer-Olkin (KMO) measure.

At the bivariate level, correlations among DASS-21 subscales, between each DASS-21

subscale and cardiovascular autonomic function measures, namely chronotropic index and

HRR1, and between the two autonomic outcomes were assessed. Pearson correlation

coefficients were used for normally distributed variable pairs, while Spearman's rank

correlation was applied for nonparametric data. Differences in chronotropic index and HRR1

across tertiles of each DASS-21 subscale were evaluated using one-way analysis of variance

(ANOVA) for normally distributed outcomes or the Kruskal-Wallis test for skewed

outcomes. Corresponding box plots were generated for graphical presentation.

For multivariable analysis, DASS-21 subscale scores were standardized to a mean of 0 and a

standard deviation of 1 to facilitate interpretable effect sizes relating to the chronotropic

index and HRR1. Primary analyses were conducted using linear regression models, adjusting

for biologically relevant covariates, following assessment of residuals and other model

assumptions. Quantile regression was fitted for the chronotropic index model due to

violations of multiple linear regression diagnostics. For HRR1, linear regression with robust

variance estimation was used to account for the presence of outliers. Secondary analyses

employed multivariable logistic regression models for categorized chronotropic index and

HRR1 based on predefined clinical cutoffs. Linearity in the logit for continuous predictors

was assessed graphically and found to be satisfactory; thus, predictors were retained in their

original continuous form. Associations were expressed per standard deviation increase in

DASS-21 scores to facilitate clinically interpretable effect sizes and comparison across

psychosocial constructs. Statistical significance was set at a two-sided p-value less than 0.05,

with a corrected threshold of 0.01 to account for multiple comparisons involving two co-

primary outcomes and three factor variables.

Ethical considerations

Approval was obtained from the Institutional Review Board (IRB) of the Korle-Bu Teaching Hospital (REF No. KBTH-IRB 000154/22). Written informed consent was obtained from each participant. Study data were completely anonymised.

Results

Demographic characteristics of respondents

A total of 268 junior doctors were invited to participate in this study, with a response rate of 91% (244/268). Sociodemographic and clinical characteristics are shown in Table 2. Among respondents, 123 (50.41%) were male. The mean age of participants was 31.46 (SD 4.41) years. The median body mass index (BMI) was 25.58 (IQR, 22.79-28.66) kg/m^2. Sixty-nine (69) participants reported that they consume alcoholic beverages (28.28%) at least occasionally, while 6 (2.46%) and 100 (40.98%) reported smoking (tobacco and shisha) and drinking coffee, respectively.

Table 2 Sociodemographic and clinical characteristics of study participants (N = 244)

Characteristic	Value (%)
Sex	
Male	123 (50.41%)
Female	121 (49.59%)
Age (years), mean (SD)	31.46 (4.41)

Age-group	
25 years and below	26 (10.66%)
26-30 years	81 (33.20%)
31-35 years	86 (35.25%)
36-40 years	45 (18.44%)
41-45 years	5 (2.05%)
Above 45 years	1 (0.41%)
Marital status	
Single	128 (52.46%)
Married	114 (46.72%)
Divorced	2 (0.82%)
Years worked as a doctor	
Less than 2 years	66 (27.05%)
2-5 years	67 (27.46%)
5-10 years	89 (36.48%)
More than 10 years	22 (9.02%)
Grade/Rank	
House officer	65 (26.64%)
Medical officer	59 (24.18%)

Junior resident	85 (34.84%)
Senior resident	35 (14.34%)
Years in current rank	
Less than 2 years	162 (66.39%)
2-5 years	74 (30.33%)
5-10 years	7 (2.87%)
More than 10 years	1 (0.41%)
Department	
Anaesthesia	33 (13.52%)
Child health	36 (14.75%)
Emergency medicine	18 (7.38%)
Family medicine	19 (7.79%)
Internal medicine	60 (24.59%)
Obstetrics and Gynaecology	33 (13.52%)
Surgery	45 (18.44%)
Average hours of work per week	
Less than 30 hours	4 (1.64%)
30 – 39 hours	65 (26.64%)
40 – 49 hours	85 (34.84%)
50 hours or more	90 (36.89%)

Alcohol consumption	
Yes	69 (28.28%)
No	175 (71.72%)
Smoking	
Yes	6 (2.46%)
No	238 (97.54%)
Coffee intake	
Yes	100 (40.98%)
No	144 (59.02%)
BMI (kg/m^2), median (IQR)	25.58 (22.79-28.66)

Values are presented as frequency (percentage), mean (SD), or median (IQR).

Abbreviations: BMI, body mass index; IQR, interquartile range; kg/m^2, kilogram per meter square; N, sample size; SD, standard deviation

Psychosocial factor assesment

The DASS-21 demonstrated good internal consistency, with Cronbach's alpha values of 0.86, 0.80, and 0.90 for the stress, anxiety, and depression subscales, respectively. The Kaiser-Meyer-Olkin (KMO) measure indicated excellent sampling adequacy (overall KMO = 0.92). The distribution of DASS-21 subscale scores is shown in Table 3. Stress scores were approximately normally distributed, with a mean score of 15.51 (SD 8.65), whereas anxiety and depression scores were skewed, with median anxiety and depression scores of 6 (IQR 2-14) and 4 (IQR 0-10), respectively. The prevalence of stress, anxiety, and depressive symptoms was determined to be 48.36%, 48.77%, and 29.51%, respectively, with varied degrees of symptom severity (i.e., from mild to extremely severe). Participants were

322   relatively evenly distributed across low, middle, and high score tertiles for the stress subscale,

323   while they were largely in low tertiles for anxiety and depression subscales.

325   Table 3. Distribution of DASS-21 subscale scores among study participants (N=244)

Variable	Stress	Anxiety	Depression
Mean/median score	15.51 (SD 8.65)	6 (IQR 2-14)	4 (IQR 0-10)
Severity category, n (%)			
Normal	126 (51.64)	125 (51.23)	172 (70.49)
Mild	35 (14.34)	14 (5.74)	23 (9.43)
Moderate	50 (20.49)	50 (20.49)	31 (12.70)
Severe	26 (10.66)	28 (11.48)	13 (5.33)
Extremely severe	7 (2.87)	27 (11.07)	5 (2.05)
Tertiles□, n (%)			
Low	93 (38.11)	102 (41.80)	106 (43.44)
Middle	84 (34.43)	74 (30.33)	66 (27.05)
High	67 (27.46)	68 (27.87)	72 (29.51)

Scores are presented as mean (SD) for normally distributed variables (stress) and median

(IQR) for skewed variables (anxiety and depression). Categorical variables are presented as a

number, n (percentage, %). Severity categories are based on DASS-21 subscale cut-offs.

Abbreviations and symbols: n, number; SD, standard deviation; IQR, interquartile range.

□Tertiles of the respective subscale scores.

There was a moderate positive correlation between stress and anxiety scores (Spearman's ρ =

0.55, p < 0.001), stress and depression scores (Spearman's ρ = 0.61, p < 0.001), and anxiety

and depression scores (Spearman's ρ = 0.57, p < 0.001).

Treadmill stress test and cardiovascular autonomic function

parameters

All participants underwent a symptom-limited TMT using the standard Bruce protocol.

Fatigue was the main reason for TMT termination (68.9%). The median resting heart rate was

81 (IQR, 73-88) bpm, while the median peak heart rate was 175 (IQR, 165 – 180.5) bpm. Up

to 238 participants (97.54%) achieved the target heart rate, i.e., at least 85% of the maximum

predicted heart rate (median, 94.5%; IQR, 90%-100%).

For the cardiovascular autonomic function measures (Table 4), the HRR_1 and HRR_2 were

approximately normally distributed, and the mean HRR_1 was 38.09 (SD, 9.45; 95% CI, 36.9-

39.29) bpm, while the mean HRR_2 was 58.14 (SD, 9.82; 95% CI, 56.90-59.38) bpm.

Chronotropic index data were skewed, with a median of 0.92 (IQR, 0.85 – 1.00; bootstrap

95% CI, 0.89-0.94). While an abnormal chronotropic index (chronotropic incompetence) was

observed in 23 participants (9.43%; 95% CI, 6.07% - 13.81%), HRR_1 and HRR_2 were

abnormal among 8 (3.28%; 95% CI, 1.43% - 6.36%) and 15 (6.15%; 95% CI, 3.48% -

9.94%) participants, respectively.

Table 4. Summary and categorical distribution of cardiovascular autonomic function

measures among study participants (N = 244)

Outcome measure	Value

Chronotropic index	
Median (IQR) [bootstrap◻ 95% CI]	0.92 (0.85 – 1.00) [0.89-0.94]
Chronotropic incompetence, n (%) [95% CI]	23 (9.43%) [6.07% - 13.81%]
Heart rate recovery at 1 minute, bpm	
Mean (SD) [95% CI]	38.09 (9.45) [36.9-39.29]
Abnormal HRR_1, n (%) [95% CI]	8 (3.28%) [1.43% - 6.36%]
Heart rate recovery at 2 minutes, bpm	
Mean (SD) [95% CI]	58.14 (9.82) [56.90-59.38]
Abnormal HRR_2, n (%) [95% CI]	15 (6.15%) [3.48% - 9.94%]

Values are presented as mean (SD) [95% CI], median (IQR) [bootstrap 95% CI], or n (%)

[95% CI]. Chronotropic incompetence was defined as a chronotropic index < 0.80. Abnormal

HRR_1 and HRR_2 were defined as $HRR_1 \leq$ 18bpm and $HRR_2 \leq$ 42bpm, respectively.

Abbreviations: bpm, beat per minute; CI, confidence interval; HRR_1, Heart rate recovery at 1

minute; HRR_2, Heart rate recovery at 2 minutes; IQR, interquartile range; n, number; SD,

standard deviation.

◻Bootstrap 95% CI was obtained using 1,000 replications of the sample.

Chronotropic index and HRR_1 were not correlated (ρ = -0.09, p = 0.17), while HRR_1 and

HRR_2 were strongly correlated (Pearson's r = 0.77, p < 0.001).

366 Relationship between psychosocial factors and cardiovascular

367 autonomic function

Chronotropic index was not correlated with DASS-21 stress (ρ = -0.09, p = 0.15); anxiety (ρ

= -0.05, p = 0.45); or depression (ρ = 0.00, p = 0.96). Similarly, HRR_1 was not correlated

with DASS-21 stress (r = 0.11, p = 0.09); anxiety (ρ = 0.03, p = 0.64); or depression (ρ = -

0.01, p = 0.90).

Box plots showed similar distributions of both chronotropic index and HRR_1 across tertiles of

stress, anxiety, and depression scores, with largely overlapping medians and interquartile

ranges (Figs 3a-c and 4a-c). The Kruskal-Wallis test revealed no difference in chronotropic

index across tertiles of stress (χ^2 (2) = 3.89, p = 0.14); anxiety (χ^2 (2) = 1.12, p = 0.57); or

depression (χ^2 (2) = 0.03, p = 0.99) (S1 Table). Similarly, one-way ANOVA showed no

difference in mean HRR_1 across tertiles of stress ($F_{2,241}$ = 1.90, p = 0.15); anxiety ($F_{2,241}$ =

0.37, p = 0.69); or depression ($F_{2,241}$ = 0.05, p = 0.95) (S2 Table).

Fig 3a-c. Box and whisker plots of chronotropic index across tertiles of psychosocial factor

subscales assessing group differences using the Kruskal-Wallis test. Boxes represent

interquartile ranges with median lines, and whiskers indicate 1.5 times the interquartile range.

3a, chronotropic index vs stress tertiles, χ^2 (2) = 3.89, p = 0.14. 3b) chronotropic index vs

anxiety tertiles, χ^2 (2) = 1.12, p = 0.57. 3c) chronotropic index vs depression tertiles, χ^2 (2) =

0.03, p = 0.99.

Figure 4a-c. Box and whisker plot of heart rate recovery at 1 minute (HRR_1) across tertiles

of psychosocial factor subscales assessing group differences using one-way ANOVA. Boxes

represent interquartile ranges with median lines, and whiskers indicate 1.5 times the

interquartile range. 4a), HRR_1 vs stress tertiles, F (2, 241) = 1.90, p = 0.15. 4b) HRR_1 vs

anxiety tertiles, F (2, 241) = 0.37, p = 0.69. 4c) HRR_1 vs depression tertiles, F (2, 241) = 0.05,

p = 0.95.

After adjustment for age, sex, BMI, smoking, coffee intake, and resting heart rate, higher

DASS-21 stress scores were associated with a small reduction in chronotropic index at the

median in quantile regression, although the 95% CI included the null (β per SD –0.01, 95%

CI –0.03 - 0.00; Table 5). Anxiety and depression scores showed minimal associations with

chronotropic index, with effect estimates close to zero (S3-6 Tables). In contrast, higher

stress scores were associated with higher HRR1 in adjusted linear regression (β per SD 1.53

bpm, 95% CI 0.03 - 3.02), while anxiety and depression scores were not meaningfully

associated with HRR1 (Table 5). In logistic regression using clinically defined cut-offs (S7-

10 Tables), higher stress and anxiety scores were associated with increased odds of

chronotropic incompetence, although estimates were imprecise (adjusted OR per stress score

SD 1.68, 95% CI 0.93 - 3.04; adjusted OR per anxiety score SD 1.30, 95% CI 0.73 – 2.31),

whereas higher depression scores were associated with lower odds of chronotropic

incompetence (adjusted OR 0.44, 95% CI 0.22 - 0.88; Table 6). Associations between DASS-

21 subscales and abnormal HRR1 were generally weak and characterized by wide confidence

intervals spanning clinically relevant effects in both directions (Table 6).

Table 5. Quantile and linear regression models assessing the association between

psychosocial factors and autonomic function

DASS-21 subscale	Unadjusted coefficient, β(95% CI) [p-value]	Adjusted coefficient, β(95% CI) [p-value]
Chronotropic index: Quantile regression, n= 244, Pseudo R²= 0.12		
Stress score	-0.02 (-0.03 - -0.00) [0.03]	-0.01 (-0.03 – 0.00) [0.17]
Anxiety score	-0.01 (-0.03 – 0.01) [0.19]	-0.00 (-0.02 – 0.01) [0.68]

Depression score	-0.00 (-0.02 – 0.01) [0.62]	0.01 (-0.01 – 0.02) [0.50]
HRR₁: Linear regression with vce robust, n=244, R²=0.11, p=0.03		
Stress score	1.01 (-0.18 – 2.20)bpm [0.09]	1.53 (0.03 – 3.02)bpm [0.04]
Anxiety score	0.40 (-0.76 – 1.57)bpm [0.49]	-0.08 (-1.46 – 1.31)bpm [0.91]
Depression score	0.14 (-0.90 – 1.18)bpm [0.78]	-0.60 (-1.82 – 0.61)bpm [0.33]

Chronotropic index was modelled using quantile regression at the median, while HRR_1 was

analysed using linear regression with robust variance estimation (vce). Regression

coefficients (β) represent effects per one standard deviation increase in the respective DASS-

21 subscale score. Adjusted models included age, sex, body mass index, smoking status,

coffee intake, and resting heart rate. Results are presented with 95% confidence intervals to

emphasize effect size and precision; p-values are provided for completeness.

Abbreviations: bpm, beats per minute; CI, confidence interval; HRR_1, Heart rate recovery at 1

minute; n, sample; R², R-squared of the model.

Table 6. Logistic regression models assessing the association between psychosocial factors

and autonomic function

DASS-21 subscale	Unadjusted odds ratio (95% CI) [p-value]	Adjusted odds ratio (95% CI) [p-value]
Chronotropic incompetence: n= 244, Pseudo R²= 0.09, p=0.12, area under ROC=0.74		
Stress score	1.22 (0.80 - 1.86) [0.36]	1.68 (0.93 – 3.04) [0.08]
Anxiety score	1.15 (0.76 – 1.74) [0.51]	1.30 (0.73 – 2.31) [0.38]
Depression score	0.74 (0.44 – 1.25) [0.25]	0.44 (0.22 – 0.88) [0.02]
Abnormal HRR₁: n=238, Pseudo R²=0.34 p < 0.01, area under ROC=0.87		
Stress score	0.65 (0.30 – 1.40) [0.27]	0.68 (0.17 – 2.79) [0.59]

Anxiety score	1.06 (0.53 – 2.12) [0.87]	2.29 (0.64 – 8.11) [0.20]
Depression score	0.42 (0.12 – 1.49) [0.18]	0.11 (0.01 – 1.50) [0.10]

Chronotropic incompetence and abnormal HRR_1 were defined using established clinical cut-

offs (<0.80 and ≤ 18bpm). Odds ratios represent effects per one standard deviation increase in

DASS-21 subscale scores. Adjusted models included age, sex, body mass index, smoking

status, coffee intake, and resting heart rate. Model performance is summarized using pseudo-

R^2 values and area under the receiver operating characteristic (ROC) curve. Confidence

intervals are emphasized to reflect the exploratory, cross-sectional nature of the analysis.

Abbreviations: bpm, beats per minute; CI, confidence interval; HRR_1, Heart rate recovery at 1

minute; n, sample; R^2, R-squared of the model.

Discussion

The study was conducted among junior doctors at the KBTH, with a mean age of 31.46 years.

It revealed a high prevalence of psychosocial factors which correlated moderately with one

another. Less than 10% of participants recorded abnormal parameters of cardiovascular

autonomic function, with no significant differences after comparing based on tertiles of the

psychosocial subscales. There was no relationship between psychosocial factors and

chronotropic index after adjusting for possible confounders, while higher stress scores were

associated with an increase in HRR_1 in a linear regression model. High depression scores

were however associated with decreased odds of chronotropic incompetence.

The high KMO and Cronbach's alpha across all psychosocial subscales assure adequate

sampling and internal consistency of the DASS-21 tool within the sample population. The

prevalences of stress, anxiety and depression were 48.36%, 48.77% and 29.15% respectively.

These are consistent with other studies, with a systematic review by Mata et al. reporting a

28.8% pooled prevalence of depression [29]. Early stages of professional development have

been associated with higher rates of psychosocial distress [34], consistent with the population

demographics of this study. Despite the use of different stress assessment scales, studies in

Ethiopia [26,35] and Saudi Arabia [36] all showed high levels of stress, ranging from 48-

73%. These factors also correlated with one another as already established [37], underscoring

their intricate relationship and the concern for multimorbidity. Over 75% of this study's

population comprised doctors in training positions, suggesting multiple responsibilities;

conducting daily inpatient reviews, taking responsibility for day-to-day patient care decisions,

running multiple clinics under supervision, providing daily cover for wards and the

emergency unit, seeking out opportunities to observe and practice procedural skills required

to be proficient for the next career stage, and seeking to balance these with family and

personal commitments. Similar factors were described as contributory to stress among

Nigerian junior doctors [38].

Participants with moderate psychosocial symptoms or higher across all assessed factors

represented majority of those who reported symptoms, consistent with findings from Saudi

Arabia which show a 34.9% prevalence of severe stress among interns (house officers) [36].

Similarly, a 21% and 58% prevalence of at least moderate levels of stress were reported in

studies among nurses and midwives in Northern Ghana respectively [39,40], in consonance

with our findings. The presence of these symptoms may the risk of unforced work-related

errors and the adoption of risky lifestyle behaviours [37,41] which may harm patients

inadvertently, though it has been suggested that there is an inconsistent relationship between

stress and risky behaviours such as alcohol use, smoking and physical inactivity [42]. Such

levels of psychosocial stress could progress to burnout as suggested by studies among

Ghanaian and Nigerian health workers [27,43]. Consequently, sufferers may experience

downstream cardiovascular and psychological/psychiatric effects [4,34,37,44].

There was a low prevalence of cardiovascular autonomic dysfunction with no significant

variation when analysed across tertiles of psychosocial factors. This could be due to high

resilience because of good coping mechanisms, because while psychosocial factors invoke an

autonomic response, the development of autonomic dysfunction follows long-term exposure

to stressors that overwhelm intrinsic coping mechanisms [6,8]. Alternatively, there may be an

asymptomatic stage that the tools used in this study are not sensitive to diagnose. Heart rate

variability (HRV) has been used in assessing cardiovascular autonomic function in other

studies, including a study among Ghanaian nurses which found no association with

occupational stress [40]. However, a meta-analysis on the relationship between HRV and

occupational stress identified 8 out of 10 studies across various professions that demonstrated

abnormal cardiovascular autonomic function [45]. HRR is described by Goldberger et al. as

greater in assessing the dynamic changes of the autonomic nervous system, though both HRR

and HRV have limited real-world clinical utility [17]. Though a study with a small number of

participants identified a significant correlation between HRR and HRV, other evidence

comparing them have been equivocal, with neither showing a clear diagnostic advantage [46].

Chronotropic index showed no relationship with the independent variables when adjusted for

confounding, whereas HRR_1 values were higher with higher stress scores. The associated

SNS activation in response to stress, together with the exercise-induced increase in SNS

output, can result in higher peak HR following exercise. This could lead to marked HR

reductions after exercise as the PSNS output increases, leading to increases in HRR. When

PSNS output is however diminished due to chronic exposure to the psychosocial stressors,

this may then manifest as autonomic dysfunction. As seen in Figure 1, it may be plausible

that the study participants, despite the high prevalence of psychosocial factors, have not

progressed to the chronic state responsible for autonomic dysfunction. Additionally, stressors

may rather improve efficiency and resourcefulness based on one's subjective assessment of

the situation and coping mechanisms. This is called eustress [9,26]. This is consistent with

the population mean and medians for the psychosocial constructs falling within the mild

symptom bracket, suggesting that the stressors they face overall may not be overwhelming

yet. Similar mechanisms could explain the lower odds of chronotropic incompetence related

to depression scores.

The low prevalence of cardiovascular autonomic dysfunction may be attributable to the

relatively young ages of the study population (mean age = 31.46 years), presumably shorter

exposure to psychosocial stressors, coping mechanisms, or other factors not explored in this

study. Additionally, doctors may be presumed to be more health conscious than the general

population, putting effort into controlling psychosocial stressors. The individual response to

stress is however varied and difficult to characterise distinctly [42]. The small effect sizes and

wide confidence intervals of some outcome parameters may reflect limited statistical power

to identify such associations. Nonetheless, the prevalences of depressive, anxiety and stress

symptoms have clinical significance that requires attention to safeguard the health of the

junior doctor. It is therefore recommended that the findings of this study be interpreted as

exploratory, generating hypothesis for further longitudinal investigation. These can assess

how psychosocial factors relate with cardiovascular autonomic function and potential

mechanisms for this relationship, strengthened with the use of invasive psychosocial stress

markers like cortisol and catecholamine levels to enhance objectivity.

Limitations

While the study has some strengths, there are significant limitations that need to be acknowledged. The use of the DASS-21 tool is subject to recall/self-report bias, potentially resulting in an over- or under-estimation of the true burden of stress. Secondly, the duration of exposure to the psychosocial factors may not qualify as chronic, depending on the time over which this has happened and coping mechanisms. The definition of acute versus chronic stress is individualised based on each person's threshold determined by one's perception of situations and coping mechanisms. This could influence results as the longstanding haemodynamic changes from autonomic dysfunction are due to chronic stress, rather than exposure to stress acutely. Additionally, the study could not differentiate between eustress, which improves output, and distress which causes deleterious effects. Routine exercise habits, sleep patterns and shifts before exercise were not factored into the study, which could potentially have influenced the results as autonomic function was assessed. Finally, this was a single-centre cross-sectional study, limiting the generalisability of the results as well as the inference of causality.

Conclusion

There are high prevalences of stress and anxiety symptoms among junior doctors at the KBTH, with a lower (albeit significant) prevalence of depressive symptoms. The prevalence of cardiovascular autonomic dysfunction was low, and some psychosocial factors were associated with enhanced cardiovascular autonomic function suggesting a more acute duration of these factors or durable coping mechanisms. There were no significant differences in cardiovascular autonomic function based on severity of psychosocial factors.

It is suggested that sources and effects of these psychosocial stressors be studied further with

adequate sampling to enable improved efforts at mental health protection of the junior doctor.

Acknowledgements

Dr Muhyideen Bashir assisted with the review of the statistical analysis and interpretation of

results in the revised manuscript.

The efforts of Dorcas Baiden-Amissah, Angelina Martin, Raheal Osman, Gloria Dzata and

Georgina Banahene, all staff of the National Cardiothoracic Centre at the time of the study,

are acknowledged for their assistance with the data collection process.

Data availability

All relevant data are within the manuscript and its supporting information files

Supporting information

S1 Table Comparison of chronotropic index with stress, anxiety and depressive symptoms by

tertiles using the Kruskal-Wallis test

S2 Table Comparison of heart rate recovery at 1 minute with stress, anxiety and depressive

symptoms by tertiles using one-way ANOVA

S3 Table Linear regression model (crude) assessing the association between psychosocial

factors and cardiovascular autonomic function

S4 Table Quantile regression model (adjusted) assessing the association between

psychosocial factors and chronotropic index

S5 Table Linear regression model (adjusted) assessing the association between psychosocial

factors and HRR_1

S6 Table Linear regression model (adjusted) assessing the association between psychosocial

factors and HRR_2

S7 Table Logistic regression model (crude) assessing the association between psychosocial

factors and cardiovascular autonomic function

S8 Table Logistic regression model (adjusted) assessing the association between

psychosocial factors and chronotropic index

S9 Table Logistic regression model (adjusted) assessing the association between

psychosocial factors and HRR_1

S10 Table Logistic regression model (adjusted) assessing the association between

psychosocial factors and HRR_2

S11 Table Study dataset

References

1.　Yuyun MF, Sliwa K, Kengne AP, Mocumbi AO, Bukhman G. Cardiovascular

diseases in sub-Saharan Africa compared to high-income countries: An

epidemiological perspective. Glob Heart. 2020;15(1): 1–18.

doi:10.5334/GH.403

2.　Boateng D, Wekesah F, Browne JL, Agyemang C, Agyei-Baffour P, De-Graft

Aikins A, et al. Knowledge and awareness of and perception towards

cardiovascular disease risk in sub-Saharan Africa: A systematic review. PLoS

One. 2017;12(12): 1–21. doi:10.1371/journal.pone.0189264

3.   Asamoah KT, Dei-Asamoa R. Psychiatry Meets Cardiology: A Case Report on

593   the Need for Mental Health Assessment in the Evaluation of Cardiovascular

594   Symptoms. Case Rep Psychiatry. 2022; doi:10.1155/2022/5415196

595   4.   Raggi P, Quyyumi AA, Henein MY, Vaccarino V. Psychosocial stress and

596   cardiovascular disease. Am J Prev Cardiol. 2025;22.

597   doi:10.1016/j.ajpc.2025.100968

598   5.   Bueno H, Deaton C, Farrero M, Forsyth F, Braunschweig F, Buccheri S, et al.

599   2025 ESC Clinical Consensus Statement on mental health and cardiovascular

600   disease: developed under the auspices of the ESC Clinical Practice Guidelines

601   Committee. Eur Heart J. 2025. doi:10.1093/eurheartj/ehaf191

602   6.   Osborne MT, Shin LM, Mehta NN, Pitman RK, Fayad ZA, Tawakol A.

603   Disentangling the Links between Psychosocial Stress and Cardiovascular

604   Disease. Circ Cardiovasc Imaging. 2020;13(8): e010931.

605   doi:10.1161/CIRCIMAGING.120.010931

606   7.   Kerr JI, Naegelin M, Weibel RP, Ferrario A, La Marca R, von Wangenheim F,

607   et al. The effects of acute work stress and appraisal on psychobiological stress

608   responses in a group office environment. Psychoneuroendocrinology. 2020;121:

609   104837. doi:10.1016/j.psyneuen.2020.104837

610   8.   Taouk Y, Spittal MJ, Lamontagne AD, Milner AJ. Psychosocial work stressors

611   and risk of all-cause and coronary heart disease mortality: A systematic review

612   and meta-analysis. Scand J Work Environ Health. 2020;46(1): 19–31.

613   doi:10.5271/sjweh.3854

614   9.   Hinterdobler J, Schunkert H, Kessler T, Sager HB. Impact of Acute and Chronic

615   Psychosocial Stress on Vascular Inflammation. Antioxid Redox Signal.

616   2021;35(18): 1531–1550. doi:10.1089/ars.2021.0153

617   10.   Khemani P, Mehdirad AA. Cardiovascular Disorders Mediated by Autonomic

618   Nervous System Dysfunction. Cardiol Rev. 2020;28(2): 65–72.

619   doi:10.1097/CRD.0000000000000280

620   11.   Lankhorst S, Keet SWM, Bulte CSE, Boer C. The impact of autonomic

621   dysfunction on peri-operative cardiovascular complications. Anaesthesia.

622   2015;70(3): 336–343. doi:10.1111/anae.12904

623   12.   Bustamante-Sánchez Á, Tornero-Aguilera JF, Fernández-Elías VE, Hormeño-

624   Holgado AJ, Dalamitros AA, Clemente-Suárez VJ. Effect of Stress on

625   Autonomic and Cardiovascular Systems in Military Population: A Systematic

626   Review. Cardiol Res Pract. 2020. doi:10.1155/2020/7986249

627   13.   Golbidi S, Frisbee JC, Laher IX. Chronic stress impacts the cardiovascular

628   system: Animal models and clinical outcomes. Am J Physiol Heart Circ Physiol.

629   2015;308(12): H1476–H1498. doi:10.1152/ajpheart.00859.2014

630   14.   Majidinia M, Rasmi Y, Ansaria MHK, Seyed-Mohammadzad M, Saboory E,

631   Shirpoor A. Metoprolol improves endothelial function in patients with cardiac

632   syndrome X. Iran J Pharm Res. 2016;15(3): 561–566.

633   doi:10.22037/ijpr.2016.1869

634   15.   Yusuf S, Hawken S, Ôunpuu S, Dans T, Avezum A, Lanas F, et al. Effect of

635   potentially modifiable risk factors associated with myocardial infarction in 52

636   countries (the INTERHEART study): Case-control study. Lancet.

637   2004;364(9438): 937–952. doi:10.1016/S0140-6736(04)17018-9

638   16.   O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, et al. Global

639   and regional effects of potentially modifiable risk factors associated with acute

640   stroke in 32 countries (INTERSTROKE): a case-control study. The Lancet.

641   2016;388(10046): 761–775. doi:10.1016/S0140-6736(16)30506-2

642   17.   Goldberger JJ, Arora R, Buckley U, Shivkumar K. Autonomic Nervous System

643   Dysfunction: JACC Focus Seminar. J Am Coll Cardiol. 2019;73(10): 1189–

644   1206. doi:10.1016/j.jacc.2018.12.064

645   18.   Zaim S, Schesser J, Hirsch LS, Rockland R. Influence of the Maximum Heart

646   Rate Attained during Exercise Testing on Subsequent Heart Rate Recovery.

647   Ann Noninvasive Electrocardiol. 2010;15(1): 43–48.

648   19.   Antelmi I, Chuang EY, Grupi CJ, Latorre MDRDDO, Mansur AJ. Heart rate

649   recovery after treadmill electrocardiographic exercise stress test and 24-hour

650   heart rate variability in healthy individuals. Arq Bras Cardiol. 2008;90(6): 413–

651   418. doi:10.1590/S0066-782X2008000600005

652   20.   El Agaty SM, Kirmani A, Labban E. Heart rate variability analysis during

653   immediate recovery from exercise in overweight/obese healthy young adult

654   females. Ann Noninvasive Electrocardiol. 2017;22(3): 1–5.

655   doi:10.1111/anec.12427

656   21.   Myers J, Tan SY, Abella J, Aleti V, Froelicher VF. Comparison of the

657   chronotropic response to exercise and heart rate recovery in predicting

658   cardiovascular mortality. Eur J Cardiovasc Prev Rehabil. 2007;14(4): 215–221.

659   22.   Cole CR, Blackstone EH, Pashkow FJ, Snader CE, Lauer MS. Heart-rate

660   recovery immediately after exercise as a predictor of mortality. N Engl J Med.

661   1999;341(18): 1351–1357.

662   23.   Carreira MAMQ, Nogueira AB, Pena FM, Kiuchi MG, Rodrigues RC,

663   Rodrigues RR, et al. Detection of autonomic dysfunction in hemodialysis

664   patients using the exercise treadmill test: The role of the chronotropic index,

665   heart rate recovery, and R-R variability. PLoS One. 2015;10(6): 1–11.

666   doi:10.1371/journal.pone.0128123

667   24.   Vaidya GN. Application of exercise ECG stress test in the current high cost

668   modern-era healthcare system. Indian Heart J. 2017;69(4): 551–555.

669   doi:10.1016/j.ihj.2017.06.004

670   25.   Harvey SB, Epstein RM, Glozier N, Petrie K, Strudwick J, Gayed A, et al.

671   Mental illness and suicide among physicians. Lancet 2021; 398: 920–30

672   26.   Birhanu M, Gebrekidan B, Tesefa G, Tareke M. Workload Determines

673   Workplace Stress among Health Professionals Working in Felege-Hiwot

674   Referral Hospital, Bahir Dar, Northwest Ethiopia. J Environ Public Health.

675   2018. doi:10.1155/2018/6286010

676   27.   Odonkor ST, Frimpong K. Burnout among Healthcare Professionals in Ghana:

677   A Critical Assessment. Biomed Res Int. 2020. doi:10.1155/2020/1614968

678   28.   Bolarinwa OA. Sample Size Estimation for Health and Social Science

679   Researchers: The Principles and Considerations for Different Study Designs.

680   Niger Postgrad Med J. 2020;27: 67–75. doi:10.4103/npmj.npmj

681   29.   Mata DA, Ramos MA, Bansal N, Khan R, Guille C, Di Angelantonio E, et al.

682   Prevalence of depression and depressive symptoms among resident physicians a

683   systematic review and meta-analysis. JAMA. 2015;314(22): 2373–2383.

684   doi:10.1001/jama.2015.15845

685   30.   Coker AO, Coker OO, Sanni D. Psychometric properties of the 21-item

686   Depression Anxiety Stress Scale (DASS-21). Afr Res Rev. 2018;12(2): 135.

687   doi:10.4314/afrrev.v12i2.13

688   31.   Smarż K, Jaxa-Chamiec T, Bednarczyk T, Bednarz B, Eysymontt Z, Gałaszek

689   M, et al. Electrocardiographic exercise testing in adults: Performance and

690   interpretation. An expert opinion of the Polish Cardiac Society Working Group

on Cardiac Rehabilitation and Exercise Physiology. Kardiol Pol. 2019;77(3):

399–408. doi:10.5603/KP.a2018.0241

32.   Okutucu S, Karakulak UN, Aytemir K, Oto A. Heart rate recovery: A practical

clinical indicator of abnormal cardiac autonomic function. Expert Rev

Cardiovasc Ther. 2011;9(11): 1417–1430. doi:10.1586/erc.11.149

33.   Watanabe J, Thamilarasan M, Blackstone EH, Thomas JD, Lauer MS. Heart

Rate Recovery Immediately After Treadmill Exercise and Left Ventricular

Systolic Dysfunction as Predictors of Mortality - The Case of Stress

Echocardiography. Circulation. 2001;104: 1911–1916.

34.   Bambi S, Foà C, De Felippis C, Lucchini A, Guazzini A, Rasero L. Workplace

incivility, lateral violence and bullying among nurses. A review about their

prevalence and related factors. Acta Biomedica. 2018;89(6): 51–79.

doi:10.23750/abm.v89i6-S.7461

35.   Gebeyehu S, Zeleke B. Workplace stress and associated factors among

healthcare professionals working in public health care facilities in Bahir Dar

City, Northwest Ethiopia, 2017. BMC Res Notes. 2019;12(1): 249.

doi:10.1186/s13104-019-4277-1

36.   Abdulghani HM, Irshad M, Al Zunitan MA, Al Sulihem AA, Al Dehaim MA,

Al Esefir WA, et al. Prevalence of stress in junior doctors during their internship

training: A cross-sectional study of three Saudi medical colleges' hospitals.

Neuropsychiatr Dis Treat. 2014;10: 1879–1886. doi:10.2147/NDT.S68039

37.   Ryan E, Hore K, Power J, Jackson T. The relationship between physician

burnout and depression, anxiety, suicidality and substance abuse: A mixed

methods systematic review. Front Public Health. 2023;11.

38.   Adeolu J, Yussuf O, Popoola O. Prevalence and Correlates of Job Stress Among

716   Junior Doctors in the University College Hospital, Ibadan. Ann Ib Postgrad

717   Med. 2016;14.

718   39.   Kaburi BB, Bio FY, Kubio C, Ameme DK, Kenu E, Sackey SO, et al.

719   Psychological working conditions and predictors of occupational stress among

720   nurses, Salaga Government Hospital, Ghana, 2016. Pan Afr Med J. 2019;33.

721   doi:10.11604/pamj.2019.33.320.16147

722   40.   Nutor JJ, Aborigo RA, Okiring J, Kuwolamo I, Dorzie JBK, Getahun M, et al.

723   Individual and situational predictors of psychological and physiological stress

724   and burnout among maternity providers in Northern Ghana. PLoS One.

725   2022;17. doi:10.1371/journal.pone.0278457

726   41.   Fishta A, Backé EM. Psychosocial stress at work and cardiovascular diseases:

727   an overview of systematic reviews. Int Arch Occup Environ Health. 2015;88(8):

728   997–1014. doi:10.1007/s00420-015-1019-0

729   42.   Bamert M, Schaffter Y, Bally F, O'Connor DB, Inauen J. Assessment of stress

730   and its relationship with health behaviour in daily life: A systematic review.

731   Health Psychol Rev. 2025;19: 906–936. doi:10.1080/17437199.2025.2534000

732   43.   Ezenwaji IO, Eseadi C, Okide CC, Nwosu NC, Ugwoke SC, Ololo KO, et al.

733   Work-related stress, burnout, and related sociodemographic factors among

734   nurses. Medicine (United States). 2019;98(3).

735   doi:10.1097/MD.0000000000013889

736   44.   Abdelall ES, Eagle Z, Finseth T, Mumani AA, Wang Z, Dorneich MC, et al.

737   The Interaction Between Physical and Psychosocial Stressors. Front Behav

738   Neurosci. 2020;14: 1–15. doi:10.3389/fnbeh.2020.00063

739   45.   Järvelin-Pasanen S, Sinikallio S, Tarvainen MP. Heart rate variability and

740   occupational stress-systematic review. Ind Health. 2018;56: 500–511.

741   46.   Bechke E, Kliszczewicz B, McLester C, Tillman M, Esco M, Lopez R. An

742   examination of single day vs. multi-day heart rate variability and its relationship

743   to heart rate recovery following maximal aerobic exercise in females. Sci Rep.

744   2020;10(1). doi:10.1038/s41598-020-71747-8

Obtained lists of junior doctors from various departments

Categorised by sex and rank in random order

Selected 268 junior doctors to participate based on department, rank and sex

All 268 selected junior doctors invited to participate

12 replaced by ballot based on department, rank and sex due to meeting exclusion criteria.

24 non-responders - 14 declined invitation and 10 accepted but did not proceed

244 participants enrolled in the study

Figure 3a

Figure 3b

Figure 3c

Figure4c

Figure4b

Figure4c