Predicting future cognitive impairment in preclinical Alzheimer's disease using amyloid PET and MRI: a multisite machine learning study

Author information

Braden Yang[a,b,c] (0000-0002-2558-4132), Tom Earnest[a] (0000-0001-8671-8424), Murat Bilgel[d*] (0000-0001-5042-7422), Marilyn S. Albert[e] (0000-0002-8167-8985), Sterling C. Johnson[f] (0000-0002-8501-545X), Christos Davatzikos[c] (0000-0002-1025-8561), Guray Erus[c] (0000-0001-6633-4861), Colin L. Masters[g] (0000-0003-3072-7940), Susan M. Resnick[d*] (0000-0003-1115-7145), Michael I. Miller[h] (0000-0003-4689-3855), Arnold Bakker[i,e] (0000-0003-0417-641X), John C. Morris[j] (0000-0001-9820-5618), Tammie L.S. Benzinger[a] (0000-0002-8114-0552), Brian A. Gordon[a] (0000-0003-2109-2955), Aristeidis Sotiras[a,b,c,k] (0000-0003-0795-8820), for the Alzheimer's Disease Neuroimaging Initiative[†], for the Preclinical Alzheimer's Disease Consortium

[a] Mallinckrodt Institute of Radiology, Washington University School of Medicine in St. Louis, St. Louis, MO, USA 63110

[b] Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA 19104

[c] Center for AI and Data Science for Integrated Diagnostics (AI^{2}D), Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA 19104

[d] Laboratory of Behavioral Neuroscience, National Institute on Aging Intramural Research Program, National Institutes of Health, Baltimore, MD, USA 21224

[e] Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

21205

[f] Wisconsin Alzheimer's Disease Research Center, University of Wisconsin-Madison School of

Medicine and Public Health, Madison, WI, USA 53726

[g] Florey Institute of Neuroscience and Mental Health, Parkville, VIC, Australia 3052

[h] Center of Imaging Science and Department of Biomedical Engineering, Johns Hopkins

University, Baltimore, Maryland, USA 21218

[i] Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of

Medicine, Baltimore, MD, USA 21205

[j] Department of Neurology, Washington University School of Medicine in St. Louis, St. Louis,

MO, USA 63110

[k] Institute for Informatics, Data Science and Biostatistics, Washington University School of

Medicine in St. Louis, St. Louis, MO, USA 63110

Please address all correspondence to: Braden Yang (b.y.yang@wustl.edu), Aristeidis Sotiras

(aristeidis.sotiras@pennmedicine.upenn.edu)

45 participate in the analysis or writing of this report. A complete listing of ADNI investigators can

46 be found at: http://adni.loni.usc.edu/wp-

47 content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Highlights

- Machine learning can predict future cognitive impairment in preclinical Alzheimer's

- Models achieved high out-of-sample ROC-AUC on external sites and PET tracers

- Models were able to distinguish cognitively stable from decliners in the A4 cohort

- ML cohort enrichment enhanced secondary treatment effect detection in the A4 cohort

Abstract

Predicting the likelihood of developing Alzheimer's disease (AD) dementia in at-risk individuals is important for the design of and optimal recruitment for clinical trials of disease-modifying therapies. Machine learning (ML) has been shown to excel in this task; however, there remains a lack of models developed specifically for the preclinical AD population, who display early signs of abnormal brain amyloidosis but remain cognitively unimpaired. Here, we trained and evaluated ML classifiers to predict whether individuals with preclinical AD will progress to mild cognitive impairment or dementia within multiple fixed time windows, ranging from one to five years. Models were trained on regional imaging features extracted from amyloid positron emission tomography and magnetic resonance imaging pooled across seven independent sites and from two amyloid radiotracers ([^{18}F]-florbetapir and [^{11}C]-Pittsburgh-compound-B). Out-of-sample generalizability was evaluated via a leave-one-site-out and leave-one-tracer-out cross-validation. Classifiers achieved an out-of-sample receiver operating characteristic area-under-the-curve of 0.66 or greater when applied to all except one hold-out sites and 0.72 or greater when applied to each hold-out radiotracer. Additionally, when applying our models in a retroactive cohort enrichment analysis on A4 clinical trial data, we observed increased statistical power of detecting differences in amyloid accumulation between placebo and treatment arms after enrichment by ML stratifications. As emerging investigations of new disease-modifying therapies for AD increasingly focus on asymptomatic, preclinical populations, our findings underscore the potential applicability of ML-based patient stratification for recruiting more homogeneous cohorts and improving statistical power for detecting treatment effects for future clinical trials.

Keywords

preclinical Alzheimer's disease; machine learning; support vector machine; amyloid positron emission tomography; structural magnetic resonance imaging; patient stratification; clinical trial cohort enrichment

1 Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disease marked by the accumulation of amyloid-β plaques and tau neurofibrillary tangles in the brain, which leads to neuronal cell death and eventual development of cognitive deficits such as short-term memory loss, impairments in orientation, or difficulty with written and verbal communication. According to current models of the AD pathogenesis timeline, accumulation of these pathologies typically begin years before the onset of clinically-measurable symptoms (Jack and Holtzman, 2013; Jack Jr. et al., 2024). Thus, clinical trials of AD treatments have largely focused on disease-modifying therapies, which target these pathologies in the hopes of altering the biological course of the disease and slowing or even preventing the progression of clinical symptoms (Dyck et al., 2023; Sims et al., 2023). Yet to date, there have only been a handful of approved disease-modifying therapies, with the rest failing to demonstrate a significant treatment effect (Kim et al., 2022). A potential reason for this high failure rate is that recruited participants may have been at too advanced of a disease stage for treatments to meaningfully alter its progression (Sperling et al., 2014), with the majority of previously conducted trials primarily recruiting individuals who already displayed objective cognitive impairment (Huang et al., 2020; Sims et al., 2023; Swanson et al., 2021). In response, newly emerging trials have transitioned to targeting preclinical AD, where patients exhibit abnormal brain amyloidosis while showing no signs of cognitive impairment. Furthermore, failures may have also been attributed to high heterogeneity in the phenotypic expression of the disease, particularly in the rates of cognitive decline over

time. This heterogeneity potentially leads to differential responses to administered therapies,

which may negatively impact the estimation of treatment effects (Jutten et al., 2021).

To mitigate this issue, stratification models for distinguishing fast-cognitive decliners from slow-

or non-decliners are commonly employed in AD clinical trials. These models enable cohort

enrichment by identifying and selectively enrolling candidates who are most likely to benefit from

disease-modifying therapies in order to increase the likelihood of observing a potential treatment

effect (Ballard et al., 2019; Wolz et al., 2016). Additionally, biased allocation of fast-decliners

and slow-decliners into control and treatment groups may lead to overestimation or

underestimation of a potential treatment effect. Proper stratification of clinical trial participants

ensures that an equal distribution of fast-to-slow decliners are allocated into each trial arm and

may aid in reducing uncertainties in treatment effect estimation, on top of reducing the required

sample size and cost of running a trial (Wang et al., 2024). Previous trials have used single risk

assessments such as neuropsychological testing and quantitative amyloid and tau

measurements from positron emission tomography (PET) to stratify participants into high and

low risk groups (Abdelnour et al., 2022; Sims et al., 2023; Swanson et al., 2021). However,

recent evidence has suggested that data-driven stratification models built on machine learning

(ML) can outperform single risk factor stratification in enhancing clinical trial operations

(Birkenbihl et al., 2024; Gladstein et al., 2025; Nallapu et al., 2025; Tam et al., 2022; Wang et

al., 2024). By leveraging information across multiple data modalities such as imaging, genetics

and cognitive assessments, ML models can learn complex patterns within these multiple

modalities to provide more accurate stratifications. These stratifications may be used to guide a

more optimized cohort enrichment, leading to increased power of detect hypothetical treatment

effects (Nallapu et al., 2025), reduction in random allocation bias (Wang et al., 2024), and

reduction in the sample size required to observe a treatment effect (Birkenbihl et al., 2024;

Gladstein et al., 2025; Tam et al., 2022). Among data modalities commonly used for

stratification, medical imaging data like PET and magnetic resonance imaging (MRI) are

particularly useful for probing the topographical distribution of core and associated AD

pathologies in the brain (Jack Jr. et al., 2024), which have been shown to be highly

discriminative of cognitive decliners from non-decliners (Mathotaarachchi et al., 2017; Pascoal

et al., 2020; Pfeil et al., 2021).

Despite recent progress in developing ML models for this application, previous literature has

predominantly focused on individuals who already exhibit cognitive deficits in the form of mild

cognitive impairment (MCI) (Ezzati et al., 2021; Samper-González et al., 2018), with many

studies only training on features extracted from structural MRI (Basaia et al., 2019;

Mathotaarachchi et al., 2017; Misra et al., 2009). Efforts have been made to develop models for

the cognitively unimpaired (CU) population that also incorporate amyloid PET, but most have

either imposed a single time-to-progression window (Choi et al., 2024; Dansson et al., 2021;

Pfeil et al., 2021) or don't ensure the presence of amyloid pathology in their CU cohorts

(Karaman et al., 2022). Additionally, with trials routinely involving large consortiums of

institutions, each utilizing a different set of data acquisition protocols and scanner models, this

introduces site-specific biases in the collected data which may negatively affect the ability of

models to generalize on out-of-sample data. Furthermore, disparities in the pharmacokinetics

and binding properties of different PET radiotracers lead to significant biases in quantitative

amyloid PET measurements (Landau et al., 2014; Su et al., 2019; Villemagne et al., 2012). For

these reasons, a rigorous evaluation of ML generalizability to external sites and different

amyloid PET radiotracers is needed.

Motivated by these gaps, the current study seeks to develop and evaluate ML classifiers to

predict future cognitive decline in asymptomatic, preclinical AD patients. We trained support

vector machines (SVM) on regional imaging features extracted from amyloid PET and structural

MRI to perform binary classification of whether an individual will develop cognitive deficits within

multiple follow-up times, ranging from one year to five years. We obtained data from seven independent sites and two amyloid PET radiotracers and estimated the out-of-sample performance of trained models via leave-one-site-out and leave-one-tracer-out cross-validation. Lastly, to gauge the benefit of applying these models in a clinical trial setting, we utilized data from the Anti-Amyloid Treatment in Asymptomatic Alzheimer's (A4) study (Sperling et al., 2023) to estimate the change in statistical power to detect treatment effects after retroactive cohort enrichment using our models' predictions.

2 Materials and methods

2.1 Participants and data

Participants and their respective data were obtained from the following independent studies and consortia – the A4 study (Sperling et al., 2023), the Alzheimer's Disease Neuroimaging Initiative (ADNI), the Harvard Aging Brain Study (HABS) (Dagley et al., 2017), the Mayo Clinic Study of Aging (MCSA) (Roberts et al., 2008), the Open Access Series of Imaging Studies 3 (OASIS) (LaMontagne et al., 2019), and the Preclinical Alzheimer's Disease Consortium (PAC). ADNI, launched in 2003 as a public-private partnership and led by Principal Investigator Michael W. Weiner, MD, has aimed to test whether serial MRI, PET, other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of MCI and early AD. PAC is a consortium comprising five AD-focused datasets – the Adult Children Study (ACS) (Coats and Morris, 2005; Price and Morris, 1999), the Australian Imaging Biomarkers and Lifestyle (AIBL) study (Fowler et al., 2021), the Biomarkers of Cognitive Decline Among Normal Individuals (BIOCARD) study (Albert et al., 2014), the Baltimore Longitudinal Study of Aging (BLSA) (Resnick et al., 2000), and the Wisconsin Registry for Alzheimer's Prevention (WRAP) (Johnson et al., 2018). All participants provided informed consent to participate in their respective studies in accordance with the Declaration of Helsinki, and ethics approvals were obtained by each site's respective institutional review board. From the A4 study, only

participants from the placebo arm of the clinical trial or from the Longitudinal Evaluation of

Amyloid Risk and Neurodegeneration (LEARN) study were used to train the ML models.

Participants from the Solanezumab treatment arm were excluded from training and were only

used as an additional evaluation set (detailed in section *2.4 A4 trial analyses*). Additionally, we

omitted ACS from our analyses since this dataset had significant participant overlap with

OASIS.

All participants with available [^{18}F]-florbetapir (FBP) or [^{11}C]-Pittsburgh-compound-B (PiB)

amyloid PET imaging were selected. We restricted these scans to be amyloid-positive, defined

as having a mean cortical amyloid PET signal of greater than a threshold value. We used the

dataset-specific definition of the summary cortical region mask and cutoff signal that was

optimally determined for each dataset. Further details on how amyloid positivity was determined

for each site can be found in the Supplementary Methods. All scans were matched with an

accompanying structural T1-weighted MRI scan within one year of the PET acquisition date.

Baseline age, sex and *APOE4* carriership were selected as additional predictors. We used the

global Clinical Dementia Rating® (CDR®) (Morris, 1997), which rates an individual's cognitive

performance in six domains (memory, orientation, judgment and problem solving, community

affairs, home and hobbies, and personal care), as a measure of overall dementia severity. All

PET scans were matched to a global CDR assessment within one year of the PET acquisition

date. Participants who were missing any matching data were excluded from further analyses.

The analysis was restricted to participants who were CU at the time of the amyloid PET scan,

defined as a global CDR score of zero. We labeled participants as stable if they remained CU

for at least five years from the time of the amyloid PET, and as progressor if they converted to a

global CDR of greater than zero within a follow-up time that varied from one to five years.

Progressors who reverted to CU were excluded. Additionally, stables who did not have a follow-

up CDR of zero at least five years from the time of the amyloid PET were excluded. We

observed a larger proportion of stable individuals compared to progressors; to reduce class

imbalance, we imposed the five-year stability criterion on the stable cohort to obtain the most

homogeneous group of participants, with presumably the least amount of pathology, to

represent our control group. PAC did not provide CDR scores; we instead used the clinical

diagnoses in place of CDR, with a diagnosis of either MCI or dementia being equivalent to

CDR>0. In total, 343 stable and 247 progressor participants across seven sites (A4, ADNI,

AIBL, BLSA, HABS, MCSA, OASIS) were selected for this study. We excluded BIOCARD and

WRAP from our analyses due to too few participants meeting the selection criteria. Participant

counts after exclusions are illustrated in Supplementary Fig. 1.

2.2 Image acquisition and processing

All scans underwent a standard PET-MRI processing pipeline consisting of segmentation of the

MRI into regions-of-interest (ROI) and derivation of regional amyloid standardized uptake value

ratios (SUVR) from the amyloid PET. Each participant's T1-weighted MRI was preprocessed

using a fully automated pipeline which included N4 bias correction (Tustison et al., 2010), skull-

stripping, and segmentation into 145 cortical and subcortical ROI using Multi-atlas region

Segmentation utilizing Ensembles of registration algorithms and parameters (MUSE) (Doshi et

al., 2016). Manual quality checks were performed on the outputs to ensure correctness of

segmentations. We selected 125 relevant ROI and extracted their volumes, normalized by the

total intracranial volume, to be used as the volumetric input features of our model.

FBP and PiB PET scans were downloaded either in a frame-averaged, smoothed and uniform-

resolution format (ADNI only) or as the full dynamic scan (all other datasets). For dynamic

scans, we selected frames within the 50-70 minute window post-tracer injection for FBP and the

40-60 minute window for PiB. All frames were realigned to remove patient motion, averaged,

and coregistered to the T1 image using FSL FLIRT rigid body registration (Smith et al., 2004).

Additionally, we performed an iterative smoothing operation by applying a Gaussian smoothing

kernel with progressively increasing full width at half maximum until a target resolution was

reached (estimated using AFNI's *3dFWHMx* tool). The target resolution was set to the

3dFWHMx-estimated resolution of ADNI frame-averaged scans, which was found to be

approximately 10mm isotropic. After smoothing, the PET image was converted to an SUVR

image by dividing each voxel by the average signal of the left and right cerebellar gray matter

regions. Finally, the MRI-to-PET transformation was applied to the anatomical MUSE ROI labels

to sample them to native PET space, and regional SUVRs were computed by averaging the

SUVR image over each ROI. For ADNI scans, we only performed coregistration to the T1,

SUVR image computation and derivation of regional SUVRs. Manual quality checks were

performed to ensure correctness of the frame alignment, PET-MRI coregistration and SUVR

computation steps. Of the 145 total ROI, we selected 114 cortical and subcortical gray matter

regions to be used as the amyloid input features of our model. The full list of ROIs used in this

study are listed in Supplementary Table 1.

2.3 Model development and evaluation

2.3.1 Training

The overview of the study is illustrated in Figure 1. We trained SVM binary classifiers with linear

kernel to predict whether an individual will progress to CDR greater than zero using baseline

amyloid PET SUVR, volumetric and non-imaging (i.e., age, sex, *APOE4*) features. We chose to

train SVM classifiers since it has been widely used and extensively validated for AD diagnosis

classification and prediction of future cognitive decline (Rathore et al., 2017). To account for

imbalances in class frequency, we employed class-balanced weighting in which

misclassifications of the minority class (progressor class in most cases) were more heavily

penalized during optimization. Five separate time-to-progression models were trained

independently of one another to predict progressor status, with the time window varied between

one and five years. For example, the three-year model predicts if an individual will progress to a

CDR of greater than zero within three years from baseline. Note that each selected time-to-

progression would result in a different set of progressor participants. However, we used the

same set of stable participants who remained CU for at least five years as the control group in

all five models.

2.3.2 Cross-validation

We employed a leave-one-site-out and leave-one-tracer-out cross-validation framework, in

which data from one site or from one PET tracer were held out for independent testing, while the

remaining data were used for training. Along with five times-to-progression, this resulted in 45

Figure 1. Overview of the study. Analyses were split into *model development and evaluation* (**a-c**) and *A4 trial analyses* (**d-e**). **(a)** Amyloid PET regional SUVRs, MRI regional volumes, and non-imaging features (age, sex, APOE-ε4) were inputted to train support vector machine classifiers to distinguish preclinical AD stables from progressors, where stables are individuals who remain CDR=0 for at least five years from baseline and progressors are individuals who convert to CDR>0 within a specified time window. **(b)** To evaluate model generalizability to unseen data, models were trained on all but one site or one tracer and tested on the hold-out site or tracer. **(c)** To estimate relative importance of feature modalities, nested models were trained on a subset of features and classification performance was compared to the model trained on the full feature set. To obtain individual feature-level importance, linear SVM weights were transformed using the data covariance matrix. **(d)** The SVM classifier trained on all sites except A4 was applied to the A4 cohort to predict progressor status. Only predicted progressors were retained to retroactively enrich the cohort. **(e)** To gauge the benefit of machine learning cohort enrichment, differences in PACC and global amyloid SUVR were tested between placebo and treatment arms post-enrichment. *Abbreviations*. CDR: Clinical Dementia Rating, PACC: Preclinical Alzheimer's Cognitive Composite, SUVR: standardized uptake value ratio, SVM: support vector machine.

total models. All continuous features were z-scored using the standard deviation of only the CU-

stable participants of the training fold, which was shown to enhance separation between linearly

separable groups (Linn et al., 2016). A grid search with five-fold cross-validation was conducted

using the training fold to determine the optimal value of hyperparameter C, which regulates how

strongly misclassifications impact the SVM optimization. We optimized hyperparameter C

independently for each of our 45 models. To evaluate model performance on the leave-out test

set, the receiver operating characteristic area-under-the-curve (ROC-AUC), accuracy, balanced

accuracy, F1 score, sensitivity, specificity, positive predictive value (PPV) and negative

predictive value (NPV) were reported.

2.3.3 Centiloid sensitivity analysis

To address biases in amyloid PET SUVRs due to the inclusion of multiple tracers, we performed

a sensitivity analysis where we harmonized regional SUVRs into regional Centiloids before

inputting as features to the SVM classifier. The Centiloid scale is a method of linearly

transforming raw amyloid PET SUVRs into a scale that is standardized across different tracers

(Klunk et al., 2015). We utilized the conversion equations that were uniquely calibrated to each

site to convert SUVRs to Centiloid, then retrained our models with Centiloids as the amyloid

PET features. Equations were only available for A4, ADNI, MCSA and OASIS, so we included

only these sites in the sensitivity analysis. The DeLong test was used to compare ROC-AUC

between models utilizing regional Centiloids versus regional SUVRs. Bonferroni correction was

applied to p-values to adjust for multiple comparisons.

2.3.4 Feature importance

We assessed the relative importance of each input feature for influencing the classification task

via two methods. First, to estimate the relative importance of each modality of input feature

(amyloid PET SUVR, MRI volume, or non-imaging features), we trained nested models wherein

a reduced subset of modalities was used as input. We then computed the change in ROC-AUC

on the hold-out test set relative to the full model as a measure of the relative contribution of that

modality to the prediction. The DeLong test with Bonferroni correction for multiple comparisons

was performed to test for statistical significance of ROC-AUC differences. Secondly, to derive a

feature-level measure of the strength of association between each input feature and the

progressor class, we computed the covariance-corrected linear SVM weights of the full model,

as proposed by Haufe et al. (2014). This approach transforms classifier weights into

interpretable estimates of the feature-level association with the progressor class, with high

positive values indicating positive association, high negative values indicating negative

association, and values close to zero indicating little to no association. A major advantage of

this is that feature importance estimates are less prone to artificial distortion caused by

collinearity of features, unlike impurity-based metrics for models such as random forest (Strobl

et al., 2008). For each classifier, we applied this method to estimate importance values of each

input feature, and we averaged the values across all 45 classifiers to derive an overall feature

importance. Additionally, to gauge how the magnitude of feature importance varies in relation to

the time-to-progression, we estimated the slope of the absolute value of feature importance

versus time-to-progression via an ordinary least squares linear regression.

2.4 A4 trial analyses

To further validate trained SVM classifiers, we performed three analyses using the extended A4

study cohort as the external testing set. This extended cohort consisted of participants in the

placebo arm and participants treated with Solanezumab, an immunoglobulin monoclonal

antibody designed to increase amyloid clearance in the brain. Firstly, we applied the classifier

trained on all datasets except A4 to classify A4 participants into stables and progressors using

imaging acquired during the screening phase of the trial. We then compared the Preclinical

Alzheimer's Cognitive Composite (PACC), a composite score which has been shown to better

track early AD-related cognitive changes (Donohue et al., 2014), between predicted stables and

progressors. Mirroring previous work (Donohue et al., 2023; Sperling et al., 2023), we fitted a

natural cubic spline model on longitudinal PACC scores as the target variable, using continuous

time-from-baseline as the independent variable and baseline age, years of education, *APOE4*

carriership, PACC test version administered, and baseline FBP SUVR of a cortical composite

region as covariates. We then estimated a model-adjusted mean PACC score and 95%

confidence interval at the 240-week follow-up point (corresponding to the end of the placebo-

controlled study period) and compared the estimates derived for predicted stables versus

predicted progressors using a two-sample t-test. This analysis was independently performed

within the placebo and Solanezumab arms.

Secondly, we applied the SVM classifications to conduct retroactive enrichment wherein we

restricted the cohort to only include predicted progressors. We then tested for treatment effects

post-enrichment in the primary endpoint (model-adjusted PACC) and a secondary endpoint

(change in amyloid PET SUVR). For the primary endpoint, we fitted the same natural cubic

spline model with covariates on the longitudinal PACC scores of the enriched placebo and

treatment cohorts. Model-adjusted means and 95% confidence intervals of PACC at the 240-

week follow-up point were estimated and compared between placebo and Solanezumab arms

using a two-sample t-test. For the secondary endpoint, we computed the change in FBP cortical

SUVR from baseline to the 240-week follow-up point and used this metric as the target variable.

An analysis of covariance (ANCOVA) was fitted to compare the mean change in SUVR between

placebo and Solanezumab arms, controlling for baseline age, *APOE4* carriership and baseline

SUVR as covariates.

Lastly, we performed a bootstrap analysis to estimate the statistical power of detecting a

difference in the change in FBP cortical SUVR between baseline and 240-weeks, testing

different sample sizes and sampling populations (i.e. enriched vs. unenriched). Here, we

focused only on the secondary endpoint rather than the PACC primary endpoint since no

statistically significant treatment effect was reported for the primary endpoint in the original A4

study. For a fixed sample size n, we randomly selected n participants each from the placebo

and Solanezumab arms. The pool of participants to select from was either the full, unenriched cohort or the ML-enriched cohort. We then reran the ANCOVA as conducted previously to test for significant differences in the adjusted change in SUVR between the randomly selected cohorts. We repeated this procedure 1000 times. Power was estimated as the percentage of significant findings across all 1000 iterations. Samples sizes per trial cohort were varied from 10 to 80 in increments of 10.

3 Results

3.1 Cohort characteristics

Table 1 summarizes the cohort characteristics of each site and of the pooled data; p-values were derived from two-sample t-tests for all continuous variables and Fisher's exact test for all categorical variables. Across all sites, progressors were older on average (75.9 ± 6.4) than stables (72.4 ± 5.6, *p < 0.001*) and displayed similar proportions of females-to-males and *APOE4* carriers-to-noncarriers compared to stables (*p > 0.05*). Progressors had a mean time to progression of 2.4 ± 1.3 years and stables remained CU for an average of 7.1 ± 1.6 years. Stables were scanned with a roughly equal proportion of FBP to PiB radiotracer, whereas progressors were more often scanned with FBP (60%). Within individual sites, only age in the A4 cohort displayed a statistically significant difference between stables and progressors across all demographics reported (*p < 0.001*).

3.2 Model evaluation

We evaluated the external site generalizability of the SVM classifier using leave-one-site-out cross-validation. The classifier generalized well to most leave-out sites and times-to-progression, achieving an ROC-AUC of 0.66 or greater, with a majority exceeding 0.75 or above (Figure 2, Table 2). The exception was HABS, for which the ROC-AUC fell below 0.50. We also computed aggregate binary classification metrics by combining the out-of-sample predictions of

Table 1. Cohort characteristics of stables and progressors from all individual sites and the pooled dataset.

	A4		ADNI		AIBL		BLSA		HABS		MCSA		OASIS		pooled	
	stable	progressor	stable	progressor	stable	progressor	stable	progressor	stable	progressor	stable	progressor	stable	progressor	stable	progressor
number of participants	128	111	22	29	28	14	13	11	4	15	94	43	54	24	343	247
age, mean (SD)	70.05 (3.47)	73.47 (5.17)**	74.25 (6.79)	78.14 (6.00)	72.14 (5.78)	74.23 (6.83)	77.08 (7.04)	83.01 (4.77)	75.56 (1.66)	76.28 (5.33)	75.43 (5.34)	79.19 (7.09)	70.75 (6.17)	75.91 (6.68)	72.41 (5.62)	75.89 (6.43)**
female, count (%)	91 (71)	63 (57)	13 (59)	18 (62)	13 (46)	5 (36)	6 (46)	3 (27)	1 (25)	13 (87)	45 (48)	20 (47)	38 (70)	11 (46)	207 (60)	133 (54)
APOE4 carriership, count (%)	78 (61)	63 (57)	6 (27)	11 (38)	12 (43)	10 (71)	2 (15)	5 (45)	2 (50)	9 (60)	38 (40)	26 (60)	29 (54)	12 (50)	167 (49)	136 (55)
amyloid PET tracer, count (%)																
FBP	128 (100)	111 (100)	22 (100)	29 (100)	-	-	-	-	-	-	-	-	20 (37)	10 (42)	170 (50)	150 (61)
PIB	-	-	-	-	28 (100)	14 (100)	13 (100)	11 (100)	4 (100)	15 (100)	94 (100)	43 (100)	34 (63)	14 (58)	173 (50)	97 (39)
years to CDR>0 progression, mean (SD)	-	2.35 (1.23)	-	2.37 (1.26)	-	3.04 (1.37)	-	2.99 (1.34)	-	2.75 (1.54)	-	2.27 (1.05)	-	2.42 (1.49)	-	2.44 (1.27)
years of CU-stable, mean (SD)	6.68 (0.90)	-	7.55 (1.74)	-	8.68 (2.50)	-	7.31 (2.21)	-	5.38 (0.32)	-	6.84 (1.31)	-	7.61 (1.97)	-	7.10 (1.61)	-
years to CDR>0 progression, count (%)																
0-1 year	-	5 (4.5)	-	6 (21)	-	-	-	-	-	3 (20)	-	4 (9.3)	-	7 (29)	-	25 (10)
1-2 years	-	37 (33)	-	6 (21)	-	4 (29)	-	4 (36)	-	4 (27)	-	13 (30)	-	2 (8.3)	-	70 (28)
2-3 years	-	33 (30)	-	8 (28)	-	2 (14)	-	1 (9.1)	-	1 (6.7)	-	17 (40)	-	8 (33)	-	70 (28)
3-4 years	-	20 (18)	-	4 (14)	-	3 (21)	-	2 (18)	-	4 (27)	-	8 (19)	-	2 (8.3)	-	43 (17)
4-5 years	-	16 (14)	-	5 (17)	-	5 (36)	-	4 (36)	-	3 (20)	-	1 (2.3)	-	5 (21)	-	39 (16)

*$p < 0.05$
**$p < 0.01$
***$p < 0.001$

every model on their respective leave-out site and comparing them to the combined ground

truth labels from all sites (Table 2). Classifiers achieved an aggregate accuracy of >0.69 and

balanced accuracy of >0.58. Classifiers were also highly specific (>0.74) and had high negative

predictive value (>0.74). While models also achieved relatively high sensitivity, PPV and F1

score, these metrics were particularly sensitive to time-to-progression. Notably, smaller times-

to-progression resulted in poorer sensitivity, PPV and F1 score, likely due to the fact that

classifiers were trained on a fewer number of positive cases and were prone to overfitting to the

Figure 2. Receiver operating characteristic curves of SVM classifiers trained on each of five times-to-progression. Curves were obtained by applying each trained model to data from its respective hold-out group. Solid lines represent leave-out sites and dotted lines represent leave-out tracers. Across all times-to-progression, SVM classifiers achieved area-under-the-curve of 0.66 or greater in all leave-out sites except for HABS and 0.72 or greater in both leave-out tracers.

control cases. Classifiers trained on higher times-to-progression exhibited a better balance

between sensitivity/PPV to specificity/NPV.

We additionally performed leave-one-tracer-out validation to assess the generalizability of the

SVM classifier to unseen tracers. The classifier achieved ROC-AUC of >0.72 across all times-

to-progression for both leave-out tracers (Table 2). However, the model trained on PiB and

tested on FBP had a low sensitivity and high specificity while the opposite was true for the

model trained on FBP and tested on PiB. This may reflect bias in regional SUVRs between FBP

and PiB tracers. To address this, we retrained the classifiers using Centiloid-harmonized

regional amyloid PET SUVRs and reported the resulting changes in performance

(Supplementary Table 2). After harmonization, models achieved a better balance of sensitivity

to specificity for both leave-out tracers: sensitivity increased when testing on FBP, and

specificity increased when testing on PiB. This was also accompanied by increased aggregate

balanced accuracy across all times-to-progression. ROC-AUC was slightly decreased in some

cases, but none of these changes were statistically significant via the DeLong test (*p > 0.05*).

3.3 Feature importance

The performances of nested models on each hold-out site are shown in Figure 3 and

Supplementary Table 3. When amyloid PET was omitted from the input set, decreases of up to

0.223 in ROC-AUC were observed particularly at time windows of three years or greater (with

the exception of A4), although these differences were not statistically significant ($p > 0.05$).

Table 2. Evaluation metrics of SVM classifiers of all five times-to-progression, tested on each site and tracer as the independent testing set.

		leave-out site/tracer										
	progression time-horizon (yrs)	A4	ADNI	AIBL	BLSA	HABS	MCSA	OASIS	aggregated (site)	FBP	PIB	aggregated (tracer)
Receiver Operating Characteristic Area-under-the-curve (ROC-AUC)	1	0.8062	0.8030	-	-	0.2500	0.8298	0.9074	-	0.7235	0.8282	-
	2	0.7487	0.7576	1.000	0.7500	0.2143	0.8004	0.9630	-	0.7897	0.8352	-
	3	0.7230	0.8364	0.9881	0.6615	0.3750	0.7078	0.8758	-	0.7537	0.7918	-
	4	0.7573	0.8049	0.9603	0.7473	0.4583	0.7538	0.8489	-	0.7647	0.7679	-
	5	0.7478	0.7743	0.8622	0.7902	0.5000	0.7640	0.8395	-	0.7569	0.7904	-
Accuracy	1	0.8872	0.8214	0.9286	0.5385	0.2857	0.6837	0.9016	0.8098	0.9239	0.6630	0.7935
	2	0.7941	0.7353	0.8125	0.5294	0.2727	0.6126	0.8571	0.7306	0.7807	0.6095	0.6986
	3	0.6897	0.8095	0.8235	0.5000	0.5000	0.6719	0.8310	0.7126	0.6960	0.6553	0.6772
	4	0.7265	0.7826	0.8108	0.6000	0.5000	0.6471	0.8082	0.7169	0.6835	0.6102	0.6497
	5	0.7155	0.7451	0.7619	0.6667	0.5263	0.6204	0.7692	0.6983	0.6469	0.6407	0.6441
Balanced Accuracy	1	0.6531	0.5833	0.9286	0.5385	0.2500	0.7154	0.6336	0.5827	0.5328	0.6931	0.5925
	2	0.6953	0.7197	0.8929	0.6923	0.3214	0.6990	0.8704	0.6948	0.6144	0.7205	0.6325
	3	0.6601	0.8114	0.8929	0.5308	0.5000	0.6264	0.7881	0.6865	0.6162	0.7038	0.6273
	4	0.7061	0.7822	0.8373	0.6593	0.5000	0.6722	0.7510	0.7045	0.6419	0.6810	0.6316
	5	0.7051	0.7429	0.7500	0.6853	0.5167	0.6540	0.7176	0.6901	0.6280	0.6812	0.6270
F1 Score	1	0.2105	0.2857	-	-	0	0.1622	0.4000	0.1860	0.1250	0.2051	0.1915
	2	0.5455	0.6400	0.5714	0.5000	0.2000	0.3944	0.6400	0.5042	0.3902	0.4459	0.4261
	3	0.5655	0.8095	0.6667	0.4000	0.5714	0.4615	0.6667	0.5805	0.4196	0.5525	0.4938
	4	0.6391	0.7917	0.6957	0.6000	0.6000	0.5636	0.6316	0.6355	0.4891	0.5892	0.5459
	5	0.6458	0.7719	0.6667	0.7143	0.6400	0.5517	0.6087	0.6397	0.4645	0.6226	0.5513
Sensitivity	1	0.4000	0.1667	-	-	0	0.7500	0.2857	0.3200	0.0714	0.7273	0.3600
	2	0.5000	0.6667	1.000	1.000	0.1429	0.8235	0.8889	0.6316	0.2759	0.8919	0.5158
	3	0.5467	0.8500	1.000	0.6000	0.5000	0.5294	0.7059	0.6121	0.2913	0.8065	0.4848
	4	0.5684	0.7917	0.8889	0.8571	0.5000	0.7381	0.6316	0.6538	0.3543	0.8765	0.5577
	5	0.5586	0.7586	0.7143	0.9091	0.5333	0.7442	0.5833	0.6397	0.3267	0.8247	0.5223
Specificity	1	0.9062	1.000	0.9286	0.5385	0.5000	0.6809	0.9815	0.8455	0.9941	0.6590	0.8251
	2	0.8906	0.7727	0.7857	0.3846	0.5000	0.5745	0.8519	0.7580	0.9529	0.5491	0.7493
	3	0.7734	0.7727	0.7857	0.4615	0.5000	0.7234	0.8704	0.7609	0.9412	0.6012	0.7697
	4	0.8438	0.7727	0.7857	0.4615	0.5000	0.6064	0.8704	0.7551	0.9294	0.4855	0.7055
	5	0.8516	0.7273	0.7857	0.4615	0.5000	0.5638	0.8519	0.7405	0.9294	0.5376	0.7318
Positive Predictive Value (PPV)	1	0.1429	1.000	-	-	0	0.0909	0.6667	0.1311	0.5000	0.1194	0.1304
	2	0.6000	0.6154	0.4000	0.3333	0.3333	0.2593	0.5000	0.4196	0.6667	0.2973	0.3630
	3	0.5857	0.7727	0.5000	0.3000	0.6667	0.4091	0.6316	0.5519	0.7500	0.4202	0.5031
	4	0.7297	0.7917	0.5714	0.4615	0.7500	0.4559	0.6316	0.6182	0.7895	0.4438	0.5346
	5	0.7654	0.7857	0.6250	0.5882	0.8000	0.4384	0.6364	0.6397	0.8033	0.5000	0.5837
Negative Predictive Value (NPV)	1	0.9748	0.8148	1.000	1.000	0.4000	0.9846	0.9138	0.9446	0.9286	0.9744	0.9465
	2	0.8444	0.8095	1.000	1.000	0.2500	0.9474	0.9787	0.8814	0.7941	0.9596	0.8482
	3	0.7444	0.8500	1.000	0.7500	0.3333	0.8095	0.9038	0.8031	0.6867	0.8966	0.7564
	4	0.7248	0.7727	0.9565	0.8571	0.2500	0.8382	0.8704	0.7825	0.6583	0.8936	0.7246
	5	0.6899	0.6957	0.8462	0.8571	0.2222	0.8281	0.8214	0.7405	0.6100	0.8455	0.6802

385   When MRI volumetric features were omitted from the input set, we observed less consistent

386   trends across leave-out sites. For some sites such as A4 and OASIS, slight decreases in ROC-

387   AUC were observed, whereas for other sites such as ADNI and HABS slight increases were

388   observed. All differences were not statistically significant ($p > 0.05$). In many cases, the

389   reduction of model generalizability when amyloid features were omitted was greater than when

Figure 3. Difference in hold-out ROC-AUC of nested models relative to the full model trained on all feature modalities. Dotted lines represent imaging models that omit non-imaging features. In many hold-old sites, omission of amyloid PET features (red) at high times-to-progression led to a decrease in ROC-AUC compared to the full model. Omission of volumetric features (blue) resulted in less consistent trends. Except for BLSA and HABS, the non-imaging only model (purple) performs worse than the full model for nearly all times-to-progression. Omission of non-imaging features (green) led to nearly identical performance compared to the full model.

volumetric features were omitted, especially at later time windows, suggesting a higher relative

importance of amyloid PET in these tasks. In all leave-out sites except for BLSA and HABS, the

non-imaging only model consistently performed worse than the model integrating features from

both imaging modalities, with the difference in ROC-AUC of the one-year model on MCSA being

statistically significant ($p < 0.05$). Finally, omission of non-imaging features led to nearly

identical performance compared to the full model.

Figure 4. Covariance-corrected linear SVM weights per amyloid SUVR (**a**), volumetric (**b**), and non-imaging (**c**) feature. The values shown are the mean importance across all 45 trained models. The ten features with the highest magnitude of mean importance within each imaging modality are shown on the left. Among amyloid SUVR features, temporal, middle & superior frontal and middle occipital gyri had the highest magnitude mean importance. Among volumetric features, lateral ventricle, hippocampus and amygdala had the highest magnitude mean importance. Age was the only non-imaging feature with a high mean importance. Abbreviations. Inf Lat Vent: inferior lateral ventricle, ITG: inferior temporal gyrus, MFG: middle frontal gyrus, MOG: middle occipital gyrus, MTG: middle temporal gyrus, SFG: superior frontal gyrus, STG: superior temporal gyrus

The average importance of each input feature across all trained models is visualized in Figure 4

and Supplementary Fig. 2. For amyloid SUVR features, widespread contributions for predicting

the progressor class were observed throughout the cortex, with bilateral inferior and middle

temporal gyri exhibiting the strongest mean importance, followed by middle frontal, superior

frontal and middle occipital gyri. In contrast, subcortical amyloid SUVR generally had much

lower importance. For volumetric features, bilateral inferior lateral ventricular volumes had the

highest absolute value of feature importance out of all features (including amyloid SUVR and

non-imaging variables). This was followed by lateral ventricle, amygdala and hippocampus

volume. Ventricular volumes all had positive importance values, whereas all other volumetric

features had negative importances. Among the three non-imaging features, baseline age was

most strongly associated with the progressor class. Neither sex nor *APOE4* carriership showed

strong associations.

We also estimated the rate-of-change of the absolute value of feature importance versus the

time-to-progression via a linear regression (Supplementary Fig. 3 & 4). For amyloid SUVR

features, regions in the orbitofrontal, temporal and occipital cortices had the highest rate of

increase in importance as the time-to-progression window grew. Among volumetric regions, the

right lateral inferior ventricles, right frontal operculum and bilateral posterior cingulate gyri had

the highest rate of increase. Age had a modest rate of increase, while sex and *APOE4* did not

show substantial change. Amyloid SUVR features displayed much greater rates-of-change in

feature importance compared to either volumetric features or non-imaging features, suggesting

that their influence on model outputs becomes more pronounced as the time-to-progression

window expands to capture earlier disease progression. This was also consistent with the

trends observed in the nested model analysis.

3.4 A4 trial analysis

We conducted retroactive analyses on the A4 clinical trial dataset to further validate our SVM

classifiers and demonstrate their potential benefit for clinical trial cohort enrichment. Table 3

Table 3. Cohort characteristics of the placebo and Solanezumab treatment arms of the A4 study. The original, unenriched cohort and ML-enriched cohorts are shown.

	cohort					
	original cohort	1-year progressor	2-year progressor	3-year progressor	4-year progressor	5-year progressor
Placebo						
number of participants	444	72	132	158	149	150
participants with FBP PET at 240-week follow-up, count (%)	325 (73)	43 (60)	79 (60)	108 (68)	94 (63)	95 (63)
baseline age, mean (SD)	72.0 (5.0)	75.2 (5.5)***	75.3 (5.2)***	74.6 (5.4)***	75.2 (5.2)***	74.9 (5.3)***
female, count (%)	270 (61)	25 (35)**	60 (45)	86 (54)	74 (50)	78 (52)
APOE4 carriership, count (%)	253 (57)	46 (64)	78 (59)	90 (57)	85 (57)	87 (58)
baseline cortical FBP SUVR, mean (SD)	1.36 (0.17)	1.47 (0.20)**	1.49 (0.18)***	1.46 (0.18)***	1.48 (0.18)***	1.49 (0.18)***
baseline PACC, mean (SD)	0.0 (2.6)	-1.5 (3.1)*	-1.5 (2.9)***	-1.0 (3.1)	-1.3 (2.9)***	-1.2 (3.0)**
annualized PACC rate-of-change, mean (SD)	-0.25 (1.24)	-0.48 (1.44)	-0.49 (1.51)	-0.49 (1.28)	-0.48 (1.45)	-0.48 (1.40)
Solanezumab						
number of participants	404	61	110	140	150	143
participants with FBP PET at 240-week follow-up, count (%)	305 (75)	41 (67)	74 (67)	103 (74)	107 (71)	100 (70)
baseline age, mean (SD)	72.2 (4.5)	74.6 (5.1)*	75.1 (4.4)***	74.3 (4.6)***	74.5 (4.5)***	74.6 (4.5)***
female, count (%)	227 (56)	18 (30)**	56 (51)	74 (53)	75 (50)	71 (50)
APOE4 carriership, count (%)	254 (63)	39 (64)	71 (65)	93 (66)	98 (65)	90 (63)
baseline cortical FBP SUVR, mean (SD)	1.37 (0.16)	1.44 (0.17)	1.51 (0.17)***	1.46 (0.18)***	1.48 (0.17)***	1.49 (0.17)***
baseline PACC, mean (SD)	0.05 (2.77)	-1.53 (2.89)**	-1.25 (2.57)***	-0.96 (2.83)*	-1.01 (2.79)**	-1.11 (2.74)**
annualized PACC rate-of-change, mean (SD)	-0.31 (1.17)	-0.76 (1.43)	-0.78 (1.48)	-0.71 (1.37)	-0.69 (1.40)	-0.73 (1.45)

*p < 0.05
**p < 0.01
***p < 0.001

422   summarizes the characteristics of the unenriched A4 cohort and the retroactively enriched

423   cohorts using each of our five time-to-progression models. In both placebo and Solanezumab

Figure 5. Natural cubic spline models of longitudinal PACC scores of predicted stables and progressors from the A4 study cohort. The model-adjusted PACC trajectory and 95% confidence interval are shown for predicted stables (blue), predicted progressors (red), and the overall cohort (black). The vertical dotted line marks the 240-week follow-up point. The effect size of the contrast between stables and progressors and corresponding p-value are also displayed. Asterisks indicate the significance level (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$). In all cases except for the 1-year progressors of the Solanezumab arm, significant differences were observed between stables and progressors in the model-adjusted PACC at 240-weeks.

424 arms, significant differences between the unenriched and enriched cohorts were observed in

425 baseline age, baseline cortical FBP SUVR (except for the one-year progressors of the

426 Solanezumab arm) and baseline PACC (except for the three-year progressors of the placebo

427   arm) ($p < 0.05$). A lower percentage of females were selected for the enriched cohort in both

428   trial arms compared to the unenriched cohort; this difference was only significant for the one-

429   year progressors ($p < 0.01$). No significant differences were observed in *APOE4* carriership or

430   annualized PACC rate-of-change ($p > 0.05$).

Figure 6. Power to detect differences in change in FBP cortical SUVR between placebo and Solanezumab arms of the A4 study cohort, estimated by bootstrap analysis. Power before (black) and after (colored) retroactive cohort enrichment using SVM classifier predictions are shown. Cohort enrichment by the two-year, four-year and five-year models resulted in increased power at most sample sizes compared to the original, unenriched cohort.

Cohort	Arm	N	Mean ± SE	Lower CI	Upper CI	Effect size	p
1-year progressor	Placebo	43	0.0961 ± 0.02	0.05629	0.1360		
	Solanezumab						
	Contrast	153	-0.0648 ± 0.0201	-0.1043	-0.02535	-0.523 ± 0.165	0.009**
3-year progressor	Placebo	108	0.089 ± 0.0116	0.06612	0.1119		
	Solanezumab	103	0.044 ± 0.0121	0.02016	0.06780		
	Contrast	211	-0.045 ± 0.0164	-0.07713	-0.01291	-0.379 ± 0.139	0.04*
4-year progressor	Placebo	94	0.0782 ± 0.013	0.05267	0.1038		
	Solanezumab	107	0.0215 ± 0.0123	-0.002737	0.04579		
	Contrast	201	-0.0567 ± 0.0173	-0.09064	-0.02278	-0.464 ± 0.143	0.007**
5-year progressor	Placebo	95	0.076 ± 0.0131	0.05015	0.1018		
	Solanezumab	100	0.0175 ± 0.0128	-0.007682	0.04272		
	Contrast	195	-0.0585 ± 0.0178	-0.09328	-0.02367	-0.473 ± 0.146	0.007**

*$p < 0.05$
**$p < 0.01$
***$p < 0.001$

To validate that the SVM classifier accurately distinguish cognitive stables from decliners, we

fitted a natural cubic spline model to longitudinal PACC scores of predicted stables and

progressors and estimated the model-adjusted mean PACC score at the 240-week follow-up. In

all cases except for the one-year progressor model in the Solanezumab arm, predicted stables

had a significantly higher mean estimate of the model-adjusted PACC score at 240 weeks than

predicted progressors ($p < 0.01$) (Figure 5, Supplementary Table 4). When using the five-year

model, which roughly aligns with the duration of the placebo-controlled period of the A4 trial,

predicted progressors of the Solanezumab arm had a mean adjusted PACC score of -3.92 ±

0.57, which was significantly worse (*p < 0.001*) than -0.84 ± 0.26 for predicted stables

(Supplementary Table 4). Similarly, in the placebo group, predicted progressors had a mean

adjusted PACC score of -3.39 ± 0.43 compared to -0.96 ± 0.22 for predicted stables (*p < 0.001*).

To assess whether cohort enrichment by the trained classifiers improves our ability to detect

treatment effects, we tested for significant effects in the primary (PACC) and secondary (FBP

cortical SUVR) endpoints after retroactive enrichment. For PACC, we observed no significant

difference in the model-adjusted mean PACC score between placebo and Solanezumab arms

when the A4 cohort remained unenriched (Table 4), consistent with the conclusions of the

original A4 study (Sperling et al., 2023). The difference in model-adjusted mean PACC

remained non-significant after enrichment by any model (Table 4), indicating no clear benefit of

retroactive enrichment for recovering an underlying treatment effect in the primary outcome.

For FBP cortical SUVR, a significant difference between trial arms was observed in the mean

change in SUVR of the unenriched cohort, with a mean effect size of -0.407 ± 0.08 (*p < 0.001*)

(Table 4), suggesting that a reduction in the rate of amyloid accumulation was associated with

the treatment. After enrichment, this significant difference was preserved in all but the one-year

progressor model. Enrichment by the two, four and five-year models resulted in a modest

increase in the mean effect size of the contrast, albeit with an increase in the standard error due

to reduced sample sizes. Power analysis revealed that for sample sizes between 20 and 80, an

increased power to detect this effect was observed after enrichment by the two, four and five-

year models (Figure 6, Supplementary Fig. 5). However, enrichment by the one-year model

and, paradoxically, the three-year model showed a decrease in power. This pattern aligned with

the results of the ANCOVA in Table 4, wherein mean effect size was increased after enrichment

by the two, four and five-year models, but not the one or three-year models.

4 Discussion

We conducted a rigorous evaluation of machine learning classifiers to predict future cognitive impairment among individuals with preclinical Alzheimer's disease using imaging features from amyloid PET and MRI. Through leave-one-site-out and leave-one-tracer-out cross-validation, we demonstrated that our models consistently achieved strong generalizability to unseen sites and amyloid PET radiotracers, supporting the viability of these models towards enhancing large, multi-consortia clinical trials for AD. Additionally, we demonstrated that these classifiers may be useful in improving the statistical power of detecting treatment effects using real clinical trial data from the A4 study. The present study represents an important and clinically relevant step towards developing and validating stratification models tailored to the asymptomatic, preclinical AD population. This is of particular importance for future investigations of disease-modifying therapies for AD, given that many are trending towards targeting the disease at an early stage. Furthermore, as amyloid PET is commonly utilized for screening participants into clinical trials (Rafii et al., 2023; Sims et al., 2023; Sperling et al., 2023), our findings suggest that there is added value in quantifying the spatial distribution of amyloid accumulation, beyond a simple dichotomization of global amyloid burden, for stratifying and enriching trial cohorts. In terms of clinical applications, current use cases of amyloid PET include establishing the presence of abnormal pathology before administering anti-amyloid therapy (Cummings et al., 2023), although the use of PET in asymptomatic individuals remains an area of active debate (Leuzy et al., 2025; Rabinovici et al., 2025; Rafii and Aisen, 2023). However, we envision, on the premise that treatments intended for asymptomatic preclinical AD individuals are approved in the future, that amyloid PET coupled with predictive models such as ours would serve as valuable prognostic tools for guiding individualized treatment plans. Such applications are anticipated to expand as mechanisms of reimbursement for amyloid PET are relaxed ("Beta Amyloid Positron Emission Tomography in Dementia and Neurodegenerative Disease," 2023).

Several prior studies have demonstrated the ability of ML and imaging features extracted from

amyloid PET to predict progression to cognitive impairment in CU cohorts with high accuracy

(Choi et al., 2024; Dansson et al., 2021; Karaman et al., 2022). The model developed by

Karaman *et al.* achieved an ROC-AUC of 0.82 or higher when predicting the future diagnosis of

CU individuals from ADNI up to 5-years from baseline. Choi *et al.* reported a mean ROC-AUC of

0.87 when applying their DL-ADprob model to an external CU cohort from HABS to predict

future decline at the 4-year follow-up point. In the current study, our classifiers achieved slightly

lower performances at comparable follow-up windows. For most leave-out sites, SVM classifiers

achieved an out-of-sample ROC-AUC of 0.66 or higher across all times to progression,

including ROC-AUC of 0.74 or higher in identifying 4-year progressors or beyond. The notable

exception to this was HABS, which scored an ROC-AUC below 0.5. This likely led to high

sampling variability that potentially accentuated any existing site-specific biases which the

model may not have been robust to. Despite lower comparative performance, we highlight an

important distinction of our study compared to previous literature in that we restricted our cohort

of interest to CU individuals who showed evidence of abnormal brain amyloidosis, which has

been proposed as a core criterion for the biological definition of AD (Jack Jr. et al., 2024).

Discrepancies in performance may potentially be attributable to reduced amyloid signal

available for differentiating stables from progressors. Additionally, unlike the previously reported

classification performance on a HABS CU cohort by (Choi et al., 2024), which was not restricted

to be amyloid-positive, our amyloid-positive CU cohort suffered from low sample sizes and

extreme imbalances in the stables-to-progressors ratio. Nonetheless, our findings still suggest

considerable robustness of SVM classifiers to out-of-sample data for the task of interest, which

is crucial for the viability of such models in clinical trial settings for preclinical AD.

In addition to site generalizability, we also showed good generalizability to different amyloid PET

radiotracers. Models trained on a single amyloid radiotracer achieved ROC-AUC of 0.72 or

greater when applied to the other tracer, even when no explicit tracer harmonization was

performed. This, along with previous studies which have shown ML models to accurately predict

amyloid positivity (Fan et al., 2024) or future cognitive decline (Choi et al., 2024) without explicit

harmonization, may suggest that ML models are capable of finding low-dimensional

representations of high-dimensional PET data which are tracer-agnostic. However, we also

observed that in the context of a binary classification task, the cutoff thresholds (i.e. the

separating hyperplane of the linear SVM) used to determine a binary class decision may not

generalize to out-of-sample tracers, as evidenced by high imbalances in the sensitivity and

specificity. Converting regional amyloid SUVRs into Centiloids produced models that achieved a

much better balance of these metrics, suggesting that explicit harmonization of amyloid PET

measurements still holds value for ML models.

While the global amyloid PET signal has been known to be associated with future cognitive

decline (Landau et al., 2018; Parent et al., 2023; Small et al., 2012), recent studies have

suggested that regional patterns of amyloid accumulation may serve as a more sensitive marker

for the risk of cognitive decline, particularly for preclinical AD (Pascoal et al., 2020; Pfeil et al.,

2021). By leveraging amyloid PET biomarkers at the regional level, our models were able to

discern spatial patterns which discriminate stables from progressors in the preclinical stage.

Among the most important amyloid SUVR features were the temporal, superior and middle

frontal, and middle occipital cortices. Many of these regions were previously identified as

important correlates to increased risk of cognitive decline (Pascoal et al., 2020; Pfeil et al.,

2021). Notably, these regions comprise the intermediate stage of amyloid accumulation

according to PET staging models (Mattsson et al., 2019). In contrast, our models assigned a

lower average feature importance to regions comprising the early accumulation stage, such as

the precuneus and orbitofrontal cortices, suggesting that past a global amyloid threshold, the

intermediate stage regions become more important for discriminating future cognitive decline.

Additionally, low importances were assigned to subcortical structures such as hippocampus and

amygdala, which tend to accumulate amyloid at much later stages of the disease (Hanseeuw et

al., 2018; Thal et al., 2002). In terms of MRI features, lateral ventricular volume and the volumes

of select subcortical regions such as the hippocampus and amygdala had the highest feature

importance. Atrophy in these regions has consistently been identified as robust markers of AD

disease progression (Laakso et al., 1995; Mungas et al., 2002; Nestor et al., 2008). Additionally,

we found that the importance of amyloid SUVR features increased at a higher rate than

volumetric features as the time-to-progression was extended. Our nested model analysis was

consistent with this pattern, wherein at higher times-to-progression, the omission of amyloid

PET features led to greater decreases in model performance and generalizability compared to

either MRI or non-imaging features. These findings may suggest that the influence of amyloid

PET on model predictions becomes increasingly important as the cohort includes more early-

stage progressors. This aligns with current models of AD pathogenesis (Jack Jr. et al., 2024)

where amyloid is expected to rise to abnormality well before neurodegeneration occurs, and

thus may serve as a better indicator of future cognitive decline in the earlier stages of the

disease. Our study notably did not utilize tau PET, which has been shown to closely track

longitudinal cognitive decline (Bejanin et al., 2017; Johnson et al., 2016), and may serve as a

better predictor of the rate of cognitive decline than either amyloid PET or MRI (Ossenkoppele

et al., 2021). Despite this, amyloid PET imaging alone has been shown to provide strong

prognostic value for predicting cognitive decline even within amyloid-positive, cognitively normal

cohorts (Farrell et al., 2017; Pfeil et al., 2021; van der Kall et al., 2021), a notion which is further

supported by our own findings. This is particularly of value for clinical trials for asymptomatic,

preclinical AD, where amyloid PET is more often utilized as a screening criterion compared to

tau PET.

In the context of cohort recruitment for clinical trials, the clinical presentation of AD may remain

highly heterogeneous even after stratification by single risk factors such as global amyloid PET

signal or *APOE4* genotype. Alternatively, ML stratification models trained on a combination of

multimodal risk factors have been shown to produce more accurate forecasts of longitudinal

cognitive decline, leading to greater enhancements in clinical trial operations (Wang et al.,

2024). We successfully identified two distinct subtypes within A4 trial participants that exhibited

different rates of cognitive decline, where predicted progressors experienced a significantly

faster rate of decline compared to stables. However, when applied to cohort enrichment, our

models failed to demonstrate a measurable benefit for enhancing the treatment effect in the

primary cognitive outcome, PACC. This could be attributed to a true lack of an underlying

treatment effect of Solanezumab therapy. Nonetheless, assuming the presence of a ground-

truth drug effect, ML models may still provide a boost in treatment effect detection as evidenced

by previous literature (Birkenbihl et al., 2024; Nallapu et al., 2025; Tam et al., 2022).

Interestingly, in terms of the secondary outcome, cortical amyloid deposition, the two, four and

five-year classifiers identified enriched cohorts that resulted in an increased power for detecting

differences in longitudinal amyloid accumulation. However, this was not the case for either the

one-year model, likely due to insufficient sample size selected, or the three-year model, perhaps

due to selection of a more heterogeneous cohort that consisted of slightly younger participants

with slightly lower cortical SUVRs compared to other enriched cohorts. Given these

contradictory findings and the fact that models were not directly trained to predict longitudinal

amyloid accumulation, our results should be interpreted with caution. Nonetheless, these results

suggest that ML stratification models hold potential to identify trial candidates who are more

likely to respond to treatments, and as more clinical trial data become available, future

investigations should continue to validate ML models for use within this context.

We note several key limitations which represent areas of future improvement. Firstly, our study

did not utilize tau PET, as previously mentioned, and it did not utilize metabolic PET, such as

[^{18}F]-fluorodeoxyglucose PET, which has also been found to be predictive of cognitive decline in

preclinical AD (Mayblyum et al., 2021). For a more comprehensive assessment of AD

biomarkers and their prognostic value, future predictive models should integrate these additional

modalities. Secondly, due to the strict inclusion criteria employed for cohort selection, the

sample sizes in the present study are relatively low compared to other studies employing ML

techniques for AD risk prediction. We combatted this by obtaining data from seven independent

sites and including PET imaging from two tracers. Additionally, our selected cohort suffered

from a high amount of class imbalance, with stable participants heavily outnumbering

progressors especially in low times-to-progression. In an effort to reduce the effect of class

imbalance, SVM models were optimized with a balanced class weighting, and class-balance-

invariant metrics such as balanced accuracy and F1 score were reported. Nonetheless, a larger

sample of preclinical AD individuals, particularly those who eventually progress to clinical

symptoms, is desirable especially for developing more sophisticated ML models such as ones

built upon deep learning. Thirdly, given issues of class imbalance, the stable cohort was limited

to those who remain CN for at least five years from baseline, representative of a control group

with presumably the least amount of pathology. However, a more clinically interpretable model

would ideally match the time of stability to the time-to-progression (i.e. an individual predicted

not to progress within three years would necessarily remain stable for three years). Lastly, each

time-to-progression model was trained independently of the rest, ignoring the constraint that

predicted classifications should be consistent across time windows. Future work will explore

alternate models such as time-to-event models like the survival SVM or the Mirai model for

breast cancer prediction (Yala et al., 2021) which constrain predictions at multiple follow-up

times to be temporally consistent.

5 Conclusion

We performed an extensive validation of machine learning for predicting future cognitive impairment in cognitively unimpaired individuals at the preclinical stage of AD. Our classifiers maintained strong generalizability across most independent testing sites as well as on unseen amyloid PET radiotracers. We further evaluated our models on the A4 clinical trial cohort and observed potential increases in statistical power to detect a secondary treatment effect, indicating potential utility for cohort enrichment. As emerging investigations of new disease-modifying therapies for AD increasingly focus on asymptomatic, preclinical populations, our findings underscore the potential applicability of ML-based patient stratification for recruiting more homogeneous cohorts and improving statistical power for detecting treatment effects for future clinical trials.

6 Data availability

Data used in the preparation of this study are openly available and can be requested at https://www.a4studydata.org, https://adni.loni.usc.edu, https://www.mayo.edu/research/centers-programs/alzheimers-disease-research-center/research-activities/mayo-clinic-study-aging/overview, and https://sites.wustl.edu/oasisbrains. HABS data release 1.10, obtained September 2019, was used in the preparation of this article, and can be requested via https://habs.mgh.harvard.edu. A separate application process is involved for accessing PAC. Investigators interested in accessing the data should contact the PAC Coordinating Center at Johns Hopkins University for details.

SVM was implemented in scikit-learn 1.6.0 using python v3.12.8. All statistical analyses were performed in R v4.4.0. Accompanying code will be made publicly available at https://github.com/sotiraslab.

7 Acknowledgements

This work was supported by the donors of the Alzheimer's Disease Research (ADR) program A2021042S, a program of the BrightFocus Foundation. Research reported in this publication was supported by National Institute on Aging of the National Institutes of Health under award number R01AG067103. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The A4 Study was a secondary prevention trial in preclinical Alzheimer's disease, aiming to slow cognitive decline associated with brain amyloid accumulation in clinically normal older individuals. The A4 Study was funded by a public-private-philanthropic partnership, including funding from the National Institutes of Health-National Institute on Aging, Eli Lilly and Company, Alzheimer's Association, Accelerating Medicines Partnership, GHR Foundation, an anonymous foundation, and additional private donors, with in-kind support from Avid Radiopharmaceuticals, Cogstate, Albert Einstein College of Medicine and the Foundation for Neurologic Diseases. The companion observational Longitudinal Evaluation of Amyloid Risk and Neurodegeneration (LEARN) Study was funded by the Alzheimer's Association and GHR Foundation. The A4 and LEARN Studies were led by Dr. Reisa Sperling at Brigham and Women's Hospital, Harvard Medical School, and Dr. Paul Aisen at the Alzheimer's Therapeutic Research Institute (ATRI) at the University of Southern California. The A4 and LEARN Studies were coordinated by ATRI at the University of Southern California, and the data are made available under the auspices of Alzheimer's Clinical Trial Consortium through the Global Research & Imaging Platform (GRIP). The complete A4 Study Team list is available on: https://www.actcinfo.org/a4-study-team-lists/. We would like to acknowledge the dedication of the study participants and their study partners who made the A4 and LEARN Studies possible. Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012).

ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging

and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's

Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen;

Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.;

Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy

Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development

LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx

Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal

Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The

Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in

Canada. Private sector contributions are facilitated by the Foundation for the National Institutes

of Health (www.fnih.org). The grantee organization is the Northern California Institute for

Research and Education, and the study is coordinated by the Alzheimer's Therapeutic

Research Institute at the University of Southern California. ADNI data are disseminated by the

Laboratory for Neuro Imaging at the University of Southern California. Part of the data used in

the preparation of this article were obtained from the Harvard Aging Brain Study (HABS-

P01AG036694; https://nam10.safelinks.protection.outlook.com/?url=https%3A%2F%2Fhabs.m

gh.harvard.edu%2F&data=05%7C02%7Ctom.earnest%40wustl.edu%7C70bcf8c15ffb44a98484

08dccc4c749f%7C4ccca3b571cd4e6d974b4d9beb96c6d6%7C0%7C0%7C6386098759785146

19%7CUnknown%7CTWFpbGZsb3d8eyJWIjoiMC4wLjAwMDAiLCJQIjoiV2luMzIiLCJBTiI6Ik1ha

WwiLCJXVCI6Mn0%3D%7C0%7C%7C%7C&sdata=uZIOX1693oz1jyd49mvRaqaaIHC%2BXK

DKPHUdotwp%2B8I%3D&reserved=0). The HABS study was launched in 2010, funded by

the National Institute on Aging. and is led by principal investigators Reisa A. Sperling MD and

Keith A. Johnson MD at Massachusetts General Hospital/Harvard Medical School in Boston,

MA. Part of the data contained in this analysis were obtained under one of the following

research grants from the National Institutes of Health to the Mayo Clinic Study of Aging (U01

AG06786, Ronald Petersen, PI) or the Mayo Alzheimer's Disease Research Center (P50

AG16574, Ronald Petersen, PI). Part of the data were provided by OASIS-3: Longitudinal

Multimodal Neuroimaging (Principal Investigators: T. Benzinger, D. Marcus, J. Morris). OASIS-3

was supported by the following funding sources: NIH P30 AG066444, P30 NS09857781, P01

AG026276, P01 AG003991, R01 AG043434, UL1 TR000448, R01 EB009352. AV-45 doses

were provided by Avid Radiopharmaceuticals, a wholly owned subsidiary of Eli Lilly. Avid

Radiopharmaceuticals, Inc., a wholly owned subsidiary of Eli Lilly and company, enabled use of

the 18F-florbetapir tracer, but did not provide direct funding and was not involved in data

analysis or interpretation. The Preclinical AD Consortium is supported by the National Institutes

of Health (NIH), USA [grant number RF1-AG059869]. The individual studies in the consortium

are funded, in part, by the following grants: U19-AG033655, P01-AG026276, RF1-AG027161

and the Australian Commonwealth Scientific Industrial Research Organization (CSIRO); as well

as the National Institute on Aging Intramural Research Program.

8 CRediT authorship contribution statement

Braden Yang: Conceptualization, Investigation, Formal analysis, Methodology, Software,

Visualization, Writing - original draft. **Tom Earnest**: Methodology, Software, Data curation,

Writing - review & editing. **Murat Bilgel**: Methodology, Software, Data curation, Writing - review

& editing. **Marilyn S. Albert**: Data curation. **Sterling C. Johnson**: Data curation. **Christos

Davatzikos**: Data curation. **Guray Erus**: Data curation. **Colin L. Masters**: Data curation. **Susan

M. Resnick**: Data curation, Writing - review & editing. **Michael I. Miller**: Data curation. **Arnold

Bakker**: Data curation. **John C. Morris**: Data curation, Writing - review & editing. **Tammie L.S.

Benzinger**: Conceptualization, Supervision, Data curation, Writing - review & editing. **Brian A.

Gordon**: Conceptualization, Supervision, Writing - review & editing. **Aristeidis Sotiras**:

Conceptualization, Supervision, Project administration, Funding acquisition, Writing - review &

editing.

9 Funding

BY was supported by the Imaging Science Pathways NIH T32 EB014855 and BrightFocus

Foundation grant ADR A2021042S. AS was supported by NIH award R01 AG067103 and

BrightFocus Foundation grant ADR A2021042S.

Computations were performed using the facilities of the Washington University Research

Computing and Informatics Facility, which were partially funded by NIH grants S10OD025200,

1S10RR022984-01A1 and 1S10OD018091-01. Additional support is provided by The

McDonnell Center for Systems Neuroscience.

10 Competing interests

719 AS reported receiving personal fees from BrightFocus for serving as a grant reviewer and stock

720 from TheraPanacea outside the submitted work. All remaining authors have no conflicting

721 interests to report.

11 References

Abdelnour, C., Agosta, F., Bozzali, M., Fougère, B., Iwata, A., Nilforooshan, R., Takada, L.T.,
Viñuela, F., Traber, M., 2022. Perspectives and challenges in patient stratification in
Alzheimer's disease. Alzheimers Res. Ther. 14, 112. https://doi.org/10.1186/s13195-
022-01055-y
Albert, M., Soldan, A., Gottesman, R., McKhann, G., Sacktor, N., Farrington, L., Grega, M.,
Turner, R., Lu, Y., Li, S., Wang, M.-C., Selnes, O., 2014. Cognitive changes preceding
clinical symptom onset of mild cognitive impairment and relationship to ApoE genotype.
Curr. Alzheimer Res. 11, 773–784.
Ballard, C., Atri, A., Boneva, N., Cummings, J.L., Frölich, L., Molinuevo, J.L., Tariot, P.N., Raket,
L.L., 2019. Enrichment factors for clinical trials in mild-to-moderate Alzheimer's disease.
Alzheimers Dement. Transl. Res. Clin. Interv. 5, 164–174.
https://doi.org/10.1016/j.trci.2019.04.001
Basaia, S., Agosta, F., Wagner, L., Canu, E., Magnani, G., Santangelo, R., Filippi, M., 2019.
Automated classification of Alzheimer's disease and mild cognitive impairment using a
single MRI and deep neural networks. NeuroImage Clin. 21, 101645.
https://doi.org/10.1016/J.NICL.2018.101645

Bejanin, A., Schonhaut, D.R., La Joie, R., Kramer, J.H., Baker, S.L., Sosa, N., Ayakta, N.,
Cantwell, A., Janabi, M., Lauriola, M., O'Neil, J.P., Gorno-Tempini, M.L., Miller, Z.A.,
Rosen, H.J., Miller, B.L., Jagust, W.J., Rabinovici, G.D., 2017. Tau pathology and
neurodegeneration contribute to cognitive impairment in Alzheimer's disease. Brain 140,
3286–3300. https://doi.org/10.1093/brain/awx243
Beta Amyloid Positron Emission Tomography in Dementia and Neurodegenerative Disease,
2023.
Birkenbihl, C., de Jong, J., Yalchyk, I., Fröhlich, H., for the Alzheimer's Disease Neuroimaging
Initiative, 2024. Deep learning-based patient stratification for prognostic enrichment of
clinical dementia trials. Brain Commun. 6, fcae445.
https://doi.org/10.1093/braincomms/fcae445
Choi, Y.S., Ngam, P.I., Lee, J.R., Hwang, D., Tan, E.-K., for the Harvard Aging Brain Study, for
the Australian Imaging Biomarkers and Lifestyle Study of Aging, for the Japanese
Alzheimer's Disease Neuroimaging Initiative, for the Alzheimer's Disease Neuroimaging
Initiative, 2024. Deep learning-based amyloid PET harmonization to predict cognitive
decline in non-demented elderly. Radiol. Adv. 1, umae019.
https://doi.org/10.1093/radadv/umae019
Coats, M., Morris, J.C., 2005. Antecedent biomarkers of Alzheimer's disease: The Adult
Children Study. J. Geriatr. Psychiatry Neurol. 18, 242–244.
https://doi.org/10.1177/0891988705281881
Cummings, J., Apostolova, L., Rabinovici, G.D., Atri, A., Aisen, P., Greenberg, S., Hendrix, S.,
Selkoe, D., Weiner, M., Petersen, R.C., Salloway, S., 2023. Lecanemab: appropriate use
recommendations. J. Prev. Alzheimers Dis. 10, 362–377.
https://doi.org/10.14283/jpad.2023.30
Dagley, A., LaPoint, M., Huijbers, W., Hedden, T., McLaren, D.G., Chatwal, J.P., Papp, K.V.,
Amariglio, R.E., Blacker, D., Rentz, D.M., Johnson, K.A., Sperling, R.A., Schultz, A.P.,
2017. Harvard Aging Brain Study: Dataset and accessibility. NeuroImage, Data Sharing
Part II 144, 255–258. https://doi.org/10.1016/j.neuroimage.2015.03.069
Dansson, H.V., Stempfle, L., Egilsdóttir, H., Schliep, A., Portelius, E., Blennow, K., Zetterberg,
H., Johansson, F.D., for the Alzheimer's Disease Neuroimaging Initiative (ADNI), 2021.
Predicting progression and cognitive decline in amyloid-positive patients with
Alzheimer's disease. Alzheimers Res. Ther. 13, 151. https://doi.org/10.1186/s13195-
021-00886-5
Donohue, M.C., Langford, O., Insel, P.S., van Dyck, C.H., Petersen, R.C., Craft, S.,
Sethuraman, G., Raman, R., Aisen, P.S., Initiative, F. the A.D.N., 2023. Natural cubic
splines for the analysis of Alzheimer's clinical trials. Pharm. Stat. 22, 508–519.
https://doi.org/10.1002/pst.2285
Donohue, M.C., Sperling, R.A., Salmon, D.P., Rentz, D.M., Raman, R., Thomas, R.G., Weiner,
M., Aisen, P.S., 2014. The Preclinical Alzheimer Cognitive Composite: measuring
amyloid-related decline. JAMA Neurol. 71, 961–970.
https://doi.org/10.1001/JAMANEUROL.2014.803
Doshi, J., Erus, G., Ou, Y., Resnick, S.M., Gur, R.C., Gur, R.E., Satterthwaite, T.D., Furth, S.,
Davatzikos, C., 2016. MUSE: MUlti-atlas region Segmentation utilizing Ensembles of
registration algorithms and parameters, and locally optimal atlas selection. NeuroImage
127, 186–195. https://doi.org/10.1016/j.neuroimage.2015.11.073
Dyck, C.H. van, Swanson, C.J., Aisen, P., Bateman, R.J., Chen, C., Gee, M., Kanekiyo, M., Li,
D., Reyderman, L., Cohen, S., Froelich, L., Katayama, S., Sabbagh, M., Vellas, B.,
Watson, D., Dhadda, S., Irizarry, M., Kramer, L.D., Iwatsubo, T., 2023. Lecanemab in
early Alzheimer's disease. N. Engl. J. Med. 388, 9–21.
https://doi.org/10.1056/NEJMoa2212948

Ezzati, A., Abdulkadir, A., Jack, C.R., Thompson, P.M., Harvey, D.J., Truelove-Hill, M.,
Sreepada, L.P., Davatzikos, C., Lipton, R.B., 2021. Predictive value of ATN biomarker
profiles in estimating disease progression in Alzheimer's disease dementia. Alzheimers
Dement. 17, 1855–1867. https://doi.org/10.1002/ALZ.12491

Fan, S., Ponisio, M.R., Xiao, P., Ha, S.M., Chakrabarty, S., Lee, J.J., Flores, S., LaMontagne,
P., Gordon, B., Raji, C.A., Marcus, D.S., Nazeri, A., Ances, B.M., Bateman, R.J., Morris,
J.C., Benzinger, T.L.S., Sotiras, A., Haller, S., Atzen, S., 2024. AmyloidPETNet:
Classification of amyloid positivity in brain PET imaging using end-to-end deep learning.
Radiology 311, e231442. https://doi.org/10.1148/radiol.231442

Farrell, M.E., Kennedy, K.M., Rodrigue, K.M., Wig, G., Bischof, G.N., Rieck, J.R., Chen, X.,
Festini, S.B., Devous, M.D., Sr, Park, D.C., 2017. Association of longitudinal cognitive
decline with amyloid burden in middle-aged and older adults: evidence for a dose-
response relationship. JAMA Neurol. 74, 830–838.
https://doi.org/10.1001/jamaneurol.2017.0892

Fowler, C., Rainey-Smith, S.R., Bird, S., Bomke, J., Bourgeat, P., Brown, B.M., Burnham, S.C.,
Bush, A.I., Chadunow, C., Collins, S., Doecke, J., Doré, V., Ellis, K.A., Evered, L.,
Fazlollahi, A., Fripp, J., Gardener, S.L., Gibson, S., Grenfell, R., Harrison, E., Head, R.,
Jin, L., Kamer, A., Lamb, F., Lautenschlager, N.T., Laws, S.M., Li, Q.-X., Lim, L., Lim,
Y.Y., Louey, A., Macaulay, S.L., Mackintosh, L., Martins, R.N., Maruff, P., Masters, C.L.,
McBride, S., Milicic, L., Peretti, M., Pertile, K., Porter, T., Radler, M., Rembach, A.,
Robertson, J., Rodrigues, M., Rowe, C.C., Rumble, R., Salvado, O., Savage, G., Silbert,
B., Soh, M., Sohrabi, H.R., Taddei, K., Taddei, T., Thai, C., Trounson, B., Tyrrell, R.,
Vacher, M., Varghese, S., Villemagne, V.L., Weinborn, M., Woodward, M., Xia, Y.,
Ames, D., 2021. Fifteen years of the Australian Imaging, Biomarkers and Lifestyle (AIBL)
Study: progress and pbservations from 2,359 older adults spanning the spectrum from
cognitive normality to Alzheimer's disease. J. Alzheimers Dis. Rep. 5, 443–468.
https://doi.org/10.3233/ADR-210005

Gladstein, S., Yang, L., Wooten, D., Huang, X., Comley, R., Guo, Q., Initiative, the A.D.N.,
2025. Machine learning–enhanced screening funnel for clinical trials in Alzheimer's
disease. Alzheimers Dement. Transl. Res. Clin. Interv. 11, e70084.
https://doi.org/10.1002/trc2.70084

Hanseeuw, B.J., Betensky, R.A., Mormino, E.C., Schultz, A.P., Sepulcre, J., Becker, J.A.,
Jacobs, H.I.L., Buckley, R.F., LaPoint, M.R., Vannini, P., Donovan, N.J., Chhatwal, J.P.,
Marshall, G.A., Papp, K.V., Amariglio, R.E., Rentz, D.M., Sperling, R.A., Johnson, K.A.,
Initiative, for the A.D.N., Study, and the H.A.B., 2018. PET staging of amyloidosis using
striatum. Alzheimers Dement. 14, 1281–1292. https://doi.org/10.1016/j.jalz.2018.04.011

Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., Bießmann, F.,
2014. On the interpretation of weight vectors of linear models in multivariate
neuroimaging. NeuroImage 87, 96–110.
https://doi.org/10.1016/j.neuroimage.2013.10.067

Huang, L.-K., Chao, S.-P., Hu, C.-J., 2020. Clinical trials of new drugs for Alzheimer disease. J.
Biomed. Sci. 27, 18. https://doi.org/10.1186/s12929-019-0609-7

Jack, C.R., Holtzman, D.M., 2013. Biomarker modeling of Alzheimer's disease. Neuron 80,
1347–1358. https://doi.org/10.1016/j.neuron.2013.12.003

Jack Jr., C.R., Andrews, J.S., Beach, T.G., Buracchio, T., Dunn, B., Graf, A., Hansson, O., Ho,
C., Jagust, W., McDade, E., Molinuevo, J.L., Okonkwo, O.C., Pani, L., Rafii, M.S.,
Scheltens, P., Siemers, E., Snyder, H.M., Sperling, R., Teunissen, C.E., Carrillo, M.C.,
2024. Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's
Association Workgroup. Alzheimers Dement. 20, 5143–5169.
https://doi.org/10.1002/alz.13859

Johnson, K.A., Schultz, A., Betensky, R.A., Becker, J.A., Sepulcre, J., Rentz, D., Mormino, E.,
Chhatwal, J., Amariglio, R., Papp, K., Marshall, G., Albers, M., Mauro, S., Pepin, L.,
Alverio, J., Judge, K., Philiossaint, M., Shoup, T., Yokell, D., Dickerson, B., Gomez-Isla,
T., Hyman, B., Vasdev, N., Sperling, R., 2016. Tau positron emission tomographic
imaging in aging and early Alzheimer disease. Ann. Neurol. 79, 110–119.
https://doi.org/10.1002/ana.24546
Johnson, S.C., Koscik, R.L., Jonaitis, E.M., Clark, L.R., Mueller, K.D., Berman, S.E., Bendlin,
B.B., Engelman, C.D., Okonkwo, O.C., Hogan, K.J., Asthana, S., Carlsson, C.M.,
Hermann, B.P., Sager, M.A., 2018. The Wisconsin Registry for Alzheimer's Prevention:
A review of findings and current directions. Alzheimers Dement. Diagn. Assess. Dis.
Monit. 10, 130–142. https://doi.org/10.1016/j.dadm.2017.11.007
Jutten, R.J., Sikkes, S.A.M., Flier, W.M.V. der, Scheltens, P., Visser, P.J., Tijms, B.M., Initiative,
for the A.D.N., 2021. Finding treatment effects in Alzheimer trials in the face of disease
progression heterogeneity. Neurology 96, e2673–e2684.
https://doi.org/10.1212/WNL.0000000000012022
Karaman, B.K., Mormino, E.C., Sabuncu, M.R., Initiative, for the A.D.N., 2022. Machine
learning based multi-modal prediction of future decline toward Alzheimer's disease: An
empirical study. PLOS ONE 17, e0277322. https://doi.org/10.1371/journal.pone.0277322
Kim, C.K., Lee, Y.R., Ong, L., Gold, M., Kalali, A., Sarkar, J., 2022. Alzheimer's disease: key
insights from two decades of clinical trial failures. J. Alzheimer's Dis. 87, 83–100.
https://doi.org/10.3233/JAD-215699
Klunk, W.E., Koeppe, R.A., Price, J.C., Benzinger, T.L., Devous Sr., M.D., Jagust, W.J.,
Johnson, K.A., Mathis, C.A., Minhas, D., Pontecorvo, M.J., Rowe, C.C., Skovronsky,
D.M., Mintun, M.A., 2015. The Centiloid Project: Standardizing quantitative amyloid
plaque estimation by PET. Alzheimers Dement. 11, 1-15.e4.
https://doi.org/10.1016/j.jalz.2014.07.003
Laakso, M.P., Soininen, H., Partanen, K., Helkala, E.-L., Hartikainen, P., Vainio, P., Hallikainen,
M., Hänninen, T., Riekkinen Sr, P.J., 1995. Volumes of hippocampus, amygdala and
frontal lobes in the MRI-based diagnosis of early Alzheimer's disease: Correlation with
memory functions. J. Neural Transm. - Park. Dis. Dement. Sect. 9, 73–86.
https://doi.org/10.1007/BF02252964
LaMontagne, P.J., Benzinger, T.LS., Morris, J.C., Keefe, S., Hornbeck, R., Xiong, C., Grant, E.,
Hassenstab, J., Moulder, K., Vlassenko, A.G., Raichle, M.E., Cruchaga, C., Marcus, D.,
2019. OASIS-3: longitudinal neuroimaging, clinical, and cognitive dataset for normal
aging and Alzheimer disease. medRxiv 2019.12.13.19014902.
https://doi.org/10.1101/2019.12.13.19014902
Landau, S.M., Horng, A., Jagust, W.J., For the Alzheimer's Disease Neuroimaging Initiative,
2018. Memory decline accompanies subthreshold amyloid accumulation. Neurology 90,
e1452–e1460. https://doi.org/10.1212/WNL.0000000000005354
Landau, S.M., Thomas, B.A., Thurfjell, L., Schmidt, M., Margolin, R., Mintun, M., Pontecorvo,
M., Baker, S.L., Jagust, W.J., the Alzheimer's Disease Neuroimaging Initiative, 2014.
Amyloid PET imaging in Alzheimer's disease: a comparison of three radiotracers. Eur. J.
Nucl. Med. Mol. Imaging 41, 1398–1407. https://doi.org/10.1007/s00259-014-2753-3
Leuzy, A., Bollack, A., Pellegrino, D., Teunissen, C.E., La Joie, R., Rabinovici, G.D.,
Franzmeier, N., Johnson, K., Barkhof, F., Shaw, L.M., Arkhipenko, A., Schindler, S.E.,
Honig, L.S., Moscoso Rial, A., Schöll, M., Zetterberg, H., Blennow, K., Hansson, O.,
Farrar, G., 2025. Considerations in the clinical use of amyloid PET and CSF biomarkers
for Alzheimer's disease. Alzheimers Dement. 21, e14528.
https://doi.org/10.1002/alz.14528
Linn, K.A., Gaonkar, B., Satterthwaite, T.D., Doshi, J., Davatzikos, C., Shinohara, R.T., 2016.
Control-group feature normalization for multivariate pattern analysis of structural MRI

data using the support vector machine. NeuroImage 132, 157–166.
https://doi.org/10.1016/J.NEUROIMAGE.2016.02.044
Mathotaarachchi, S., Pascoal, T.A., Shin, M., Benedet, A.L., Kang, M.S., Beaudry, T., Fonov,
V.S., Gauthier, S., Rosa-Neto, P., 2017. Identifying incipient dementia individuals using
machine learning and amyloid imaging. Neurobiol. Aging 59, 80–90.
https://doi.org/10.1016/J.NEUROBIOLAGING.2017.06.027
Mattsson, N., Palmqvist, S., Stomrud, E., Vogel, J., Hansson, O., 2019. Staging β-amyloid
pathology with amyloid positron emission tomography. JAMA Neurol. 76, 1319–1329.
https://doi.org/10.1001/jamaneurol.2019.2214
Mayblyum, D.V., Becker, J.A., Jacobs, H.I.L., Buckley, R.F., Schultz, A.P., Sepulcre, J.,
Sanchez, J.S., Rubinstein, Z.B., Katz, S.R., Moody, K.A., Vannini, P., Papp, K.V., Rentz,
D.M., Price, J.C., Sperling, R.A., Johnson, K.A., Hanseeuw, B.J., 2021. Comparing PET
and MRI biomarkers predicting cognitive decline in preclinical Alzheimer disease.
Neurology 96, e2933–e2943. https://doi.org/10.1212/WNL.0000000000012108
Misra, C., Fan, Y., Davatzikos, C., 2009. Baseline and longitudinal patterns of brain atrophy in
MCI patients, and their use in prediction of short-term conversion to AD: Results from
ADNI. NeuroImage 44, 1415–1422. https://doi.org/10.1016/j.neuroimage.2008.10.031
Morris, J.C., 1997. Clinical Dementia Rating: a reliable and valid diagnostic and staging
measure for dementia of the Alzheimer type. Int. Psychogeriatr. 9, 173–176.
https://doi.org/10.1017/S1041610297004870
Mungas, D., Reed, B.R., Jagust, W.J., DeCarli, C., Mack, W.J., Kramer, J.H., Weiner, M.W.,
Schuff, N., Chui, H.C., 2002. Volumetric MRI predicts rate of cognitive decline related to
AD and cerebrovascular disease. Neurology 59, 867–873.
https://doi.org/10.1212/WNL.59.6.867
Nallapu, B.T., Petersen, K.K., Qian, T., Demirsoy, I., Ghanbarian, E., Davatzikos, C., Lipton,
R.B., Ezzati, A., for the Alzheimer's Disease Neuroimaging Initiative, 2025. A machine
learning approach to predict cognitive decline in Alzheimer disease clinical trials.
Neurology 104, e213490. https://doi.org/10.1212/WNL.0000000000213490
Nestor, S.M., Rupsingh, R., Borrie, M., Smith, M., Accomazzi, V., Wells, J.L., Fogarty, J.,
Bartha, R., 2008. Ventricular enlargement as a possible measure of Alzheimer's disease
progression validated using the Alzheimer's disease neuroimaging initiative database.
Brain 131, 2443–2454. https://doi.org/10.1093/brain/awn146
Ossenkoppele, R., Smith, R., Mattsson-Carlgren, N., Groot, C., Leuzy, A., Strandberg, O.,
Palmqvist, S., Olsson, T., Jögi, J., Stormrud, E., Cho, H., Ryu, Y.H., Choi, J.Y., Boxer,
A.L., Gorno-Tempini, M.L., Miller, B.L., Soleimani-Meigooni, D., Iaccarino, L., La Joie,
R., Baker, S., Borroni, E., Klein, G., Pontecorvo, M.J., Devous, M.D., Sr, Jagust, W.J.,
Lyoo, C.H., Rabinovici, G.D., Hansson, O., 2021. Accuracy of tau positron emission
tomography as a prognostic marker in preclinical and prodromal Alzheimer disease: a
head-to-head comparison against amyloid positron emission tomography and magnetic
resonance imaging. JAMA Neurol. 78, 961–971.
https://doi.org/10.1001/jamaneurol.2021.1858
Parent, C., Rousseau, L.-S., Predovan, D., Duchesne, S., Hudon, C., 2023. Longitudinal
association between ß-amyloid accumulation and cognitive decline in cognitively healthy
older adults: A systematic review. Aging Brain 3, 100074.
https://doi.org/10.1016/j.nbas.2023.100074
Pascoal, T.A., Therriault, J., Mathotaarachchi, S., Kang, M.S., Shin, M., Benedet, A.L.,
Chamoun, M., Tissot, C., Lussier, F., Mohaddes, S., Soucy, J.P., Massarweh, G.,
Gauthier, S., Rosa-Neto, P., 2020. Topographical distribution of Aβ predicts progression
to dementia in Aβ positive mild cognitive impairment. Alzheimers Dement. Diagn.
Assess. Dis. Monit. 12, e12037. https://doi.org/10.1002/DAD2.12037

Pfeil, J., Hoenig, M.C., Doering, E., van Eimeren, T., Drzezga, A., Bischof, G.N., 2021. Unique regional patterns of amyloid burden predict progression to prodromal and clinical stages of Alzheimer's disease. Neurobiol. Aging 106, 119–129. https://doi.org/10.1016/J.NEUROBIOLAGING.2021.06.014

Price, J.L., Morris, J.C., 1999. Tangles and plaques in nondemented aging and "preclinical" Alzheimer's disease. Ann. Neurol. 45, 358–368. https://doi.org/10.1002/1531-8249(199903)45:3%253C358::AID-ANA12%253E3.0.CO;2-X

Rabinovici, G.D., Knopman, D.S., Arbizu, J., Benzinger, T.L.S., Donohoe, K.J., Hansson, O., Herscovitch, P., Kuo, P.H., Lingler, J.H., Minoshima, S., Murray, M.E., Price, J.C., Salloway, S.P., Weber, C.J., Carrillo, M.C., Johnson, K.A., 2025. Updated appropriate use criteria for amyloid and tau PET: A report from the Alzheimer's Association and Society for Nuclear Medicine and Molecular Imaging Workgroup. Alzheimers Dement. 21, e14338. https://doi.org/10.1002/alz.14338

Rafii, M.S., Aisen, P.S., 2023. Detection and treatment of Alzheimer's disease in its preclinical stage. Nat. Aging 3, 520–531. https://doi.org/10.1038/s43587-023-00410-4

Rafii, M.S., Sperling, R.A., Donohue, M.C., Zhou, J., Roberts, C., Irizarry, M.C., Dhadda, S., Sethuraman, G., Kramer, L.D., Swanson, C.J., Li, D., Krause, S., Rissman, R.A., Walter, S., Raman, R., Johnson, K.A., Aisen, P.S., 2023. The AHEAD 3–45 Study: Design of a prevention trial for Alzheimer's disease. Alzheimers Dement. J. Alzheimers Assoc. 19, 1227–1233. https://doi.org/10.1002/alz.12748

Rathore, S., Habes, M., Iftikhar, M.A., Shacklett, A., Davatzikos, C., 2017. A review on neuroimaging-based classification studies and associated feature extraction methods for Alzheimer's disease and its prodromal stages. NeuroImage 155, 530–548. https://doi.org/10.1016/J.NEUROIMAGE.2017.03.057

Resnick, S.M., Goldszal, A.F., Davatzikos, C., Golski, S., Kraut, M.A., Metter, E.J., Bryan, R.N., Zonderman, A.B., 2000. One-year Age Changes in MRI Brain Volumes in Older Adults. Cereb. Cortex 10, 464–472. https://doi.org/10.1093/cercor/10.5.464

Roberts, R.O., Geda, Y.E., Knopman, D.S., Cha, R.H., Pankratz, V.S., Boeve, B.F., Ivnik, R.J., Tangalos, E.G., Petersen, R.C., Rocca, W.A., 2008. The Mayo Clinic Study of Aging: design and sampling, participation, baseline measures and sample characteristics. Neuroepidemiology 30, 58–69. https://doi.org/10.1159/000115751

Samper-González, J., Burgos, N., Bottani, S., Fontanella, S., Lu, P., Marcoux, A., Routier, A., Guillon, J., Bacci, M., Wen, J., Bertrand, A., Bertin, H., Habert, M.O., Durrleman, S., Evgeniou, T., Colliot, O., 2018. Reproducible evaluation of classification methods in Alzheimer's disease: Framework and application to MRI and PET data. NeuroImage 183, 504–521. https://doi.org/10.1016/J.NEUROIMAGE.2018.08.042

Sims, J.R., Zimmer, J.A., Evans, C.D., Lu, M., Ardayfio, P., Sparks, J., Wessels, A.M., Shcherbinin, S., Wang, H., Monkul Nery, E.S., Collins, E.C., Solomon, P., Salloway, S., Apostolova, L.G., Hansson, O., Ritchie, C., Brooks, D.A., Mintun, M., Skovronsky, D.M., TRAILBLAZER-ALZ 2 Investigators, 2023. Donanemab in early symptomatic Alzheimer disease: the TRAILBLAZER-ALZ 2 randomized clinical trial. JAMA 330, 512–527. https://doi.org/10.1001/jama.2023.13239

Small, G.W., Siddarth, P., Kepe, V., Ercoli, L.M., Burggren, A.C., Bookheimer, S.Y., Miller, K.J., Kim, J., Lavretsky, H., Huang, S.-C., Barrio, J.R., 2012. Prediction of cognitive decline by positron emission tomography of brain amyloid and tau. Arch. Neurol. 69, 215–222. https://doi.org/10.1001/archneurol.2011.559

Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg, H., Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niazy, R.K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances in functional and structural MR image analysis and implementation as FSL. NeuroImage,

Mathematics in Brain Imaging 23, S208–S219.
https://doi.org/10.1016/j.neuroimage.2004.07.051

Sperling, R.A., Donohue, M.C., Raman, R., Rafii, M.S., Johnson, K., Masters, C.L., Dyck, C.H. van, Iwatsubo, T., Marshall, G.A., Yaari, R., Mancini, M., Holdridge, K.C., Case, M., Sims, J.R., Aisen, P.S., 2023. Trial of Solanezumab in preclinical Alzheimer's disease. N. Engl. J. Med. 389, 1096–1107. https://doi.org/10.1056/NEJMoa2305032

Sperling, R.A., Rentz, D.M., Johnson, K.A., Karlawish, J., Donohue, M., Salmon, D.P., Aisen, P., 2014. The A4 study: Stopping AD before symptoms begin? Sci. Transl. Med. 6. https://doi.org/10.1126/SCITRANSLMED.3007941/ASSET/1A7D4501-F70E-47E3-B98C-C19B558DDE40/ASSETS/GRAPHIC/6228FS13-F1.JPEG

Strobl, C., Boulesteix, A.-L., Kneib, T., Augustin, T., Zeileis, A., 2008. Conditional variable importance for random forests. BMC Bioinformatics 9, 307. https://doi.org/10.1186/1471-2105-9-307

Su, Y., Flores, S., Wang, G., Hornbeck, R.C., Speidel, B., Joseph-Mathurin, N., Vlassenko, A.G., Gordon, B.A., Koeppe, R.A., Klunk, W.E., Jack Jr., C.R., Farlow, M.R., Salloway, S., Snider, B.J., Berman, S.B., Roberson, E.D., Brosch, J., Jimenez-Velazques, I., van Dyck, C.H., Galasko, D., Yuan, S.H., Jayadev, S., Honig, L.S., Gauthier, S., Hsiung, G.-Y.R., Masellis, M., Brooks, W.S., Fulham, M., Clarnette, R., Masters, C.L., Wallon, D., Hannequin, D., Dubois, B., Pariente, J., Sanchez-Valle, R., Mummery, C., Ringman, J.M., Bottlaender, M., Klein, G., Milosavljevic-Ristic, S., McDade, E., Xiong, C., Morris, J.C., Bateman, R.J., Benzinger, T.L.S., 2019. Comparison of Pittsburgh compound B and florbetapir in cross-sectional and longitudinal studies. Alzheimers Dement. Diagn. Assess. Dis. Monit. 11, 180–190. https://doi.org/10.1016/j.dadm.2018.12.008

Swanson, C.J., Zhang, Y., Dhadda, S., Wang, J., Kaplow, J., Lai, R.Y.K., Lannfelt, L., Bradley, H., Rabe, M., Koyama, A., Reyderman, L., Berry, D.A., Berry, S., Gordon, R., Kramer, L.D., Cummings, J.L., 2021. A randomized, double-blind, phase 2b proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-Aβ protofibril antibody. Alzheimers Res. Ther. 13, 80. https://doi.org/10.1186/s13195-021-00813-8

Tam, A., Laurent, C., Gauthier, S., Dansereau, C., 2022. Prediction of cognitive decline for enrichment of Alzheimer's disease clinical trials. J. Prev. Alzheimers Dis. 9, 400–409. https://doi.org/10.14283/jpad.2022.49

Thal, D.R., Rüb, U., Orantes, M., Braak, H., 2002. Phases of Aβ-deposition in the human brain and its relevance for the development of AD. Neurology 58, 1791–1800. https://doi.org/10.1212/WNL.58.12.1791

Tustison, N.J., Avants, B.B., Cook, P.A., Zheng, Y., Egan, A., Yushkevich, P.A., Gee, J.C., 2010. N4ITK: Improved N3 bias correction. IEEE Trans. Med. Imaging 29, 1310–1320. https://doi.org/10.1109/TMI.2010.2046908

van der Kall, L.M., Truong, T., Burnham, S.C., Doré, V., Mulligan, R.S., Bozinovski, S., Lamb, F., Bourgeat, P., Fripp, J., Schultz, S., Lim, Y.Y., Laws, S.M., Ames, D., Fowler, C., Rainey-Smith, S.R., Martins, R.N., Salvado, O., Robertson, J., Maruff, P., Masters, C.L., Villemagne, V.L., Rowe, C.C., 2021. Association of β-Amyloid level, clinical progression, and longitudinal cognitive change in normal older individuals. Neurology 96, e662–e670. https://doi.org/10.1212/WNL.0000000000011222

Villemagne, V.L., Mulligan, R.S., Pejoska, S., Ong, K., Jones, G., O'Keefe, G., Chan, J.G., Young, K., Tochon-Danguy, H., Masters, C.L., Rowe, C.C., 2012. Comparison of 11C-PiB and 18F-florbetaben for Aβ imaging in ageing and Alzheimer's disease. Eur. J. Nucl. Med. Mol. Imaging 39, 983–989. https://doi.org/10.1007/s00259-012-2088-x

Wang, C., Tachimori, H., Yamaguchi, H., Sekiguchi, A., Li, Y., Yamashita, Y., 2024. A multimodal deep learning approach for the prediction of cognitive decline and its effectiveness in clinical trials for Alzheimer's disease. Transl. Psychiatry 14, 1–11. https://doi.org/10.1038/s41398-024-02819-w

Wolz, R., Schwarz, A.J., Gray, K.R., Yu, P., Hill, D.L.G., For the Alzheimer's Disease
Neuroimaging Initiative, 2016. Enrichment of clinical trials in MCI due to AD using
markers of amyloid and neurodegeneration. Neurology 87, 1235–1241.
https://doi.org/10.1212/WNL.0000000000003126
Yala, A., Mikhael, P.G., Strand, F., Lin, G., Smith, K., Wan, Y.-L., Lamb, L., Hughes, K.,
Lehman, C., Barzilay, R., 2021. Toward robust mammography-based models for breast
cancer risk. Sci. Transl. Med. 13, eaba4373.
https://doi.org/10.1126/scitranslmed.aba4373
1049