The use of Mendelian randomization to explore the causal consequences of childhood
maltreatment: consideration of assumptions and challenges

Ka Kei Sum,[1,2] Amanda M. Hughes,[1] Alexandra Havdahl,[3,4,5] George Davey Smith,[1,2] Laura
D. Howe[1,2]

[1]MRC Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, Bristol,
UK

[2]Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

[3]PsychGen Centre for Genetic Epidemiology and Mental Health, Norwegian Institute of
Public Health, Oslo, Norway

[4]PROMENTA Research Centre, Department of Psychology, University of Oslo, Oslo, Norway

[5]Nic Waals Institute, Lovisenberg Diaconal Hospital, Oslo, Norway

<u>Abstract</u>

Mendelian randomization (MR) uses genetic variants as instrumental variables to enhance
causal inference. Studies have identified genetic variants related to childhood maltreatment,
but interpreting the effects of these variants or assessing the plausibility of MR assumptions
is complex. We aim to investigate the feasibility of applying MR to complex social traits
using the association between childhood maltreatment and mental health and behavioral
outcomes as an example. We explore four potential key concerns: confounding by population
phenomena, horizontal and vertical pleiotropy, reverse causality, and selection. For each
concern, we demonstrate scenarios where MR studies of childhood maltreatment may be
biased using DAGs and critical appraisal of previous MR analyses. For confounding by
population phenomena, we further perform within-family genetic analyses in 42,101 parent-
offspring trios from the Norwegian Mother, Father and Child Cohort Study (MoBa) to
address bias due to family-level processes since childhood maltreatment often occurs within
households. Our results showed same-trait shrinkage (11% attenuation of the association
between children's polygenic risk scores of childhood maltreatment (PRS_{CM}) and mothers'
report of children's physical abuse) but not cross-trait shrinkage (children's PRS_{CM} and
children's mental health and behavioral outcomes) after adjusting for parental PRS_{CM}. The
lack of cross-trait shrinkage suggests that genetic variants related to childhood maltreatment
may be capturing other child-level phenotypes, after adjusting for family-level processes.
Mothers' PRS_{CM} were also associated with mothers' own maltreatment experiences in
childhood and adulthood with similar magnitudes, suggesting these genetic effects are not
specific to childhood maltreatment. Due to the complexity involved in the causal chain of
childhood maltreatment and it being reported, the interpretation of MR studies for childhood
maltreatment is challenging. Other causal approaches should be considered for observational
studies of complex social traits.

Introduction

Childhood maltreatment encompasses physical, emotional or sexual abuse, and neglect. Maltreatment can occur in the home, in an institution, in the community, or, more rarely, by people unknown to the child. It is most commonly perpetrated by parents or caregivers within the household,[1,2] but in most population-based research, a combined measure of maltreatment is considered, regardless of the perpetrator. Childhood maltreatment is common (global prevalence up to 36%),[2,3] and is strongly associated with worse physical and mental health outcomes[4] across the life course. However, given the strong influence of poverty on childhood maltreatment,[5] it is possible that some of the association between childhood maltreatment and later outcomes reflects confounding by socioeconomic processes. Similarly, parental mental health problems are a risk factor for childhood maltreatment,[6] and given the known genetic component to mental health, some degree of genetic confounding is possible. Moreover, children with neurodevelopmental conditions such as autism and ADHD may be more likely to experience harsh responses from adults due to evocative gene-environment correlations.[7,8] Applying causal inference approaches may therefore help us better understand the causal effects of childhood maltreatment itself, over and above the effects of confounding influences. However, causal inference within the field of childhood maltreatment research is challenging.

Mendelian randomization (MR) is an approach that leverages genetic variation to understand phenotypic causation.[9] It is commonly implemented as an instrumental variable (IV) analysis that uses genetic variants, most commonly single nucleotide polymorphisms (SNPs), as IVs to estimate the causal effect of an exposure on an outcome using observational data.[10] It has been widely used in the field of epidemiology to improve causal inference because of its potential to overcome one of the major barriers to establishing causality in observational studies, namely unmeasured and residual confounding.[11] To infer a causal effect of exposure on outcome, three main MR assumptions need to be met.[10] The assumptions of MR include relevance (genetic IVs are robustly associated with the exposure of interest), independence (there are no common causes between genetic IVs and the outcome of interest), and exclusion restriction (genetic IVs influence the outcome only through the exposure). To quantify the magnitude of a causal estimate, an additional fourth assumption is required.[12–14] There are two versions of this assumption, each providing slightly different interpretations of causal effects. These include homogeneity (the exposure-outcome association does not depend on the value of the IV) and monotonicity (the direction of IV-exposure association is consistent across all participants). Under the homogeneity assumption, MR estimates the average treatment effect, whereas under the monotonicity assumption (a weaker version of homogeneity), MR estimates the local average treatment effect. Depending on the research question, researchers should adopt the version that is most plausible in their context. MR also makes implicit assumption about consistency, a key criterion in establishing causal inference, which states that the outcome under a given exposure value is the same regardless of how that exposure is achieved.[15] In MR, this assumption is known as the gene-environment equivalence principle.[14,16,17] It requires changes in the exposure due to genetic variation to

have the same effect on the outcome as changes induced by modifiable or environmental factors.

MR was initially conceived to study exposures with genetic proxies that have a well-defined biological function. However, due to the proliferation of large genome-wide association studies (GWAS), SNPs have been identified relating to a wide variety of phenotypes, and MR is increasingly used for complex traits where the functional relevance of genetic proxies is less well understood. This makes it more likely that MR assumptions are violated. While the relevance assumption can be empirically tested, independence, exclusion restriction, and gene-environment equivalence assumptions are challenging to falsify.[18]

Many[12,13,19,20] have previously explored scenarios in which genetic instruments are associated with multiple traits in MR and discussed when these scenarios may bias MR results and what are the appropriate causal interpretations. Some of these scenarios include confounding, horizontal and vertical pleiotropy, reverse causation, and selection. Complex social traits are often interlinked and clustered. This means genetic variants associated with one complex social trait are often associated with others. Genetic associations of complex social traits in unrelated individuals such as educational attainment and family socioeconomic position (SEP) have been shown to reflect more than direct genetic effects alone and also include indirect effects due to population phenomena such as population stratification, assortative mating, and dynastic effects.[21–24] Population stratification happens when the genotype-phenotype associations are confounded by population structure, which is systematic differences in allele frequencies between ancestral subpopulations, often after being segregated by geographical location.[21] Dynastic effects refer to the indirect influence of parental genotype on offspring phenotype through its expression in the parental phenotype,[21] sometimes called "the nature of nurture."[25] This phenomenon is also a form of gene-environment correlation,[21] most closely resembling passive gene-environment correlation,[22,26] which states that the parents' genotypes shape both the child's genotype and the environment they provide, thus affecting the child's phenotype.[27] Other than parental indirect genetic effects, the term is sometimes used more broadly to reflect the social stratification processes within or across generations, for example, effects from siblings, relatives, or grandparents.[28] Meanwhile, assortative mating arises when couples choose partners who are more similar to themselves than would be expected by chance with respect to phenotypes such as height and education.[29] When assortative mating happens related to phenotypes that are genetically influenced, couples are likely to be more genetically similar than is typical across the population.[21] In the context of MR, these population phenomena can induce spurious correlations through confounding between genotypes and phenotypes, violating the independence assumption, in turn biasing MR estimates.[14,24,28] This potential bias is illustrated by results of within-family MR studies of complex traits, which by holding parental genotype constant are robust to the influence of population stratification, assortative mating, and dynastic effects. MR studies of unrelated individuals suggest a causal effect of body mass index (BMI) on educational attainment, but within-family MR studies show a null effect, suggesting the association in a standard MR analysis reflects bias from population structure and family-level processes.[28]

MR analyses have previously been used to investigate the relationships between childhood
maltreatment and mental health and behavioral outcomes using SNPs identified from a
GWAS of childhood maltreatment.[30] However, SNPs identified for complex social traits such
as childhood maltreatment are likely to be correlated with other social, environmental, and
mental health traits, making it highly improbable that the independence, exclusion restriction,
and gene-environment equivalence assumptions are met. Since childhood maltreatment is
something that happens to a child, perpetrated by another person, it is implausible that a
child's genetic variants directly cause maltreatment. Rather, they may influence individual
traits (e.g. impulsivity, aggression, emotional regulation) that alter the likelihood of
maltreatment occurring through social interactions, or that alter its reporting/recollection.
These mechanisms align with evocative gene-environment correlation, where the child's
genotype influences his / her behavior, which evokes reactions from others. Moreover,
estimates from 171 countries indicate that the most common perpetrators of childhood
maltreatment are household members of both sexes and various ages, except for sexual
abuse.[1] Similarly in high income countries, approximately 80% of maltreatment cases involve
a parent or other caregiver, meaning maltreatment can be considered a family-level process
for many children.[2] This suggests MR studies of childhood maltreatment in unrelated
individuals may be confounded by parental genotype (*i.e.*, dynastic effects), reflecting passive
gene-environment correlation.

Hence, we aim to explore the feasibility of applying MR to study complex social traits, using
the association between childhood maltreatment and mental health and behavioral outcomes
as an example. We explore four potential key concerns: confounding by population structure,
parental genotype (*i.e.*, dynastic effects, passive gene-environment correlation), and
assortative mating; horizontal and vertical pleiotropy; reverse causality; and selection. For
each of these concerns we (1) demonstrate scenarios where MR studies of childhood
maltreatment and mental health and behavioral outcomes may be biased using directed
acyclic graphs (DAGs), and (2) critically appraise previous MR analyses of childhood
maltreatment.[30] For confounding by population structure, parental genotype, and assortative
mating, we additionally perform within-family analysis using genotype data from parent-
offspring trios in the Norwegian Mother, Father, and Child Cohort Study (MoBa) to explore
bias due to family-level processes.

<u>Key concerns that may bias MR studies of childhood maltreatment</u>

1. Confounding by population structure, parental genotype, and assortative mating

a. Confounding by population structure

160 *b. Confounding by parental genotype (i.e., dynastic effects, passive gene-environment*
161 *correlation)*

162
163

164
165
166
167

169 *c. Confounding by single-trait assortative mating through vertical pleiotropy*

170
171

179 *d. Confounding by single-trait assortative mating through horizontal pleiotropy*

180
181

190   *e. Confounding by cross-trait assortative mating*

Figure 1. *Directed acyclic graph (DAG) representations of confounding by population*
phenomena. The arrow from child genotype to childhood maltreatment is shown to illustrate
potential causal structures, but we do not assume a true direct genetic effect of the child
genotype on maltreatment. (a) Child genotype-outcome association is confounded by
population structure such as ancestral or geographical variation. (b) Child genotype-
outcome association is confounded by parental genotype and parental characteristics / the
environment they provide. (c), (d) and (e) Child genotype-outcome association is confounded
by single-trait and cross-trait assortative mating. Single-trait assortative mating does not
always bias MR estimates. Bias by single-trait assortative mating only happens when the trait
under assortment is genetically correlated with both the exposure and outcome, either
through (c) vertical or (d) horizontal pleiotropy. All the mechanisms described are not
mutually exclusive and multiple mechanisms can happen simultaneously. Unmeasured
confounders are not illustrated.

Previous literature has demonstrated how population phenomena such as population
stratification between ancestral groups, dynastic effects, and assortative mating may bias
genotype-phenotype associations[21,24] and estimates from MR analysis of unrelated
individuals.[13,24,28,31] All these phenomena may bias MR analysis conducted in samples of
unrelated individuals by inducing spurious association between the genetic IV and the
outcome through confounding.[21] MR analysis in family-based samples of related individuals
– either sibling pairs, or mother-father-child 'trios' - is more robust to the impact of these
family-level phenomena, because the genotype of other family members can be controlled
for.[24,28,31]

Figure 1 demonstrates how population structure, dynastic effects, and assortative mating may
bias observed associations in MR studies of childhood maltreatment and mental health and
behavioral outcomes.

1.1. Previous MR analyses of childhood maltreatment

There have been several MR studies of childhood maltreatment published on physical and
mental health outcomes.[30,32–39] Here, we consider the first of these studies by Warrier et al.[30]
A GWAS meta-analysis of childhood maltreatment (N=185,414) was first performed by the
authors using data from five datasets. Variables used to define childhood maltreatment varied
across the five datasets and information about perpetrators was usually not specified. They
subsequently did a bidirectional MR study investigating the impact of childhood
maltreatment on mental health and behavioral outcomes, and vice versa. A 'two-sample' MR,
it drew on summary statistics from the GWAS of both childhood maltreatment and mental
health and behavioral outcomes. Standard inverse-variance weighted (IVW) analysis
suggested an impact of childhood maltreatment on depression (beta=0.59, SE=0.15, p=3.63 x
10^{-5}) and bidirectional associations of childhood maltreatment with schizophrenia (forward
MR of childhood maltreatment on schizophrenia: beta=1.17, SE=0.27, p=1.35 x 10^{-5}; reverse
MR of schizophrenia on childhood maltreatment: beta=0.05, SE=0.006, p=4.83 x 10^{-15}) and
ADHD (forward MR of childhood maltreatment on ADHD: beta=1.04, SE=0.36, p=4.02 x
10^{-3}; reverse MR of ADHD on childhood maltreatment: beta=0.08, SE=0.02, p=1.22 x 10^{-4}).

As summary statistics are usually derived from GWAS in different populations, failure to
account for population phenomena in those GWAS will violate the independence assumption
and bias MR analyses. While all the GWAS included by the authors for both SNP-exposure
and SNP-outcome associations found minimal evidence of genomic inflation due to
population structure after considering the intercept from linkage disequilibrium score
regression (LDSC), removing genetic outliers, and adjusting for principal components, bias
due to dynastic effects and assortative mating may still have been present and can only be
accounted for using a within-family design. For this reason, the authors derived polygenic
risk scores (PRS) of childhood maltreatment based on the genome-wide significant SNPs
from the population-based childhood maltreatment GWAS and performed within-family
analyses of the PGS on retrospective, self-reported childhood maltreatment (total scores
range from 0-20) using sibling data from the UK Biobank in a linear mixed effects model.
The model included two fixed effects, (1) between-family effect (corresponds to the
coefficient of the mean family PRS, *i.e.*, mean PRS of the siblings) and (2) between-sibling
effect (corresponds to the coefficient of the difference in PRS between the individual sibling
and the mean family PRS). They identified a between-sibling (*i.e.*, within-family) effect
(beta=0.053, SE=0.02, p=0.015) of the PRS on retrospective, self-reported childhood
maltreatment, interpreted as the direct genetic effect that is free from familial effects which
maps onto active/evocative gene-environment correlations.[26] This between-sibling effect
accounted for 58% of the total effect of the PRS (*i.e.*, between-family effect) on childhood
maltreatment. However, there was little statistical evidence for passive gene-environment
correlation quantified by the difference in between-family and between-sibling PRS effects
on childhood maltreatment (beta=0.039, SE=0.026, p=0.13; 42% of the total effect). They

also found little attenuation in the variance in prospectively reported childhood maltreatment
explained by a PRS for retrospectively reported childhood maltreatment in a separate cohort
after adjusting for prenatal parental risk factors including smoking, alcohol consumption,
depression, and parental maltreatment, which should partly account for potential dynastic
effects. Thus, the authors concluded active or evocative gene-environment correlation to be
the main mechanism at play, but they could not completely rule out passive gene-
environment correlation.

While a sibling-based design accounts for parental indirect effects, between-sibling estimates
can still be impacted by indirect genetic effects of siblings.[24,40] Also, the two analyses only
considered confounding by parental genotype in the PRS-exposure (same-trait) association
but not the PRS-outcome (cross-trait) association, in this instance, mental health and
behavioral outcomes. This means if dynastic processes affect both childhood maltreatment
and mental health and behavioral outcomes (DAG 1b), bias due to dynastic effects would not
be corrected for in the analyses performed by Warrier et al.[30] since there are still open paths
from parental genotype to the outcomes.[28] Previous literature has also shown cross-trait
shrinkage in within-sibship MR estimates for body mass index (BMI) genetic variants on
educational attainment, age at first birth, and cognitive ability when compared to the
population estimates even in the absence of same-trait shrinkage in the GWAS estimates of
BMI.[24] Currently, there are no other MR studies on childhood maltreatment that have used
study designs to correct for bias due to population phenomena.

1.2. Empirical study using parent-offspring trio genetic data from the Norwegian
Mother, Father, and Child Cohort Study (MoBa)

To account for possible family processes influencing childhood maltreatment and mental
health and behavioral outcomes, we performed within-family analysis using genetic data on
mother-father-offspring trios in the Norwegian Mother, Father, and Child cohort study[41,42]
(MoBa; see detailed methods in supplementary materials; Supp. Fig. 1). Using a trio-based
design to correct for indirect genetic effects has been shown to be less biased compared to
other within-family designs.[40] Briefly, we constructed z-standardized childhood maltreatment
PRS for mothers (PRS_M), fathers (PRS_F), and children (PRS_C) using 12 independent genome-
wide significant SNPs derived from the GWAS of childhood maltreatment performed by
Warrier et al.[30] We then assessed the association between PRS_M and the mothers' reports of
their own childhood and adulthood experiences of maltreatment (physical, sexual, emotional,
verbal abuse) and their own traits of anxiety, depression, and ADHD. The same regressions
were performed for fathers using PRS_F. Lastly, mothers' reports of their children's exposure
to physical abuse, and children's traits of anxiety, depression, and ADHD were regressed on
PRS_C. We then assessed whether associations of PRS_C with the children's mental health and
behavioral outcomes were attenuated after adjustment for parental genotype.

We found statistical evidence of associations between PRS_M and mothers' reports of almost
all maltreatment experiences in both childhood and adulthood (all RRs between 1.04-1.09 per
SD higher of PRS; Supp. Fig. 2), with effect sizes similar for maternal experiences of
maltreatment during childhood and adulthood. However, we did not find statistical evidence

of an association between PRS_F and fathers' reports of their own sexual (RR=1.14 [95%

CI=0.79, 1.63], p=0.49) and physical abuse (RR=1.09 [95% CI=0.96, 1.24], p=0.17)

experiences in the past year during mothers' mid pregnancies. Fathers' reports of sexual and

physical abuse measured during their partners' pregnancies were rare, only 0.18% and 1.42%

respectively, compared to mothers' reports (17.81% and 13.95%) during the same period.

PRS_C was also associated with mothers' reports of their children's exposure to physical abuse

at age 8 years (RR=1.27 [95% CI=1.02, 1.57], p=0.029; Supp. Fig. 3). After adjusting for

parental genotype, the association between PRS_C and mothers' reports of their children's

exposure to physical abuse was partly attenuated (RR=1.13 [95% CI=0.84, 1.53], p=0.42; 11%

attenuation; Supp. Fig. 3), showing same-trait shrinkage.

Mental health and behavioral outcomes were log-transformed because of their right-skewed

residual distributions and were dichotomized if log transformation did not result in normally

distributed residuals. In terms of PRS associations with mental health and behavioral

outcomes (Supp. Fig. 4), there was statistical evidence of associations between PRS_M and

mothers' reports of their own depression and anxiety traits during pregnancy week 30

(RR=1.05 [95%CI=1.00, 1.11], p=0.03), PRS_C and mothers' reports of children's depression

(RR=1.25 [95%CI=1.09, 1.45], p=0.002) and ADHD traits (expected ratio of geometric

means=1.02 [95%CI=1.004, 1.03], p=0.01) at age 8 years, and PRS_C and children's reports of

their own depression traits (expected ratio of geometric means=1.02 [95%CI=1.003, 1.03],

p=0.02) and mothers' reports of children's ADHD traits (expected ratio of geometric

means=1.04 [95%CI=1.03, 1.06], p=2.38 x 10^{-6}) at age 14 years.

Nevertheless, we found little to no attenuation in the estimates between PRS_C and children's

mental health and behavioral outcomes after adjusting for parental genotype (*i.e.*, absence of

cross-trait shrinkage; Supp. Fig. 5). We also found no strong statistical evidence of

associations between PRS_M or PRS_F and children's mental health and behavioral outcomes

before or after accounting for PRS_C, suggesting minimal parental indirect effects and that the

child genotype does not reflect parental genotype (Supp. Fig. 6). We performed sensitivity

analyses by regressing mothers' childhood maltreatment experiences on PRS_C as negative

control and found that PRS_C did not predict mothers' childhood maltreatment experiences

with or without adjusting for PRS_M (Supp. Fig. 7). We also repeated our analyses using a PRS

that was constructed with the inclusion of all SNPs from the childhood maltreatment GWAS

(as opposed to restricting to genome-wide significant SNPs) and observed similar results

(Supp. Fig. 8).

2. Pleiotropy

a. Uncorrelated horizontal pleiotropy

347   *b. Horizontal pleiotropy*

353   *c. Correlated horizontal pleiotropy due to heritable confounding*

362   *d. Vertical pleiotropy through misspecification of the primary phenotype*

Figure 2*. Directed acyclic graph (DAG) representations of non-exhaustive scenarios of*
horizontal pleiotropy. The arrow from child genotype to childhood maltreatment is shown to
illustrate potential causal structures, but we do not assume a true direct genetic effect of the
child genotype on maltreatment. (a) Horizontal pleiotropy happens when genetic variants
believed to influence childhood maltreatment are also associated with child mental and
behavioral outcomes, but there is no causal relationship between childhood maltreatment and
the outcomes of interest. This represents uncorrelated pleiotropy, where the effects of genetic
variants on the exposure and the outcome are independent. (b) Horizontal pleiotropy happens
through the same mechanism as (a) but there is causal relationship between childhood
maltreatment and child mental health and behavioral outcomes. This can be uncorrelated or

 correlated depending on whether the genotype-outcome path is independent of or has shared
mechanism with the genotype-exposure path. (c) Genetic variants believed to influence
childhood maltreatment are associated with a confounder (i.e., child's behavior / personality).
This represents correlated horizontal pleiotropy, where the pleiotropic effect is correlated
with the genetic variants' association with the exposure. It is a case of heritable confounding
because the confounder is a shared heritable factor that is genetically driven. (d) This is
essentially (c) but without the arrow from child genotype to childhood maltreatment. Here,
the genetic variants are primarily associated with the confounder but not with childhood
maltreatment. This represents a subtype of vertical pleiotropy, which biases MR estimates.
Misspecification of primary phenotype is also a case of heritable confounding. Unmeasured
confounders are not illustrated.

One of the biggest threats to the validity of MR studies is pleiotropy, where genetic variants
are associated with multiple phenotypes. There are different types of pleiotropy, mainly
categorized into horizontal and vertical pleiotropy. Horizontal pleiotropy happens when there
is an alternative pathway from the genetic IV to the outcome that is not through the exposure,
thereby violating the exclusion restriction assumption and causing bias in MR estimates.
Horizontal pleiotropy is further divided into uncorrelated and correlated pleiotropy.
Uncorrelated pleiotropy occurs when genetic variants influence the exposure and the outcome
through distinct pathways and the pleiotropic effect on the outcome is independent from the
genetic effect on the exposure.[43] Meanwhile, correlated pleiotropy occurs when genetic
variants influence the exposure and the outcome through a confounder such as a shared
pathway and the pleiotropic effect is correlated with the effect of the genetic variants on the
exposure.[43] Correlated pleiotropy is a consequence of heritable confounding, because the
confounder is driven by the same genetic variants affecting the exposure, and thus
"heritable."[20] On the other hand, vertical pleiotropy happens when genetic variants are
associated with phenotypes that are on the same causal pathway as the exposure and the
outcome. It has been shown that vertical pleiotropy does not usually bias MR estimates
except when it happens through misspecification of the primary phenotype.[14] When genetic
variants are associated with a phenotype that is upstream of the exposure (*i.e.,*
misspecification of the primary phenotype), bias in the MR estimates of the exposure on the
outcome happens through the alternative path from the true exposure to the outcome.
Misspecification of the primary phenotype is a case of heritable confounding, which leads to
correlated pleiotropy.[20]

We present four scenarios of how the genetic IV of childhood maltreatment may influence
child mental health and behavioral outcomes via independent pathways and potentially bias
MR analysis (Figure 2). In DAGs 2a and 2b, child genotype influences both childhood
maltreatment and child mental health and behavioral outcomes independently but in DAG 2b,
child genotype also influences child mental health and behavioral outcomes via childhood
maltreatment. In DAG 2a, there is no true causal effect of childhood maltreatment on child
mental health and behavioral outcomes but they both share the same genetic predictor. DAG
2c and 2d could arise in the situation when a heritable confounder exists between the
exposure and the outcome. In DAG 2c, some of the genetic variants in the child genotype are

associated with the childhood maltreatment through the confounder, whereas in DAG 2d, all
genetic variants in the child genotype are associated with childhood maltreatment through the
confounder. DAG 2d demonstrates a scenario that childhood maltreatment is a proxy for a
separate underlying causal trait that affects mental health and behavioral outcomes, for
instance relating to the child's own characteristics that predispose the child to mental health
conditions. When this underlying causal trait is related to child behavior or personality, this
scenario represents evocative gene-environment correlation, where child genotype influences
child behavior or personality which in turns influences the child's risk of being maltreated,
thus affecting their risk of developing mental health and behavioral outcomes. For example,
children with ADHD and autism may be more likely to experience harsh parenting because of
behaviors perceived as challenging.[7,8] The evocative responses may even be mediated by an
active gene-environment correlation, where the genetically influenced child characteristics
shape the family environment that affects their risk of maltreatment. It is also possible that the
genetic variants may be related to the child's characteristics associated with differential
reporting of childhood maltreatment instead of childhood maltreatment itself. Even if
maltreatment may not be self-reported, parental or caregiver response may differ based on
what the child has disclosed to them. In DAG 2d, the testing of the null hypothesis of no
causal relationship between the exposure and the outcome remains valid because some
aspects of the underlying causal trait are being captured by the mis-specified primary
phenotype. However, the causal effect estimate will be biased.[12]

2.1. Previous MR analyses of childhood maltreatment

Performing multiple pleiotropy-robust MR methods that have different assumptions has been
recommended to increase the credibility of causal claims.[44] Weighted median MR relaxes the
IV assumption by allowing less than half of the variants to be invalid instruments. MR-Egger
allows all the genetic instruments to have pleiotropic effects and includes an intercept term to
estimate the average pleiotropic effect. However, it relies on the Instrument Strength
Independent of Direct Effects (InSIDE) assumption, where variant-exposure associations
should not be associated with the pleiotropic paths, which is often not plausible. Meanwhile,
MR-PRESSO is a variation of inverse-variance weighted (IVW) regression which involves
sequential removal of variants that may have substantially different ratio estimates (*i.e.*,
potential pleiotropic variants) from the rest of the variants. Weighted median MR and MR-
PRESSO are robust to outliers but not MR-Egger. While MR-PRESSO is efficient with valid
IVs, it is likely to have high false positive rate in the presence of invalid IVs. Another
important caveat to all the pleiotropy robust methods is that they can only account for
uncorrelated pleiotropy but less so for correlated pleiotropy.[20,43]

Warrier et al.[30] replicated the results of their standard IVW analysis using both median
weighted MR and MR-PRESSO, whereas MR-Egger only showed statistical evidence for
bidirectional effect for schizophrenia. Evidence of associations between childhood
maltreatment and major depressive disorder and ADHD were not replicated in MR-Egger.
Statistical power in MR-Egger analyses is generally lower than for other MR methods, which
could explain these discrepancies.

Furthermore, their leave-one-out analysis (repeating the MR analysis omitting each SNP in
turn) did not suggest a particular genetic variant was driving the result. The authors also
attempted to understand if childhood maltreatment SNPs might be related to other phenotypes
by identifying previous GWAS associations of those SNPs and performing genetic
correlations between childhood maltreatment SNPs and other health related phenotypes. They
found childhood maltreatment SNPs were previously associated with mental health and
neurodevelopment (depression, ADHD, autism, schizophrenia), personality (neuroticism),
risk-taking behavior, smoking and cannabis use, cognitive performance, and educational
attainment. In genetic correlation analyses, they also found positive genetic correlations with
mental health and behavioral outcomes and negative genetic correlations with intelligence
and educational attainment. The largest magnitudes of genetic correlations were those with
mental health and neurodevelopmental phenotypes including depressive symptoms (r=0.65),
ADHD (r=0.56), and autism (r=0.41).

The authors also performed polygenic transmission disequilibrium tests (pTDT),[45] a within-
family PRS-based method that assesses the likelihood of allele transmission of a certain trait
from unaffected parents to their affected offspring, by quantifying how much the offspring
PRS deviates from the parentally derived expected value. They hypothesized that siblings
who are discordant for neurodevelopmental conditions such as autism or ADHD may prompt
differential parental responses, thus indicating evocative gene-environment correlation. They
found over-transmission of the childhood maltreatment PRS (*i.e.*, where offspring inherited a
higher value of the childhood maltreatment PRS than expected by chance) in autistic children
in two cohorts, suggesting the presence of evocative gene-environment correlation. Yet, there
was no statistical evidence of a relationship between childhood maltreatment and autism in
their MR results. The authors suggested this might have been due to weak instrument bias
and underlying heterogeneity in autism. They also mentioned the observed over-transmission
could arise in the event where childhood maltreatment and autism were driven by a common
cause even in the absence of a true causal relationship between the two.

Despite Warrier et al.'s best effort to comprehensively consider potential instrument validity
and pleiotropic effects by performing pleiotropy-robust MR methods and additional
sensitivity analyses, it is challenging to completely tease apart the scenarios depicted by the
DAGs presented in Figure 2 since they may be statistically indistinguishable. In addition, a
recent simulation study[20] has shown that all standard pleiotropy-robust methods will be
biased in the presence of correlated pleiotropy caused by heritable confounding, especially if
a large proportion of the genetic variants for an exposure are associated with a confounder of
the exposure and the outcome. Given the large magnitudes of genetic correlations observed
between childhood maltreatment and mental health and behavioral outcomes,[46,47] it is also
possible that childhood maltreatment may have shared etiology with these outcomes, even in
the absence of a true causal effect.[12,44]

3. Reverse causation

a. Reverse causation due to heritable confounding

510   *b. Reverse causation due to feedback mechanism*

Figure 3. *Directed acyclic graph (DAG) representations of reverse causation. The subscript*
T refers to timepoint. The arrow from child genotype to childhood maltreatment is shown to
illustrate potential causal structures, but we do not assume a true direct genetic effect of the
child genotype on maltreatment. (a) Reverse causation happens due to genetic variants
primarily affect child mental health and behavioral outcomes rather than childhood
maltreatment. This is essentially the structure of heritable confounding as presented in DAG
2d. The heritable confounder here is the outcome itself. However, it can also be some
psychopathological processes involved before the outcome is developed (dashed box). (b)
Reverse causation occurs as a result of feedback mechanism, where there is a bidirectional
relationship between childhood maltreatment and child mental health and behavioral
outcomes. The variables in the dashed box can be replaced by other psychopathological
processes involved before the outcome is developed. Unmeasured confounders are not
illustrated.

Reverse causation can primarily happen under two scenarios, (1) when the genetic
instruments used for the exposure primarily influences the outcome, which in turn influences
the exposure of interest (DAG 3a) and (2) due to a feedback mechanism, where the exposure
affects the outcome and the outcome affects the exposure at a later time point (DAG
3b).[48] DAG 3a is essentially the same representation of DAG 2d but here the heritable
confounder can be the outcome itself or the psychopathological processes preceding the
outcome. For example, some of the psychopathological processes that affect individuals who
experience childhood adversity are emotional dysregulation and cognitive impairment.[49]
These processes are known to be central to self-regulation and decision making. As discussed

in section 2.1, childhood maltreatment SNPs were highly correlated with mental health and
behavioral outcomes and processes involved in psychopathology. Children who have
impairment in these processes to begin with may influence their risk of being maltreated and
developing mental health and behavioral outcomes. Under the feedback mechanism scenario,
it is also possible that there exists a bidirectional relationship between childhood
maltreatment and these psychopathological processes. Children who are experienced
maltreatment may experience impairment in emotional and cognitive processing,
perpetuating their vulnerabilities to be maltreated. Bidirectional relationships between
childhood maltreatment and mental health and behavioral consequences can be explained by
passive or evocative gene-environment correlations. Under both scenarios, since the outcome
or development process influences the exposure, genetic variants for the outcome or
development process are also associated with the exposure. In large enough sample sizes,
genetic variants associated with the outcome or development process will be detected as
significantly associated with the exposure in a GWAS of the exposure. In such cases of
reverse causation, MR effect estimates are expected to be biased.

3.1. Previous MR analyses of childhood maltreatment

Bidirectional MR can be used to orient the causal direction of effect between two traits using
two corresponding genetic instruments. To correctly identify causal directionality, the two
genetic instruments should be independent from each other. This is as indicated by a Steiger
directionality test, which compares the proportion of variance or pseudo-variance (for
categorical factors) in the exposure and outcome and removes genetic variants that have
greater association with the outcome compared to the exposure. Warrier et al.[30] performed
bidirectional MR to assess the effect of childhood maltreatment on mental health and
behavioral outcomes and vice versa. Steiger directionality tests were also performed by the
authors in their bidirectional MR, supporting the causal direction indicated by all MR
analyses. However, results from a Steiger test may be erroneous in the presence of differential
measurement error, unmeasured confounding, and large differences in sample sizes between
the exposure and outcome GWAS[50,51] and detection of the primary influence of genetic
variants is not guaranteed.[48] In most cases, Steiger test is over-permissive of the genetic
variants included because it is generally implausible for a genetic variant to statistically
explain more than 10% of the variance in the outcome compared with the exposure. For
example, even for a strong risk factor such as smoking accounts for only around 10% of the
pseudo-variance in lung cancer mortality in a cohort.[52] Sensitivity analyses to assess the
impact of a range of samples sizes, varying levels of measurement error, and unmeasured
confounding can be carried out to further understand these sources of bias.[50,51] Given the
challenges of differential measurement error and unmeasured confounding, even in the event
where bidirectional MR and Steiger directionality tests may detect some associations in both
directions, it is also possible that the observed effects may be indicating shared etiological
pathways in the absence of a true causal relationship, *i.e.*, heritable confounding. Besides the
MR analysis done by Warrier et al., only two[35,37] of the other eight MR studies on childhood
maltreatment performed Steiger tests. However, none performed the sensitivity analyses
suggested above to assess the potential biases that may invalidate Steiger test.

4. Selection

Figure 4. *Directed acyclic graph (DAG) representations of selection. The arrow from child genotype to childhood maltreatment is shown to illustrate potential causal structures, but we do not assume a true direct genetic effect of the child genotype on maltreatment. Both childhood maltreatment and child mental health and behavioral outcomes influence the child's selection into the study. Conditioning on the selection factors can cause spurious association between the child genotype and the outcome even if there is no true causal effect. Unmeasured confounders are not illustrated.*

Conditioning on common effects (or descendants) of both exposure and outcome can create spurious exposure-outcome associations, which is also known as collider bias.[53] Selection bias is a specific case of collider bias when the characteristics of participants selected into the study are colliders, *i.e.* are caused by both the exposure of interest (here, childhood maltreatment) and the outcome of interest (here, child mental health and behavioral outcomes).[53] Selection bias is common in most epidemiological studies due to challenges in recruiting participants who are representative of the target population, loss to follow-up, and missing information. Moreover, it is especially problematic in biobanks since participants who volunteered biospecimens are often a subset of people who are fundamentally different from the general population.[54] As such, analyses that involve genetic data are likely to suffer from selection bias. In the case of MR, selection bias invalidates the independence assumption, where the genetic IV and confounders become conditionally dependent, thus biasing MR results by inducing a spurious association between the genetic IV and outcome even in the absence of a true causal effect between exposure and outcome. Selection bias has been shown to bias MR estimates especially for socio-behavioral traits such as education, diet, smoking, and BMI[55] since these traits are also more likely to be associated with the broader socio-economic factors, which influence selection. The magnitude of bias is also shown to be more severe in the presence of strong selection effects, *i.e.*, strong exposure-selection, exposure-confounder, confounder-outcome, and confounder-selection relationships but less so when selection effects are moderate.[56]

People with complete data on childhood maltreatment and mental health and behavioral outcomes have been shown to have higher socioeconomic position (more educational qualifications, higher household income, less material deprivation) in multiple cohorts,[57–59] including those included in Warrier et al.[30] analysis and our empirical analysis in MoBa. Given the strong effects of poverty on the risk of childhood adversity,[60] selection due to social patterning will likely underestimate the MR effect sizes of childhood maltreatment and mental health and behavioral outcomes. However, despite the likely underestimation of estimates, the directionality of estimates is likely to be consistent as previously shown.[55]

Hence, we expect the true association estimates between childhood maltreatment and mental
health and behavioral outcomes to be in the same direction as estimates affected by selection,
but with greater effect sizes.

<u>Potential interpretations and implications of MR studies of childhood maltreatment</u>

Mendelian randomization is a useful tool to uncover causal relationship between two
variables using genetic variants when the IV assumptions are adequately met. In most cases,
these assumptions are more likely to be met when a trait is primarily influenced by intrinsic
biological factors. We attempted to describe scenarios in which MR estimates might be
biased and invalid in the context of a complex social trait, exemplified by childhood
maltreatment. We demonstrated strategies to overcome potential biases by critically
appraising previous MR studies of childhood maltreatment and conducting an empirical study
using trio-based data from a large-scale birth cohort in Norway of childhood maltreatment
and its association with mental health and behavioral outcomes. Our within-family analyses
showed same-trait shrinkage (PRS$_C$ and child physical abuse) but not cross-trait shrinkage
(PRS$_C$ and child mental health and behavioral outcomes) after accounting for dynastic effects
by adjusting for parental PRS. This means that dynastic effects (*i.e.*, passive gene-
environment correlation) partially explained PRS-childhood maltreatment association but not
PRS-child mental health and behavioral outcome associations). Contrary to our results,
Warrier et al.[30] found no strong statistical evidence of passive gene-environment correlation
in their between-sibling study for the PRS-childhood maltreatment association but only
statistical evidence for active / evocative gene-environment correlation. The lack of cross-
trait shrinkage in our empirical analysis suggests that genetic variants related to childhood
maltreatment may be capturing other child-level phenotypes, after adjusting for family-level
processes. Mothers' PRS were also associated similarly with mothers' own maltreatment
experiences in both childhood and adulthood, suggesting these genetic effects are not specific
to childhood maltreatment, consistent with previous finding.[46] Multiple MR pleiotropy-robust
methods and directionality tests led the authors of the original MR study to conclude potential
causal effects of childhood maltreatment on mental health and behavioral traits including
depression, ADHD, and schizophrenia. However, we argue that the interpretation of such MR
findings is extremely challenging due to potential pleiotropic pathways, reverse causality, and
selection even in the context of within-family studies that are more robust to population
confounding (one of the biggest threats to genetic studies of social traits).

As previously mentioned, childhood maltreatment is a complex social phenomenon that
involves multiple people and forces spanning multiple socio-ecological levels. As with most
complex social traits including educational attainment, some genetic variants are likely
associated with childhood maltreatment because of their associations with other individual
and societal factors along the causal chain.[61] Our analysis only provided snapshots of
potential pleiotropic pathways (Figure 2a-2d) that are parts of a bigger social ecosystem. The
complexity involved in the development of a social trait has two implications for MR studies.

First, the interpretation of MR findings becomes extremely difficult. In the absence of
comprehensive biological knowledge about the genetic variants, positive MR evidence may

reflect a combination of causal and non-causal pathways. In particular, the assumption of
gene-environment equivalence is unlikely to hold. It is implausible that genetic variants exert
a direct effect on childhood maltreatment. Instead, they may operate indirectly through
behavioral tendencies or social dynamics that affect how maltreatment occurs or is reported.
Consequently, the pathways through which genetic liability gives rise to maltreatment differ
from those through which maltreatment arises due to social, economic, or contextual factors.
The genetically influenced form of maltreatment may thus differ qualitatively in type, timing,
or severity, violating the assumption that the exposure induced by genetic variation is
equivalent to that experienced in the broader population. Even if genetic liability increases
the risk of maltreatment in one context (e.g., high-stress households), it may be less likely to
do so in another (e.g., supportive environments), meaning gene-environment mapping can be
highly context dependent. This pattern mirrors the findings for *ALDH2* variants in women,
where the expected genotype-exposure association was effectively "zero" where cultural
factors result in low alcohol consumption in this group.[62] Moreover, gene-environment
correlations further confound the interpretation of causal estimates. Together, these issues
make it difficult to satisfy gene-environment equivalence in MR studies of socially mediated
exposures like childhood maltreatment.

Second, the IV assumption of exclusion restriction may rarely or never be met. Since most
childhood maltreatment cases are likely to be to be influenced by some upstream processes,
heritable confounding is unavoidable. Despite the availability of MR pleiotropy-robust
methods to assess instrument validity, many of them require additional assumptions that are
untestable and often implausible such as the InSIDE assumption. This means evidence even
from MR pleiotropy-robust studies is undermined if additional sensitivity analyses[63] such as a
heterogeneity test, leave-one-out analyses, use of positive and negative control outcomes,
colocalization, subgroup analyses, associations with potentially pleiotropic variables,
correcting for selection bias, and investigating reverse causation are not performed. Certainly,
as described previously, in the presence of heritable confounding that leads to correlated
pleiotropy, all standard pleiotropy-robust methods will be biased if a large proportion of the
genetic variants are associated with the heritable confounder.[20] The guidelines developed to
perform[63] and report[16] MR studies can be useful to help researchers consider whether the
research question at hand can be appropriately addressed using MR by thinking through the
plausibility of assumptions and challenges in interpretations.

Moreover, the ethical implications of conducting MR studies of childhood maltreatment can
be far-reaching. Debates surrounding the risks and benefits of the wider research of social and
behavioral genomics (SBG) have been ongoing since such studies began. Recently, The
Hastings Center[61] further classifies certain SBG studies (largely focused on GWAS and PRS)
to be of heightened concern especially those that involve sensitive phenotypes, *i.e.*,
"phenotypes influencing one's social standing, historically have been part of harmful
stereotypes about minoritized groups and threaten to reify the biologization of social
identities, or central to a minoritized group's identity." They recommended that such studies
should be conducted and communicated responsibly to reduce potential harms. We argue that
genetic research specifically surrounding GWAS[30,64] and creation of PRS of childhood

maltreatment should fall under the category of sensitive phenotypes since childhood
maltreatment is not only highly damaging but can lead to social stigma for individuals who
have experienced it. Even in the context of well-conducted and carefully communicated
studies, there is a risk that studying the individual-level genetic correlates of childhood
maltreatment could add to stigmatization, discrimination, and fatalism at the individual,
group, and policy levels.[61]

That said, while using MR to instrument complex social exposures may not be the best
approach to give us causal answers for the reasons stated above, MR can sometimes be
valuable to assess the extent of confounding. In particular, MR can be used to instrument
potential confounders instead of the social exposure itself. An example is the application of
MR to understand the determinants of selection into shift work, where genetic instruments for
cardiometabolic risk factors are used to estimate the selection effect of these risk factors on
shift work.[65] These estimates are useful to inform the extent to which these risk factors may
confound the association between shift work and adverse health outcomes. A second example
of using germline genetics to understand confounding relates to interleukin 6 (IL-6) activity
effects on high-density lipoprotein (HDL) cholesterol, and how these confound the
associations of HDL cholesterol with various health outcomes.[66] Other genetic methods may
be useful to help disentangle causal effects of childhood maltreatment from confounding.
These include family-based designs such as adoption, children of twin, and sibling
comparison studies. Some of these designs can also help us to better understand mechanisms
of transmission and gene-environment correlation without the use of GWAS and PRS.[67]
However, PRS can also be useful in cases when it is used in a within family design such as
the pTDT approach mentioned earlier or sensitivity analyses to account for genetic
confounding.[68] Other non-genetic designs such as marginal structural models and structural
nested models that allow modelling of time-varying exposures and confounders are useful for
increasing evidence credibility.[69] Natural experiments that examine the effects of childhood
maltreatment through wider social or political events can also help overcome biases and
limitations of traditional observational studies.[70] This is exemplified by the English and
Romanian Adoptees study[71] that examines the long term mental health outcomes of
institutional neglect experienced by young children as a result of extreme early life
deprivation that occurred during the Ceau□escu regime in Romania before being adopted by
UK families.
736
Conclusion

The explosion of GWAS has enabled us to study the genetic predisposition of many complex
traits and use genetic variants as instruments for MR studies. MR studies are only valid when
the instrumental variable assumptions are met, and they are less likely to be met for traits that
are extrinsic to an individual, complex, and socially or environmentally driven. In the context
of childhood maltreatment, the underlying causal mechanism linking a person's genotype to
their risk of exposure is likely to be complex and multi-factorial. As demonstrated in our
study, even in a well-executed MR analysis, the interpretation is extremely challenging due to
potential pleiotropic pathways and gene-environment correlations. Established MR

pleiotropy-robust methods and sensitivity analyses have their own limitations as certain
assumptions remain untestable and often implausible for complex social traits. There are also
important ethical considerations regarding GWAS and PRS research on traits which are more
common in vulnerable groups or may be subject to stigmatization. We urge researchers to be
mindful when implementing MR for complex social traits such as childhood maltreatment
and consider other approaches that may help strengthen causal inference in observational
studies.

Acknowledgements

This study includes data from the Norwegian Mother, Father and Child Cohort Study (MoBa)
conducted by the Norwegian Institute of Public Health. We are grateful to all the participating
families in Norway who take part in this on-going cohort study. For generating high-quality
genomic data, we thank the Norwegian Institute of Public Health (NIPH), the HARVEST
collaboration, the NORMENT Centre at the University of Oslo, the Center for Diabetes
Research at the University of Bergen, deCODE Genetics, the Research Council of Norway,
the South-Eastern and Western Norway Regional Health Authorities, the ERC AdG, Stiftelsen
KG Jebsen, the Trond Mohn Foundation, and the Novo Nordisk Foundation. This work was
performed on the Tjeneste for Sensitive Data (TSD) facilities, owned by the University of
Oslo, operated and developed by the TSD service group at the University of Oslo, IT-
Department (USIT), using resources provided by Sigma2—the National Infrastructure for
High Performance Computing and Data Storage in Norway (UNINETT).

Funding statement

This research was funded in whole, or in part, by the Wellcome Trust. KKS is supported by a
Wellcome PhD studentship in Molecular, Genetic, and Lifecourse Epidemiology
(218495/Z/19/Z). LDH is supported by Wellcome Trust (309183/Z/24/Z). KKS, AMH, GDS,
and LDH work in a unit that receives funding from the University of Bristol and the UK
Medical Research Council (MC_UU_00032/1, MC_UU_00032/2). AH is supported by the
Norwegian Research Council (336085) and the Norwegian Health Authority (2020022,
#2022029 and #2022083). The Norwegian Mother, Father and Child Cohort Study is
supported by the Norwegian Ministry of Health and Care Services and the Ministry of
Education and Research. For the purpose of Open Access, the author has applied a CC BY
public copyright licence to any Author Accepted Manuscript version arising from this
submission.

Data availability

Data from the Norwegian Mother, Father and Child Cohort Study is managed by the
Norwegian Institute of Public Health. Access requires approval from the Regional
Committees for Medical and Health Research Ethics (REC), compliance with GDPR, and

data owner approval. Participant consent does not allow individual-level data storage in
repositories or journals. Researchers seeking access for replication must apply via
www.helsedata.no.

<u>References</u>

1. Devries K, Knight L, Petzold M, et al. Who perpetrates violence against children? A
systematic analysis of age-specific and sex-specific data. *BMJ Paediatr Open*.
2018;2(1):e000180. doi:10.1136/bmjpo-2017-000180

2. Gilbert R, Widom CS, Browne K, Fergusson D, Webb E. Burden and consequences of
child maltreatment in high-income countries. *The Lancet*. 2009;373:68-81.
doi:10.1016/S0140

3. Stoltenborgh M, Bakermans☐Kranenburg MJ, Alink LRA, van IJzendoorn MH. The
prevalence of child maltreatment across the globe: review of a series of meta☐analyses.
Child Abuse Review. 2015;24(1):37-50. doi:10.1002/car.2353

4. Carr A, Duff H, Craddock F. A systematic review of reviews of the outcome of
noninstitutional child maltreatment. *Trauma Violence Abuse*. 2020;21(4):828-843.
doi:10.1177/1524838018801334

5. Bywaters P, Skinner G, Cooper A, Eilis K, Malik A. *The Relationship between Poverty*
and Child Abuse and Neglect: New Evidence.; 2022.

6. Ayers S, Bond R, Webb R, Miller P, Bateson K. Perinatal mental health and risk of
child maltreatment: a systematic review and meta-analysis. *Child Abuse Negl*.
2019;98:104172. doi:10.1016/j.chiabu.2019.104172

7. Hamshere ML, Langley K, Martin J, et al. High loading of polygenic risk for ADHD in
children with comorbid aggression. *American Journal of Psychiatry*. 2013;170(8):909-
916. doi:10.1176/appi.ajp.2013.12081129

8. Warrier V, Baron-Cohen S. Childhood trauma, life-time self-harm, and suicidal
behaviour and ideation are associated with polygenic scores for autism. *Mol Psychiatry*.
2019;26(5):1670-1684. doi:10.1038/s41380-019-0550-x

9. Davey Smith G. Mendelian randomization for strengthening causal inference in
observational studies: application to gene × environment interactions. *Perspectives on*
Psychological Science. 2010;5(5):527-545. doi:10.1177/1745691610383505

10. Lawlor DA, Harbord RM, Sterne JAC, Timpson N, Davey Smith G. Mendelian
randomization: using genes as instruments for making causal inferences in
epidemiology. *Stat Med*. 2008;27(8):1133-1163. doi:10.1002/sim.3034

11. Davey Smith G, Ebrahim S. Mendelian randomization: prospects, potentials, and
limitations. *Int J Epidemiol*. 2004;33(1):30-42. doi:10.1093/ije/dyh132

12.   Burgess S, Butterworth AS, Thompson JR. Beyond Mendelian randomization: how to
interpret evidence of shared genetic predictors. *J Clin Epidemiol*. 2016;69:208-216.
doi:10.1016/j.jclinepi.2015.08.001

13.   Yang Q, Sanderson E, Tilling K, Borges MC, Lawlor DA. Exploring and mitigating
potential bias when genetic instrumental variables are associated with multiple non-
exposure traits in Mendelian randomization. *Eur J Epidemiol*. 2022;37(7):683-700.
doi:10.1007/s10654-022-00874-5

14.   Sanderson E, Glymour MM, Holmes M V., et al. Mendelian randomization. *Nature*
Reviews Methods Primers. 2022;2(1):6. doi:10.1038/s43586-021-00092-5

15.   VanderWeele TJ. Concerning the consistency assumption in causal inference.
Epidemiology. 2009;20(6):880-883. doi:10.1097/EDE.0b013e3181bd5638

16.   Skrivankova VW, Richmond RC, Woolf BAR, et al. Strengthening the reporting of
observational studies in epidemiology using Mendelian randomisation (STROBE-MR):
explanation and elaboration. *BMJ*. Published online October 26, 2021:n2233.
doi:10.1136/bmj.n2233

17.   Ebrahim S, Davey Smith G. Mendelian randomization: can genetic epidemiology help
redress the failures of observational epidemiology? *Hum Genet*. 2008;123(1):15-33.
doi:10.1007/s00439-007-0448-6

18.   Davies NM, Holmes M V., Davey Smith G. Reading Mendelian randomisation studies:
a guide, glossary, and checklist for clinicians. *BMJ (Online)*. 2018;362.
doi:10.1136/bmj.k601

19.   Hemani G, Bowden J, Davey Smith G. Evaluating the potential role of pleiotropy in
Mendelian randomization studies. *Hum Mol Genet*. 2018;27(R2):R195-R208.
doi:10.1093/hmg/ddy163

20.   Sanderson E, Rosoff D, Palmer T, Tilling K, Davey Smith G, Hemani G. Heritable
confounding in Mendelian randomization studies: structure, consequences and
relevance for gene-environment equivalence. *medRxiv*. Published online August 28,
2025. doi:10.1101/2024.09.05.24312293

21.   Morris TT, Davies NM, Hemani G, Davey Smith G. Population phenomena inflate
genetic associations of complex social traits. *Sci Adv*. 2020;6(16).
doi:10.1126/sciadv.aay0328

22.   Lin Y, Procopio F, Keser E, et al. Polygenic score prediction within and between
sibling pairs for intelligence, cognitive abilities, and educational traits from childhood
to early adulthood. *Intelligence & Cognitive Abilities*. 2025;1(1):5-19.
https://www.teds.ac.uk/datadictionary/home.htm

23.   Tan T, Jayashankar H, Guan J, et al. Family-GWAS reveals effects of environment and
mating on genetic associations. *medRxiv*. Published online October 4, 2024.
doi:10.1101/2024.10.01.24314703

24. Howe LJ, Nivard MG, Morris TT, et al. Within-sibship genome-wide association
analyses decrease bias in estimates of direct genetic effects. *Nat Genet*.
2022;54(5):581-592. doi:10.1038/s41588-022-01062-7

25. Kong A, Thorleifsson G, Frigge ML, et al. The nature of nurture: effects of parental
genotypes. *Science (1979)*. 2018;359(6374):424-428. doi:10.1126/science.aan6877

26. Selzam S, Ritchie SJ, Pingault JB, Reynolds CA, O'Reilly PF, Plomin R. Comparing
within- and between-family polygenic score prediction. *Am J Hum Genet*.
2019;105(2):351-363. doi:10.1016/j.ajhg.2019.06.006

27. Jaffee SR, Price TS. Genotype–environment correlations: implications for determining
the relationship between environmental exposures and psychiatric illness. *Psychiatry*.
2008;7(12):496-499. doi:10.1016/j.mppsy.2008.10.002

28. Brumpton B, Sanderson E, Heilbron K, et al. Avoiding dynastic, assortative mating,
and population stratification biases in Mendelian randomization through within-family
analyses. *Nat Commun*. 2020;11(1). doi:10.1038/s41467-020-17117-4

29. Yengo L, Robinson MR, Keller MC, et al. Imprint of assortative mating on the human
genome. *Nat Hum Behav*. 2018;2(12):948-954. doi:10.1038/s41562-018-0476-3

30. Warrier V, Kwong ASF, Luo M, et al. Gene–environment correlations and causal
effects of childhood maltreatment on physical and mental health: a genetically
informed approach. *Lancet Psychiatry*. 2021;8(5):373-386. doi:10.1016/S2215-
0366(20)30569-1

31. Hartwig FP, Davies NM, Davey Smith G. Bias in Mendelian randomization due to
assortative mating. *Genet Epidemiol*. 2018;42(7):608-620. doi:10.1002/gepi.22138

32. Konzok J, Gorski M, Winkler TW, et al. Child maltreatment as a transdiagnostic risk
factor for the externalizing dimension: a Mendelian randomization study. *Mol
Psychiatry*. 2025;30(2):567-573. doi:10.1038/s41380-024-02700-8

33. Kim MH, Turecki G, Orri M. Investigating the contribution of childhood maltreatment
to suicide attempt: a multivariable Mendelian randomization study. *Psychiatry Res*.
2024;342:116278. doi:10.1016/j.psychres.2024.116278

34. Zhang Z, Ren H, Han R, et al. Impact of childhood maltreatment on aging: a
comprehensive Mendelian randomization analysis of multiple age-related biomarkers.
Clin Epigenetics. 2024;16(1):103. doi:10.1186/s13148-024-01720-z

35. Alkema A, Marchi M, van der Zaag JAJ, et al. Childhood abuse v. neglect and risk for
major psychiatric disorders. *Psychol Med*. 2024;54(8):1598-1609.
doi:10.1017/S0033291723003471

36. Baltramonaityte V, Karhunen V, Felix JF, et al. Biological pathways underlying the
relationship between childhood maltreatment and multimorbidity: a two-step,

multivariable Mendelian randomisation study. *Brain Behav Immun*. Published online
February 2025. doi:10.1016/j.bbi.2025.01.024

37. Baltramonaityte V, Pingault JB, Cecil CAM, et al. A multivariate genome-wide
association study of psycho-cardiometabolic multimorbidity. *PLoS Genet*.
2023;19(6):e1010508. doi:10.1371/journal.pgen.1010508

38. Zavitsanou G, Waldren LH, Walton E, Baltramonaityte V. The role of loneliness and
social isolation in mediating the relationship between childhood maltreatment and
schizophrenia: a genetically informed approach. *Journal of Psychopathology and*
Clinical Science. 2024;133(5):392-402. doi:10.1037/abn0000902

39. Chen TT, Chen CY, Liu CY, et al. Genetic architectures of childhood maltreatment and
causal influence of childhood maltreatment on health outcomes in adulthood. *Mol*
Psychiatry. Published online February 20, 2025. doi:10.1038/s41380-025-02928-y

40. Demange PA, Hottenga JJ, Abdellaoui A, et al. Estimating effects of parents' cognitive
and non-cognitive skills on offspring education using polygenic scores. *Nat Commun*.
2022;13(1). doi:10.1038/s41467-022-32003-x

41. Corfield EC, Frei O, Shadrin AA, et al. The Norwegian Mother, Father, and Child
cohort study (MoBa) genotyping data resource: MoBaPsychGen pipeline v.1.
doi:10.1101/2022.06.23.496289

42. Brandlistuen RE, Kristjansson D, Alsaker E, et al. Cohort profile update: the
Norwegian Mother, Father and Child Cohort (MoBa). *Int J Epidemiol*. 2025;54(5).
doi:10.1093/ije/dyaf139

43. Morrison J, Knoblauch N, Marcus JH, Stephens M, He X. Mendelian randomization
accounting for correlated and uncorrelated pleiotropic effects using genome-wide
summary statistics. *Nat Genet*. 2020;52(7):740-747. doi:10.1038/s41588-020-0631-4

44. Burgess S, Woolf B, Mason AM, Ala-Korpela M, Gill D. Addressing the credibility
crisis in Mendelian randomization. *BMC Med*. 2024;22(1):374. doi:10.1186/s12916-
024-03607-5

45. Weiner DJ, Wigdor EM, Ripke S, et al. Polygenic transmission disequilibrium
confirms that common and rare variation act additively to create risk for autism
spectrum disorders. *Nat Genet*. 2017;49(7):978-985. doi:10.1038/ng.3863

46. Ter Kuile AR, Hübel C, Cheesman R, et al. Genetic decomposition of the heritable
component of reported childhood maltreatment. *Biological Psychiatry Global Open*
Science. 2023;3(4):716-724. doi:10.1016/j.bpsgos.2023.03.003

47. Nielsen TT, Bali P, Grove J, et al. Genetic architecture and risk of childhood
maltreatment across 5 psychiatric diagnoses. *JAMA Psychiatry*. 2025;82(8):790.
doi:10.1001/jamapsychiatry.2025.0828

48.   Burgess S, Swanson SA, Labrecque JA. Are Mendelian randomization investigations
931   immune from bias due to reverse causation? *Eur J Epidemiol.Springer Science and*
932   *Business Media B.V.* 2021;36(3):253-257. doi:10.1007/s10654-021-00726-8

933   49.   McLaughlin KA. Future directions in childhood adversity and youth psychopathology.
934   *Journal of Clinical Child and Adolescent Psychology*. 2016;45(3):361-382.
935   doi:10.1080/15374416.2015.1110823

936   50.   Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between
937   imprecisely measured traits using GWAS summary data. *PLoS Genet*. 2017;13(11).
938   doi:10.1371/journal.pgen.1007081

939   51.   Richmond RC, Davey Smith G. Commentary: orienting causal relationships between
940   two phenotypes using bidirectional Mendelian randomization. *Int J Epidemiol.Oxford*
941   *University Press*. 2019;48(3):907-911. doi:10.1093/ije/dyz149

942   52.   Davey Smith G. Epidemiology, epigenetics and the 'Gloomy Prospect': embracing
943   randomness in population health research and practice. *Int J Epidemiol*.
944   2011;40(3):537-562. doi:10.1093/ije/dyr117

945   53.   Munafò MR, Tilling K, Taylor AE, Evans DM, Davey Smith G. Collider scope: when
946   selection bias can substantially influence observed associations. *Int J Epidemiol*.
947   2018;47(1):226-235. doi:10.1093/ije/dyx206

948   54.   Fry A, Littlejohns TJ, Sudlow C, et al. Comparison of sociodemographic and health-
949   related characteristics of UK Biobank participants with those of the general population.
950   *Am J Epidemiol*. 2017;186(9):1026-1034. doi:10.1093/aje/kwx246

951   55.   Schoeler T, Speed D, Porcu E, Pirastu N, Pingault JB, Kutalik Z. Participation bias in
952   the UK Biobank distorts genetic associations and downstream analyses. *Nat Hum*
953   *Behav*. 2023;7(7):1216-1227. doi:10.1038/s41562-023-01579-9

954   56.   Gkatzionis A, Burgess S. Contextualizing selection bias in Mendelian randomization:
955   how bad is it likely to be? *Int J Epidemiol*. 2019;48(3):691-701.
956   doi:10.1093/ije/dyy202

957   57.   Baldwin JR, Sallis HM, Schoeler T, et al. A genetically informed registered report on
958   adverse childhood experiences and mental health. *Nat Hum Behav*. 2023;7(2):269-290.
959   doi:10.1038/s41562-022-01482-9

960   58.   Soares ALG, Hammerton G, Howe LD, Rich-Edwards J, Halligan S, Fraser A. Sex
961   differences in the association between childhood maltreatment and cardiovascular
962   disease in the UK Biobank. *Heart*. 2020;106(17):1310-1316. doi:10.1136/heartjnl-
963   2019-316320

964   59.   Biele G, Gustavson K, Czajkowski NO, et al. Bias from self selection and loss to
965   follow-up in prospective cohort studies. *Eur J Epidemiol*. 2019;34(10):927-938.
966   doi:10.1007/s10654-019-00550-1

60. Walsh D, McCartney G, Smith M, Armour G. Relationship between childhood
socioeconomic position and adverse childhood experiences (ACEs): a systematic
review. *J Epidemiol Community Health (1978)*. 2019;73(12):1087-1093.
doi:10.1136/jech-2019-212738

61. Meyer MN, Appelbaum PS, Benjamin DJ, et al. Wrestling with social and behavioral
genomics: risks, potential benefits, and ethical responsibility. *Hastings Center Report*.
2023;53(S1):S2-S49. doi:10.1002/hast.1477

62. Chen L, Davey Smith G, Harbord RM, Lewis SJ. Alcohol intake and blood pressure: a
systematic review implementing a Mendelian randomization approach. *PLoS Med*.
2008;5(3):e52. doi:10.1371/journal.pmed.0050052

63. Burgess S, Davey Smith G, Davies NM, et al. Guidelines for performing Mendelian
randomization investigations: update for summer 2023. *Wellcome Open Res*.
2023;4:186. doi:10.12688/wellcomeopenres.15555.3

64. Dalvie S, Maihofer AX, Coleman JRI, et al. Genomic influences on self-reported
childhood maltreatment. *Transl Psychiatry*. 2020;10(1). doi:10.1038/s41398-020-
0706-0

65. Daghlas I, Richmond RC, Lane JM, et al. Selection into shift work is influenced by
educational attainment and body mass index: a Mendelian randomization study in the
UK Biobank. *Int J Epidemiol*. 2021;50(4):1229-1240. doi:10.1093/ije/dyab031

66. Hamilton F, Pedersen KM, Ghazal P, Nordestgaard BG, Davey Smith G. Low levels of
small HDL particles predict but do not influence risk of sepsis. *Crit Care*.
2023;27(1):389. doi:10.1186/s13054-023-04589-1

67. Jami ES, Hammerschlag AR, Bartels M, Middeldorp CM. Parental characteristics and
offspring mental health and related outcomes: a systematic review of genetically
informative literature. *Transl Psychiatry*. 2021;11(1):197. doi:10.1038/s41398-021-
01300-2

68. Pingault JB, Rijsdijk F, Schoeler T, et al. Genetic sensitivity analysis: adjusting for
genetic confounding in epidemiological associations. *PLoS Genet*. 2021;17(6).
doi:10.1371/journal.pgen.1009590

69. VanderWeele TJ, Jackson JW, Li S. Causal inference and longitudinal data: a case
study of religion and mental health. *Soc Psychiatry Psychiatr Epidemiol*.
2016;51(11):1457-1466. doi:10.1007/s00127-016-1281-9

70. Baldwin JR, Wang B, Karwatowska L, et al. Childhood maltreatment and mental
health problems: a systematic review and meta-analysis of quasi-experimental studies.
American Journal of Psychiatry. 2023;180(2):117-126. doi:10.1176/appi.ajp.20220174

71. Sonuga-Barke EJS, Kennedy M, Kumsta R, et al. Child-to-adult neurodevelopmental
and mental health trajectories after early life deprivation: the young adult follow-up of

the longitudinal English and Romanian Adoptees study. *The Lancet*.

2017;389(10078):1539-1548. doi:10.1016/S0140-6736(17)30045-4

1006