INCREASED PREVALENCE OF AUTISM IN THE CHILDREN OF FIBROMYALGIA PATIENTS RELATIVE TO OTHER TYPES OF CHRONIC PAIN.

Hirst A[1]., Mountford, R[1]., Berwick, R[1]., Christiansen, P[1]., Sandin, S[2,3]., Yin, W[2]., Andersson, D, A[4].,

Happé, F[4]., Fallon, N*[1]., Goebel, A*[1].

*Joint senior authors.

Abstract

Fibromyalgia syndrome (FMS) is a chronic primary pain condition characterised by widespread pain and hypersensitivities, and symptoms of fatigue, poor sleep and impaired cognition. The current studies were prompted by clinical observations made at a tertiary pain centre, where patients with severe FMS frequently reported having biological children on the autism spectrum. Using multimodal data collection techniques, including a clinical service evaluation (N = 589), an exploratory phenotyping study (N= 59), an online patient survey (N= 710), and a clinical prospective investigation (N = 66), the present study aimed to investigate links that may point to a connection between maternal FMS and autistic offspring. For the clinical and online data collection, the studies used comparator patients with complex regional pain syndrome (CRPS) and a mix of "other chronic pain conditions". Maternal factors of interest, including a familial history of neurodivergence and psychological variables, were considered in the analysis. Maternal FMS was consistently found to be associated with an increased likelihood of children with an autism diagnosis, even after accounting for established maternal factors associated with autism. Meeting and advocating for the support needs of autistic children may require enhanced care, which can lead to greater demands on patients already suffering from FMS or CRPS. The causes for the

observed association between maternal FMS and autistic children require further investigation. These findings highlight the importance of being aware of potential neurodivergence among the offspring of patients with FMS and suggest that this should be routinely asked about and documented.

Author affiliations:

1 University of Liverpool, Liverpool, UK.

2 Karolinska Institutet, Sweden.

3 Icahn School of Medicine at Mount Sinai, New York, USA.

4 Kings College London, London, UK.

Introduction

Fibromyalgia syndrome (FMS) is a chronic primary pain condition characterised by symptoms of widespread musculoskeletal pain, fatigue, stiffness, and "brain fog" [11; 35]. It is frequently associated with other somatic (e.g., irritable bowel syndrome) and psychological (e.g., anxiety, depression, PTSD) comorbidities [20]. The estimated global prevalence of FMS is commonly reported at 2-3% [36], with UK estimates reaching up to 5.4% [17]. Sixty-90% of cases are women [12; 46]. The pathogenesis of FMS is currently unknown, however, accumulating evidence has suggested that immune system dysfunction could underlie characteristic symptoms [19; 24; 32]. Several features of FMS are suggestive of autoimmunity, including the female predominance and mid-age onset [12], the notable prevalence of autoimmune disease comorbidity [2; 18], gene expression profiles which are typical of autoimmune disease [14], the relationship between the onset of FMS symptoms and trauma and infection [47], and the presence of autoreactive antibodies [5; 19].

Autism spectrum disorder (hereafter referred to as autism) is a neurodevelopmental condition characterised by social interaction and communication difficulties along with a range of restricted and/or repetitive patterns of behaviour, interests, or activities [3]. Moreover, autism presents with atypical sensorimotor function and often co-occurs with chronic pain [7; 45]. The estimated prevalence of autism in the general UK population is between 1% and 3% [8; 33], however, it is known to be underdiagnosed in older adults and, historically, in women. Although the exact aetiology of autism is not yet known, a complex interplay involving genetic and environmental factors is proposed [27]. Immune system dysregulation has been suggested as a non-genetic contributing factor and has been described in autistic people and their family members [15; 40]. Previous research has also implicated the prenatal immune environment as a factor in autism development [25; 37; 38], and several

studies have demonstrated an increased likelihood of autistic children of mothers, but not fathers, with autoimmune diseases [10; 13; 48].

In clinical observations made by a consultant in pain medicine at a tertiary pain centre (AG), it was noted that patients with severe FMS frequently reported having biological children on the autism spectrum. Caring and advocating for an autistic child can place additional demands on the parent [34], which may be particularly challenging for people suffering from severe chronic pain. It was thought, therefore, that understanding any such association between FMS and autistic offspring would be relevant for practitioners in this field, enabling them to gather information pertinent to gauge support requirements.

Methods

The study used a design in which data from multiple, independent sources were compiled to investigate a potential link between maternal FMS and children with an autism diagnosis. This comprised data from a clinical pain centre service evaluation (Study A1), a previously established research study (Study A2), and two newly conducted online surveys – one using a wider geographical region in the UK (Study B) and one using a subset of patients included in the clinical service evaluation (Study C).

A1) Clinical Service Evaluation

Participants

Following initial observations about autistic children of patients with FMS at a tertiary pain centre, we systematically documented the frequency of autism diagnoses in the offspring of patients with FMS and other chronic pain conditions at the same institution (the Walton Centre, a UK neuro-care centre).

Chronic pain patients were included in a service evaluation if they had attended their appointment as a new patient at the tertiary care outpatient pain clinic with a consultant in pain medicine (AG) between 26th October 2020 and 10th May 2023. All patients for whom a clinic letter was dictated during this period were included (N=589). As part of their routine consultation, the consultant had asked new patients clinical questions, which for the majority included a question about their children's health. The consultant did not specifically enquire about neurodivergent diagnoses in children in any of the available clinic letters, and no questionnaires were used. The service evaluation was approved by the Walton Centre NHS Foundation Trust (NS451).

Diagnostic categorisation

The recruiting centre provides health care in matters of pain from the north-west region of England and is also a supra-regional referral centre for persistent Complex Regional Pain Syndrome (CRPS) and other uncommon, complex chronic pain conditions. Patients were categorised by their chronic pain condition: FMS, persistent CRPS, and other chronic pain conditions. In the "other chronic pain condition" group, a majority suffered from chronic widespread pain or back and leg pain conditions. A smaller group suffered from neuropathic pain conditions or other chronic pain (for details, see Supplementary Table 1; Appendix A).

Patients who reported their children as having an autism diagnosis were classified as cases in the analysis. Patients who reported their child as having *"suspected"* or *"probable"* autism were not classified as a case of offspring autism in the analysis.

Statistical analysis

Penalised likelihood logistic regression analysis was conducted to test the association between FMS and CRPS, compared to other chronic pain conditions, and the likelihood of

offspring with autism diagnoses. Firth's penalised logistic regression is a method for modelling binary outcomes, used to mitigate issues that traditional logistic regression models may have when facing small sample sizes, rare events, and separation [43]. The categorical variable "chronic pain condition" had 3 levels (FMS, CRPS, and other chronic pain). Other chronic pain was chosen as the baseline category to observe any change in log-odds for FMS and CRPS, which are particularly severe chronic primary pain conditions, previously implicated as having an autoimmune mechanism [19; 23]. At present, no such evidence has emerged for the other chronic pain group. Only mothers with confirmed biological children were included in the analysis. Odds ratios (OR) with 95% confidence intervals (CIs) were calculated, corresponding to a two-sided test on the 5% level of significance. RStudio software (Version 4.3.1; R Core Team, 2023) was used for all statistical analyses, and logistic regression models were fit using the logistf package [22].

A2) Exploratory phenotyping study

To explore phenotypic differences between FMS mothers with and without autistic children, we utilised an established research study involving detailed phenotyping of 59 patients with moderate or severe FMS who had been identified from the pain management program registry hosted by the same hospital as the service evaluation (Study A1). Study methods and phenotyping results from a subgroup have been published [30]. With few exceptions, these patients were recruited by a senior trainee in pain medicine (RB), and none of these patients' data had been part of the service evaluation. Recruitment details are summarised in the supplementary Appendix A. The trainee asked participants about the health of their offspring as part of the assessment procedure, and, like the consultant in the service evaluation, did not specifically ask about neurodivergence. If patients volunteered the

presence of a neurodivergent condition in offspring, this was documented. Details of the source of the autism diagnosis were not requested. Patients also completed an array of questionnaires and underwent mechanical quantitative sensory testing (Supplementary Materials; Appendix A).

Statistical analysis

The sample was split into 3 groups: FMS mothers with autistic children, FMS mothers with non-autistic children, and FMS females without children. To explore differences in pain phenotype between these groups, Kruskal-Wallis tests were conducted, and where a statistically significant difference was found (two-sided 5% level), uncorrected post-hoc comparisons were performed. As this was an exploratory investigation, no correction for testing multiple parameters was applied.

RStudio software (Version 4.3.1; R Core Team, 2023) was used for all statistical analyses.

B) Online Survey

We conducted an online survey amongst participants of UK pain online support groups aiming to i) record the frequency of autistic children of patients with FMS and other chronic pain conditions in a wider geographical region in the UK, and ii) explore which maternal factors are associated with having an autistic child. We hypothesised that a) there would be an increased rate of autistic children of mothers with FMS compared to mothers with other chronic pain conditions; and b) this relationship would remain even after accounting for maternal or familial history of neurodivergent conditions. The survey was hosted by Qualtrics Survey Software (Qualtrics, 2024) and was advertised first between May 2023 and June 2023, and then between November 2024 and February 2025, through online chronic pain

support groups and social media platforms targeting individuals suffering from FMS and general chronic pain. This included *The Pain Relief Foundation* (https://painrelieffoundation.org.uk) and *Pain UK* (https://painuk.org). The advertisement briefly outlined that the study was investigating *neurodivergent* conditions in people with chronic pain and their families, without specifically mentioning autism.

Participants

The inclusion criteria were adults with a chronic pain diagnosis who were also parents of children of any age, based in the UK, and fluent in English. Individuals who were pregnant at the time were excluded from participation. Respondents were asked to confirm a chronic pain diagnosis from a specified clinician, although no proof of diagnosis was required. All participants were requested to provide consent and confirm eligibility before starting the online study. Upon completion, participants had the choice to anonymously enter a prize draw to win a £50 Amazon voucher (one for every 50 respondents who entered). The study obtained ethical approval from the University of Liverpool Research Ethics Committee (25th April 2023; Ref: 12195).

Questionnaire

The questionnaire was created by the researchers (AH, NF, AG). It recorded mothers' current pain and maternal/family factors that may be associated with child neurodiversity. It also assessed the presence of neurodivergence in biological children, and for each neurodiverse child, the presence and nature of birth complications. Mother's pain at the time of the pregnancy with any neurodiverse children was also enquired. See the supplementary materials for the full questionnaire (Appendix B).

Data diagnostics

The study was advertised online, therefore, a screening process was undertaken to remove fake responders from the analysis. The exclusion criteria removed: duplicate responses, bots identified by a function of the survey software, nonsensical timelines, non-English responses, and reporting diagnoses which were not related or relevant to the concept (e.g., reporting "headaches" as a neurodivergent condition, and simply "sore" as a chronic pain condition).

Respondents who reported having a child who had *suspected* autism, or was *on a waiting list* for diagnosis, were recorded but not classified as a case of autism. These cases were treated as a control in the analysis.

Statistical analysis

Penalised likelihood hierarchical logistic regression analyses were conducted to quantify the association between maternal chronic pain (FMS/Other chronic pain) and autistic children, whilst accounting for the mother's neurodivergence and the incidence of neurodivergent conditions in the mother's first-degree family members. A hierarchical regression model was chosen to determine whether the variables of interest improved the model's ability to predict the binary outcome, after accounting for all other predictor variables. Maternal neurodivergence (diagnosis/no diagnosis) and family neurodivergence (yes/no) were entered into the first model, as inherited and genetic factors are known to play a major role in the aetiology of autism [21]. A second model was created by adding the variable maternal chronic pain condition (FMS/Other chronic pain) to the first model. The association between the individual predictors and the outcome of having a child with an autism diagnosis was determined using ORs and the associated 95% CIs. The model fit was

assessed by conducting a model likelihood ratio (LR) chi-square test, and Akaike Information Criterion (AIC) values are also reported.

RStudio software (Version 4.3.1; R Core Team, 2023) was used for all statistical analyses, and logistic regression models were fit using the logistf package [22].

C) Clinical Prospective Investigation

A second online study invited all patients whose clinic letters had been included in the service evaluation (Study A1) to complete an online questionnaire. Our objectives for this study were to i) expand upon initial data from Study A1 by collecting more comprehensive information on respondents with autistic children, including any maternal contributing factors, and ii) gain further information regarding children's autistic traits to supplement the parents' binary report (diagnosis/no diagnosis).

Participants

All chronic pain patients included in the clinical service evaluation (N= 589, Study A1), except those who were pregnant at the time of approach, were invited to participate in an online questionnaire. Patients had the choice to enter a prize draw to win a £75 Amazon voucher upon completing the questionnaire. The study received ethical approval from the NHS Health Research Authority Research Ethics Committee Wales REC 2 (Ref: 24/WA/0056).

Materials

An online questionnaire was created jointly with an autism specialist (FH) while incorporating Patient and Public Involvement (PPI) in its development. Chronic pain patients involved in an existing FMS PPI group at our research institution advised on the content of

the survey, its clarity, and sensitivity. Several validated scales were included in the questionnaire. Internal reliability was assessed using McDonald's ω, which was calculated from the current study sample. Data were collected using Qualtrics Survey Software (Qualtrics, 2024). See the supplementary materials for the full questionnaire (Appendix C).

1. The General Anxiety Disorder (GAD-7; [42]) is a 7-item *self-report* scale which was developed to measure general anxiety disorder. This scale was included based on the consideration that higher maternal anxiety may relate to an increased desire to have clarity about their child's diagnosis, potentially increasing the probability of obtaining an autism diagnosis. Responses to all items are scored on a 4-point Likert scale ranging from *0 (not at all)* to *3 (nearly every day)* in reference to the previous 2 weeks. Total scores range from 0 to 21, with higher scores indicating higher severity of general anxiety. A score of ≥10 is considered the cut-off for moderate anxiety. The overall scale had excellent internal reliability (McDonald's ω_t = 0.97).

2. The Autism Spectrum Quotient Adult (AQ-10 Adult; [1]) is a 10-item *self-report* scale developed as a brief screening tool to capture autistic traits in adults. The scale consists of 5 subscales, each consisting of 2 items, including attention to detail, attention switching, communication, imagination, and social behaviours. Total scores range from 0 to 10. A cut-off value of ≥6 indicates a possible case of autism. The overall scale had poor internal reliability in the present sample (McDonald's ω_h = 0.48).

3. The Ritvo Autism Asperger Diagnostic Scale (RAADS-14; [16]) is a 14-item *self-report* scale which is designed to assess traits associated with autism in adults. The scale consists of 3 subscales including mentalising deficits, social anxiety, and sensory reactivity. All items, except item 6, which is reversed, are scored *3 (true now and when I was young), 2 (true only now), 1 (true only when I was young),* and *0*

(never true). Total scores range from 0 to 42 with a cut-off value of ≥14 being indicative of possible autism. The overall scale had acceptable internal reliability (McDonald's ω_h = 0.76).

4. Parent-Report Autism Spectrum Quotient (Parent-report AQ-10; [1]) is a 10-item scale developed for parents to complete *referring to their child*. Total scores range from 0 to 10, and a cut-off value of ≥6 is indicative of possible autism. The overall scale had questionable internal reliability (McDonald's ω_h = 0.67).

5. The Social Communication Questionnaire – Lifetime (SCQ; [41]) is a 40-item, parent-report measure for screening symptomology associated with autism. All items are answered dichotomously *(yes/no)*. Item 1 records whether the referred child has the ability to speak in short phrases or sentences, and items 2 – 40 are scored and summed. Total scores range from 0 to 39 with a cut-off value of ≥15 suggesting autism. The overall scale had excellent internal reliability (McDonald's ω_t = 0.91).

Procedure

After providing consent and passing eligibility checks, patients were asked about their chronic pain history and general health, and to complete 3 standardised questionnaires referring to themselves (GAD-7, AQ-10 Adult, RAADS-14). After this, patients were asked to complete questions referring to the health of all their children, in birth order. This included the parent-report AQ10 *for each of their children*. The SCQ was then only completed for children who respondents either reported as having an autism diagnosis, or about whom they had concerns regarding their social communication, behaviour, and/or development. See supplementary materials for details about children's AQ10 and SCQ scores (Supplementary Table 14 and 15; Appendix C).

Patients with autistic children were asked further questions regarding any

complications during pregnancy, delivery or after birth, co-occurring child health conditions,

child intellectual disability, and the level of any child support needs assigned by a doctor in

reference.

Statistical analysis

Hierarchical penalised likelihood logistic regression analyses were conducted to test the

association of maternal chronic pain condition (FMS/ Other chronic pain) whilst accounting

for family history variables (autistic mothers and the incidence of neurodivergent conditions

in first-degree family members) and psychological variables (GAD-7 scores). Maternal

autism diagnosis status (diagnosis/ no diagnosis) and family neurodivergence (yes/no) were

entered into the first model, GAD-7 scores were added into the second model, and maternal

chronic pain condition was entered into the third model. The association between these

individual parameters and the outcome of having a child with an autism diagnosis was

determined using ORs with 95% CIs. The model fit was determined by conducting a model

LR chi-square test and AIC values were reported. RStudio software (Version 4.3.1; R Core

Team, 2023) was used for all statistical analyses. Logistic regression models were fit using

the logistf package [22], and internal reliability was assessed using the psych package [39].

Results

A1) Clinical Service Evaluation

The cohort included 589 total patients, 398 (67.57%) females and 191 (32.43%) males.

The consultant had documented an enquiry about the health of the respective patient's

offspring in 89% of cases. Among the patients with confirmed biological children, 31

(8.68%) patients had at least one child with an autism diagnosis. Thirteen (16.25%) mothers

with FMS, 4 mothers with CRPS (14.81%) and 6 (4.05%) mothers with other chronic pain

conditions reported having at least one autistic child. Table 1 displays the full sample by chronic pain condition.

Information about the onset of mothers' pain in relation to their children's ages/ birth was not consistently documented; of the available data, 3 mothers with FMS (30.00%) and 2 mothers with CRPS (100.00%) reported that the onset of their FMS or CRPS symptoms, respectively, occurred before childbirth. Two mothers with other chronic pain (33.3%) reported that they had pain before childbirth (Table 2). Table 2 describes further available information pertaining to children with an autism diagnosis.

Although not classified as a case of autism in the analysis, 7 patients reported that they had one child who they *suspected* was autistic, but no diagnosis (2 FMS; 1 CRPS; 4 other chronic pain).

Table 1.

The total sample of chronic pain patients by condition and the frequency of reporting

children with an autism diagnosis.

Chronic pain condition	Total N of patients (females, %)	Patients with confirmed biological children (females, %)	Patients with at least one autistic child N[%][a]	Female patients with at least one autitic child N[%][b]	Male patients with at least one autistic child N [%][c]
FMS	119 (107 89.92% [d])	86 (80, 93.02% [d])	13 [15.12%]	13 [16.25%]	0 [0.00%]
CRPS	67 (42, 62.69%)	38 (26, 68.42%)	7 [18.42%]	4 [14.81%]	3 [25.00%]
Other chronic pain conditions	403 (249, 61.79%)	239 (148, 61.92%)	11 [4.60%]	6 [4.05%]	5 [5.56%]

[a] Patients with at least one autistic child as a percentage of those with confirmed biological children within the chronic pain condition category.

[b] Female patient with at least one autistic child as a percentage of females with confirmed biological children within the chronic pain condition category.

[c] Male patient with at least one autistic child as a percentage of males with confirmed biological children within the chronic pain condition category.

[d] 3 of these female patients had both FMS and CRPS (not listed in the CRPS group), 1 of whom had an autistic child.

Table 2.

Characteristics of autistic children, by parental chronic pain condition in the clinical service evaluation.

	FMS	CRPS	Other chronic pain
Characteristics of autistic children	N [%] or mean (SD)	N [%] or mean (SD)	N [%] or mean (SD)
N (of parents with that chronic pain condition)	13	7	11
Parental age at birth *(N available data)*	31.38 (5.97) *8*	21.50 (3.54) *2*	26.00 (3.37) *5*
Onset of mothers' chronic pain condition before childbirth *(N available data)*	3 [30.00%] *10*	2 [100%] *2*	2 [18.18%] *2*
More than 1 biological child with autism *(N available data)*	1 [7.69%] *13*	3 [42.86%] *7*	1 [9.09%] *11*
Sex of children			
Male:	10 [83.33%]	0 [0.00%]	8 [88.89%]
Female:	2 [16.67%]	3 [75.00%]	1 [11.11%]
Male and Female:	0 [0.00%]	1 [25.00%]	0 [0.00%]
(N available data)	*12*	*4*	*9*
Birth order of autistic child: *			
1st born:	5 [38.46%]	4 [57.14%]	4 [36.36%]
2nd born:	3 [23.08%]	3 [42.86%]	1 [9.09%]
3rd born:	0 [0.00%]	2 [28.57%]	3 [27.27%]
4th born:	0 [0.00%]	1 [14.29%]	0 [0.00%]
Unknown:	5 [38.46%]	1 [14.29%]	3 [27.27%]

Note. The table is derived from data available in the clinic letters, which were incomplete in this respect. Percentages are derived from the available data for each variable.

*Some values are duplicates for patients who have more than one biological child with an autism diagnosis.

Figure 1a shows the results of the penalised logistic regression model examining the association between maternal chronic pain condition and the likelihood of children with autism diagnoses. The overall model was statistically significant $\chi^2(2) = 11.00$, $p=.004$, McFadden $R^2 =0.01$. The likelihood of autistic offspring was estimated at OR: 4.38 (95% CI: 1.70 –12.45) for maternal FMS and at OR: 4.38 (1.14 - 15.70) for maternal CRPS, compared to other chronic pain.

Figure 1.

(a) Penalised logistic regression analysis showing the association of maternal FMS and CRPS, relative to other chronic pain conditions, to children with an autism diagnosis.

(b) Forest plot showing ORs and 95% CIs.

Maternal chronic pain condition	B	SE	OR	95% CIs	*p* value
FMS	1.48	0.50	4.38	1.70 – 12.45	**.002**
CRPS	1.48	0.65	4.38	1.14 – 15.70	**.033**

A2) Exploratory phenotyping study

Five/44 (11.36%) FMS participant mothers reported having an autistic child. In comparison to the mothers with non-autistic children, mothers of autistic children had statistically significantly lower scores on the Widespread Pain Index (i.e., their pain was less widespread, median 10.00, IQR 2.00 vs. median 15.00, IQR 5.00; $p = .027$, $\eta^2 = 0.09$ uncorrected Kruskal-Wallis test) and on the Short-Form McGill Pain Questionnaire *intermittent* subscale (i.e., lower reported severity of intermittent pain, median 3.40, IQR 2.02 vs. median 5.80, IQR 4.25; $p = .049$, $\eta^2 = 0.07$). There were no other significant differences between these 2 groups on any of the measured demographic, impact, pain, or sensory parameters (Supplementary Table 2; Appendix A).

B) Online Survey

This cohort of parents included 710 participants (589 FMS, 121 other chronic pain), of whom 622 were mothers (87.61%) and 80 were fathers (11.27%). Eight respondents (one of whom reported having a child with an autism diagnosis) did not provide their sex and, therefore, were not included in the analysis. One hundred and thirty-seven (25.95%) mothers with FMS and 10 (10.64%) mothers with other chronic pain self-reported a child with an autism diagnosis. Fifteen FMS mothers (2.84%) and 3 other chronic pain mothers (3.19%) reported having more than one autistic child.

 The most commonly reported "other chronic pain" conditions were back pain, osteoarthritis, joint pain, and neck/shoulder pain (Supplementary Table 3; Appendix B).

Table 3 describes the participant characteristics of the mothers-only sample, stratified by maternal chronic pain condition and the autism diagnosis status of children. Although not classified as a case of autism in the analysis, 7 mothers with FMS reported having a child they *suspected* was autistic or indicated that they were on a waiting list for diagnosis.

Three (5.26%) fathers with FMS and 1 (4.35%) father with other chronic pains reported having an autistic child. Additional analyses showed a statistically significant main effect of parental sex on the likelihood of autism diagnoses in children. The likelihood of autistic offspring was estimated at OR 4.82 (95% CI: 2.03 – 14.85, p <.001) for mothers compared to fathers. There was no significant interaction between parental sex and chronic pain condition (p = .341). Supplementary Table 5 shows further details about the fathers-only sample (Appendix B).

Patients also reported other neurodivergent conditions in their children. Overall, ADHD was the second most reported at a rate of 21.78% for FMS mothers and 14.89% for mothers with other chronic pain (non-significant difference, supplementary Table 6; Appendix B). Other neurodivergent conditions were rarely noted (Supplementary Table 4; Appendix B).

Table 3.

Characteristics of mothers with FMS or other chronic pain, stratified by the autism diagnosis status of children.

Patient characteristics	Mothers with FMS		Mothers with other chronic pain [a]	
	Autistic child	**Non-autistic child**	**Autistic child**	**Non- autistic child**
	N [%] or M (SD)	N [%] or M (SD)	N [%] or M (SD)	N [%] or M (SD)
N	137 [25.95%]	391 [74.05%]	10 [10.64%]	85 [90.43%]
Maternal age yrs	40.8 (8.83)	42.4 (12.10)	54.6 (12.40)	44.0 (16.20)
Respondents with a diagnosed neurodivergent condition	47 [34.31%]	102 [26.09%]	6 [60.00%]	35 [41.18%]
1st degree family with diagnosed neurodivergent condition [b]	34 [23.36%]	76 [19.44%]	4 [40.00%]	27 [31.76%]
Weekly average pain score (0-10)	6.87 (1.75)	6.66 (1.75)	5.30 (2.00)	5.46 (1.99)
High lifetime-stress estimate (much more/extremely above average)	93 [67.89%]	227 [58.06%]	4 [40.00%]	27 [31.76%]

Abbreviations: M, mean; SD, standard deviation.
[a] None of these patients had CRPS.
[b] First-degree family include the respondent's parents and siblings, but not children.
Note. There was a significant interaction between chronic pain condition and autism diagnosis status of offspring on patient age (p=.004, ηp^2=0.01; 2x2 ANOVA). Mothers with other chronic pain and an autistic child were older than those without autistic children (p=.044), and FMS mothers with (p=.003) and without autistic children (p=.009).
There was a significant main effect of chronic pain condition on weekly average pain rating, with FMS mothers reporting a higher score compared to mothers with other chronic pain (p<.001, ηp^2= 0.06), however, there was no interaction between maternal chronic pain condition and autism diagnosis status of offspring on pain scores (p=.549, ηp^2= <0.01; 2x2 ANOVA).
There was a significant main effect of chronic pain condition on a high lifetime stress estimate, with FMS mothers having an increased likelihood of reporting a high lifetime stress compared to mothers with other chronic pain (OR= 2.92, 95% CIs 1.85 - 4.70, p<.001). There was no interaction between maternal chronic pain condition and autism diagnosis status of offspring on lifetime stress estimates (OR = 1.05 95% CIs 0.27 - 4.31, p=.944)

Table 4.

Characteristics of autistic children, by maternal chronic pain condition.

	FMS	Other chronic pain
Characteristics of autistic child	N [%] or mean (SD)	N [%] or mean (SD)
N (of mothers with that chronic pain condition)	137	10
Maternal age at birth	26.52 (5.34	27.22 (6.12)
Autism co-occurring with other neurodivergent conditions	7 [5.11%]	2 [20.00%]
More than one autistic child	15 [10.95%]	3 [30.00%]
Maternal (current) chronic pain <u>diagnosis</u> before/around childbirth	32 [23.36%]	2 [20.0%]
Onset of maternal (current) chronic pain <u>symptoms</u> before/around childbirth	58 [42.34%]	4 [40.0%]
If (maternal) pain before pregnancy: average pain intensity (0-10)	5.23 (1.97)	3.40 (2.88)
If (maternal) pain during pregnancy: average pain intensity (0-10)	6.22 (2.12)	4.40 (2.97)
Child's father had a known chronic pain condition around conception/pregnancy	26 [18.98%]	3 [30.00%]
Birth/ pregnancy complications	64 [46.72%]	2 [20.0%]
- *Gestational age <35 wks*	12 [8.76%]	1 [10.00%]
- *Low birth weight*	17 [12.41%]	0 [0.00%]
- *Birth injury/trauma*	24 [17.52%]	0 [0.00%]
- *Birth defects associated with the CNS*	4 [2.92%]	0 [0.00%]
- *Meconium aspiration syndrome*	3 [2.19%]	1[10.00%]
- *Neonatal or epileptic encephalopathy*	1 [0.73%]	0 [0.00%]
- *Genetic disorder*	3 [2.19%]	0 [0.00%]

Note. Maternal chronic pain diagnosis and the presence of pain around pregnancy were calculated from the child's age and the respondent's reported year of diagnosis and pain onset.

Figure 2a shows the results of the penalised hierarchical logistic regression model examining the association between maternal variables and the likelihood of children with autism diagnoses. The first model, including the family-history factors, maternal neurodivergence and first-degree family neurodivergence, was non-significant $\chi^2(2) = 2.93$, $p = .231$, AIC = 672.78, McFadden's $R^2 = 0.01$.

Adding maternal chronic pain condition into the second model significantly improved the model fit relative to the family-history only model $\chi^2(1) = 13.57$, $p < .001$. With this, the overall model was statistically significant $\chi^2(3) = 16.50$, $p < .001$, AIC = 672.12, McFadden's $R^2 = 0.02$. The likelihood of autistic offspring was estimated at OR 3.12 (95% CI: 1.65 – 6.50) for mothers with FMS compared to mothers with other chronic pain.

Figure 2.

(a) Penalised logistic regression analysis showing the association of maternal variables to children with an autism diagnosis. (b) Forest plot showing ORs and 95% CIs.

Maternal Predictors	Model 1:			Model 2:		
	B (SE)	OR (95% CIs)	*p* value	B (SE)	OR (95% CIs)	*p* value
Step 1: Family history variables						
Maternal neurodivergence	0.27 (0.21)	1.32 (0.87-1.98)	.197	0.29 (0.21)	1.42 (0.93 – 2.15)	.106
Family neurodivergence	0.15 (0.23)	1.16 (0.73 – 1.81)	.528	0.14 (0.23)	1.24 (0.78 – 1.95)	.366
Step 2: Chronic pain variables						
FMS vs other chronic pain	-	-	-	1.14 (0.34)	3.12 (1.65 – 6.50)	**<.001**

Odds Ratio (95% CI)

C) Clinical Prospective Investigation

The cohort included a total of 66 patients (50 [75.76%] female, 16 [24.24%] male); the overall response rate was 11.2% (24.6% FMS, 17.7% CRPS, 6.3% other chronic pain). Nineteen of the 24 mothers with FMS (79.16%) had an FMS diagnosis during the service evaluation, all 7 mothers with CRPS (100%), and 8 out of the 11 mothers with other chronic pain (72.73%) were categorised as other chronic pain in the service evaluation.

Among the maternal sample, 8 mothers with FMS reported having an autistic child (33.33%), whereas no cases were reported among mothers with other chronic pain. Table 5 shows the characteristics of the maternal sample, stratified by the autism diagnosis status of offspring. Responding mothers with CRPS were not included in the analysis due to a small sample size (N= 7), see supplementary materials for descriptive details about CRPS mothers (Supplementary Table 10; Appendix C).

In all cases, patients reported having only one autistic child. All autistic children included in the analysis scored above the recommended cut-off values on the parent-report AQ10 (score of ≥6) and SCQ (score of ≥15). See Table 6 for further information pertaining to autistic children.

Table 5.

Characteristics of mothers with FMS and other chronic pain, stratified by the autism diagnosis status of children.

	Mothers with FMS		Mothers with other chronic pain
	Autistic child	**Non-autistic child**	**Non-autistic child**
Patient characteristics	N [%] or median (IQR)	N [%] or median (IQR)	N [%] or median (IQR)
N	8 [33.33%]	16 [66.67%]	11 [100%]
Maternal age yrs (mean (SD)	48.8 (7.34)	49.3 (8.93)	49 (14.2)
Autistic respondent	0 [0.00%]	2 [12.5%]	0 [0.00%]
1st degree family with diagnosed neurodivergent condition [a]	3 [37.50%]	1 [6.25%]	2 [18.18%]
GAD-7 (0-21) *Scored above cutoff (≥10)*	7 (8) 3 [37.50%]	7.5 (11.2) 7 [43.75%]	13 (8.5) 7 [63.64%]
RAADS-14 Total (0-42) *Scored above cutoff (≥14)*	25 (14.5) 6 [75.00%]	20.5 (20) 10 [62.50%]	24 (16) 9 [81.82%]
AQ-10 (0-10) *Scored above cutoff (≥6)*	4.5 (1.25) 2 [25.00%]	4.5 (1.25) 4 [25.00%]	5 (1.5) 3 [27.27%]

Abbreviations: IQR, Interquartile Range; SD, standard deviation.
[a] First-degree family include the respondent's parents and siblings, but not children.
Note. There were no significant differences in age (p=.969 η2= <0.01; ANOVA), AQ-10 scores (p=.976, η^2= -0.06), or RAADS-14 scores (p=.757, η^2= -0.05; Kruskal-Wallis) between groups.

Table 6.

Characteristics of autistic children, of mothers with FMS.

Characteristics of autistic child	Mothers with FMS N [%] or mean (SD)
N (of mothers with FMS)	8
Parental age at birth *(N available data)*	25.90 (6.62) *8*
AQ-10 score median (IQR)(0-10; cutoff ≥6) *(N available data)*	8.50 (1.50) *8*
SCQ score median (IQR) (0-39; cut off≥15) *(N available data)*	24.00 (6.50) *7*
Age at diagnosis yrs *(N available data)*	14.29 (4.82) *7*
Sex = male *(N available data)*	4 [50.00%] *8*
Birth order:	
1ˢᵗ child:	3 [37.50%]
2ⁿᵈ child:	2 [25.00%]
3ʳᵈ child:	3 [37.50%]
(N available data)	*8*
Any maternal chronic pain during pregnancy *(N available data)*	4 [50.00%] *8*
Maternal FMS or widespread pain during pregnancy *(N available data)*	2 [28.57%] *7*
Birth/ pregnancy complications *(N available data)*	2 [28.57%] *7*
	• Respiratory distress syndrome • Low blood sugar
Co-occurring intellectual disability *(N available data)*	3 [42.86%] *7*
Level of support needs:	
Level 1 (requiring support):	0 [0.00%]
Level 2 (requiring substantial support):	3 [100.00%]
Level 3 (requiring very substantial support):	0 [0.00%]
(N available data)	*3*
Child's other co-occurring conditions	• Pulmonary stenosis • Cerebral Palsy • Asthma (x2) • ADHD

Note. Parents did not know or report answers for all the presented variables. Percentages are derived from the available data for each variable.

Figure 3a shows the results of the hierarchical penalised likelihood logistic regression model examining the association between maternal variables and the offspring with autism diagnoses. The first model, including family history variables: maternal autism diagnosis and first-degree family neurodivergence, was not statistically significant $\chi^2(2) = 3.53$, $p=.171$, AIC = 40.99, McFadden's R^2 = -0.03.

The model LR chi-square test indicated that adding the psychological variable, GAD-7 scores, did not significantly improve the model fit $\chi^2(1) = 0.54$ $p=.539$, and this second model was not statistically significant $\chi^2(3) = 3.91$, $p=.271$, AIC = 39.11, McFadden's R^2 = 0.12.

The third model included maternal chronic pain condition (FMS and other chronic pain). The model LR chi-square test suggested that adding these variables significantly improved the model fit $\chi^2(1) = 6.27$, $p=.012$. With this, the overall model was statistically significant $\chi^2(4) = 10.18$, $p=.038$, AIC = 42.60, McFadden's R^2 = 0.13. The likelihood of autistic offspring was estimated at OR 24.94 (95% CI: 1.76 – 6952.87) for mothers with FMS compared to mothers with other chronic pain. The likelihood for mothers with first-degree neurodivergent relatives was estimated at OR 13.55 (1.16 – 1876.57). However, the precision of these estimates is poor due to a small sample size.

Figure 3.

(a) Penalised logistic regression analysis showing the association of maternal variables to children with an autism diagnosis. (b) Forest plot showing ORs and 95% CIs.

Maternal predictors	Model 1:			Model 2:			Model 3:		
	B (SE)	OR (95% CIs)	p value	B (SE)	OR (95% CIs)	p value	B (SE)	OR (95% CIs)	p value
Step 1: Family history variables									
Maternal autism diagnosis	-1.39 (1.72)	0.25 (0.00 − 4.93)	.400	-1.19 (1.67)	0.30 (0.00 − 5.16)	.455	-2.68 (2.00)	0.07 (0.00 − 2.23)	.155
Family neurodivergence	1.74 (0.93)	5.70 (0.90 − 43.09)	.064	1.63 (0.91)	5.11 (0.83 − 37.26)	.078	2.61 (1.34)	13.55 (1.16 − 1876.57)	**.036**
Step 2: Psychological variables									
GAD7 score	-	-	-	-0.05 (0.06	0.96 (0.83 − 1.08)	.480	-0.02 (0.06)	0.98 (0.85 − 1.13)	.803
Step 3: Chronic pain variables									
FMS vs other chronic pain	-	-	-	-	-	-	3.22 (1.65)	24.94 (1.76 − 6952.87)	**.011**

Discussion

To our knowledge, no previous report has explored the occurrence of autism specifically in the children of patients suffering from FMS. We found an increased likelihood of autistic children with mothers with FMS compared to mothers with other chronic pain conditions in each of 4 cohorts, with ORs ranging from 3.12 to 24.94. CRPS in mothers was also associated with an increased likelihood of autistic children in comparison to other chronic pains.

A clinical service evaluation (Study A1) included all chronic pain patients seen by a consultant at a tertiary centre over 2 years. Mothers with FMS and CRPS reported a high prevalence of autistic children (16% and 15%, respectively) versus patients presenting with other pain conditions (4%). This relationship was found to be significant, with 4.38 increased odds in mothers with FMS, and 4.38 increased odds for mothers with CRPS.

An exploratory phenotyping study (Study A2) analysed the responses of FMS patients who had also been routinely asked about their children's health. Eleven percent of mothers with FMS reported an autistic child. Together, these findings suggest a much higher prevalence of autistic children of FMS mothers compared to the UK general population estimate of 1%-3% [8; 33]. Comparison of demographic and FMS characteristics between those with and without autistic children in Study A2 revealed a lower reported widespread pain score and less intermittent pain (e.g., shooting, stabbing pain) for mothers with an autistic child. These results should be considered exploratory, however, the moderate effect sizes raise the possibility that there might be a difference in the FMS phenotype in mothers who have an autistic child.

An online survey (Study B) was distributed among members of online pain support groups to assess the frequency of autistic children in FMS mothers, beyond both tertiary

healthcare settings and the geographical restriction of the previous studies (NW-England). The aim was to explore mothers' personal and first-degree family members' self-reported neurodiversity, as genetic factors play a major role in the aetiology of autism [21]. Consistent with the service evaluation, a higher incidence of autism diagnoses in offspring was reported for mothers with FMS compared to mothers with other chronic pain conditions (26% and 11%, respectively). FMS in mothers remained statistically significantly associated with 3.12 increased odds of autistic children compared to mothers with other chronic pain after accounting for maternal and first-degree familial history of neurodivergence. In contrast, FMS fathers reported autistic children much less frequently than FMS mothers, and at a similar prevalence (4-5%) as fathers suffering from the other chronic pain conditions. Unexpectedly, mothers with other chronic pain reported diagnosed neurodivergent conditions in themselves and their first-degree family members more frequently than FMS mothers. The high reported neurodivergence prevalences across groups are likely affected by a selection bias whereby those with a known occurrence of neurodivergence in themselves or their family may have been more likely to participate in the study. This high prevalence is consistent with current literature suggesting an overlap between neurodivergence and chronic pain [9; 29]. However, the available literature evidence does not appear to account for the increased likelihood of children with autism diagnoses of FMS mothers specifically. Biological fathers might provide a greater genetic contribution in this group – we did not assess the fathers' or paternal first-degree family members' neurodiversity in this study, but this may be unlikely given the low offspring autism rates reported by participating fathers in the same study. Alternatively, this finding might suggest a lower genetic propensity for the autistic children of FMS mothers than of mothers with other chronic pain conditions. Correspondingly, non-genetic factors, including perinatal factors, may be more relevant in these children of FMS mothers. It has been noted that autoantibody-mediated autoimmune

conditions may be linked to autistic children through transplacental transfer of pathogenic autoantibodies [6; 13; 25]. Additional studies will be required to understand this area.

In this online survey, 47% of FMS mothers reported having birth/pregnancy complications with their autistic child, with birth injury or trauma (18%), low birth weight (12%), and a gestational age of <35 weeks (9%) being the most reported. This can be aligned with previous research which demonstrates that such parameters are associated with autistic children [4; 28]. Moreover, FMS in pregnancy has been found to be associated with several adverse maternal and neonatal outcomes, including premature birth [31]. In our study, data on birth and pregnancy outcomes were not collected for mothers without neurodivergent children, therefore, it is unclear at present whether the adverse outcomes are specific to FMS mothers with autistic children.

A final clinical prospective study (Study C) invited all patients from the initial service evaluation to complete a survey capturing detailed data on maternal factors and children's autistic traits. Consistent with the online survey, mothers with FMS were statistically significantly associated with an increased likelihood of autistic children. In this study, we additionally found that a first-degree family history of neurodivergence was associated with autism in children, which aligns with the well-documented role of genetics in the development of autism [21]. However, these findings should be interpreted with caution as the sample size is small, and therefore, the confidence intervals are wide. Maternal psychological variables – general anxiety (GAD-7) and autistic traits (AQ10 and RAADS-14) – were assessed in this study as potential factors related to autism diagnoses in offspring. Previous evidence suggests that FMS patients display high health anxiety [44], which in turn could lead to a heightened awareness regarding their children's health. However, the lack of association of maternal GAD-7 scores to autism diagnoses in children appears to suggest that psychological distress impacting parental behaviours does not explain the group differences

seen in the prevalence of autistic children. Moreover, no significant differences were found between mothers on the AQ-10 and RAADS-14 autistic trait measures, which indicates that FMS mothers did not display higher autistic traits compared to mothers with other chronic pain. However, this should be interpreted with caution as the AQ-10 displayed poor reliability in this cohort, and the sample size was small.

This is the first study to demonstrate a statistically significant association between maternal FMS and autism specifically in children, when compared to other chronic pain conditions. Our results are consistent with those from Kelly [26] who reported an increased prevalence of neurodivergent conditions among the *relatives* (first- and second-degree) of 13 patients with FMS and concomitant hypermobility, as compared to a control group of osteoarthritis patients. However, whether the mechanisms underpinning increased autism rates in children of patients with FMS are identical to those explaining increased autism rates in relatives overall is uncertain. The findings of our online survey hint at the possibility that autistic children of FMS mothers may be explained, at least in part, by the perinatal environment rather than solely by genetic factors.

There were several limitations to the current study. Firstly, self-selection bias may have influenced the high reported prevalence of diagnosed autism in both the online survey and the clinical prospective investigation. Individuals who had a known incidence of autism in themselves or their families may have been more interested in participating in the study. However, this alone is unlikely to account for the disparity between chronic pain conditions. Secondly, the research relied on parental reports for autism diagnoses. Although parental confirmation of a diagnosis from a clinician was required in the prospective study, this could not be formally verified. Nevertheless, all autistic children in the analysis exceeded the recommended cut-off values on the parent-reported AQ10 and SCQ autistic trait measures. The online survey also relied on self-reports of chronic pain, and diagnoses could not be

independently verified. Thirdly, there was a small discrepancy in the prospective investigation between the self-reported chronic pain diagnosis and the clinic letter diagnosis in the service evaluation. This could be due to misreporting, or it is possible that, between the time of the clinic letters and completing the study, some patients' diagnoses had changed. Finally, the clinical prospective investigation is limited by its small sample size and the fact that no mothers classified in the other chronic pain group had an autistic child. As a result, the confidence intervals in some of the statistical analysis models are wide, which suggests a high degree of uncertainty in the estimates. To identify the magnitude and specificity of the association between FMS in mothers and children on the autism spectrum, future research with larger samples or population cohort data should be utilised to improve understanding of the effect size.

For the first time, the present study demonstrated an increased likelihood of autistic children of mothers with FMS (and CRPS) compared to mothers with other chronic pain conditions across multiple, independent cohorts. We suggest asking mothers with severe chronic pain, particularly FMS and CRPS, in clinical practice about autistic children to better understand the patient's home and life situation. More research is required to quantify this correlation and to understand its cause.

[1] Allison C, Auyeung B, Baron-Cohen S. Toward brief "red flags" for autism screening: The short autism spectrum quotient and the short quantitative checklist in 1,000 cases and 3,000 controls. Journal of the American Academy of Child & Adolescent Psychiatry 2012;51(2):202-212. e207.

[2] Applbaum E, Lichtbroun A. Novel Sjögren's autoantibodies found in fibromyalgia patients with sicca and/or xerostomia. Autoimmunity Reviews 2019;18(2):199-202.

[3] Association AP. Diagnostic and statistical manual of mental disorders: American psychiatric association, 2013.

[4] Banerjee N, Adak P. Birth related parameters are important contributors in autism spectrum disorders. Scientific Reports 2022;12(1):14277.

[5] Berwick RJ, Sahbaie P, Kenny G, Guo T-Z, Neiland H, Andersson DA, Clark JD, Mallon P, Goebel A. Postacute COVID-19 syndrome and fibromyalgia syndrome are associated with anti-satellite glial cell IgG serum autoantibodies but only fibromyalgia syndrome serum-IgG is pronociceptive. PAIN 2025;166(10):e397-e408.

[6] Brimberg L, Sadiq A, Gregersen P, Diamond B. Brain-reactive IgG correlates with autoimmunity in mothers of a child with an autism spectrum disorder. Molecular psychiatry 2013;18(11):1171-1177.

[7] Brown CO, Uy J, Singh KK. A mini-review: Bridging the gap between autism spectrum disorder and pain comorbidities. Canadian Journal of Pain 2020;4(4):37-44.

[8] Brugha TS, McManus S, Bankart J, Scott F, Purdon S, Smith J, Bebbington P, Jenkins R, Meltzer H. Epidemiology of autism spectrum disorders in adults in the community in England. Archives of general psychiatry 2011;68(5):459-465.

[9] Casale R, Atzeni F, Bazzichi L, Beretta G, Costantini E, Sacerdote P, Tassorelli C. Pain in women: a perspective review on a relevant clinical issue that deserves prioritization. Pain and therapy 2021;10(1):287-314.

[10] Chen S-w, Zhong X-s, Jiang L-n, Zheng X-y, Xiong Y-q, Ma S-j, Qiu M, Huo S-t, Ge J, Chen Q. Maternal autoimmune diseases and the risk of autism spectrum disorders in offspring: a systematic review and meta-analysis. Behavioural brain research 2016;296:61-69.

[11] Clauw DJ. Fibromyalgia: a clinical review. Jama 2014;311(15):1547-1555.

[12] Cooksey R, Choy E. Exploring gender differences, medical history, and treatments used in patients with fibromyalgia in the UK using primary-care data: a retrospective, population-based, cohort study. The Lancet Rheumatology 2022;4:S20.

[13] Croen LA, Qian Y, Ashwood P, Daniels JL, Fallin D, Schendel D, Schieve LA, Singer AB, Zerbo O. Family history of immune conditions and autism spectrum and developmental disorders: Findings from the study to explore early development. Autism Research 2019;12(1):123-135.

[14] Dolcino M, Tinazzi E, Puccetti A, Lunardi C. Gene expression profiling in fibromyalgia indicates an autoimmune origin of the disease and opens new avenues for targeted therapy. Journal of clinical medicine 2020;9(6):1814.

[15] Erbescu A, Papuc SM, Budisteanu M, Arghir A, Neagu M. Re-emerging concepts of immune dysregulation in autism spectrum disorders. Frontiers in psychiatry 2022;13:1006612.

[16] Eriksson JM, Andersen LM, Bejerot S. RAADS-14 Screen: validity of a screening tool for autism spectrum disorder in an adult psychiatric population. Molecular Autism 2013;4(1):49.

[17] Fayaz A, Croft P, Langford R, Donaldson L, Jones G. Prevalence of chronic pain in the UK: a systematic review and meta-analysis of population studies. BMJ open 2016;6(6):e010364.

[18] Gau S-Y, Leong P-Y, Lin C-L, Tsou H-K, Wei JC-C. Higher risk for Sjögren's syndrome in patients with fibromyalgia: a nationwide population-based cohort study. Frontiers in Immunology 2021;12:640618.

[19] Goebel A, Krock E, Gentry C, Israel MR, Jurczak A, Urbina CM, Sandor K, Vastani N, Maurer M, Cuhadar U. Passive transfer of fibromyalgia symptoms from patients to mice. The Journal of clinical investigation 2021;131(13).

[20] Häuser W, Perrot S, Clauw DJ, Fitzcharles M-A. Unravelling fibromyalgia—steps toward individualized management. The Journal of Pain 2018;19(2):125-134.

[21] Havdahl A, Niarchou M, Starnawska A, Uddin M, van der Merwe C, Warrier V. Genetic contributions to autism spectrum disorder. Psychological medicine 2021;51(13):2260-2273.

[22] Heinze GP, Meinhard; Jiricka, Lena; Steiner, Gregor. logistf: Firth's Bias-Reduced Logistic Regression, 2023.

[23] Helyes Z, Tékus V, Szentes N, Pohóczky K, Botz B, Kiss T, Kemény Á, Környei Z, Tóth K, Lénárt N. Transfer of complex regional pain syndrome to mice via human autoantibodies is mediated by interleukin-1–induced mechanisms. Proceedings of the National Academy of Sciences 2019;116(26):13067-13076.

[24] Israel MR, Berwick R, Vastani N, Zheng Q, Moore W, Maurer M, Gentry C, Marshall A, Sun H, Neiland H. Aβ low threshold mechanoreceptors contribute to sensory abnormalities in fibromyalgia. Brain 2025:awaf321.

[25] Jones KL, Van de Water J. Maternal autoantibody related autism: mechanisms and pathways. Molecular psychiatry 2019;24(2):252-265.

[26] Kelly C. The association between fibromyalgia, hypermobility and neurodivergence extends to families: Brief report. International Journal of Psychiatry Research 2023;6(3):1-5.

[27] LaSalle JM. Epigenomic signatures reveal mechanistic clues and predictive markers for autism spectrum disorder. Molecular psychiatry 2023;28(5):1890-1901.

[28] Laverty C, Surtees A, O'Sullivan R, Sutherland D, Jones C, Richards C. The prevalence and profile of autism in individuals born preterm: a systematic review and meta-analysis. Journal of Neurodevelopmental Disorders 2021;13(1):41.

[29] Lipsker CW, Bölte S, Hirvikoski T, Lekander M, Holmström L, Wicksell RK. Prevalence of autism traits and attention-deficit hyperactivity disorder symptoms in a clinical sample of children and adolescents with chronic pain. Journal of Pain Research 2018:2827-2836.

[30] MA RJB. After-Sensations and Lingering Pain Following Examination in Patients with Fibromyalgia Syndrome. 2022.

[31] Magtanong GG, Spence AR, Czuzoj-Shulman N, Abenhaim HA. Maternal and neonatal outcomes among pregnant women with fibromyalgia: a population-based study of 12 million births. The Journal of Maternal-Fetal & Neonatal Medicine 2019;32(3):404-410.

[32] Merriwether EN, Agalave NM, Dailey DL, Rakel BA, Kolker SJ, Lenert ME, Spagnola WH, Lu Y, Geasland KM, Allen L-AH. IL-5 mediates monocyte phenotype and pain outcomes in fibromyalgia. Pain 2021;162(5):1468-1482.

[33] O'Nions E, Petersen I, Buckman JE, Charlton R, Cooper C, Corbett A, Happé F, Manthorpe J, Richards M, Saunders R. Autism in England: assessing underdiagnosis in a population-based cohort study of prospectively collected primary care data. The Lancet Regional Health–Europe 2023;29.

[34] Ooi KL, Ong YS, Jacob SA, Khan TM. A meta-synthesis on parenting a child with autism. Neuropsychiatric disease and treatment 2016:745-762.

[35] Plazier M, Ost J, Stassijns G, De Ridder D, Vanneste S. Pain characteristics in fibromyalgia: understanding the multiple dimensions of pain. Clinical Rheumatology 2015;34(4):775-783.

[36] Queiroz LP. Worldwide epidemiology of fibromyalgia. Current pain and headache reports 2013;17(8):356.

[37] Ramirez-Celis A, Becker M, Nuño M, Schauer J, Aghaeepour N, Van de Water J. Risk assessment analysis for maternal autoantibody-related autism (MAR-ASD): a subtype of autism. Molecular psychiatry 2021;26(5):1551-1560.

[38] Ramirez-Celis A, Croen LA, Yoshida CK, Alexeeff SE, Schauer J, Yolken RH, Ashwood P, Van de Water J. Maternal autoantibody profiles as biomarkers for ASD and ASD with co-occurring intellectual disability. Molecular psychiatry 2022;27(9):3760-3767.

[39] Revelle W. psych: Procedures for Psychological, Psychometric, and Personality Research: Northwestern University, 2025.

[40] Robinson-Agramonte MdlA, Noris García E, Fraga Guerra J, Vega Hurtado Y, Antonucci N, Semprún-Hernández N, Schultz S, Siniscalco D. Immune dysregulation in autism spectrum disorder: what do we know about it? International journal of molecular sciences 2022;23(6):3033.

[41] Rutter M, Le Couteur A, Lord C. Autism diagnostic interview-revised. Los Angeles, CA: Western Psychological Services 2003;29(2003):30.

[42] Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. Archives of internal medicine 2006;166(10):1092-1097.

[43] Suhas S, Manjunatha N, Kumar CN, Benegal V, Rao GN, Varghese M, Gururaj G. Firth's penalized logistic regression: A superior approach for analysis of data from India's National Mental Health Survey, 2016. Indian journal of psychiatry 2023;65(12):1208-1213.

[44] Uçar M, Sarp Ü, Karaaslan Ö, Gül AI, Tanik N, Arik HO. Health anxiety and depression in patients with fibromyalgia syndrome. Journal of International Medical Research 2015;43(5):679-685.

[45] Whitney DG, Shapiro DN. National prevalence of pain among children and adolescents with autism spectrum disorders. JAMA pediatrics 2019;173(12):1203-1205.

[46] Wolfe F, Walitt B, Perrot S, Rasker JJ, Häuser W. Fibromyalgia diagnosis and biased assessment: Sex, prevalence and bias. PloS one 2018;13(9):e0203755.

[47] Yavne Y, Amital D, Watad A, Tiosano S, Amital H. A systematic review of precipitating physical and psychological traumatic events in the development of fibromyalgia, Proceedings of the Seminars in arthritis and rheumatism, Vol. 48: Elsevier, 2018. pp. 121-133.

[48] Yin W, Norrbäck M, Levine SZ, Rivera N, Buxbaum JD, Zhu H, Yip B, Reichenberg A, Askling J, Sandin S. Maternal rheumatoid arthritis and risk of autism in the offspring. Psychological medicine 2023;53(15):7300-7308.