Systemic burden predicts patient-reported health status independently of left

ventricular ejection fraction in older adults with heart failure

Nguyen Le Huy Hoang[1*], Pham Thi Ngoc Anh[1], Trinh Cong Tuan[1], Than Ha Ngoc The[1,2],

Nguyen Ngoc Hoanh My Tien[1,2], and Pham Huy Tung[3]

Work performed at Thong Nhat Hospital, Ho Chi Minh City, Vietnam.

[1]School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi

Minh City, Vietnam.

[2]Department of Geriatrics and Palliative Care, University Medical Centre Ho Chi Minh City,

Ho Chi Minh City, Vietnam.

[3]Department of Radiology, Thong Nhat Hospital, Ho Chi Minh City, Vietnam.

*Corresponding author:

Nguyen Le Huy Hoang, MD, MSc

Position: cardiologist and geriatrician

Institution: School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City

Address: 217 Hong Bang Street, Cho Lon Ward, Ho Chi Minh City, Vietnam

Phone number: +84 946 639 943

Fax: not available

Email: lucian.nguyen@icloud.com

ORCID ID: 0009-0002-2791-2838

Other authors:

Pham Thi Ngoc Anh, MD, MSc

Position: physician

Institution: School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City

Address: 217 Hong Bang Street, Cho Lon Ward, Ho Chi Minh City, Vietnam

Email: phamthingocanh798@gmail.com

Trinh Cong Tuan, MD, MSc

Position: physician

Institution: School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City

Address: 217 Hong Bang Street, Cho Lon Ward, Ho Chi Minh City, Vietnam

Email: tringcongtuan@gmail.com

Than Ha Ngoc The, MD, PhD

Position: lecturer and physician

Institution: School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City

and Department of Geriatrics and Palliative Care, University Medical Centre Ho Chi Minh

City

Address: 217 Hong Bang Street, Cho Lon Ward, Ho Chi Minh City, Vietnam

Email: the.thn@umc.edu.vn

Nguyen Ngoc Hoanh My Tien, MD, MSc, Specialist Level 2

Position: lecturer and physician

Institution: School of Medicine, University of Medicine and Pharmacy at Ho Chi Minh City

and Department of Geriatrics and Palliative Care, University Medical Centre Ho Chi Minh

City

Address: 217 Hong Bang Street, Cho Lon Ward, Ho Chi Minh City, Vietnam

Email: tien.nnhm@umc.edu.vn

Pham Huy Tung, MD, MSc

Position: radiologist

Institution: Department of Radiology, Thong Nhat Hospital

Address: 1 Ly Thuong Kiet Street, Tan Son Nhat Ward, Ho Chi Minh City, Vietnam

Email: tunghuy516@gmail.com

Total word count: 2,969 words

Abstract

In heart failure (HF), management priorities are shifting to include health-related quality of

life, measured by patient-reported outcome measures (PROMs) like the Kansas City

Cardiomyopathy Questionnaire-12 (KCCQ-12). While left ventricular ejection fraction

(LVEF) guides therapy, it correlates poorly with patient-reported symptoms (the 'LVEF

paradox'). The contributions of systemic factors, such as New York Heart Association

(NYHA) functional class, comorbidity, and frailty, remain poorly quantified against LVEF in

older, ambulatory patients. This analytical, cross-sectional, observational study aimed to

identify independent predictors of patient-reported health status. We enrolled 150 consecutive

patients aged 60 years or older with chronic HF from a single cardiology centre. The primary

outcome was the KCCQ-12 overall summary score. A multivariable linear regression model

quantified associations of LVEF, NYHA functional class (I-II vs. III-IV), Charlson

Comorbidity Index (CCI), frailty (Clinical Frailty Scale 5 or higher), age, and sex with

KCCQ-12 scores. The cohort (median age 75 years) had a median KCCQ-12 score of 68,

which did not differ significantly across HF phenotypes. The model explained 76.7% of the

variance (R-squared = 0.767). Significant independent predictors of poorer health status were:

advanced NYHA functional class (Coefficient -15.88), higher comorbidity burden

(Coefficient -3.17 per 1-point CCI increase), and frailty (Coefficient -6.82) (all $p < 0.001$).

Male sex predicted higher health status (Coefficient 3.21, $p = 0.030$). LVEF ($p = 0.561$) and

age ($p = 0.909$) were not significant predictors. In this cohort, patient-reported health status

was independently determined by systemic factors (NYHA class, comorbidity, and frailty),

not LVEF. These findings underscore the 'LVEF paradox' and support a broader approach to

HF assessment incorporating functional, comorbidity, and frailty measures to guide patient-

centred management.

Introduction

Heart failure (HF) is a major global health issue, affecting over 64 million people, with its prevalence increasing due to ageing populations.[1,2] Consequently, management priorities have expanded from reducing mortality to improving functional capacity and health-related quality of life (HRQoL).[3–5] Clinical guidelines now recommend routine assessment of health status using validated patient-reported outcome measures (PROMs).[4,5] The Kansas City Cardiomyopathy Questionnaire (KCCQ), including its validated 12-item short form Kansas City Cardiomyopathy Questionnaire-12 (KCCQ-12), is the benchmark disease-specific instrument for this purpose.[6]

Pharmacological therapy for HF is guided by left ventricular ejection fraction (LVEF), which classifies patients into distinct phenotypes.[4] However, LVEF correlates poorly with patient symptoms, a phenomenon known as the 'LVEF paradox'.[7,8] Instead, patient-reported health status appears determined by systemic factors.[9] These include clinician-assessed functional status, quantified by New York Heart Association (NYHA) functional class, comorbidity burden, and frailty, a multisystem syndrome of decreased physiological reserve, all of which are established predictors of poor outcomes and reduced HRQoL.[10]

The relative contributions of NYHA functional class, multimorbidity, and frailty to health status, particularly in comparison to LVEF, remain poorly quantified. This uncertainty is pronounced in older, ambulatory patients who are frequently underrepresented in clinical trials.[11] However, no equivalent framework exists for predicting patient-reported health status, despite the established role of LVEF in guiding pharmacological therapy.[12]

This study aimed to identify and quantify the independent predictors of patient-reported health status, measured by the KCCQ-12, in older ambulatory adults with chronic HF. We hypothesised that factors reflecting systemic burden, including NYHA functional class,

108   comorbidity, frailty, age, and sex, would predict KCCQ-12 scores independently of LVEF.

109   The inclusion of sex is supported by established differences in HF pathophysiology and

110   outcomes between men and women.[13]

Methods

Study design and participants

This analytical observational, cross-sectional study followed Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for reporting.[14] The source population comprised consecutive patients aged 60 years or older with chronic HF attending Thong Nhat Hospital cardiology clinic between August 2024 and March 2025. Patients were excluded if they had acute decompensated heart failure or severe cognitive impairment preventing informed consent or questionnaire completion. Sample size calculations used two approaches: first, estimating mean KCCQ-12 scores with 5-point precision required 97 participants; second, Green's formula for testing regression coefficients with six predictors required 110 participants. The final cohort of 150 participants exceeded both requirements. Ethical approval was obtained from the University of Medicine and Pharmacy at Ho Chi Minh City (2118/HDDD-DHYD) and Thong Nhat Hospital (121/2024/CN-BVTN-HDDD). The study complied with the Declaration of Helsinki,[15] and all participants provided written informed consent before enrolment. Participant flow is shown in Figure 1.

Data collection and measurements

All procedures were completed during a single outpatient visit at Thong Nhat Hospital cardiology clinic. During the visit, trained staff conducted face-to-face interviews in Vietnamese and administered several validated instruments, including the Vietnamese KCCQ-12[16,17] and the Clinical Frailty Scale (CFS),[18] using structured case report forms. A cardiologist performed clinical assessments and assigned NYHA functional class.[19] Data for the Charlson Comorbidity Index (CCI), N-terminal pro–B-type natriuretic peptide (NT-proBNP) levels, and baseline LVEF were abstracted from medical records. Staff received protocol training with competency assessments. Data

were entered into an electronic case report form with validation checks. Analyses used

anonymised data.

Variable definitions

Primary and secondary outcomes

Participants completed a validated Vietnamese version during a baseline visit before

clinical assessment to minimise information bias. Trained staff provided standardised

instructions and scored the questionnaires according to the official manual, blinded

where feasible to clinical data. Scores for the four domains (physical limitation,

symptom frequency, quality of life, and social limitation) were transformed to a 0–100

scale, with higher scores indicating better health. The primary outcome was the overall

summary score, calculated as the mean of the available domain scores. The individual

domain scores served as secondary outcomes. We prespecified a minimal clinically

important difference of 5 or more points for the overall score, which is a conservative

estimate consistent with published data.[6]

Key explanatory variables

LVEF was derived from echocardiograms performed within three months of the study visit.

Functional capacity was defined by the NYHA functional class, which we dichotomised for

regression into mild (classes I–II) and advanced (classes III–IV) limitation.[19] The CCI

represented the comorbidity burden.[20] While treated as a continuous variable in the primary

regression model, for descriptive analyses, scores were stratified into low burden (scores 3–5),

medium burden (6–7), and high burden (≥8). This categorisation was based on the

distribution of scores within the study cohort rather than pre-specified a priori cut-points.

Frailty was defined as a score of 5 or higher on the 9-point CFS. This threshold is widely

used and validated for identifying at least mild frailty in older adults and is associated with

adverse outcomes.

Other covariates

For descriptive purposes, we classified HF phenotype according to 2021 European Society

of Cardiology guidelines based on LVEF: reduced (HFrEF; ≤40%), mildly reduced

(HFmrEF; 41–49%), or preserved (HFpEF; ≥50%).[4] In the primary model, LVEF was

treated as a continuous variable. We recorded demographics (age, sex), lifestyle factors

(smoking, alcohol use), and clinical measurements, including body mass index, averaged

resting heart rate and blood pressure, and NT-proBNP concentrations. Polypharmacy was

defined as using five or more concurrent medications. We also counted the number of

prescribed guideline-directed HF drug classes (range 0–4). Functional status was assessed

using the Katz Index of Independence in Activities of Daily Living,[21] and nutritional status

was evaluated with the Mini Nutritional Assessment Short-Form.[22]

Statistical analysis

All statistical analyses were performed using Stata version 19.5 (StataCorp, College Station,

Texas, USA). Categorical variables were summarised as counts and percentages. Continuous

variables were assessed for normality using the Shapiro-Wilk test and visual inspection of

histograms, then reported as either mean and standard deviation or median and interquartile

range as appropriate.

The primary analysis employed a multivariable linear regression model to identify predictors

of the KCCQ-12 overall summary score. To disentangle the effects of chronological age

from the cumulative comorbidity burden, the model included both age and the CCI as

separate continuous predictors. In addition to age and CCI, the model included the other key

explanatory variables: LVEF, dichotomised NYHA functional class, and binary frailty status,

along with sex. Results were expressed as adjusted regression coefficients with 95%

confidence intervals, which were calculated using robust standard errors as a default method

to account for potential heteroscedasticity. Model diagnostics involved assessing residual

plots for homoscedasticity and normality, whilst variance inflation factors were used to

evaluate multicollinearity.

For descriptive analyses, trends across ordered groups were tested using the Jonckheere-

Terpstra test for continuous or ordinal outcomes and the Cochran-Armitage test for binary

outcomes. KCCQ-12 scores were compared across categories using the Mann-Whitney U

test for two-group comparisons and the Jonckheere-Terpstra test for three or more ordered

groups.

Exploratory analyses involved stratifying the multivariable regression model by HF

phenotype (HFrEF, HFmrEF, HFpEF) to examine whether associations between the

predictor variables and KCCQ-12 scores differed across these subgroups. As the study was

not powered for formal subgroup analyses, these results require cautious interpretation.

Additional exploratory analyses examined whether the associations between predictor

variables and health status differed between HFrEF and HFmrEF, with the full results

provided in the Supplementary Material.

All analyses used complete-case data, meaning participants with missing data for any

variable were excluded from that specific analysis, and the proportions of missing data are

reported in the baseline characteristics table. A two-sided p-value below 0.05 was considered

statistically significant. No adjustment was made for multiple comparisons in the

exploratory analyses.

Results

Cohort and baseline description

The study cohort comprised 150 older adults with HF, who were categorised into HFrEF,

HFmrEF, and HFpEF groups (Figure 1). The median age was 75 years, with patients in the

HFpEF group being significantly older than those in the HFrEF group.

Geriatric syndromes such as malnutrition, frailty, and limitations in daily activities were

prevalent across the cohort. Most patients had a high comorbidity burden and were on

multiple medications. While the proportion of patients with a high comorbidity burden

appeared to increase from the HFrEF to the HFpEF phenotype, this trend was not statistically

significant. Common comorbidities included hypertension, coronary artery disease, and

dyslipidaemia. Notably, the prevalence of atrial fibrillation increased significantly from the

HFrEF to the HFpEF group (Table 1).

Among patients with HFrEF, the prescription of foundational therapies was common. The

vast majority of these patients were on combination therapy, with most receiving two or three

different drug classes (Supplementary Table 1).

Primary endpoint

The median KCCQ-12 summary score was 68 and did not differ significantly across HF

phenotypes. However, health status was significantly poorer in certain subgroups, including

patients in advanced NYHA functional classes, those with frailty or an increasing

comorbidity burden, and females (Table 2).

A multivariable linear regression model was developed to identify independent predictors of

health status, explaining 76.7% of the variance in KCCQ-12 scores. The analysis identified

higher NYHA functional class, increased comorbidity burden, and the presence of frailty as

significant predictors of poorer health status. In contrast, male sex was associated with a

higher health status score. After adjusting for other variables, age and LVEF were not found

to be significant predictors (Table 3 and Figure 2).

Diagnostic checks confirmed the validity of the model's assumptions. There was no evidence

of multicollinearity among the predictor variables (Supplementary Table 2). Visual

inspection of the quantile-quantile plot confirmed that the residuals were approximately

normally distributed (Supplementary Figure 2). Furthermore, a plot of residuals versus fitted

values showed a random scatter, satisfying the assumption of homoscedasticity

(Supplementary Figure 1).

Secondary endpoints

Consistent with the primary outcome, no significant trend was observed across HF

phenotypes for any KCCQ-12 domain scores, including physical limitation, symptom

frequency, quality of life, and social limitation. Subgroup analyses revealed consistent

patterns of poorer health status, with patients in advanced NYHA functional classes, frail

individuals, those with an increasing comorbidity burden, and females reporting

significantly lower scores across all four domains (Table 4).

Exploratory analysis

Multivariable linear regression models, stratified by HF phenotype, explained a substantial

portion of the variance in KCCQ-12 scores for all three groups (Supplementary Table 3).

In both the HFrEF and HFpEF cohorts, advanced NYHA functional class, frailty, and higher

comorbidity burden were independent predictors of poorer health status. For patients with

HFpEF, male sex also predicted higher scores. In contrast, among patients with HFmrEF,

only an increasing comorbidity score independently predicted lower health status.

A significant interaction was observed between comorbidity burden and phenotype when

comparing HFrEF and HFmrEF, indicating that health status declined more steeply with

increasing comorbidities in the HFmrEF group (Supplementary Table 5, Supplementary

Figure 5).

Diagnostic checks for the stratified models confirmed that there was no evidence of

multicollinearity and that the assumptions of homoscedasticity and normally distributed

residuals were met (Supplementary Table 4, Supplementary Figure 3, Supplementary Figure

4).

Discussion

Principal findings of the study

In this study of 150 older, ambulatory Vietnamese adults with chronic HF, patient-reported

health status was determined by systemic and functional factors, not LVEF. The multivariable

analysis identified that a higher NYHA functional class, increased comorbidity burden, and

frailty were independent predictors of poorer health status, while male sex was associated

with a higher health status score. LVEF showed no association with KCCQ-12 scores. These

results indicate that, in older adults, patient-reported health status aligns more closely with

functional capacity, comorbidity burden, and physiological reserve than with ventricular

function alone.

Contextualising health status assessment in geriatric heart failure

HF management increasingly emphasises functional capacity and HRQoL alongside mortality

reduction, reflecting demographic changes in ageing populations. Clinical guidelines now

recommend the routine use of PROMs like the KCCQ.[3–5] These findings support

incorporating geriatric assessment domains into routine HF evaluation. The results suggest

that assessment of NYHA functional class, comorbidity, and frailty complements LVEF-

based pharmacotherapy in addressing patient-centred outcomes recommended by current

guidelines.

The left ventricular ejection fraction paradox and phenotype-independent health status

The dissociation between LVEF and patient-reported health status has been widely reported.

For instance, a 2020 study that followed patients for 12 months found that the proportions of

patients experiencing clinically significant changes in KCCQ scores did not differ between

HFpEF and HFrEF groups, which suggests that the trajectory of health status is independent

of the underlying phenotype.[23] Similarly, research by Pokharel et al. found that the prognostic

power of the current KCCQ score for predicting death and hospitalisation holds true for

patients irrespective of their ejection fraction, supporting a combined analysis approach.[24]

LVEF reflects cardiac mechanics rather than systemic physiological reserve or patient-

perceived well-being, consistent with previous evidence. The model identified systemic

factors as the strongest predictors of health status. These findings suggest that patient-

reported symptoms correlate with systemic factors (frailty, comorbidity) rather than with

isolated measures of cardiac function.

The primacy of functional class and the 'discordance' problem

This finding relates to clinician-patient discordance, a recognised clinical challenge. A 2022

study found that one-third of HF patients experience a discordance between their clinician-

assessed functional status and their self-reported health (KCCQ).[25] The largest discordant

group (27.4% of patients) comprised those with 'discordant worse KCCQ'.[25] These patients

were classified as NYHA I/II but reported low KCCQ scores. This discordant group had an

80% higher risk of hospitalisation or death compared to the 'concordant good' group (low

NYHA, high KCCQ).[26] This pattern may be explained by frailty and comorbidity burden,

which independently lower KCCQ scores even at lower NYHA functional classes. For

instance, a patient assessed as low-risk (e.g., NYHA II) may still report very poor health if

they are also frail and have a high comorbidity burden, as these factors independently lower

health status scores. These are systemic factors that the NYHA classification alone does not

capture. A three-predictor model incorporating NYHA functional class, CCI, and frailty may

identify high-risk discordant patients more effectively than NYHA functional class alone.

The independent, dose-dependent impact of comorbidity burden

Health status declined progressively with increasing comorbidity burden, with median KCCQ

scores decreasing from 81 in the low-burden group to 58 in the high-burden group. These

findings are consistent with Yang et al. (2023), who reported associations between cardiac

and non-cardiac comorbidities and reduced health status in HFrEF and HFpEF.[27] The present

study in a Vietnamese cohort demonstrates similar associations. The exploratory analysis

suggested a hypothesis regarding the HFmrEF phenotype. In the HFmrEF group, only

comorbidity burden remained a significant predictor, with a larger effect size than observed in

other phenotypes. Interaction testing confirmed this difference was statistically significant,

suggesting that comorbidity burden may be a particularly important determinant of health

status in HFmrEF. The small HFmrEF sample size limits interpretation; however, the finding

suggests that non-cardiac disease burden may substantially influence health status in this

phenotype and warrants validation in larger studies.

Validating geriatric syndromes as a core determinant

This study demonstrates an association between frailty assessment and patient-reported

outcomes (KCCQ). The CFS is a practical bedside assessment tool that has been shown to

predict mortality independently in HF patients. Notably, frailty emerged as an independent

predictor distinct from other clinical variables. Frailty is sometimes conflated with older age,

multimorbidity, or higher NYHA functional class. The multivariable model did not support

this assumption. After adjusting for age, comorbidity, and functional class, frailty remained

an independent predictor. Frailty may capture loss of physiological reserve, a dimension not

reflected in other clinical measures. Frailty assessment may provide prognostic information

beyond that obtained from LVEF, NYHA functional class, and comorbidity indices.

Strengths and limitations

This study has several strengths, including its consecutive enrolment of patients from a

dedicated ambulatory clinic, focus on older adults, and simultaneous modelling of clinician-

assessed function, comorbidity, and frailty against a validated patient-reported outcome

measure. The inclusion of a Vietnamese cohort adds valuable, non-Western data to a global

literature dominated by North American and European studies. However, several limitations

must be acknowledged. The single-centre design limits the generalisability of the findings,

and the cross-sectional analysis can only establish association, not causation. The small size

of the HFmrEF subgroup renders the finding of the CCI as the sole driver in this phenotype

speculative and requires cautious interpretation. Finally, our model did not include other

potentially important factors, such as socioeconomic status or psychosocial health.

Future perspectives and clinical implications

These findings have potential clinical implications. Clinically, LVEF alone may be

insufficient for assessing, managing, or predicting patient-reported burden. Combined

assessment of NYHA functional class, CCI, and CFS can be completed rapidly in clinical

practice. This combined assessment may improve risk stratification and support

individualised care planning consistent with guideline recommendations. For future research,

longitudinal studies are required to determine if changes in frailty and comorbidity burden

over time correlate with changes in KCCQ scores. Randomised clinical trials are warranted to

test whether interventions targeting these factors improve KCCQ scores as a primary

endpoint. The hypothesis that comorbidity burden is a predominant determinant of health

status in HFmrEF requires validation in larger, multicentre cohorts.

Conclusion

The findings suggest that patient-reported health status is more closely associated with

systemic factors than with cardiac mechanics alone. These findings support a broader

approach to HF assessment beyond LVEF alone. Incorporating assessment of functional class,

comorbidity, and frailty into routine cardiovascular care may improve identification of high-

risk patients and support patient-centred management focused on quality of life.

Author contributions

All authors approved the final submitted version and any revised versions. Nguyen Le Huy Hoang is the guarantor with full data access and responsibility for accuracy.

Funding: None declared.

Acknowledgements

We thank the patients and their families for participating in this study. We are grateful to the clinicians, echocardiography technologists, and nursing staff at Thong Nhat Hospital for their support with patient recruitment, clinical assessments, and echocardiographic evaluations. We acknowledge the use of the KCCQ-12, licensed from CV Outcomes, Inc. (Kansas City, Missouri, USA).

Declaration of interest: None declared.

Data availability statement

The data that support the findings of this study are available from the corresponding author, Nguyen Le Huy Hoang, upon reasonable request. The data are not publicly available due to restrictions protecting patient privacy.

Reference

1. Savarese G, Becher PM, Lund LH, Seferovic P, Rosano GMC, Coats AJS. Global burden of heart failure: a comprehensive and updated review of epidemiology. Cardiovasc Res 2022; 118 :3272–87. doi: 10.1093/cvr/cvac013.

2. Groenewegen A, Rutten FH, Mosterd A, Hoes AW. Epidemiology of heart failure. European Journal of Heart Failure 2020; 22 :1342–56. doi: 10.1002/ejhf.1858.

3. Volterrani M, Halasz G, Adamopoulos S, Agostoni PG, Butler J, Coats AJS, et al. Quality of life in heart failure. The heart of the matter. A scientific statement of the Heart Failure Association and the European Association of Preventive Cardiology of the European Society of Cardiology. European Journal of Heart Failure 2025; 27 :1159–73. doi: 10.1002/ejhf.3440.

4. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: Developed by the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) With the special contribution of the Heart Failure Association (HFA) of the ESC. European Heart Journal 2021; 42 :3599–726. doi: 10.1093/eurheartj/ehab368.

5. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation 2022; 145 :e895–1032. doi: 10.1161/CIR.0000000000001063.

6.   Spertus JA, Jones PG, Sandhu AT, Arnold SV. Interpreting the Kansas City

Cardiomyopathy Questionnaire in Clinical Trials and Clinical Care. JACC 2020;

76 :2379–90. doi: 10.1016/j.jacc.2020.09.542.

7.   Bristow MR, Kao DP, Breathett KK, Altman NL, Gorcsan J, Gill EA, et al. Structural and

Functional Phenotyping of the Failing Heart. JACC: Heart Failure 2017; 5 :772–81. doi:

10.1016/j.jchf.2017.09.009.

8.   Gorodeski EZ, Chu EC, Reese JR, Shishehbor MH, Hsich E, Starling RC. Prognosis on

Chronic Dobutamine or Milrinone Infusions for Stage D Heart Failure. Circulation: Heart

Failure 2009; 2 :320–4. doi: 10.1161/CIRCHEARTFAILURE.108.839076.

9.   Johansson I, Balasubramanian K, Bangdiwala S, Mielniczuk L, Hage C, Sharma SK, et al.

Factors associated with health-related quality of life in heart failure in 23,000 patients

from 40 countries: results of the global congestive heart failure research program. Eur

Heart J 2022; 43 :ehac544.879. doi: 10.1093/eurheartj/ehac544.879.

10. Wu L, Liu S, Zhang M, Xiong X. Prevalence and Associated Factors of Frailty in Patients

with Chronic Heart Failure: A Systematic Review and Meta-Analysis. RCM 2025;

26 :26854. doi: 10.31083/RCM26854.

11. Van Spall HGC, Toren A, Kiss A, Fowler RA. Eligibility Criteria of Randomized

Controlled Trials Published in High-Impact General Medical JournalsA Systematic

Sampling Review. JAMA 2007; 297 :1233–40. doi: 10.1001/jama.297.11.1233.

12. Bristow MR, Kao DP, Breathett KK, Altman NL, Gorcsan J, Gill EA, et al. Structural and

Functional Phenotyping of the Failing Heart. JACC: Heart Failure 2017; 5 :772–81. doi:

10.1016/j.jchf.2017.09.009.

13. Lam CSP, Arnott C, Beale AL, Chandramouli C, Hilfiker-Kleiner D, Kaye DM, et al. Sex

differences in heart failure. Eur Heart J 2019; 40 :3859–3868c. doi:

10.1093/eurheartj/ehz835.

14. Elm E von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The

Strengthening the Reporting of Observational Studies in Epidemiology (STROBE)

Statement: Guidelines for Reporting Observational Studies. PLOS Medicine 2007;

4 :e296. doi: 10.1371/journal.pmed.0040296.

15. World Medical Association. World Medical Association Declaration of Helsinki: Ethical

Principles for Medical Research Involving Human Subjects. JAMA 2013; 310 :2191–4.

doi: 10.1001/jama.2013.281053.

16. Thành NT, Thanh NX, Hương NTT, Thắng P, Huyền VTT. 23. TÍNH TIN CẬY VÀ GIÁ

TRỊ CỦA THANG ĐO KCCQ-12 ĐỂ ĐÁNH GIÁ CHẤT LƯỢNG CUỘC SỐNG Ở

BỆNH NHÂN SUY TIM MẠN TÍNH.

17. Spertus JA, Jones PG. Development and Validation of a Short Version of the Kansas City

Cardiomyopathy Questionnaire. Circulation: Cardiovascular Quality and Outcomes 2015;

8 :469–76. doi: 10.1161/CIRCOUTCOMES.115.001958.

18. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A

global clinical measure of fitness and frailty in elderly people. CMAJ 2005; 173 :489–95.

doi: 10.1503/cmaj.050051.

19. Caraballo C, Desai NR, Mulder H, Alhanti B, Wilson FP, Fiuzat M, et al. Clinical

Implications of the New York Heart Association Classification. Journal of the American

Heart Association 2019; 8 :e014240. doi: 10.1161/JAHA.119.014240.

20. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying

prognostic comorbidity in longitudinal studies: Development and validation. Journal of

Chronic Diseases 1987; 40 :373–83. doi: 10.1016/0021-9681(87)90171-8.

21. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of Illness in the Aged:

The Index of ADL: A Standardized Measure of Biological and Psychosocial Function.

JAMA 1963; 185 :914–9. doi: 10.1001/jama.1963.03060120024016.

22. Kaiser MJ, Bauer JM, Ramsch C, Uter W, Guigoz Y, Cederholm T, et al. Validation of the

Mini Nutritional Assessment short-form (MNA®-SF): A practical tool for identification

of nutritional status. The Journal of nutrition, health and aging 2009; 13 :782–8. doi:

10.1007/s12603-009-0214-7.

23. Huang W, Teng THK, Tay WT, Richards AM, Kadam U, Lawson CA, et al. Patient-

reported outcomes in heart failure with preserved vs. reduced ejection fraction: focus on

physical independence. ESC Heart Failure 2020; 7 :2051–62. doi: 10.1002/ehf2.12950.

24. Pokharel Y, Khariton Y, Tang Y, Nassif ME, Chan PS, Arnold SV, et al. Association of

Serial Kansas City Cardiomyopathy Questionnaire Assessments With Death and

Hospitalization in Patients With Heart Failure With Preserved and Reduced Ejection

Fraction: A Secondary Analysis of 2 Randomized Clinical Trials. JAMA Cardiol 2017;

2 :1315–21. doi: 10.1001/jamacardio.2017.3983.

25. Greene SJ, Butler J, Spertus JA, Hellkamp AS, Vaduganathan M, DeVore AD, et al.

Comparison of New York Heart Association Class and Patient-Reported Outcomes for

Heart Failure With Reduced Ejection Fraction. JAMA Cardiol 2021; 6 :522–31. doi:

10.1001/jamacardio.2021.0372.

26. Michelis KC, Grodin JL, Zhong L, Pandey A, Toto K, Ayers CR, et al. Discordance

Between Severity of Heart Failure as Determined by Patient Report Versus

Cardiopulmonary Exercise Testing. Journal of the American Heart Association 2021;

10 :e019864. doi: 10.1161/JAHA.120.019864.

27. Yang M, Kondo T, Adamson C, Butt JH, Abraham WT, Desai AS, et al. Impact of

comorbidities on health status measured using the Kansas City Cardiomyopathy

Questionnaire in patients with heart failure with reduced and preserved ejection fraction.

European Journal of Heart Failure 2023; 25 :1606–18. doi: 10.1002/ejhf.2962.

Legends

Figure 1. Patient flow diagram

STROBE-style diagram illustrating the screening and enrolment process. Of 300 patients

screened, 150 were excluded for reasons including incomplete questionnaire data (n = 51),

missing left ventricular ejection fraction measurement (n = 30), acute decompensated heart

failure (n = 23), cognitive impairment (n = 23), and declined participation (n = 15), resulting

in a final cohort of 150 patients

*Other reasons for exclusion included severe non-cardiac illness (n = 5) and planned

relocation (n = 3).

Abbreviations: ADHF, acute decompensated heart failure; HFmrEF, heart failure with mildly

reduced ejection fraction; HFpEF, heart failure with preserved ejection fraction; HFrEF,

heart failure with reduced ejection fraction.

Figure 2. Independent predictors of KCCQ-12 overall summary score

The plot shows the adjusted regression coefficients (blue dots) and 95% confidence intervals

(horizontal lines) from the primary multivariable linear regression model (n = 150). The

dashed vertical line indicates a coefficient of zero, representing no effect. Coefficients and

confidence intervals that do not cross this line are statistically significant.

Abbreviations: CCI, Charlson Comorbidity Index; LVEF, left ventricular ejection fraction;

NYHA, New York Heart Association.

Tables

Table 1. Baseline characteristics by heart failure phenotype

Characteristic	Total (n = 150)	HFrEF (n = 54)	HFmrEF (n = 23)	HFpEF (n = 73)	p-value
Demographics, behavioural factors, and anthropometric measures					
Age (years), median (IQR)	75 (66–84)	70 (65–80)	71 (66–82)	77 (69–85)	0.001
Age group, n (%)					0.005
60-69 years	56 (37.3)	27 (50.0)	10 (43.5)	19 (26.0)	
70-79 years	41 (27.3)	13 (24.1)	6 (26.1)	22 (30.1)	
≥80 years	53 (35.3)	14 (25.9)	7 (30.4)	32 (43.8)	
Male sex, n (%)	89 (59.3)	34 (63.0)	15 (65.2)	40 (54.8)	0.336
Current smoker, n (%)	51 (34.0)	19 (35.2)	9 (39.1)	23 (31.5)	0.642
Current drinker, n (%)	57 (38.0)	24 (44.4)	9 (39.1)	24 (32.9)	0.183
BMI (kg/m^2), mean (SD)	21.6 (2.2)	21.9 (2.3)	21.4 (2.2)	21.4 (2.1)	0.156
BMI stratification, n (%)					0.072
Underweight	15 (10.0)	3 (5.6)	2 (8.7)	10 (13.7)	
Normal	102 (68.0)	35 (64.8)	17 (73.9)	50 (68.5)	
Overweight	26 (17.3)	15 (27.8)	2 (8.7)	9 (12.3)	
Obese	7 (4.7)	1 (1.9)	2 (8.7)	4 (5.5)	
Vital signs and biomarkers					
HR (bpm), median	79 (65–91)	79 (65–91)	65 (63–89)	85 (68–91)	0.420

(IQR)				
SBP (mmHg), mean (SD)	139 (12.0)	140 (12.4)	138 (12.6)	140 (11.6)	0.815
DBP (mmHg), mean (SD)	85 (8.1)	85 (7.9)	86 (9.1)	85 (8.1)	0.875
NT-proBNP (pg/mL), median (IQR)	604 (154–2122)	573 (213–2112)	654 (124–2125)	623 (158–2122)	0.973
Cardiac and functional status					
LVEF (%), median (IQR)	49 (37–66)	35 (31–38)	45 (43–48)	67 (58–72)	<0.001
NYHA functional class, n (%)					0.613
I and II	107 (71.3)	38 (70.4)	20 (87.0)	49 (67.1)	
III and IV	43 (28.7)	16 (29.6)	3 (13.0)	24 (32.9)	
Geriatric syndromes and functional assessment					
Frailty, n (%)	54 (36.0)	19 (35.2)	5 (21.7)	30 (41.1)	0.438
ADL limitation, n (%)	38 (25.3)	12 (22.2)	3 (13.0)	23 (31.5)	0.203
Nutrition status, n (%)					0.240
Malnourished	87 (58.0)	28 (51.9)	13 (56.5)	46 (63.0)	
At risk of malnutrition	52 (34.7)	21 (38.9)	10 (43.5)	21 (28.8)	
Normal	11 (7.3)	5 (9.3)	0 (0.0)	6 (8.2)	
Comorbidity burden					
CCI, median (IQR)	7 (5–8)	6 (5–8)	6 (5–8)	7 (5–9)	0.257

CCI stratification, n (%)					0.217
Low burden	47 (31.3)	16 (29.6)	10 (43.5)	21 (28.8)	
Medium burden	41 (27.3)	21 (38.9)	3 (13.0)	17 (23.3)	
High burden	62 (41.3)	17 (31.5)	10 (43.5)	35 (47.9)	
Polypharmacy, n (%)	132 (88.0)	50 (92.6)	23 (100.0)	59 (80.8)	0.033
Comorbidity, n (%)					
Hypertension	137 (91.3)	49 (90.7)	22 (95.7)	66 (90.4)	0.910
CAD	116 (77.3)	43 (79.6)	16 (69.6)	57 (78.1)	0.882
Atrial fibrillation	43 (28.7)	7 (13.0)	4 (17.4)	32 (43.8)	<0.001
Diabetes	72 (48.0)	29 (53.7)	9 (39.1)	34 (46.6)	0.460
Dyslipidaemia	115 (76.7)	42 (77.8)	15 (65.2)	58 (79.5)	0.761
CKD	43 (28.7)	17 (31.5)	4 (17.4)	22 (30.1)	0.930
Stroke	9 (6.0)	3 (5.6)	1 (4.3)	5 (6.8)	0.746

Abbreviations: ADL, activities of daily living; BMI, body mass index; CAD, coronary artery

disease; CCI, Charlson Comorbidity Index; CKD, chronic kidney disease; DBP, diastolic

blood pressure; HFmrEF, heart failure with mildly reduced ejection fraction; HFpEF, heart

failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction;

HR, heart rate; IQR, interquartile range; LVEF, left ventricular ejection fraction; n, number;

NT-proBNP, N-terminal pro-B-type natriuretic peptide; NYHA, New York Heart Association;

SBP, systolic blood pressure; SD, standard deviation.

Table 2. KCCQ-12 overall summary scores by subgroup

Subgroup	KCCQ-12 overall summary scores	p-value
Total	68 (57–82)	--
HF phenotype		0.619
HFrEF	70 (56–81)	
HFmrEF	72 (60–84)	
HFpEF	66 (56–81)	
NYHA functional class		<0.001
I and II	78 (65–85)	
III and IV	52 (41–56)	
CCI stratification		<0.001
Low burden	81 (77–89)	
Medium burden	70 (63–83)	
High burden	58 (43–63)	
Frailty		<0.001
No	79 (65–85)	
Yes	55 (42–62)	
Age group		<0.001
60-69 years	79 (65–85)	
70-79 years	76 (64–85)	
≥80 years	56 (50–64)	
Sex		0.001
Female	63 (55–74)	
Male	76 (60–83)	

495   *Data are reported as median (IQR).*

Table 3. Multivariable linear regression of health status

Total (n = 150)		
Predictor	**Coefficient (95% CI)**	**p-value**
LVEF (per 1% increase)	0.02 (-0.06 to 0.11)	0.561
NYHA functional class		
I and II	ref	ref
III and IV	-15.88 (-19.03 to -12.73)	<0.001
CCI (per 1 score increase)	-3.17 (-4.13 to -2.22)	<0.001
Frailty (yes vs. no)	-6.82 (-10.31 to -3.32)	<0.001
Age (per 1 increase)	-0.01 (-0.19 to 0.17)	0.909
Sex (male vs. female)	3.21 (0.32 to 6.10)	0.030
Model statistics		
R-squared	0.767	
F-statistic	67.94	<0.001

502 *Abbreviations: CCI, Charlson Comorbidity Index; CI, confidence interval; LVEF, left*

503 *ventricular ejection fraction; n, number; NYHA, New York Heart Association; ref, reference.*

Table 4. KCCQ-12 domain scores by subgroup

Subgroup	Physical limitation scores	p-value	Symptom frequency scores	p-value	Quality of life scores	p-value	Social limitation scores	p-value
Total	67 (42–83)	--	73 (65–85)	--	50 (50–75)	--	75 (67–83)	--
HF phenotype		0.565		0.262		0.539		0.556
HFrEF	75 (42–75)		74 (65–85)		75 (50–75)		71 (67–83)	
HFmrEF	67 (50–83)		73 (70–94)		50 (50–75)		75 (67–92)	
HFpEF	58 (42–75)		73 (60–79)		50 (50–75)		67 (67–83)	
NYHA functional class		<0.001		<0.001		<0.001		<0.001
I and II	75 (58–83)		79 (70–92)		75 (50–75)		75 (67–92)	
III and IV	42 (25–42)		54 (42–67)		50 (38–50)		58 (50–67)	
CCI stratification		<0.001		<0.001		<0.001		<0.001
Low burden	83 (75–100)		85 (73–94)		75 (50–88)		83 (75–92)	
Medium burden	67 (58–75)		73 (67–85)		75 (50–75)		75 (67–83)	
High burden	42 (33–58)		65 (50–73)		50 (50–50)		67 (50–67)	

Frailty		<0.001		<0.001		<0.001		<0.001
No	75 (58–83)		79 (70–92)		75 (50–88)		75 (67–92)	
Yes	42 (33–58)		60 (44–70)		50 (50–50)		63 (50–67)	
Age group		<0.001		<0.001		<0.001		<0.001
60-69 years	75 (58–83)		79 (70–93)		75 (50–75)		83 (67–92)	
70-79 years	75 (58–83)		73 (67–92)		75 (50–88)		75 (67–83)	
≥80 years	42 (42–58)		65 (50–70)		50 (50–75)		67 (58–67)	
Sex		0.011		0.001		0.002		<0.001
Female	58 (42–75)		70 (58–73)		50 (50–75)		67 (58–75)	
Male	75 (50–83)		75 (67–92)		75 (50–75)		75 (67–83)	

505 *Data are reported as median (IQR).*

506 *Abbreviations: CCI, Charlson Comorbidity Index; HFmrEF, heart failure with mildly reduced ejection fraction; HFpEF, heart failure with*

507 *preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; IQR, interquartile range; KCCQ-12, Kansas City*

508 *Cardiomyopathy Questionnaire-12; NYHA, New York Heart Association.*

509