Divergence Between Net Fluid and Weight-Based Evaluation in Calculating Cumulative Fluid Balance

Finley J. Shinnick,[1] Denise C. Hasson, MD,[2] Ulka Kothari, MD,[3] Ami Shah, MD,[2] James D. Odum, MD,[4] Chloe G. Braun, MD,[5] Celeste G. Dixon, MD,[6] Julie C. Fitzgerald, MD, PhD, MSCE,[6] Susan D. Martin, MD,[7] Nina Terry, BSN,[7] Adam C. Dziorny, MD, PhD,[7] on behalf of the Evidence-Driven Evaluation and Management of fluid Accumulation (EDEMA) Collaborative

[1] Department of Pediatrics, Golisano Children's Hospital, University of Rochester School of Medicine, Rochester, NY

[2] Department of Pediatrics, Hassenfeld Children's Hospital at NYU Langone Health, New York, NY

[3] Department of Pediatrics, NYU Long Island School of Medicine, Mineola, NY

[4] Department of Pediatrics, University of Alabama at Birmingham, Birmingham, AL

[5] Department of Pediatrics, University of Louisville School of Medicine, Louisville, KY

[6] Department of Anesthesiology and Critical Care Medicine, Children's Hospital of Philadelphia, The University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

[7] Department of Pediatrics, Golisano Children's Hospital, University of Rochester School of Medicine, Rochester, NY

Corresponding Author:

Adam Dziorny, MD, PhD

adam_dziorny@urmc.rochester.edu

601 Elmwood Ave, Box 667

Rochester, NY 14642

Keywords: PICU, Cumulative fluid balance, Fluid overload, Fluid balance Assessment

Financial support for this study: Funding for this work was provided by OKRA (Grant Number: 5U24DK137318).

ABSTRACT

Objective: Although efforts have been made to standardize fluid balance calculations in the ICU, there is a limited understanding of how different calculation methods relate to one another across admission. We quantify the agreement between the cumulative fluid balance calculated from serial weights (CFBw) and the cumulative fluid balance calculated from fluid intake and output (CFBf) in critically ill children during the first week of intensive care unit (PICU) admission.

Design: Retrospective, multicenter, federated observational study.

Setting: We included four PICUs and two cardiac PICUs from four tertiary care centers.

Patients: Analysis included 8,895 pediatric patients (<19 years old) representing 12,388 PICU encounters from 2023-2024.

Interventions: None.

Measurements and Main Results: A patient's anchor weight was determined as their weight closest to ICU admission, and CFBw and CFBf were calculated at the time of each new weight measurement. We assessed agreement between CFBf and CFBw using Bland-Altman analyses stratified by ICU day and patient subgroups (neonates, early anchor weights [weight on ICU day 0], and encounters with unmeasured urine events). Across all units and subgroups, CFBf exceeded CFBw (mean difference: all patients = 4.7 %CFB, early anchor weight = 4.7 %CFB, neonates = 5.9 %CFB). The mean difference increased significantly over time (days 0–3: 2.7 %CFB vs. days 4–7: 8.1 %CFB, $p<0.05$), with greater divergence observed in neonates and patients with early anchor weights.

Conclusions: CFB derived from fluid intake and output consistently exceeded weight-based calculations, with a greater divergence on ICU days 4-7 in all subgroups. Early anchor weights, which may be potentially truer starting weights, increased the mean difference between CFBf and CFBw. Future studies should explore each method's association with patient outcomes to identify the most clinically informative CFB method.

Research in Context

- Net fluid (intake minus output) and weight measurements are used to evaluate PICU patients' fluid balance, but few studies have directly compared these methods across multiple institutions, unit types, and throughout PICU admission.

- CFB estimated by net fluid consistently exceeds weight-based estimates, with divergence increasing later in admission and for patients with anchor weights recorded closer to PICU admissions.

- Recording PICU admission weights early and maintaining accurate weight and fluid monitoring are essential for reliable fluid balance assessment, as CFB calculations diverge between methods.

At the Bedside

- This multicenter study demonstrates a consistent positive divergence between CFB calculated by net fluid compared to weight-based calculations across diverse PICU and PCICU populations.

- Differences between methods diverged later in admission (e.g., after day 3), particularly among neonates and patients with earlier recorded anchor weights, while CFB estimates before the first unmeasured urine differed little from the overall sample.

- Clinicians should understand these differences and the factors that influence them, prioritizing early and frequent patient weights throughout ICU admission to recognize when net fluid-based CFB diverges from a weight-based CFB.

INTRODUCTION

Fluid overload, or the pathologic increase in cumulative fluid balance (CFB), occurs in more than one-third of children admitted to the pediatric intensive care unit (PICU) (1) and is associated with multiple morbidities (2). Due to the dose-dependent association between fluid accumulation and poor outcomes in PICU populations, precise tracking of CFB across the course of admission is essential for guiding clinical decision-making (3–7). Although efforts have been made to standardize fluid balance definitions and calculations in the ICU (8), there is a limited understanding of how different calculation methods relate to one another across admission (9,10).

Two commonly accepted methods are used to assess CFB through ICU admission, each with its own advantages and disadvantages (11). The first method compares net fluid intake and output relative to ICU admission weight (CFBf) (12). While this method aims to capture fluid that remains in a patient, there is uncertainty in quantifying immeasurable fluid losses, and it is prone to error when certain outputs cannot be measured (e.g., flushed urine, emesis, unweighed stool) (13). Alternatively, physicians track fluid balance by serial weight changes (CFBw), which assumes that all weight change is solely attributable to fluid accumulation (14). Weighing critically ill children can be cumbersome, resource-intensive, and inaccurate due to lines, drains, and tubes, and is often not done routinely. Both methods require an anchor weight, which serves as the reference point for changes in weight or fluid volume. Despite limitations, each method aims to capture a patient's true CFB. While both methods are commonly used in clinical practice, their degree of agreement beyond the first 48 hours of admission across diverse patient populations remains unclear.

The objective of this multi-institutional study was to quantify the agreement between CFBw and CFBf, stratified by patient populations over the first 7 days of ICU and PCICU admission. We report the variability in obtaining daily weights across units and the general features of CFB evaluation by each method, and specifically how they differ in high-risk populations (e.g., neonates) and with different methodologies (e.g., definition of anchor weight). We hypothesize that CFBf and CFBw will diverge as a patient gets later into their ICU stay, and divergence will be greater in neonates and those in whom anchor weight is recorded after the first day of ICU admission.

METHODS

Study Design

This is a retrospective observational cohort study conducted within our Evidence-Driven Evaluation and Management of fluid Accumulation (EDEMA) collaborative (4, 15). This study followed procedures in accordance with ethical standards as delineated by the Helsinki Declaration of 1975. Each unit's institutional review board (IRB) deemed the project exempt from IRB review.

Characteristics of our collaborative member units have been described previously (4). We included data from six pediatric ICUs within four tertiary care centers: 132 beds from mixed medical/surgical units (PICUs) and 53 beds from cardiac ICUs (PCICUs), totaling 185 beds. All units use the same electronic health record (EHR), Epic (Epic Systems Corporation, Verona, WI). Our patient cohort included all pediatric patients <19 years old admitted to their PICU/PCICU between each unit's respective date ranges, generally spanning January 1, 2023, to

July 1, 2024. (**Table 1**). Similar to our collaborative's prior federated studies, each unit maintained the privacy of its data, but identical data structures allowed for the sharing of data queries and analysis scripts (4).

Measures and Outcomes

We collected baseline demographics, patients' weights, total intake and output fluid volume by route and subtype, and the number of unmeasured urine occurrences. Consistent with prior studies, we defined Day 0 of ICU stay as admission time through 7 AM the following day (2). We defined Day 1 as the subsequent 24 hours (7 AM to 7 AM), and other days in the following 24 hours until Day 7 or the patient's end of ICU admission. We defined the anchor weight as the measurement temporally closest to ICU admission, whether obtained before or during admission. Due to the federated nature of this study, we could not distinguish potential bias between pre- and post-admission weights. If a patient's anchor weight was recorded before ICU admission, we considered that weight to be the one used for ICU clinical decision-making, and we use that weight for all calculations.

Acknowledging the variability in timing of weights obtained during ICU admission, we calculated CFBw at the time of each additional recorded weight as the change from the anchor weight, divided by the anchor weight. At each non-admission weight measurement timestamp, we calculated CFBf as cumulative fluid intake minus output divided by anchor weight (Equation 1). Granularity of fluid intake and output measurements was to the day, while weights were reported as seconds since ICU admission. To identify fluid values at specific weight timestamps, we proportionally distributed each subtype of fluid intake and output over the ICU day, and we calculated net fluid intake relative to anchor weight (Equation 2). We also calculated a truncated

CFBf (Pre-Occur), to exclude any CFB value that followed a patient's first unmeasurable urine

occurrence.

$$CFBw = \frac{\text{Additional Weight (kg) - Anchor Weight (kg)}}{\text{Anchor Weight (kg)}} \times 100\% \qquad (1)$$

$$CFBf = \frac{\text{Cumulative Intake (L) - Cumulative Output (L)}}{\text{Anchor Weight (kg)}} \times 100\% \qquad (2)$$

We conducted subgroup analyses for patients who had a weight measurement taken on

their ICU day 0 (Early Anchor Weights) and patients younger than one month (Neonate). Early

anchor weight allows a truer assessment of changes in CFB over the full ICU admission, while

neonates, who typically have smaller body weights and may experience nutritional weight gain,

are more sensitive to the impact of changes in fluid volume or weight on CFB calculations. We

also segregated all CFB calculations into ICU days 0-3 and ICU days 4-7 to understand the

relationship between CFBf and CFBw through ICU admission.

Statistical Analysis

Statistical analysis was performed using R (16). We could not compute regression

analyses across units due to the federated nature of this study. We used Shapiro-Wilk tests to

assess normality between the mean difference between CFBf and CFBw in both early versus late

ICU timeframes. We subsequently applied Whitney-Mann tests or Student t-tests to assess

differences in the mean difference of CFB methods between time in ICU admission. We used

linear models to assess unit variation in obtaining and recording weights, based on patient age

group (Table 1) and ventilator status (e.g., active order placed for invasive ventilation). We

calculated raw CFBf and CFBw values across admission to inform the general direction of the

mean difference between methods (Supplemental eFigure 1). We performed Bland-Altman

analysis to assess the relationship between CFBf and CFBw (mean difference +/- standard deviation). Bland–Altman plots for each unit across ICU days 0–7 showed greater scatter later in admission, prompting further evaluation of the relationship between CFBf and CFBw on ICU days 0–3 versus 4–7 (Supplemental eFigure 2). We defined a significant difference as $p < 0.05$.

RESULTS

We included 9,832 PICU and 2,534 PCICU encounters for 8,895 unique patients. Median age of patients at ICU admission varied across units from 3.9 to 7.1 years (Table 1). Across all units, 42-45% of patients were female, and 42-59% of patients were Caucasian (Table 1). The median length of PICU stay was 3-4 days, and the median length of PCICU stay was 4-5 days (Table 1). The ratio of additional non-anchor weights per ICU admissions ranged from 0.2 to 1.8, significantly differing across units (Table 1, $p < 0.005$). Across all units, we failed to find a significant association between patients' ventilator status and the number of repeated weights (Table 1, $p > 0.10$). The proportion of encounters with anchor weights recorded on ICU day 0 ranged from 12% to 62%, significantly differing across units (Table 1, $p < 0.05$). Unit C was significantly associated with both the greatest repeated weight frequency and the greatest proportion of ICU patients with anchor weight on ICU day 0 ($p < 0.001$). At Units B and D, neonates were associated with significantly more additional weights compared to all other age categories ($p < 0.001$). Additionally, the frequency of encounters with unmeasured urine occurrences ranged from 17% to 78%, significantly differing by unit ($p < 0.001$; Table 1). The mean number of non-anchor weights per patient was higher in the PCICUs (Units A and B: 1.7 weights per patient) compared to the PICU units (0.8 weights per patient). Additionally, across

all sites, there were 1,064 patients (8.6%) with duplicated weights throughout ICU admission (i.e., that varied by less than 0.001 kg to the patient's anchor weight).

We calculated a total of 8,634 (unit variation: 224 to 3154) CFB values by each method to match and compare (Supplemental eTable 1), determined by the number of additional (new) weights provided by each unit. Both CFBf and CFBw were generally positive throughout ICU admission, with CFBf increasing faster than CFBw (Supplemental eFigure 1). Across all units, the mean difference and 95% confidence interval between CFBf and CFBw were positive (mean differences from 3.6 to 5.3 % CFB across units, Figure 1), indicating that the calculation of CFB using fluid measurements was consistently of a greater magnitude than the calculation using repeated weight measurements. When comparing patients with early anchor weights to all patients, Units B, C, and D showed increased mean difference between CFBf and CFBw, but this increased mean difference was only statistically significant at Unit C (Figure 1, supplemental eTable 1). Similarly, Units A, C, D, and E demonstrated increased mean difference for neonates compared to all patients, though this increase was only statistically significant in Units A and C (Figure 1) and was limited by the low number of encounters for neonates in Units E and F.

Early versus Late ICU days

When grouping CFB values calculated from ICU days 0-3 and ICU days 4-7, the mean difference increased, or CFBf became more divergent from CFBw, across all units. Each unit saw a significant change in mean difference between timepoints (p<0.05, Figure 1). A sensitivity analysis focused on encounters with early recorded anchor weights maintained this statistically significant increase in mean difference from ICU days 0-3 to ICU days 4-7 (p<0.001) at all but

Unit F (Figure 1). Neonates also showed a significant difference in the mean difference between CFB methods calculated on ICU days 0-3 and ICU days 4-7 ($p < 0.05$).

Impact of Unmeasured Urine Occurrences

Excluding values calculated after a patient's first occurrence resulted in a slight decrease in the mean difference between CFBf and CFBw. This exclusion reduced available CFB comparisons by between 19% and 68%; however, all units still had a positive mean difference between CFBf (Pre-Occur) and CFBw (Supplemental eTable 2). All units, except Unit F, showed a decrease in the average mean difference across ICU days 0-7 when excluding CFB values calculated after a patient's first occurrence (mean (standard deviation, SD) decrease in mean difference: all patients = −0.65% (SE 0.42), early anchor = −1.2% (SE 0.74), neonates = −0.70% (SE 0.54), Supplemental eTable 3). This decrease was only significant for patients with early anchor weights on ICU days 0-3 (Supplemental eTable 2). Only Unit B showed a significant difference between values calculated up until a patient's first occurrence and values calculated over 7 days of ICU admission (Supplemental eTable 2). Both neonates and patients with early anchor weights showed greater mean difference between CFBf and CFBw than CFBf (Pre-Occur) and CFBw on ICU days 4-7 compared to ICU days 0-3 (Supplemental eTable 2).

DISCUSSION

Across all patients and units, cumulative fluid balance calculated from net intake and output (CFBf) was greater in magnitude than that calculated from serial weight measurements (CFBw). Differences between methods increased later in ICU admission (days 4-7), especially for neonates and patients with anchor weights recorded early in ICU admission. Excluding

values calculated after a patient's first unmeasured urine occurrence still maintained a statistically significant mean difference between methods, meaning this difference is likely not solely a result of unmeasured occurrences. To our knowledge, this is the first multicenter, pragmatic observational study to measure these mean differences across children admitted to medical/surgical PICUs and PCICUs over 7 ICU days.

Although there was a relationship between CFBf and CFBw, both methods have inherent clinical complications that might confound the calculation of fluid balance (2). Neonates demonstrate nutritional weight gain during ICU admission, which may be incorrectly attributed to fluid balance (17). Our work, however, showed a greater divergence between CFBf and CFBw for neonates, suggesting neonates' weight did not increase as quickly as net fluid through ICU day 7. There are also well-documented challenges in obtaining accurate weights for critically ill children, including practical difficulties in measurement and standard error inherent to recording weights (18). In this study, however, units range from a 35%-77% frequency of additional weights per cumulative ICU day, likely reflecting a perceived clinical challenge that varies between units. Additionally, across all units, there were 1,064 patients with weights throughout ICU admission identical to 0.001 kg to the patient's anchor weight, which might reflect documentation shortcuts. Past research also highlights challenges in charting complete fluid records (19). Inherent to CFBf calculations, a large amount of "insensible" fluid is known to be lost through respiration or perspiration, which is unaccounted for in CFBf calculations (20). Because of these apparent barriers to true CFB assessment, despite a consistent albeit discrepant relationship between CFBw and CFBf, we cannot confidently determine which method best approximates the patient's true fluid balance.

A similar mean difference between CFBf and CFBw was observed within the first 48 hours of PICU admission (13, 21) and over 72 hours with non-critically ill neonates and with adults (22, 12), all in single-center settings. Similarly, the mean difference between methods increases when anchor weights are taken closer to ICU admission, likely reflecting the positive trend in patient weight from hospital admission through ICU admission (23). Both findings suggest an early fluid accumulation following hospital and ICU admission, which later anchor weights may fail to capture (24). Because CFBf consistently exceeds CFBw, the relative difference is proportionally larger in neonates and patients with early, and likely smaller anchor weights. With this finding, using an earlier anchor weight to calculate CFB results is likely a more accurate reflection of fluid status. In previous studies of patients receiving renal replacement therapy, predicted dry ICU admission weights correlated more closely with organ dysfunction than original anchor weights (25). In our study, we did not adjust for theoretical dry weight.

Excluding CFB values after a patient's first unmeasured urine occurrence decreases the mean difference across all patient groups. Previous studies quantified the threshold of clinically relevant differences in fluid measurements (13), but inconsistent trends in our unit's relationship between CFBf (pre-occur) and CFBf with CFBw limited direct quantitative comparison. No consistent mean difference change was seen on ICU days 4–7, and fewer values were excluded, which may reflect differences in recording urine output for patients with conditions associated with longer ICU stays (Supplemental eTable 2). Some studies consider stool volume when evaluating fluid output, which we did not do in this study, unless they were measured as a

urine/stool mixture with an associated volume; inclusion of this volume may have improved agreement between the two CFB methods (27). Importantly, we did not evaluate the contribution of urine occurrences to individual patients' CFB due to limited weight measurements. Units that had infrequent weight measurements might not have shown a change in mean difference due to fewer weights taken after the patient's first occurrence.

There were several important limitations to this study. The count of daily weights present in the data set limited our comparison between CFBf and CFBw to a small subset of measures per patient. As a result, we evaluated differences between values calculated on early versus late ICU days and pre-/post-occurrence on a per-patient basis. Additionally, none of our analyses evaluated the risks associated with increased CFB via each method; rather, we evaluated the relationship between the two methods. Prior studies from this collaborative have reported CFBf quantification (4). Due to limitations in data extraction, we assumed an approximately linear distribution of each intake and output subtype across an ICU day. This minimizes errors at weight times close to the beginning and end of each day (7 AM) but maximizes potential errors at weights furthest from this day switch (e.g., 7 PM). Federated analysis did not allow for cross-unit regression comparison, and no severity of illness measures were available. Lastly, we did not correlate either methodology to ICU outcomes, which will be done in future studies to further evaluate which method may be most clinically meaningful.

CONCLUSION

We report a consistent positive difference between CFB calculated using fluid intake-output versus weights, which generally increased throughout ICU admission. Neonates with

relatively small ICU weights and patients with early anchor weights likely contribute to the increased difference. Determining a patient's starting anchor weight close to ICU admission also affected this relationship, likely providing a more accurate baseline for change in fluid balance throughout admission. These findings underscore the limitations of both methods and highlight the importance of recording accurate admission weights and frequent weight measurements in critically ill children. Future work will assess the prognostic value of each CFB method in relation to patient outcomes for all critically ill patients, controlling for severity of illness, to better evaluate the clinical significance of the difference between methods.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Sheel Chitre and Adrian La Duca (NYU), Amina Khan (CHOP), and Heather Dutton (UAB) for their assistance with clinical data extraction. We would also like to thank Dr. Benjamin Kozyak, attending physician in the Cardiac Intensive Care Unit at the Children's Hospital of Philadelphia, for his assistance with CHOP CICU data analysis.

CITATIONS

1. Alobaidi, R., Morgan, C., Basu, R. K., Stenson, E., Featherstone, R., Majumdar, S. R., & Bagshaw, S. M. (2018). Association Between Fluid Balance and Outcomes in Critically Ill Children: A Systematic Review and Meta-analysis. *JAMA Pediatrics*, *172*(3), 257–268. https://doi.org/10.1001/jamapediatrics.2017.4540

2. Selewski, D. T., Gist, K. M., Basu, R. K., Goldstein, S. L., Zappitelli, M., Soranno, D. E., Mammen, C., Sutherland, S. M., Askenazi, D. J., Ricci, Z., Akcan-Arikan, A., Gorga, S. M., Gillespie, S. E., Woroniecki, R., & Assessment of the Worldwide Acute Kidney Injury, Renal Angina and Epidemiology (AWARE) Investigators. (2023). Impact of the Magnitude and Timing of Fluid Overload on Outcomes in Critically Ill Children: A Report From the Multicenter International Assessment of Worldwide Acute Kidney Injury, Renal Angina, and Epidemiology (AWARE) Study. *Critical Care Medicine*, *51*(5), 606–618. https://doi.org/10.1097/CCM.0000000000005791

3. Van Regenmortel, N., Verbrugghe, W., Roelant, E., Van Den Wyngaert, T., & Jorens, P. G. (2018). Maintenance fluid therapy and fluid creep impose more significant fluid, sodium, and chloride burdens than resuscitation fluids in critically ill patients: a retrospective study in a tertiary mixed ICU population. *Intensive Care Medicine*, *44*(4), 409–417. https://doi.org/10.1007/s00134-018-5147-3

4. Hasson, D., Shah, A., Braun, C. G., Kothari, U., Drury, S., Dapul, H., Fitzgerald, J. C., Dixon, C., Barbera, A., Odum, J., Terry, N., Weiss, S. L., Martin, S. D., & Dziorny, A. C. (2025). Identifying Opportunities for Fluid Balance Optimization in Critically Ill Children. *J Intensive Care Med*. 2025 Jul 15:8850666251359543. https://doi.org/10.1177/08850666251359543.

5. Barhight, M. F., Nelson, D., Chong, G., Basu, R. K., & Sanchez-Pinto, L. N. (2022). Non-resuscitation fluid in excess of hydration requirements is associated with higher mortality in critically ill children. *Pediatric Research*, *91*(1), 235–240. https://doi.org/10.1038/s41390-021-01456-z

6. Pierce, R., Luckett, P., & Faustino, E. (2017). A Survey of Pediatric Critical Care Providers on the Presence, Severity, and Assessment of Capillary Leak in Critically Ill Children. *Journal of Pediatric Intensive Care*, *06*(03), 145–151. https://doi.org/10.1055/s-0036-1593388

7. Castañuela-Sánchez, V., Hernández-Suárez, A., García-Benítez, L., Díaz-García, L., Martínez-Jasso, G., & Palacios Macedo-Quenot, A. (2022). Fluid overload as a predictor of morbidity and mortality in pediatric patients following congenital heart surgery. *Boletín Médico del Hospital Infantil de México*, *79*(3), 7985. https://doi.org/10.24875/BMHIM.21000183

8. Selewski, D. T., Barhight, M. F., Bjornstad, E. C., Ricci, Z., de Sousa Tavares, M., Akcan-Arikan, A., Goldstein, S. L., Basu, R., Bagshaw, S. M., & Pediatric the Acute Disease Quality Initiative (ADQI) Consensus Committee Members (2024). Fluid assessment, fluid balance, and fluid overload in sick children: a report from the Pediatric Acute Disease Quality Initiative (ADQI) conference. *Pediatric nephrology (Berlin, Germany)*, *39*(3), 955–979. https://doi.org/10.1007/s00467-023-06156-w

9. Hopwood, A., Schade Willis, T., Starr, M., Hughes, K., & Malin, S. (2025). 1340: Development and implementation of a fluid management strategy in the pediatric ICU. *Critical Care Medicine, 53*(1), 1340. https://doi.org/10.1097/01.ccm.0001104024.50214.d4

10. Raina, R., Sethi, S. K., Wadhwani, N., Vemuganti, M., Krishnappa, V., & Bansal, S. B. (2018). Fluid overload in critically ill children. *Frontiers in Pediatrics, 6,* 306. https://doi.org/10.3389/fped.2018.00306

11. Gorga, S. M., Selewski, D. T., Goldstein, S. L., & Menon, S. (2023). An update on the role of fluid overload in the prediction of outcome in acute kidney injury. *Pediatric Nephrology (Berlin, Germany)*. https://doi.org/10.1007/s00467-023-06161-z

12. Antonio, A. C. P., Fernandes, V. R., & Azzolin, K. D. O. (2021). The Correspondence between Fluid Balance and Body Weight Change Measurements in Critically Ill Adult Patients. *The Journal of Critical Care Medicine*, *7*(1), 46–53. https://doi.org/10.2478/jccm-2020-0048

13. Bontant, T., Matrot, B., Abdoul, H., Aizenfisz, S., Naudin, J., Jones, P., & Dauger, S. (2015). Assessing fluid balance in critically ill pediatric patients. *European Journal of Pediatrics*, *174*(1), 133–137. https://doi.org/10.1007/s00431-014-2372-9

14. Bem, R. A., & Lemson, J. (2024). Evaluating fluid overload in critically ill children. *Current Opinion in Pediatrics*, *36*(3), 266–273. https://doi.org/10.1097/MOP.0000000000001347

15. Dixon, C. G., Odum, J., Kothari, U., Martin, S. D., Fitzgerald, J. C., Shah, A., Dapul, H., Braun, C. G., Barbera, A., Terry, N., Weiss, S. L., Hasson, D. C., & Dziorny, A. C. (2025). Post-Hurricane Fluid Conservation Measures Fail to Reduce IV Fluid Use in Critically Ill Children. *Pediatr Nephrol*. 2025 Aug 19. https://doi.org/10.1007/s00467-025-06931-x.

16. R Core Team. (2025). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. https://www.R-project.org/

17. Dijkhuizen, E. I., Dulfer, K., De Munck, S., Van Haren, N. E. M., De Jonge, R. C. J., Vanhorebeek, I., Wouters, P. J., Van Den Berghe, G., Verbruggen, S. C. A. T., & Joosten, K. F. M. (2023). Early weight measures and long-term neuropsychological outcome of critically ill neonates and infants: a secondary analysis of the PEPaNIC trial. *European Journal of Pediatrics*, *183*(2), 649–661. https://doi.org/10.1007/s00431-023-05298-1

18. Gelbart, B., Masterson, K., Serratore, A., Zampetti, M., Veysey, A., Longstaff, S., Bellomo, R., Butt, W., & Duke, T. (2021). Precision of weight measurement in critically ill infants: a technical report. *Critical Care and Resuscitation*, *23*(4), 414–417. https://doi.org/10.51893/2021.4.TN

19. Leinum, L. R., Krogsgaard, M., Tantholdt-Hansen, S., Gögenur, I., Baandrup, A. O., & Azawi, N. (2023). Quality of fluid balance charting and interventions to improve it: a systematic review. *BMJ Open Quality*, *12*(4), e002260. https://doi.org/10.1136/bmjoq-2023-002260

20. McNeil-Masuka, J., & Boyer, T. J. (2025). Insensible Fluid Loss. In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK544219/

21. Gelbart, B., Marchesini, V., Kapalavai, S. K., Veysey, A., Serratore, A., Appleyard, J., Bellomo, R., Butt, W., & Duke, T. (2023). Agreement Between Measured Weight and Fluid Balance in Mechanically Ventilated Children in Intensive Care. *Pediatric Critical Care Medicine*, *24*(10), e459–e467. https://doi.org/10.1097/PCC.0000000000003258

22. Van Asperen, Y., Brand, P. L., & Bekhof, J. (2012). Reliability of the fluid balance in neonates. *Acta Paediatrica*, *101*(5), 479–483. https://doi.org/10.1111/j.1651-2227.2012.02591.x

23. Selewski, D. T., Cornell, T. T., Lombel, R. M., Blatt, N. B., Han, Y. Y., Mottes, T., Kommareddi, M., Kershaw, D. B., Shanley, T. P., & Heung, M. (2011). Weight-based determination of fluid overload status and mortality in pediatric intensive care unit patients requiring continuous renal replacement therapy. *Intensive Care Medicine*, *37*(7), 1166–1173. https://doi.org/10.1007/s00134-011-2231-3

24. Malbrain, M. L. N. G., Martin, G., & Ostermann, M. (2022). Everything you need to know about deresuscitation. *Intensive Care Medicine*, *48*(12), 1781–1786. https://doi.org/10.1007/s00134-022-06761-7

25. Lombel, R. M., Kommareddi, M., Mottes, T., Selewski, D. T., Han, Y. Y., Gipson, D. S., Collins, K. L., & Heung, M. (2012). Implications of different fluid overload definitions in pediatric stem cell transplant patients requiring continuous renal replacement therapy. *Intensive Care Medicine*, *38*(4), 663–669. https://doi.org/10.1007/s00134-012-2503-6

26. Brandewie, K., Alten, J., Winder, M., Mah, K., Holmes, K., Reichle, G., Smith, A., Zang, H., & Bailly, D. (2024). Neonatal Chylothorax and Early Fluid Overload After Cardiac Surgery: Retrospective Analysis of the Neonatal and Pediatric Heart and Renal Outcomes Network Registry (2015–2018)*. *Pediatric Critical Care Medicine*, *25*(3), 231–240. https://doi.org/10.1097/PCC.0000000000003415

27. Chen, J., Li, X., Bai, Z., Fang, F., Hua, J., Li, Y., Pan, J., Wang, J., Feng, X., & Li, Y. (2016). Association of Fluid Accumulation with Clinical Outcomes in Critically Ill Children with Severe Sepsis. *PLOS ONE*, *11*(7), e0160093. https://doi.org/10.1371/journal.pone.0160093

Unit	Unit A	Unit B	Unit C	Unit D	Unit E	Unit F
ICU Type	Cardiac ICU	Cardiac ICU	PICU	PICU	PICU	PICU
ICU Encounters	1902	632	715	6390	1200	1527

Female, n	823 (43%)	285 (45%)	307 (43%)	2655 (42%)	531 (44%)	688 (45%)
Age, n						
<1 month	550 (29%)	100 (16%)	12 (1.68%)	118 (2%)	23 (1.92%)	24 (1.57%)
1 mo–<4 y	771 (41%)	301 (48%)	260 (36%)	1813 (28%)	429 (36%)	625 (41%)
4 yr -< 14 yr	394 (21%)	165 (26%)	265 (37%)	2480 (39%)	455 (38%)	551 (36%)
14–<19 y	154 (8%)	65 (10%)	178 (25%)	1855 (29%)	291 (24%)	309 (20%)
Weight (kg), n						
< 10kg	1091 (57%)	326 (52%)	146 (20%)	1351 (21%)	241 (20%)	359 (24%)
10-30 kg	514 (27%)	175 (28%)	270 (38%)	2961 (46%)	473 (39%)	604 (40%)
>= 30 kg	264 (14%)	129 (20%)	299 (42%)	1946 (30%)	484 (40%)	546 (36%)
Race, n						
White	899 (47%)	334 (53%)	401 (56%)	2671 (42%)	712 (59%)	893 (59%)
Black or African American	278 (15%)	59 (9%)	248 (35%)	1697 (27%)	245 (20%)	225 (15%)
Asian	73 (4%)	23 (4%)	2 (0.28%)	272 (4%)	26 (2%)	83 (5%)
Native Hawaiian or Pacific Islander	1 (0.05%)	1 (0.16%)	0 (0%)	0 (0%)	0 (0%)	5 (0.33%)
American Indian or Alaskan Native	10 (0.53%)	1 (0.16%)	2 (0.28%)	4 (0.06%)	1 (0.08%)	12 (0.79%)
Other/Multiple	497 (26%)	119 (19%)	24 (3%)	1480 (23%)	162 (14%)	104 (7%)
Unknown/Refused	111 (6%)	94 (15%)	38 (5%)	140 (2%)	51 (4%)	187 (12%)
Ethnicity, n						
Hispanic/Latino	416 (22%)	35 (6%)	67 (9%)	1092 (17%)	112 (9%)	221 (14%)
Non-Hispanic/Latino	1327 (70%)	412 (65%)	604 (84%)	4956 (78%)	920 (77%)	495 (32%)
Other/Unknown/Refused	126 (7%)	184 (29%)	44 (6%)	216 (3%)	166 (14%)	793 (52%)
Clinical Characteristics						
Date Range	01/01/2023 - 07/01/2024	01/01/2023 - 07/01/2024	12/01/2023- 05/01/2024	01/01/2023 - 07/01/2024	01/01/2023 - 07/01/2024	01/01/2023 - 07/01/2024
Cum. ICU Days	9918	3414	2635	27370	4609	6311
Encounters	1902	632	715	6390	1200	1527
Median LOS [IQR]	4 [2, 6]	5 [2, 7]	2 [1, 3]	2 [2, 5]	2 [1, 4]	2 [2, 4]
Frequency of Additional Wt.	2352 (1.2)	1131 (1.8)	1321 (1.8)	3260 (0.5)	231 (0.2)	733 (0.5)
Additional Wt. on Vent.	1544 (66%)	267 (24%)	565 (43%)	1636 (50%)	126 (55%)	466 (64%)
Encounters with Early Anchor Wt.	1686 (89%)	596 (94%)	678 (95%)	5910 (92%)	317 (98%)	1489 (98%)
Encounters with Urine Occur.	324 (17%)	496 (78%)	261 (37%)	2720 (43%)	828 (69%)	540 (35%)

Table 1. Patient demographics, weight measurements, and occurrence frequency at each unit. The date range indicates the period of data extraction for each unit. Percentages for patient demographics indicate the proportion of patients in a given demographic at a given unit. Cum. ICU days include the sum of all encounters' days 0-7 in the PICU. LOS indicates the median and interquartile range (IQR) length of ICU stay (in days). The frequency of additional weights gives the count of additional weights and the ratio to each unit's cumulative ICU days. The additional weight on ventilator shows how many extra weights were taken for patients on a ventilator and what percentage of all additional weights that represents.

Encounters with early anchor weights had an anchor weight on ICU day 0, and proportions relative to total encounters at each unit.