Comparing Wastewater-Based and Case-Based R_t Estimates of SARS-CoV-2 Transmission in Georgia, USA Using Generalised Linear Mixed Models

2025-03-05

Authors:

Seth Edmunds[1*], Doug Landsittel[1,3], Marco Ajelli[2], Maria Litvinova[1]

Affiliations:

[1] Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA

[2] Laboratory for Computational Epidemiology and Public Health, Indiana University School of Public Health, Bloomington, IN, USA

[3] Department of Biostatistics, State University of New York at Buffalo, Buffalo, NY, USA

[*]Corresponding Author

Email: shedmund@iu.edu

Abstract

The COVID-19 pandemic has highlighted limitations in case-based surveillance due to inconsistent testing and reporting. Wastewater-based epidemiology (WBE) has emerged as a complementary surveillance approach for tracking SARS-CoV-2 transmission, capturing both symptomatic and asymptomatic infections. The aim of this study is to evaluate the effectiveness of WBE in estimating the effective reproduction number (R_t) of SARS-CoV-2 in Georgia, USA. We used a Generalised Linear Mixed Model to analyse viral concentration data from multiple wastewater treatment plants (WWTPs) collected between June 1, 2022 and December 15, 2022. After controlling for flow rates and site-level heterogeneity, model residuals were transformed into a non-negative incidence-like series used to estimate wastewater-based R_t. Wastewater-based R_t was compared with case-based R_t estimates using Spearman correlation. The two R_t estimates showed concordant temporal patterns across most sites, with stronger correlations in areas with higher case counts (Spearman correlations ranging from 0.39 to 0.84, $p < 0.001$). Wastewater-based R_t tracked increases and decreases in transmission over similar time scales as case-based estimates, while exhibiting reduced sensitivity to short-term changes in clinical testing and reporting behaviour. These findings suggest that WBE can support estimation of

34 transmission trends and complement traditional case-based surveillance for public health
35 monitoring.

36 ───

37 Introduction

The COVID-19 pandemic has underscored the need for effective surveillance systems to
monitor disease transmission at the population level. Initially, case-based surveillance
was critical for tracking the spread of COVID-19. However, fluctuations in testing rates and
the changes in reporting practices gradually undermined its reliability. By late 2022, the
shift by the Centers for Disease Control and Prevention (CDC) from daily to weekly
aggregated case and death reporting reduced the temporal resolution of data available for
public health monitoring. This change was intended to reduce the reporting burden and
improve operational efficiency, but it limited the ability to assess localised and short-term
transmission dynamics. After the expiration of the public health emergency on May 11,
2023, the U.S. Department of Health and Human Services (HHS) could no longer require
laboratories to submit COVID-19 testing results; subsequently, hospital COVID-19
reporting requirements were discontinued effective May 1, 2024 (1). Together, these
changes highlight the need for complementary surveillance approaches capable of
monitoring population-level transmission when clinical reporting is incomplete or delayed.

In response to these challenges, wastewater-based epidemiology (WBE) has re-emerged
as a promising complementary approach, providing a non-invasive, scalable method to
track viral RNA in community wastewater, capturing both symptomatic and asymptomatic
infections. WBE offers a unique opportunity to supplement traditional case-based
surveillance, particularly by providing a population-level signal that is less dependent on
individual testing behaviour and can reflect changes in community transmission. Because
wastewater surveillance occurs at the community level, it is less sensitive to biases arising
from uneven access to testing and underreporting, making WBE especially valuable in
settings with limited healthcare access or inconsistent clinical testing (2, 3).

With the rise of at-home COVID-19 testing, which is largely not reported, comparing
wastewater-based and clinical estimates of the effective reproduction number (R_t) has
become increasingly important for understanding transmission dynamics while
recognizing that each data stream reflects different processes and sources of spatio-
temporal heterogeneity. Recent studies have begun to leverage wastewater data alongside
clinical surveillance to infer transmission dynamics and develop early-warning systems
under varying assumptions about shedding, reporting, and observation processes (4). By
capturing SARS-CoV-2 RNA at the community level, wastewater-based epidemiology
(WBE) provides a population-scale signal that is less influenced by biases in clinical data,
such as uneven testing access, changing individual behaviour, or reporting delays, though
it is subject to its own sources of variability (5

Several studies have estimated wastewater-based SARS-CoV-2 effective reproduction
numbers using various assumptions, preprocessing, and model complexities (2, 6) While
county-level estimations have shown high concordance with traditional case-based
estimates (7), there is a need to validate these methods at finer scales. Using a unique
dataset that includes both reported cases and wastewater surveillance at the level of
individual sewersheds, we demonstrate the concordance of R_t estimations on a
disaggregated level. This study provides evidence for the operational utility of wastewater-
based R_t estimations and offers a practical pipeline accounting for sewershed-related
heterogeneity.

The aim of this study is to evaluate the effectiveness of WBE in estimating R_t of SARS-CoV-
2 at a sewershed level. We focus on data collected between June 1 and December 15,
2022 - a period during which routine case reporting via electronic laboratory and provider
reports remained available, enabling a contemporaneous comparison of wastewater-
based and case-based R_t estimates. To compare these signals, we apply a Generalised
Linear Mixed Model (GLMM) to wastewater concentrations, adjusting for temporal trends,
flow-related dilution, and persistent sewershed differences. This approach allows us to
assess the degree to which wastewater-derived transmission dynamics align with
contemporaneous case-based R_t estimates at the sewershed level, and to evaluate the
role of WBE as a complementary surveillance signal for clinical data at finer scales. (1, 8).

91 Methods

92 Data Collection

93 Wastewater Data Sources

WWTP employees collected twice-weekly wastewater samples, which were shipped to two
university laboratories for analysis. Sampling frequency was typically twice weekly across
sites, though exact schedules varied by facility and operational constraints (summarized in
Appendix Table A1). Viral RNA concentrations were quantified using PCR, specifically
targeting the N2 gene for its widespread use and comparable performance to the N1 target
(Huisman et al., 2022). RT-qPCR was used at WWTP 9 following the method detailed by
Lott et al. (2023) due to logistical constraints, while dPCR, preferred for its higher
sensitivity to low viral concentrations, was used at all other sites following the protocol
described by Sablon et al. (2022) (Sablon et al. 2022). Differences in analytical platform
across sites were treated as time-invariant site characteristics and accounted for through
site-level random effects in downstream modelling. Flow rate data were obtained from
routine facility records and aligned with wastewater sampling dates; depending on facility
practice, measurements corresponded to composite (time- or flow-weighted) or grab
sampling approaches, consistent with standard wastewater surveillance reporting
practices (11). Concurrent flow rates were incorporated to account for dilution-related
variability in wastewater volume.

Case Data Sources

Case data were obtained from routine provider and electronic laboratory reports submitted to the Department of Public Health (DPH) and stored within an electronic disease reporting system. Residential addresses were geocoded to obtain XY coordinates. Sewershed shapefiles were provided by the facilities. Case-to-sewershed assignment assumes accurate residential geocoding and alignment with sewershed boundaries, consistent with standard public health surveillance practice. Case data were aggregated daily by sewershed geography using symptom onset date when available; specimen collection date was used otherwise.

Data Processing

Case Data Processing

Cases were assigned to sewersheds based on geocoded residential addresses and the spatial overlap with sewershed boundaries. Epidemic curves were generated using either the date of symptom onset or, if unavailable, the date of the first positive specimen (including asymptomatic cases). Missing daily case counts were imputed using Kalman filtering to provide smoothed estimates, ensuring the preservation of temporal sequences while handling reporting fluctuations. Imputation primarily reflects completion of a daily time series from routine non-daily reporting rather than filling extended gaps in surveillance and the Kalman approach was selected based on comparative performance in preserving temporal structure, as summarized in the Appendix. Data were restricted to the study period from June 1, 2022, to December 15, 2022, and the processed case data were used in conjunction with wastewater viral concentrations for analysis.

Wastewater Data Processing

All data are from 8 sewersheds in Georgia, USA. Outliers in wastewater viral concentrations and flow rates were detected using the Interquartile Range (IQR) method, with outliers replaced by the median to reduce their influence on downstream analysis. Flagged values were treated as transient spikes for robustness rather than assumed measurement errors, and the number of flagged observations by site is reported in Appendix Table A2. While Appendix Table A1 summarizes the characteristics of all 10 facilities participating in the broader state surveillance program, two facilities, WWTP 4 and WWTP 7, lacked sufficient data during the study period window (June 1 – December 15, 2022). Because only observations exceeding the IQR threshold were modified, minimum and maximum values could remain unchanged if those extremes did not meet the outlier criterion. This method was selected based on a sensitivity analysis comparing multiple outlier detection techniques, including Median Absolute Deviation (MAD), using agreement with case-based R_t (Spearman correlation and RMSE) as evaluation criteria. Samples were predominantly 24-hour composite, collected twice weekly. Population served by individual sewersheds varied from 10,000 (WWTP 10) to 650,000 (WWTP 1) Sewershed locations spanned urban, suburban, and rural areas across Georgia; specific

geographic coordinates are not disclosed to protect facility identities, but site-level
population, flow, and sample characteristics are summarized in Appendix Table A1.
Concentrations were not pre-normalised. Instead, by including flow as a fixed effect in the
GLMM, the model inherently adjusts for flow-related dilution, with the resulting residuals
representing viral signal fluctuations independent of flow variation.

Preprocessing method selection was guided by comparative evaluation across multiple
approaches. Outlier detection using the IQR method flagged 27 concentration values and
57 flow values across sites, compared with 27, and 123 respectively for the MAD approach.
Missing data imputation using Kalman filtering (71.8% of the daily time series) reflected
completion to daily resolution from approximately twice-weekly sampling; inter-sample
gaps had a median of 2 days. Sensitivity analyses across rolling window lengths showed
that 14-day windows achieved the highest concordance with case-based R_t estimates
(median Spearman ρ = 0.608, IQR = 0.121), comparable to 10-day windows (ρ = 0.612, IQR
= 0.184) and superior to 7-day windows (ρ = 0.593, IQR = 0.254). The selected
preprocessing combination (IQR outlier detection, Kalman imputation, 14-day rolling
average) balanced noise reduction with preservation of transmission trends and minimized
data loss (flagging fewer than half as many flow outliers as the MAD approach), as
assessed by correlation with case-based estimates (median RMSE = 0.098). Complete
site-level characteristics, outlier counts, missingness patterns, and sensitivity results are
provided in Appendix Tables A1-A4.

Missing data were interpolated using Kalman filtering, chosen for its performance relative
to linear and spline interpolation methods in preserving temporal structure during routine
non-daily sampling. (Miyazawa et al. 2024; Narci et al. 2021) Both viral concentrations and
flow rates were log-transformed to stabilise variance and normalise the data distribution.
Comparisons of raw and processed data are presented in Appendix Table A5. These
preprocessing steps were implemented to stabilise the time series for downstream
transmission trend estimation rather than to reconstruct true incidence or shedding
processes.

177   Statistical Modelling

178   Model Specification

A Generalised Linear Mixed Model (GLMM) was employed (significance level set to α=0.05)
to analyse viral concentrations in wastewater, with both fixed and random effects to
capture the hierarchical and temporal structure of the data. The fixed effects included time
and flow rate, while the random effects accounted for site-level variability across WWTPs.
Model diagnostics were used to assess statistical fit and stability of the mixed-effects
model, including residual behaviour and random-effect variance. While these diagnostics
ensure the statistical robustness of the framework, they do not, by themselves, imply the
biological validity of the resulting residual time series as a direct and comprehensive
measure of the number of infections. By accounting for both site-specific and time-based

variability, the model provided a flexible framework for analyzing transmission dynamics
signal. (14, 15)

$$\log(Y_{it}) = \beta_0 + \beta_1 \cdot t + \beta_2 \cdot \log(\text{flow}_{it}) + \gamma_{0i} + \epsilon_{it}$$

Where:

- Y_{it} is the viral concentration in wastewater at time t for WWTP i.
- β_0 is the fixed intercept representing the global baseline concentration.
- β_1 is the coefficient for the linear temporal trend.
- β_2 is the coefficient adjusting for the log-transformed wastewater flow rate.
- t is a daily time variable to account for temporal trends.
- flow_{it} is the flow rate of wastewater at time t for WWTP i.
- γ_{0i} is the random intercept for WWTP i, capturing time-invariant site-specific
differences where $\gamma_{0i} \sim N(0, \sigma_\gamma^2)$.
- ϵ_{it} is the residual error term.

The primary specification includes a fixed effect for temporal trend and a site-specific
random intercept, implying a common temporal trend with persistent site-specific
differences. By utilizing a GLMM instead of manual flow-normalization, we provide a
unified framework to adjust simultaneously for flow-related dilution, shared temporal
trends, and site-level heterogeneity (e.g., variations in measurement and sampling
approaches) that are captured by the random intercept (γ_{0i}). By intentionally omitting site-
specific random slopes for time, we ensure that localised deviations from the regional
trend are preserved within the residuals (ϵ_{it}) rather than being absorbed into the model
parameters. This approach allows the residuals to more accurately reflect the site-level
dynamics necessary for subsequent R_t estimations while accommodating the nested data
structure.

It is important to note that the linear time covariate is specified as a global fixed effect
across a geographically diverse set of sewersheds. It is intended to control for shared
baseline trend without absorbing localised epidemic waves. However, researchers
applying this framework should note that if applied to highly synchronized, homogeneous
sites over a short period, the global time effect may absorb the epidemic signal, especially
during the period of consistent epidemic growth or decline. In such cases, the approach
should be implemented without the time trend fixed effects. Empirical demonstrations of
the model's robustness across varying observation windows and distinct transmission
phases are provided in Appendix Figures A1 and A2, respectively.

Residual Extraction and Smoothing

Residuals ϵ_{it} from the GLMM were extracted to capture variability in viral concentrations
not explained by the fixed effects of time, flow rate, and site-specific differences in
wastewater concentrations, including measurement differences. The inclusion of site-
specific random intercepts accounts for persistent between-site differences in baseline

viral concentration, allowing within-site temporal deviations to be examined on a
comparable scale. To reduce high-frequency noise, a 14-day rolling average was applied to
the residual series. While rolling averages introduce additional temporal smoothing and
may attenuate short-term changes, this window length reflects a pragmatic balance
between noise reduction and trend interpretability and was evaluated through sensitivity
analyses reported in Appendix Table A4. This approach ensures that critical transmission
patterns are preserved while minimizing the impact of daily fluctuations. Accordingly,
residual-based analyses are intended to support inference on transmission trends rather
than to reconstruct individual-level infection, shedding processes, case or hospitalization
incidence.

Estimation of the Effective Reproduction Number (Rt)

Rt Calculation

The effective reproduction number (R_t) was estimated using the EpiEstim package in R.
(16, 17) A parametric serial interval distribution was specified with a mean of 2.75 days and
a standard deviation of 2.53 days, utilizing updated parameters specific to the SARS-CoV-2
Omicron variant. (Kremer et al., 2022) To align with the requirements for R_t estimation, the
GLMM residuals were exponentiated to back-transform the values from the log-link model
and ensure the non-negativity required for an incidence proxy. Two estimates of R_t were
calculated: one based on case-derived incidence, and one based on a transformed
wastewater-derived residual series. Both case-based and wastewater-based series were
smoothed using a 14-day rolling average prior to R_t estimation to reduce short-term noise,
with the understanding that this additional smoothing further convolves the underlying
signal; the impact of window length on R_t concordance was evaluated through sensitivity
analyses reported in the Appendix Table A4. (2, 3) As a result, wastewater-derived R_t
estimates are interpreted as a signal of relative transmission dynamics rather than as
direct estimates of intrinsic incidence-based reproduction parameters.

Sensitivity Analysis: Rolling Average Window

A sensitivity analysis was performed to assess the impact of rolling average window length
on both case-based and wastewater-based R_t estimates. Rolling windows ranging from 3
to 30 days were evaluated, with performance assessed based on agreement with case-
based R_t (Spearman correlation and RMSE) and qualitative stability of inferred trends. The
14-day window was selected for primary analysis because it provided a pragmatic balance
between noise reduction and temporal responsiveness across sites, as summarized in
Appendix Table A4. Longer windows further reduced noise but increasingly attenuated
short-term changes during periods of rapid transmission change. (19)

Estimate Alignment

To quantify concordance between wastewater-based and case-based R_t estimates, we
computed Spearman rank correlation coefficients across sites. Analyses were restricted to
the period prior to December 2022, after which changes in case reporting requirements

substantially altered the reliability and completeness of case-based surveillance.
Alignment was further assessed by comparing the temporal co-movement of wastewater-
based R_t, case-based R_t, and reported case counts, focusing on direction and timing of
major transmission changes rather than pointwise equality. This approach emphasizes
concordance in transmission trends under differing observation processes rather than
validation against a presumed ground truth. (20)

Results

The initial dataset comprised 294 observations for both PCR target average concentration
and flow rate measurements across eight WWTPs. After data preprocessing, which
included outlier handling and missing data imputation, the number of observations
increased to 1,019. Detailed statistics on preprocessing outcomes are presented in the
Appendix. A total of 749 data points (72% of the dataset) were imputed using Kalman
smoothing reflecting routine completion of a daily time series from non-daily (typically
twice-weekly) sampling rather than extended gaps in surveillance. Kalman smoothing was
employed because it effectively estimates missing values in time series data by modelling
the underlying stochastic processes. This method accounts for variability by incorporating
both the observed data and the process noise, allowing for smoothed estimates of the
underlying temporal signal amidst measurement uncertainty and routine sampling
variability. (Narci et al. 2021)

Table 1. Core Statistics of R_t Estimates, Correlations and Sewershed Characteristics
(ordered by descending case burden)

Sewershed	Duration (days)	Total Cases	Median Rt (Case)	Median Rt (WW)	Spearman Correlation	P-Value
WWTP 1	191	15288	1	1	0.73	<0.001
WWTP 3	191	2564	1	1	0.39	<0.001
WWTP 5	177	1560	1.02	1.03	0.84	<0.001
WWTP 8	191	1100	1.02	1.03	0.58	<0.001
WWTP 2	26	952	1.11	1.17	0.95	<0.001
WWTP 9	191	503	1.06	1.02	-0.01	0.841
WWTP 6	26	126	1.06	1.23	0.38	0.052
WWTP 10	26	52	1.06	1.2	0.41	0.035

An overview of the R_t estimates derived from both case-based and wastewater-based
surveillance is provided in Table 1. This table includes the duration of data collection, total
case counts, median R_t values, and Spearman correlation coefficients across the eight
wastewater treatment plants (WWTPs). All statistically significant results illustrate notable
positive correlation between the two R_t estimations, with the highest correlation being

observed for WWTP 5 (0.84, $p < 0.001$) and the lowest for WWTP 3 (0.39, $p < 0.001$). The
results (Table 1) indicate that average case burden influenced both correlation strength
and statistical significance (Figure 1). For example, no statistically significant correlation
was observed for WWTP 9; while its data collection duration was comparable to WWTP 1,
its average case burden was substantially lower. Generally, correlation strength and
statistical significance aligned with overall case burden. Sewersheds with very low daily
case counts frequently lacked the temporal variability required to power rank-based
correlation (e.g., WWTP 9 and WWTP 6). However, statistically significant correlations were
occasionally observed even in low-incidence settings (e.g., WWTP 10, averaging
approximately 2 cases per day). The overall pattern is illustrated in Figure 1, where
R_t concordance generally increases as a function of the average number of cases.

Figure 1: Average Number of Cases and Spearman Correlation Coefficient Between Case-
Based and Wastewater-Based R_t Estimates by Wastewater Treatment Plant Sewershed

The vertical dashed line indicates approximately five average daily cases (log scale), below
which rank-based correlation was generally underpowered in our data, though exceptions
were observed (e.g., WWTP 10).

Overview of Rt Estimates Across All Sites

Table 2. Percent of Days with R_t Confidence Intervals Above or Below 1 (ordered by descending case burden)

Sewershed	Case-Based				Wastewater-Based			
	95% CI		50% CI		95% CI		50% CI	
	> 1	< 1	> 1	< 1	> 1	< 1	> 1	< 1
WWTP 1	3.14	17.28	20.94	28.8	25.13	7.85	43.46	33.51
WWTP 3	0.52	0	9.95	12.04	12.57	12.57	37.7	30.89
WWTP 5	0.56	0	14.69	10.17	31.64	15.82	54.24	33.9
WWTP 8	0	0	4.71	4.19	8.9	8.38	49.74	27.75
WWTP 2	19.23	0	96.15	0	84.62	0	100	0
WWTP 9	0	0	8.38	0	2.09	0	40.84	8.38
WWTP 6	0	0	7.69	0	88.46	0	96.15	0
WWTP 10	0	0	7.69	0	65.38	15.38	73.08	19.23

Table 2 illustrates the percentage of days on which lower and upper bounds of the R_t confidence intervals (CIs) were either above or below the epidemic threshold (1.0). This metric provides supplementary information on directional agreement relative to the epidemic threshold (R_t = 1), but was not used as a primary performance criterion because it is sensitive to smoothing choices and uncertainty width.

Detailed Analysis of Representative Sites

The wastewater-based and case-based R_t estimates were analysed across five representative sites with total case incidence greater than 1,000 cases during the study period. Each site exhibits unique patterns in transmission dynamics, as captured by the two surveillance methods. Sites were selected to represent higher-incidence sewersheds where case-based R_t estimates showed sufficient variability to support meaningful temporal comparisons. The sites are presented in the order of decreasing case burden. Two levels of uncertainty around median estimates (50% and 95% Cis) are used to represent two levels of potential risk-tolerance of public health practitioners in making decisions based on uncertain data.

Figure 2. Comparison of Wastewater-Based and Case-Based R_t Estimates for WWTP 1

At WWTP 1, the wastewater-based and case-based R_t estimates closely tracked from June
through August, both peaking in mid-July, which aligns with the peaks in the interpolated
wastewater signal and clinical case incidence (Figure 2). By late August, both estimates fell
below the epidemic threshold (Rt=1). From mid-October onward, the wastewater-based
climbed above epidemic threshold earlier than the case-based estimate in this sewershed.

334
335
336

Figure 3. Comparison of Wastewater-Based and Case-Based R_t Estimates for WWTP 3

In WWTP 3, the wastewater-based estimates sometimes lagged peaks in the case-based
R_t (Figure 3). In mid-October, the wastewater-based R_t increased to approximately
1.1−1.3, with comparatively narrower confidence intervals, while the case-based
estimates exhibited wider uncertainty that encompassed the threshold of 1.0.

Figure 4. Comparison of Wastewater-Based and Case-Based R_t Estimates for WWTP 5

For WWTP 5, the wastewater-based R_t initially aligned with case-based trends during early
summer, with values above 1.0 (Figure 4). In mid-September, the wastewater-based R_t
declined below 0.8 prior to a similar decline in the case-based estimate. A subsequent
increase in the wastewater-based R_t in mid-October preceded a rise in cases.

Figure 5. Comparison of Wastewater-Based and Case-Based R_t Estimates for WWTP 8

At WWTP 8, the wastewater-based R_t remained above 1.0 from early July, with brief dips
(Figure 5). Case-based R_t exhibited broadly similar temporal patterns but with shifts in
timing consistent with differences in smoothing and observation processes.

 Figure 6. Comparison of Wastewater-Based and Case-Based R_t Estimates for WWTP 2

364 At WWTP 2, over the shorter study period, the wastewater-based R_t remained consistently
365 above 1.0, while the case-based R_t rose above 1.0 in early December (Figure 6). An initial
366 elevation in the wastewater-based estimate was interpreted as a transient anomaly
367 consistent with preprocessing and smoothing effects rather than sustained transmission
368 change.

Discussion

Our analyses indicate that wastewater-based R_t estimates exhibited broadly similar
temporal patterns to case-based estimates across most sites, capturing transmission
shifts within approximately two weeks of clinical data. The fundamental purpose of
evaluating a clinical comparator is to determine whether pathogen detection in wastewater
translates into a tangible public health burden. However, selecting an appropriate clinical
comparator requires navigating the inherent limitations of all potential public health
signals. Because R_t estimation relies on the ratio of successive incidence values rather

than absolute counts, it remains mathematically robust to systematic underreporting
provided ascertainment does not shift abruptly. Therefore, the optimal choice must be
guided by the stability of the signal during the observation window and the specific
practical needs of public health authorities.

For this study, we utilized reported case incidence by symptom onset or collection date.
While such data inherently underestimates true infection burden, as demonstrated by
infection incidence being 2.2 to 3.7 times higher than reported cases in the United States
during the Omicron wave (21), reporting requirements during our study period (June–
December 2022) were sufficiently stable to establish a reliable baseline. The subsequent
expiration of public health mandates and the wider utilization of non-reportable at-home
testing rapidly rendered clinical ascertainment lower and highly volatile (21), necessitating
our choice of the study's end date. Therefore, for this study, reported cases provided the
most stable, geospatially appropriate comparator to address the local public health burden
at the sewershed level. However, during periods of rapid surveillance infrastructure
changes, divergence between wastewater and clinical signals should not be interpreted as
a failure of wastewater surveillance, but rather as a complementary reflection of
community transmission dynamics essential for comprehensive situational awareness.

Several limitations should be acknowledged. A substantial portion of the time series
required completion to daily resolution for R_t estimation, primarily reflecting routine non-
daily sampling rather than prolonged data gaps. Kalman smoothing was used to
interpolate missing days and stabilise temporal structure, which may attenuate short-term
variability but is necessary for renewal-based estimation frameworks. Additionally,
wastewater measurements reflect a convolution of infections over heterogeneous
shedding durations and sewer transport processes. As a result, wastewater-derived R_t
should be interpreted as a smoothed indicator of transmission trends rather than a direct
measure of incident infections. This interpretation is subject to uncertainties and
heterogeneities in SARS-CoV-2 shedding likelihood, rates, duration, and routes.
Furthermore, while we account for the flow differences, variability introduced by site-
specific shedding dynamics, viral variants, and in-sewer environmental factors impacts
the precision of these estimates. Nevertheless, these uncertainties do not preclude the
detection of major transmission shifts as evidenced by the strong concordance observed
with case data in higher-burden areas.

Finally, assignment of COVID-19 cases to sewersheds relied on residential address
geocoding, which may misclassify infections if substantial exposure or shedding occurred
outside the residential sewershed. Future studies could enhance geolocation accuracy by
incorporating additional data sources or methods, such as improved address resolution or
alternative approaches to case-to-sewershed assignment. These limitations highlight
areas for methodological improvement but do not alter the central finding that
wastewater-based R_t estimates captured transmission trends broadly consistent with
case-based surveillance. Rather than enabling direct intervention decisions, wastewater-
derived R_t provides a complementary indicator of changing transmission dynamics that
may support situational awareness when clinical surveillance is incomplete or delayed.

The weaker correlation observed at WWTP 3 ($\rho = 0.39$) despite higher total case counts compared with WWTP 5 ($\rho = 0.84$) likely reflects differences in case variability rather than absolute burden. WWTP 3 exhibited lower day-to-day variance in case incidence during the study period, reducing the discriminatory power of rank-based correlation metrics even when directional trends were broadly consistent.

Hill et al. (2025) compared eight WBE-based R_t estimation approaches and found that methodological choices substantially influenced estimates. The present study does not seek to advance a competing method but rather to assess whether a relatively simple GLMM framework applied to routine twice-weekly sampling can yield transmission trend indicators concordant with case-based surveillance at the sewershed level.

Future research may benefit from evaluating how wastewater-based surveillance scales to larger geographic units, such as counties or clusters of wastewater treatment plants, and whether multiple facilities within a region capture consistent spatiotemporal transmission patterns. Assessing how wastewater-derived indicators can be aggregated without obscuring local dynamics would inform their use in regional surveillance frameworks. As wastewater surveillance continues to be integrated into public health practice, clarifying its role next to the traditional epidemiologic data will be essential for understanding how population-level signals can complement, rather than replace, case-based reporting systems.

Acknowledgments

We thank the Georgia Department of Public Health for providing access to sewershed-level COVID-19 case data, and the participating wastewater treatment plant operators and university laboratory partners for sample collection and analysis.

Data Availability

Code used for this analysis is available at: https://github.com/sethmund/wastewater_glmm. Wastewater and case surveillance data are available upon request from the Georgia Department of Public Health subject to data use agreements.

References

1. Silk BJ. COVID-19 Surveillance After Expiration of the Public Health Emergency Declaration—United States, May 11, 2023. Morbidity and Mortality Weekly Report. 2023;72:e1.

2. Huisman JS, et al Wastewater-Based Estimation of the Effective Reproductive Number of SARS-CoV-2. Environmental Health Perspectives. 2022;130(5):057011.

3. Zheng X, et al Wastewater Surveillance Provides Spatiotemporal SARS-CoV-2 Infection Dynamics. Engineering. 2024; 40(9) pages 70-77,

4. Safford H, et al Wastewater-Based Epidemiology for COVID-19: Handling qPCR Nondetects and Comparing Spatially Granular Wastewater and Clinical Data Trends. ACS ES&T Water. 2022;2(11):2114-24.

5. Nourbakhsh S, et al A Wastewater-Based Epidemic Model for SARS-CoV-2 with Application to Three Canadian Cities. Epidemics. 2022;39:100560.

6. Hill A, et al Comparison of methods for estimating the effective reproduction number from wastewater data. Epidemics. 2025;52:100839.

7. Ravuri S, et al Estimating effective reproduction numbers using wastewater data from multiple sewersheds for SARS-CoV-2 in California counties. Epidemics. 2025;50:100803.

8. Ungar L. Pandemic Gets Tougher to Track as COVID Testing Plunges. Associated Press. 2022.

9. Lott MEJ, et al Direct Wastewater Extraction as a Simple and Effective Method for SARS-CoV-2 Surveillance and COVID-19 Community-Level Monitoring. FEMS Microbes. 2023;4:xtad004.

10. Sablon O, et al Nanotrap® KingFisher Concentration/Extraction & MagMAX KingFisher Extraction. protocols.io. 2022.

11. Antkiewicz DS, et al Wastewater-Based Protocols for SARS-CoV-2: Insights into Virus Concentration, Extraction, and Quantitation Methods from Two Years of Public Health Surveillance. Environmental Science: Water Research & Technology. 2024;10(8):1766-84.

12. Miyazawa S, et al Wastewater-Based Reproduction Numbers and Projections of COVID-19 Cases in Three Areas in Japan, November 2021 to December 2022. Eurosurveillance. 2024;29(8):2300277.

13. Narci R, et al Inference for Partially Observed Epidemic Dynamics Guided by Kalman Filtering Techniques. Computational Statistics & Data Analysis. 2021;164:107319.

14. Daza-Torres ML, et al A Mixed-Effects Model to Predict COVID-19 Hospitalizations Using Wastewater Surveillance. Journal of Environmental Chemical Engineering. 2024;12(2):112485.

15. de Graaf M, et al Capturing the SARS-CoV-2 Infection Pyramid Within the Municipality of Rotterdam Using Longitudinal Sewage Surveillance. The Science of the Total Environment. 2023;883:163599.

16. Cori A, et al A New Framework and Software to Estimate Time-Varying Reproduction
Numbers During Epidemics. American Journal of Epidemiology. 2013;178(9):1505-
12.

17. Nash RK, et al Estimating the Epidemic Reproduction Number from Temporally
Aggregated Incidence Data: A Statistical Modelling Approach and Software Tool.
PLOS Computational Biology. 2023;19(8):e1011439.

18. Kremer C, et al Serial Intervals for SARS-CoV-2 Omicron and Delta Variants,
Belgium, November 19-December 31, 2021. Emerging Infectious Diseases.
2022;28(8):1699-1702.

19. Wannigama DL, et al COVID-19 monitoring with sparse sampling of sewered and
non-sewered wastewater in urban and rural communities. iScience.
2023;26(7):107019.

20. Rezaeitavabe F, et al Beyond Linear Regression: Modelling COVID-19 Clinical Cases
with Wastewater Surveillance of SARS-CoV-2 for the City of Athens and Ohio
University Campus. Science of The Total Environment. 2024;912:169028.

21. Deng Y, et al Ratio of Infections to COVID-19 Cases and Hospitalizations in the
United States based on SARS-CoV-2 Seroprevalence Data, September 2021-
February 2022. Open Forum Infectious Diseases. 2025;12(1):ofae719.

Appendix

508