Title: Longitudinal symptom transitions predict incident tuberculosis risk in serial active

case finding

Authors: Esther Jung[1], José Victor Bortolotto Bampi[2], Alessandra Moura da Silva[2],

Yiran E. Liu[1,3,4], Daniel Henrique Tsuha[2], Argita D. Salindri[1], Andrea da Silva Santos[5],

Roberto Dias de Oliveira[5,6], Julio Croda[2,7,8#], Jason R. Andrews[1#]

[#]JC and JRA contributed equally to this manuscript

Affiliations:

1. Stanford University School of Medicine, Stanford, California

2. Federal University of Mato Grosso do Sul, Campo Grande, Brazil

3. Yale University School of Medicine, New Haven, Connecticut

4. Yale Law School, New Haven, Connecticut

5. Federal University of Grande Dourados, Dourados, Brazil

6. State University of Mato Grosso do Sul, Dourados, Brazil

7. Oswaldo Cruz Foundation, Campo Grande, Brazil

8. Yale University School of Public Health, New Haven, Connecticut

Keywords: tuberculosis; asymptomatic; prisons; active case finding; screening

Corresponding Author:

Jason Andrews, MD, SM

Stanford University School of Medicine, 240 Pasteur Dr, Stanford, CA 94304-1049

Email: jandr@stanford.edu

Phone: 415-608-9386

GENERAL SUMMARY (40 words)

In an active case-finding program for TB, developing new symptoms was associated

with over double the TB risk, while persistent symptoms were not. Monitoring symptom

changes longitudinally, especially combined with chest radiography, may improve TB

case detection in high-risk settings.

ABSTRACT (100 words)

Tuberculosis (TB) screening often relies on cross-sectional symptom assessment. To

determine the added value of longitudinal symptom monitoring, we conducted a

prospective cohort study among 2,282 incarcerated men in Brazil. Every four months,

participants were assessed for symptoms and incident TB (Xpert or culture positive).

Compared to remaining asymptomatic, developing symptoms was associated with

higher TB risk (aRR 2.31; 95% CI 1.55-3.43), strongest among those with prior

radiographic abnormalities (aRR 2.49; 95% CI 1.56-3.97), while remaining symptomatic

was not associated (aRR 1.27; 95% CI 0.72-2.24). Longitudinal symptom monitoring

may enhance TB screening in high-risk settings, particularly in complement with

radiography.

BACKGROUND

Tuberculosis (TB) continues to pose a major global health challenge[1]. Early detection and treatment are essential to interrupt transmission and improve outcomes, particularly in high-risk populations. Efforts to identify individuals living with TB often leverage symptom screening to prioritize individuals for confirmatory molecular rapid diagnostic tests[2]. However, conventional approaches to TB screening typically only assess symptoms present at the time of screening entry, which may not capture the dynamic nature of symptom changes during TB disease progression[3,4].

Symptoms that are typically classified as features of TB[5], such as cough, fever, night sweats, and weight loss, can fluctuate over the course of disease[6]. In high-risk populations such as persons deprived of liberty (PDL), these symptoms may also be caused by other common conditions, including upper respiratory tract infections, pneumonia, chronic obstructive lung disease and tobacco smoking[7]. Moreover, recent studies reveal that over 50% of individuals living with undiagnosed TB do not report symptoms at the time of diagnosis[1,8]. Together, the non-specific presentation of TB and high prevalence of asymptomatic TB may limit the ability of cross-sectional symptom screening to reliably identify individuals at imminent risk of active TB, resulting in continued transmission[8,9].

Longitudinal monitoring of symptoms offers an opportunity to overcome some of these challenges by capturing individuals' changes in symptom status over time, including new onset, persistence, or resolution. This may provide additional predictive value for

disease, improve identification of individuals at high risk of TB, or indicate higher risk for

future evaluations, thereby facilitating the prioritization of confirmatory diagnostic testing.

Prisons harbor some of the highest TB incidence and transmission rates worldwide[10–12],

and the World Health Organization recommends systematic screening be performed

among incarcerated populations at least yearly[2]. However, active case finding is not

widely implemented in most prisons in high-burden countries, partly due to resource

constraints. Understanding how TB symptoms evolve in such settings could inform

more resource-efficient screening approaches, yet longitudinal symptom trajectories in

high-risk populations remain uncharacterized.

To address this knowledge gap, we characterized longitudinal symptom trajectories in a

cohort of incarcerated individuals who were evaluated for TB symptoms and tested for

TB every four months. We assessed the relationship between symptom transitions and

risk of incident TB.

METHODS

We performed systematic screening for TB every four months from February 2023 to

September 2025 in two all-male prisons in the Brazilian state of Mato Grosso do Sul.

After providing informed consent to participate, participants completed a standardized

symptom questionnaire, administered by study nurses, to ascertain the presence of

cough, fever, night sweats, weight loss, chest pain, or dyspnea. Presence of each

symptom at each visit was assessed through yes/no questions (i.e. "Do you have a

cough?"). An individual was considered symptomatic if they confirmed the presence of

any of these symptoms.

At baseline, defined as the first visit in which a participant was enrolled in the study, all

consenting participants had a posterior-anterior chest radiograph performed and two

sputum samples collected for testing by culture and Xpert® MTB/RIF Ultra (hereinafter

"Xpert"). Chest radiographs were evaluated using automated detection software LunitTB

(South Korea; version v3.1.5.1), which outputs a numerical TB risk score on a

continuous scale between 0 and 100.

At each four-month follow-up visit, procedures were repeated: symptom assessment, a

single sputum sample collection, and chest radiography. Symptom status transitions

over sequential four-month intervals were categorized as asymptomatic-to-symptomatic,

symptomatic-to-asymptomatic, persistently symptomatic, or persistently asymptomatic.

Incident TB was defined by at least one positive sputum culture or Xpert result. Trace

Xpert results were repeated with a new sample. Those that persisted as traces were

considered negative for TB and all trace samples were cultured.

Eligibility was restricted to those with negative baseline Xpert and culture results who

did not initiate TB treatment, regardless of radiographic findings, to ensure participants

did not have prevalent TB. However, during follow-up, a small number of

microbiologically negative participants initiated TB treatment based on clinical

judgement. Only participants who completed at least one follow-up visit during the study

period were included, to permit longitudinal assessment of symptom status and incident

TB.

Given the frequent transfers and movement within the prison system, nearly 25% of

individuals enrolled in the cohort missed follow-up visits intermittently. We excluded

individuals with non-consecutive symptom information, as gaps in reporting precluded

reliable classification of symptom transitions across sequential four-month intervals into

the predefined symptom trajectory categories.

We used the chi-square and Kruskal-Wallis tests to compare baseline demographic and

clinical characteristics across symptom transition groups in the first four-month interval.

In supplementary analyses, we evaluated all four-month symptom transitions across the

study period to identify predictors associated with developing or resolving symptoms.

We performed modified Poisson regression with robust standard errors clustered at the

individual level to estimate adjusted risk ratios (aRRs)[13]. Models adjusted for the

following baseline covariates: age, prison, prior TB, exposure to symptomatic cellmates

(yes/no), TB contact (ever/never), current smoking status, any drug use, duration of

incarceration, and binary LunitTB x-ray score using a threshold of 50 as a proxy for lung

abnormalities. Separate models were constructed for individuals who were

asymptomatic at the prior visit (assessing the risk of becoming symptomatic) and for

those who were symptomatic at the prior visit (assessing the risk of becoming

asymptomatic).

Specific symptom-level transition patterns were summarized descriptively by calculating

the proportion of each individual TB-related symptom reported at visits where a

transition occurred, distinguishing between new onset, resolution, and persistent

symptoms.

We also characterized how long participants remained in their initial symptom state

during follow-up using survival analysis. For symptomatic individuals, the event was the

first resolution of symptoms, while for asymptomatic individuals, the event was the first

acquisition of symptoms. Participants were censored at TB diagnosis, loss to follow-up,

or last study visit.

For the primary analysis, we assessed whether symptom transitions during the

preceding four-month interval were associated with incident TB diagnosed at the

subsequent visit, using the same modified Poisson framework described above but

adjusting for radiographic status (Lunit score <50 vs. ≥50) at the start of each interval

rather than at baseline. Results were also stratified by radiographic status. Persistently

asymptomatic individuals served as the reference category. All analyses were

conducted in R (version 4.4.1).

Ethical approval was obtained from the Research Ethics Committee of the Federal

University of Mato Grosso do Sul, the Brazilian National Commission for Ethics in

Research, and the Stanford University Research Ethics Committee.

RESULTS

After applying inclusion criteria, our study identified 2,282 participants without TB at

baseline who were assessed for symptoms at two or more consecutive visits between

February 2023 and September 2025. The median baseline age of participants was 33

years (IQR 28-40), over 80% of participants were Black or mixed race (1,854/2,282),

and most participants did not complete high school (83%) (Table 1).

The majority of individuals reported no TB symptoms at baseline (1,747/2,282, 76.5%)

and 62.1% (1,417/2,282) remained asymptomatic at the first follow-up. Notably, 52%

(26/50) of incident TB cases occurred among participants who remained asymptomatic.

Among those asymptomatic at baseline, 14.5% developed symptoms by the first follow-

up. Of the 23.4% symptomatic at baseline, 40% remained symptomatic and 60%

reported resolution (Table 1). Several characteristics differed significantly across

transition groups, with individuals with persistent symptoms showing the highest

prevalence of these factors: previous TB, exposure to symptomatic cellmates, current

smoking, and any drug use (Table 1).

Of 2,282 total participants, 1,725 (75.6%) were assessed for at least 3 visits, 1,105

(48.4%) for at least 5, and 578 (25.3%) completed all 7 possible assessments. Among

participants symptomatic at baseline (n = 535), the probability of remaining symptomatic

declined steeply to 40.2% at the first follow-up, 19.3% at the second, and 7.6% by the

last visit. In contrast, symptom absence was more stable: among those asymptomatic at

baseline (n = 1,749), 81.1% remained asymptomatic at the first follow-up and 43.8%

were still asymptomatic at the last visit (Supplementary Figure 1).

Symptom-level transitions revealed that cough (with or without sputum production) was

the most frequently reported symptom among those developing symptoms (66%) and

among those resolving symptoms (68%). Other common symptoms associated with

symptom changes included chest pain, breathing difficulty, weight loss, and fever

(Supplementary Figure 2).

When evaluating predictors for symptom transitions longitudinally among previously

asymptomatic individuals, the onset of any TB symptom within the subsequent four-

month interval was associated with a previous TB diagnosis (aRR 1.26; 95% CI 1.08-

1.47; p=0.004), current smoking (aRR 1.81; 95% CI 1.53-2.15; p<0.001), past smoking

(aRR 1.34; 95% CI 1.09-1.66; p=0.005), any drug use (aRR 1.23; 95% CI 1.08-1.41;

p=0.002), and exposure to symptomatic cellmates (aRR 1.30; 95% CI 1.15-1.47;

p<0.001). Among previously symptomatic individuals, resolution of symptoms was less

likely among those who reported current smoking (aRR 0.82; 95% CI 0.72-0.92;

p=0.001) or any drug use (aRR 0.88; 95% CI 0.79-0.97; p=0.010) (Supplementary

Table 1). In interaction models, these predictors did not significantly modify the

association between symptom transitions and TB risk.

Longitudinal analyses showed that compared to persistently asymptomatic individuals,

transitioning from an asymptomatic to a symptomatic state within a four-month interval

was significantly associated with incident TB (aRR 2.31; 95% CI 1.55-3.43). In contrast,

remaining symptomatic (aRR 1.27; 95% CI 0.72-2.24) or resolution from symptomatic to

asymptomatic (aRR 1.13; 95% CI 0.68-1.90) were not significantly associated with TB

risk, compared to those who stayed persistently asymptomatic (Figure 1).

Stratified analyses revealed that this overall elevated risk among those developing

symptoms was primarily driven by individuals with abnormal chest radiographs (aRR

2.49; 95% CI 1.56-3.97), though a similar trend was observed among those with normal

x-rays (aRR 1.97; 95% CI 0.94-4.14). These findings indicate that symptom emergence

was associated with elevated TB risk, with the strongest effects observed among those

with radiographic abnormalities.

DISCUSSION

In our longitudinal cohort study in two high-incidence prisons, we found that TB-related

symptoms followed dynamic trajectories over four-month intervals. Between enrollment

and the first follow-up visit just four months later, nearly one in seven (14.5%)

developed new symptoms, and 60% of those symptomatic at baseline reported

resolution, underscoring the fluctuating nature of TB-related symptom reporting in this

population.

Importantly, more than half of all incident TB cases (52%, 26/50) occurred among

participants who remained asymptomatic during the first interval, highlighting the

substantial limitations of symptom-based screening in this setting. Prior studies in

prisons and other high-risk congregate settings have similarly documented that reliance

on cross-sectional symptom screening alone misses a considerable fraction of active

TB cases, supporting the need for repeated active case-finding[4,14]. Our intensive

screening program combining sputum testing and radiography every four months

successfully detected microbiologically confirmed asymptomatic cases. Although our

study was not designed to assess the additional yield of repeated symptom

assessments, longitudinal monitoring may provide value in understanding disease

progression, particularly when combined with complementary strategies such as routine

sputum testing or radiographic evaluation.

Specifically, our analyses demonstrate that participants who developed new symptoms

had more than double the risk of TB diagnosis compared to those who remained

asymptomatic, while persistent symptoms conferred no excess risk. This pattern

suggests that symptom emergence, rather than chronic symptom presence, offers

greater resolution for identifying short-term risk. This may reflect the high sensitivity of

our combined diagnostic strategy, which likely detected new TB cases rapidly, leaving

few recent-onset cases undiagnosed. In contrast, participants with persistent symptoms

may have alternative or unresolved etiologies, such as concomitant respiratory

infections, which could obscure the relationship with incident TB. The overall

association has practical implications for resource-limited settings where radiography

may be unavailable, while stratified analyses show that symptom transitions most

strongly predict TB risk among those with radiographic abnormalities, enabling more

precise targeting of diagnostic resources.

We also identified key factors associated with four-month symptom transitions. Among

previously asymptomatic individuals, previous TB, current and past smoking, and

exposure to symptomatic cellmates increased the risk of developing symptoms. In

contrast, current smoking and drug use were associated with persistent symptoms.

While these predictors shaped symptom trajectories, they did not significantly modify

the primary association between transitions and TB risk in interaction models.

Nonetheless, combining repeated symptom assessments with knowledge of exposure-

related risk factors may allow public health programs to more precisely target diagnostic

interventions toward those at highest short-term TB risk.

Our study had several limitations. Symptom reporting relied entirely on self-report,

which may be influenced by recall bias and subjective differences in symptom

perception. Individuals who typically feel well may be more attuned to new symptoms,

while those with chronic symptoms may have reduced sensitivity to incremental

changes. However, our longitudinal assessment of symptom changes uses each

individual as their own reference, potentially reducing the contextual and interviewer

variability inherent in cross-sectional symptom reporting[15]. Additionally, the four-month

interval between screenings may not have captured shorter-term or transient symptom

fluctuations. Excluding participants with intermittently missing visits may have

introduced bias, as those excluded were more likely to report drug use and lower

education, and may have had less stable follow-up due to transfer or refusal. We also

lacked reliable data on nutrition and longitudinal smoking behavior, precluding

assessment of these important TB risk factors. Despite these limitations, our findings

offer important insights into the temporal dynamics of TB symptoms and their potential

utility in prioritizing individuals at highest risk for diagnostic screening and treatment.

In conclusion, longitudinal assessment of TB-related symptoms reveals dynamic

patterns of emergence and resolution that identify individuals at elevated short-term risk

of active disease. Algorithms prioritizing individuals with either new-onset symptoms or

radiographic abnormalities for molecular testing may improve sensitivity and cost-

effectiveness in high-transmission, resource-limited settings. A recent study found that

symptom screening, while less sensitive than annual radiography, can achieve

comparable impact when applied frequently and at lower cost[16]. Our findings suggest

that monitoring symptom transitions may enhance this existing approach. Future studies

should assess how integrating longitudinal symptom monitoring, radiography, and rapid

diagnostics can be optimized to reduce transmission across high-burden settings.

Acknowledgements EJ, JRA, and JC conceived of the study. JRA, JC, and RDO

developed the study plans. AmdS and AdSS enrolled participants and collected data.

DHT performed data management. EJ, JVBB, YL, ADS, and JRA designed and

performed the data analysis. EJ and JVBB wrote the manuscript. All authors contributed

to revising the manuscript.

The authors acknowledge the use of ChatGPT (GPT-4, OpenAI) for troubleshooting

coding errors and minor grammar checking. AI was not used for generation of

conclusions or interpretation of results. All content and analyses were independently

verified by the authors, who take full responsibility for the manuscript.

Funding This work was supported by Brazil's Ministry of Health; Brazil's National

Council for Scientific and Technological Development [CNPq; 404182/2019-4 to JC] and

the U.S. National Institutes of Health [R01AI130058, K24AI182647 to JRA].

Conflicts of interest The authors do not declare any conflicts of interests in preparing

this article.

Data availability Analytic code is available at https://github.com/Andrews-Lab-

Stanford/LongitudinalTBSymptoms_BrazilPrisons.git

Table 1. Characteristics of participants by symptom transition between baseline and

first follow-up visit (N = 2,282)

Characteristics	Overall (N = 2,282)	First 4-month interval symptom transition				p-value*
		Persistently asymptomatic (n = 1,417)	Persistently symptomatic (n = 215)	Asymptomatic to Symptomatic (n = 330)	Symptomatic to Asymptomatic (n = 320)	
Median age (IQR)	33 (28, 40)	33 (28, 40)	33 (28, 40)	32 (27, 39)	33 (27, 40)	0.208
Incident TB	50 (2.2)	26 (1.8)	4 (1.9)	15 (4.5)	5 (1.6)	0.018
Ethnicity						0.204
White	419 (18.4)	251 (17.7)	40 (18.6)	57 (17.3)	71 (22.2)	
Black	217 (9.5)	127 (8.9)	27 (12.6)	33 (10.0)	30 (9.4)	
Mixed	1,637 (71.7)	1,034 (73.0)	145 (67.4)	239 (72.4)	219 (68.4)	
Asian	8 (0.4)	4 (0.3)	3 (1.4)	1 (0.3)	0 (0)	
Indigenous	1 (<0.1)	1 (<0.1)	0 (0)	0 (0)	0 (0)	
Education						0.562
Less than high school	1,893 (83.0)	1,175 (82.9)	181 (84.2)	279 (84.5)	258 (80.6)	
Completed high school	389 (17.0)	242 (17.1)	34 (15.8)	51 (15.5)	62 (19.4)	
Prison						0.256
EPJFC	1,745 (76.5)	1,089 (76.9)	165 (76.7)	239 (72.4)	252 (78.8)	
IPCG	537 (23.5)	328 (23.1)	50 (23.3)	91 (27.8)	68 (21.3)	
Previous TB	409 (17.9)	225 (15.9)	54 (25.1)	67 (20.3)	63 (19.7)	0.004
TB contact	1,335 (58.5)	818 (57.7)	135 (62.8)	204 (61.8)	178 (55.6)	0.239
Symptomatic cellmates	854 (37.4)	415 (29.3)	132 (61.4)	126 (38.2)	181 (56.6)	<0.001
Current smoking	1,342 (58.8)	741 (52.3)	165 (76.7)	220 (66.7)	216 (67.5)	<0.001
Any drug use	1,416 (62.1)	829 (58.5)	150 (69.8)	226 (68.5)	211 (65.9)	<0.001
LunitTB Score at Baseline						0.961
<50	1,584 (69.4)	984 (69.4)	146 (67.9)	230 (69.7)	224 (70.0)	
≥50	698 (30.6)	433 (30.6)	69 (32.1)	100 (30.3)	96 (30.0)	
Months incarcerated, median (IQR)	12 (2, 44)	13 (2, 41)	14 (2, 71)	11 (2, 34)	9 (1, 61)	0.213

No (% of column n) unless otherwise specified.

*Pearson's Chi-squared test; Kruskal-Wallis test

Figure 1. Adjusted risk of incident TB by symptom transitions over four-month intervals, overall

and stratified by radiographic status

N represents the total number of four-month symptom transition observations across all follow-up intervals during the study period. Adjusted risk ratios

(aRRs) represent adjusted risk of TB diagnosed by sputum culture or Xpert within the same four-month interval as the symptom transition. Models adjust

for baseline characteristics including age, prison, prior TB, smoking, drug use, duration of incarceration, symptomatic cellmates, and TB contact. The

overall model adjusts for Lunit x-ray score category at the first visit of each interval. The stratified models adjust for the baseline characteristics and are

separated by Lunit x-ray score category at the first visit of each interval.

Alt text: Forest plot showing adjusted risk of incident tuberculosis by symptom transitions over four-month intervals. Compared with participants who

remained asymptomatic, those who became symptomatic had over twice the risk of TB. Risks for persistent or resolving symptoms were similar to the

reference group.

REFERENCES

1. World Health Organization. Global Tuberculosis Report 2024.
   https://www.who.int/teams/global-programme-on-tuberculosis-and-lung-health/tb-
   reports/global-tuberculosis-report-2024 (2024).

2. World Health Organization. *WHO Consolidated Guidelines on Tuberculosis. Module
   2: Screening - Systematic Screening for Tuberculosis Disease.* (2021).

3. Kendall, E. A., Shrestha, S. & Dowdy, D. W. The Epidemiological Importance of
   Subclinical Tuberculosis. A Critical Reappraisal. *Am. J. Respir. Crit. Care Med.* **203**,
   168–174 (2021).

4. Yoon, C., Dowdy, D. W., Esmail, H., MacPherson, P. & Schumacher, S. G.
   Screening for tuberculosis: time to move beyond symptoms. *Lancet Respir. Med.* **7**,
   202–204 (2019).

5. CDC. Clinical Symptoms of Tuberculosis. *Tuberculosis (TB)*
   https://www.cdc.gov/tb/hcp/clinical-signs-and-symptoms/index.html (2024).

6. Richards, A. S. *et al.* Quantifying progression and regression across the spectrum of
   pulmonary tuberculosis: a data synthesis study. *Lancet Glob. Health* **11**, e684–e692
   (2023).

7. Viglianti, E. M., Iwashyna, T. J. & Winkelman, T. N. A. Mass Incarceration and
   Pulmonary Health: Guidance for Clinicians. *Ann. Am. Thorac. Soc.* **15**, 409–412
   (2018).

8. Emery, J. C. *et al.* Estimating the contribution of subclinical tuberculosis disease to
   transmission: An individual patient data analysis from prevalence surveys. *eLife* **12**,
   e82469 (2023).

9. Bezerra, W. da S. P. et al. Role of tuberculosis symptoms in transmission risk to cell

contacts in prisons. *BMJ Public Health* **3**, (2025).

10. Martinez, L. *et al.* Global, regional, and national estimates of tuberculosis incidence

and case detection among incarcerated individuals from 2000 to 2019: a systematic

analysis. *Lancet Public Health* **8**, e511–e519 (2023).

11. Walter, K. S. *et al.* The escalating tuberculosis crisis in central and South American

prisons. *Lancet Lond. Engl.* **397**, 1591–1596 (2021).

12. Cords, O. *et al.* Incidence and prevalence of tuberculosis in incarcerated

populations: a systematic review and meta-analysis. *Lancet Public Health* **6**, e300–

e308 (2021).

13. Zou, G. A Modified Poisson Regression Approach to Prospective Studies with

Binary Data. *Am. J. Epidemiol.* **159**, 702–706 (2004).

14. Bailey, S. L. *et al.* Missed opportunities for tuberculosis diagnosis. *Int. J. Tuberc.*

Lung Dis. Off. J. Int. Union Tuberc. Lung Dis. **15**, 205–i (2011).

15. McCreesh, N. et al. Reported Tuberculosis Symptoms: An Inadequate Classifier of

Disease State. *Clin. Infect. Dis.* ciaf611 (2025).

16. Liu, Y. E. et al. Comparative impacts and cost-effectiveness of tuberculosis active

case-finding strategies in prisons in Brazil, Colombia, and Peru: a mathematical

modeling study. *medRxiv* 2025.09.25.25336625 (2025).

APPENDIX

Supplementary Figure 1. Symptom persistence by baseline symptom status

Risk tables with raw numbers of participants at each visit are shown beneath the curves.

Alt text: Kaplan-Meier curves showing symptom persistence by baseline symptom status. Participants starting asymptomatic had a gradual decline in

remaining asymptomatic, while those starting symptomatic showed rapid symptom resolution over follow-up visits.

Supplementary Figure 2. Specific symptom changes by symptom transition group

% of total instances of symptom transition in that symptom category. Symptom changes refer to new onset (Asymptomatic to Symptomatic group), resolution

(Symptomatic to Asymptomatic), or either (Symptomatic to Symptomatic).

Alt text: Heatmap showing percentage changes in specific symptoms by symptom transition group. Color intensity reflects percent change, highest

among transitions involving cough-related symptoms.

Supplementary Figure 3. Symptom transitions among participants with at least three symptom

assessments (N = 1725)

Alt text: A Sankey diagram visualizing symptom state transitions for 1,725 participants across three timepoints: Baseline, Visit 1, and Visit 2. The chart

illustrates frequent bidirectional movement between asymptomatic and symptomatic states, highlighting the dynamic nature of symptoms over time.

Supplementary Table 1. Baseline predictors of symptom change stratified by prior symptom

status

Previously asymptomatic

Baseline predictors	Becoming symptomatic		
	aRR*	95% CI	p-value
Previous TB	1.26	(1.08-1.47)	0.004
Current smoking	1.81	(1.53-2.15)	<0.001
Past smoking	1.34	(1.09-1.66)	0.005
Any drug use	1.23	(1.08-1.41)	0.002
Symptomatic cellmates	1.30	(1.15-1.47)	<0.001
TB contact	1.03	(0.91-1.17)	0.662
LunitTB ≥50	0.90	(0.78-1.04)	0.164

393 Previously symptomatic

Baseline predictors	Becoming asymptomatic		
	aRR*	95% CI	p-value
Previous TB	0.94	(0.83-1.06)	0.340
Current smoking	0.82	(0.72-0.92)	0.001
Past smoking	0.96	(0.83-1.10)	0.563
Any drug use	0.88	(0.79-0.97)	0.010
Symptomatic cellmates	0.92	(0.84-1.00)	0.057
TB contact	0.92	(0.84-1.01)	0.077
LunitTB ≥50	0.98	(0.88-1.10)	0.784

394 *Adjusted risk ratios (aRRs) represent risk of symptom status change from prior status compared to no change in symptom status. Multivariate logistic regression

395 models with cluster-robust standard errors at the individual level to estimate odds of symptom transitions, adjusting for baseline characteristics including age,

396 prison, prior TB, smoking, drug use, prison time, symptomatic cellmates, TB contact, and categorical LunitTB x-ray score.