Atmospheric pattern correlates of human brain structure

Max Korbmacher,[1,2,3,+] Ole A. Andreassen,[4,5] Lars T. Westlye,[4-6] Ivan I. Maximov[2,*], Ivan Kuznetsov[7,*]

+ Corresponding author.

* Shared senior contribution.

Affiliations

[1]Neuro-SysMed, Department of Neurology, Haukeland University Hospital, Norway

[2]Department of Health and Functioning, Western Norway University of Applied Sciences, Norway

[3]Mohn Medical Imaging and Visualisation centre, Department of Radiology, Haukeland University Hospital, Norway

[4]Center for Precision Psychiatry, University of Oslo and Oslo University Hospital, Norway

[5]K.G. Jebsen Centre for Neurodevelopmental disorders, University of Oslo, Norway

[6]Department of Psychology University of Oslo, Norway

[7]Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany.

Abstract

Climate change increasingly impacts human health, yet its neurobiological effects remain poorly understood. Analysing structural neuroimaging data of 30,831 UK participants (4,294 with follow-up assessments), we show that ambient weather conditions (warm, sunny, low precipitation and wind speed) associate with brain structure variations that exceed contributions from Alzheimer's disease genetic risk scores and self-reported mental health. These findings establish atmospheroc patterns as measurable environmental correlates of brain structure and reveal new pathways for understanding climate-brain interactions.

Main Text

Despite growing concerns about climate change impacts on human health and well-being,[1,2]

evidence linking atmospheric conditions to brain structure remains limited. This knowledge

gap persists partly due to the challenges of integrating large-scale meteorological data with

neuroimaging cohorts. Atmospheric variability (temperature, precipitation, solar radiation,

wind) may influence brain structure through multiple pathways, including effects on

multiple bodily systems,[2] physical activity,[3] vitamin D synthesis,[4] circadian rhythms.[5] Given

projections of increased climatic variability and extreme weather events,[6] understanding

neurobiological sensitivity to atmospheric exposures could inform preventive strategies and

help anticipate neurological health burdens associated with climate change.[1] Hence, an

improved understanding of associations between atmospheric patterns and brain structure

is essential as environmental conditions continue to shift both globally and locally. Here, we

provide a first step towards investigating atmospheric-brain associations, establishing brain

structural correlates of meteorological conditions in the United Kingdom.

Fig. 1. Study design. Monthly meteorological data were linked to structural brain imaging (N = 30,831),
yielding measures of grey matter volume, white matter volume, and white matter microstructure (diffusion
metrics). Genetic risk scores and self-reported mental health data served as comparison variables.

We linked atmospheric data averaged over the United Kingdom for the month preceding

the health and MRI assessments (Fig. 1). The derived averaged atmospheric indices were

joined with individuals' brain metrics for all 30,831 participants at baseline (see Table 1 in

the Online Methods for an overview of key variables), and for n=4,294 at follow-up

(Supplemental Table 1). As a comparison to atmospheric variables as predictors of brain

structure, we also used self-reports of mental health, correspondingly polygenic risk scores

(PGRS) of Alzheimer's Disease, considering the sample age.

Our analyses, all corrected for assessment site, household income, sex, age, intracranial

volume, and surface holes, revealed that atmospheric variables presented stronger

associations (median±mean absolute deviation (MAD) $|\beta|$ = 0.013±0.013) with brain

structure than PGRS of Alzheimer's disease (AD, median±MAD $|\beta|$ = 0.006±0.001), and on

average similar to self-reports of neuroticism and depression ($|\beta|$ = 0.013±0.012; see Fig. 2;

Supplemental Data 1). However, the strongest associations were found for atmospheric

variables for associations between fractional anisotropy and a) 10 metre wind speed

(standardized effect size β=-0.059, 95% CI [-0.068, -0.049], p_{FDR}=1.78*10^{-30}), and surface

downward UV radiation (β=0.055, 95% CI [0.045, 0.064], p_{FDR}=4.42*10^{-27}).

Fig. 2. Associations between brain structural and a) atmospheric indicators, b) polygenic risk of Alzheimer's disease, and c) self-reports of recent depression and neuroticism.
PGRS = polygenic risk score, CGMV = cortical grey matter volume, DTI = diffusion tensor imaging, AD = axial diffusivity, FA = fractional anisotropy, MD = mean diffusivity, RD = radial diffusivity, WMV = white matter volume. Std. Beta = standardized beta or regression coefficients.

The associations suggest that better weather (lower precipitation, lower wind speed, higher

short- and long-wave radiation and higher surface pressure) was associated with greater

cortical volume and higher fractional anisotropy (indicating anisotropy of water diffusion

along a single direction and thereby structural integrity of axonal bundles), in addition to

lower axial, radial and mean diffusivity (indicating the magnitude of diffusivity, along or

perpendicular to the fibre bundles or their average) (Fig. 2).

Likelihood Ratio Tests suggested that adding AD PGRS or self-reported outcomes on recent

depression and neuroticism did not improve models predicting brain structure. Changes to

the model were non-significant when adding the covariates ($p>0.05$) in 65 of 90 (27.78%) of

the models with FDR-adjustment for multiple comparisons. Adding neuroticism to the

models explaining cortical volume presented the largest contribution ($26.12<F<27.16$,

$p<0.05$), followed by neuroticism ($11.29<F<11.89$, $p<0.05$; see Supplemental Data 2).

However, while statistically significant the added variance of AD PGRS and self-reported

outcomes was low ($\Delta R^2 < 0.03\%$).

Note that the pattern of associations was also found over time, using mixed linear models

considering both timepoints, in the n=4,294 participants with follow-up scans (Supplemental

Figure 1, Supplemental Data 3). Here, the effects of total precipitation were strongest,

associating negatively with cortical volume ($\beta=-0.051$, 95% CI [-0.053, -0.048], $p_{FDR}<2.23*10^{-308}$)

and fractional anisotropy ($\beta=-0.048$ [-0.051, -0.046], $p_{FDR}<2.23*10^{-308}$), and positively

associating with mean diffusivity ($\beta=0.043$ [0.039, 0.047], $p_{FDR}=2.81*10^{-108}$) and radial

diffusivity ($\beta=0.043$ [0.039, 0.046], $p_{FDR}=4.93*10^{-140}$).

Finally, considering that the strongest cross-sectional and longitudinal associations between

atmospheric conditions and whole-brain variables were found for precipitation, surface

pressure and wind speed (compare Fig.2, Supplemental Figure 1), we assessed associations

of these variables with tract- and region-level brain metrics (Supplemental Figures 2-4). For

both associations, white matter microstructure was more sensitive to weather phenomena.

Yet also the volume of the amygdala and ventricles were among the strongest correlates of

total precipitation (Supplemental Figure 2). Across associations between tracts and regional

brain characteristics, atmospheric conditions, genetic, and behavioural scores, diffusivity in

the inferior longitudinal fasciculus presented the strongest associations ($\beta < 0.08$) with

multiple atmospheric variables (for a full overview across associations see Supplemental

Data 4).

Atmospheric patterns were associated with markers of brain structure, with effect sizes

generally exceeding those of AD PGRS, as well as mental health self-reports. Ambient

atmospheric patterns, defined by lower precipitation and wind speed and higher

temperature, solar radiation and atmospheric pressure, corresponded to higher cortical

volume, greater fractional anisotropy, and lower white matter diffusivity at baseline and

over time. These findings demonstrate a measurable link between meteorological

conditions and brain structure.

Several biological mechanisms may underlie the examined weather-brain associations. Solar

radiation exposure, positively associated with structural integrity, could act through vitamin

D synthesis or circadian regulation affecting neuroplasticity and cellular maintenance.

Atmospheric pressure and precipitation may influence brain structure indirectly via

behavioural pathways, as favourable weather supports outdoor activity, social engagement,

and exposure to natural environments,[7,8] all being factors linked to general and brain

health.[9,10] High-pressure systems are also associated with low dispersion and hence air

pollution,[11] which is in turn associated with neurotoxic burden,[12] resulting in increased AD

risk.[13,14]

Confounding effects from third variables and limitations to measurements must also be

considered. Despite adjustment for multiple covariates, atmosheric patterns and brain

structure covary with unmeasured socioeconomic and lifestyle factors. Differences in

urbanicity, healthcare access, diet, occupation, and cultural practices can all influence brain

structure as well as direct and indirect exposure to atmospheric conditions. Individuals with

greater resources may mitigated adverse weather through housing, transport, or relocation.

The observed associations may therefore partly reflect interactions between meteorological

and social determinants that were not captured by our models. Moreover, unmeasured

confounders (beyond scanner site, household income, sex, age, ICV, and surface holes) may

also contribute to both weather exposure and brain outcomes. Weather data averaged

across the United Kingdom mask local microclimatic variation and individual exposure

differences. Time spent outdoors, indoor conditions, vitamin D supplementation, circadian

rhythm, and air pollution exposure were not captured. The one-month averaging window,

while pragmatic, may not reflect the most relevant temporal scale for structural effects. The

cohort's geographic and demographic composition further limits generalisability. Namely,

the UK Biobank imaging sample is composed of middle-aged and older adults in relatively

good health, not representative of the general UK population.

Non-linearity and boundary effects may also exist, potentially limiting the generalisability of

this study. Extreme weather or long-term cumulative exposure may have distinct impacts

from the monthly means analysed here. Seasonal cycles could exert neurobiological

influences distinct from stochastic weather variation. However, a previous multi-scanning

case study suggests that brain structure estimates are relatively robust to seasonal effects.[15]

Nevertheless, individual differences in weather sensitivity, shaped by genetic,

developmental, or health factors, were not examined but may modulate these relationships.

Our findings have implications for neuroscience and public health. Longitudinal designs are

needed to establish temporal precedence, assess reversibility, and quantify dose-response

relationships. Such studies could clarify whether within-person variation in weather

exposure predicts brain structural change and identify sensitive exposure windows. Future

research should also examine regional specificity, vulnerable populations, and potential

modifiers such as outdoor time, activity levels, social engagement, and baseline health.

Integrating wearable sensor data would improve exposure assessment beyond aggregated

meteorological indices.

If causal relationships are confirmed, the findings could inform interventions. Urban

planning and building design might incorporate strategies to mitigate negative weather

effects on neurological health. Individual-level interventions, including light therapy, vitamin

D supplementation, or structured activity during unfavourable weather, warrant

investigation. As global climate change alters weather variability and extremes,

understanding the neurobiological systems sensitive to meteorological variation becomes

critical for anticipating and mitigating future health burdens.

In summary, recent ambient weather patterns are associated with human brain structure,

with effect sizes exceeding those of frequently applied PGRS for common mental and brain

disorders and self-reported mental health measures. These data provide an empirical

foundation for environmental neuroscience linking climate variability to brain biology. While

causal mechanisms remain to be established, the results suggest that meteorological

conditions are neurobiologically relevant and should be integrated into models of brain

health as global environmental conditions continue to shift.

Online Methods

Sample

We extracted multimodal (T_1-weighted and diffusion-weighted images) MRI data from the

UK Biobank database.[16] We excluded participants diagnosed with any mental and

behavioural disorder (ICD-10 category F), disease of the nervous system (ICD-10 category G),

and disease of the circulatory system (ICD-10 category I), or stroke. Weather data, PGRS and

self-reported data were merged, resulting in N = 30,831 included participants (sample

overview at baseline: Table 1; at follow-up: Supplemental Table 1).

Table 1. Sample characteristics at baseline.

Characteristic	Cheadle, N = 18,018[1]	Newcastle, N = 8,260[1]	Reading, N = 4,553[1]
Demographics			
Age (years)	65 (58, 70)	66 (60, 72)	68 (61, 73)
Male (sex)	8,752 (49%)	3,888 (47%)	2,204 (48%)
Income in £1,000			
Less than 18	1,930 (11%)	905 (11%)	269 (5.9%)
18-31	3,889 (22%)	1,667 (20%)	710 (16%)
31-52	5,089 (28%)	2,308 (28%)	1,072 (24%)
52-100	4,512 (25%)	2,152 (26%)	1,425 (31%)
More than 100	928 (5.2%)	563 (6.8%)	702 (15%)
Don't know	336 (1.9%)	145 (1.8%)	92 (2.0%)
No answer	1,192 (6.7%)	502 (6.1%)	276 (6.1%)
Self-reported measures			

Recent Depression Score (range: 4-16)	5 (4, 6)	5 (4, 5)	5 (4, 5)
Neuroticism (range: 0-12)	3 (1, 5)	3 (1, 5)	2 (1, 5)
Brain-derived measures			
Intracranial Volume (litres)	1.5 (1,4, 1.6)	1.5 (1.4, 1.6)	1.5 (1.4, 1.6)
Surface Holes	53 (38, 75)	50 (36, 74)	44 (33, 62)
Fractional Anisotropy	0.46 (0.45, 0.47)	0.46 (0.44, 0.47)	0.46 (0.45, 0.47)
Mean Diffusivity	0.89 (0.87, 0.91)	0.89 (0.87, 0.91)	0.89 (0.87, 0.91)
Radial Diffusivity	0.65 (0.62, 0.67)	0.65 (0.62, 0.67)	0.64 (0.62, 0.67)
Axial Diffusivity	1.38 (1.36, 1.40)	1.37 (1.35, 1.39)	1.38 (1.36, 1.40)
Cortical Volume (litres)	0.4 (0.4, 0.5)	0.4 (0.4, 0.5)	0.4 (0.4, 0.5)
White Matter Volume (litres)	0.4 (0.4, 0.5)	0.4 (0.4, 0.5)	0.4 (0.4, 0.5)
Polygenic Risk Score (Z-scores)			
Alzheimer's Disease	-0.08 (-0.70, 0.63)	-0.04 (-0.69, 0.67)	-0.06 (-0.71, 0.63)

[1] Median (IQR); n (%)

MRI Acquisition and Processing

The MRI acquisition protocol has been described previously

(https://www.fmrib.ox.ac.uk/ukbiobank/protocol/).[16] Starting with the diffusion MRI data,

we processed these imaging data using an optimised pipeline,[17] which entails corrections of

noise, Gibbs ringing, susceptibility-induced and motion distortions, and eddy current

induced artifacts. Isotropic 1 mm^3 Gaussian smoothing was applied using fslmaths (FSL

version 6.0.1).[18] We estimated diffusion tensors at each voxel, or Diffusion Tensor Imaging

(DTI),[19] but used a signal decomposition in order to take into account kurtosis imaging as

well. This procedure has been shown to lead to more robust and reproducible DTI

estimates.[20] We employed Tract-based Spatial Statistics[21] for the analysis of white matter

integrity. The first step was to align the fractional anisotropy (FA) images to standard MNI

space using non-linear registration. A mean FA image and corresponding skeleton were then

generated from the aligned data. Each diffusion parameter map was subsequently projected

onto this mean FA skeleton. We then averaged the respective DTI metrics FA, axial, radial

and mean diffusivity across the white matter skeleton. We used the YTTRIUM method[22] for

quality control of diffusion MRI scalars. For YTTRIUM global diffusion MRI scalar metrics are

converted into 2-dimensional format using a structural similarity extension of each scalar
map to their mean image to create a 2D distribution of image and diffusion parameters.
Non-clustering values are then excluded.

For T_1-weighted MRI data, we used FreeSurfer version 5.3.0 for surface-based
reconstruction and estimation of brain grey and white matter and intra-cranial volumes, in
addition surface holes, which have previously been shown to be crucial to avoid bias in large
sample brain image analyses.[23] We used Euler numbers[24] to exclude images when the Euler
number exceeded three standard deviations from the mean.

Finally, to allow for more spatially specific assessment of brain structure, we parcellated
both grey and white matter. For the T_1-weighted MRI-derived volumes, we used the
Desikan-Killiany Atlas[25] to obtain regional estimates of brain volumes, leading to a total of
68 brain features based on the 34 regions of interest for each brain hemisphere. We used
the 20 tracts from the John Hopkins University atlas[26] which are based on a probabilistic
WM atlas. We averaged across the 20 tracts for each of the 4 DTI parameters, totalling 80
values per individual.

Polygenic risk
We estimated the polygenic risk score (PGRS) for each participant with available genomic
data, using LDPred2[27] with default settings. As input for the PGRS, we used the summary
statistics from a recent genome-wide association study (GWAS) of Alzheimer's Disease
(AD),[28] using a minor allele frequency of 0.05. We selected AD PGRS as it is based on a well-
powered GWAS[28] and has, as the most common neurodegenerative disease, real-world
applicability, particularly considering the examined ageing sample. A previous study also
presented relatively strong associations between a PGRS of late onset AD and brain
structure.[29]

Self-reported measures of mental health
We assessed two self-reported measures reflecting state- and trait-level characteristics of
mental health: recently experienced depression and the personality trait neuroticism.[30] To
assess depression, we used the Recent Depressive Symptoms (RDS-4) score (range: 4-16),

which contains four questions covering four dimensions of depression: mood, disinterest, restlessness, and tiredness. The score is computed as a sum of the 4-point Likert-like responses from the four items. The Eysenck Personality Questionnaire-Revised Short Form, which includes 12 items, was used to assess neuroticism. A sum score was computed from the binary responses to the 12 items, where symptoms were either present or absent. Both scores were previously validated against other corresponding scales using test-retest data of the UK Biobank imaging subsample and recommended to be used to assess imaging biomarkers in the context of mental health.[30]

Atmospheric data

Monthly meteorological data were obtained from the ERA5 reanalysis dataset,[31] specifically the monthly averaged single-level product prior scanning, accessed via the Copernicus Climate Data Store.[32] These data are based on historical observations with a numerical weather prediction model to produce a spatially and temporally complete reconstruction of the atmosphere over extended periods (~70+ years). Unlike raw observational records, which are irregularly distributed in space and time, reanalysis provides globally gridded fields on a regular mesh, serving as a proxy for the actual atmospheric state. This dataset captures both long-term climate trends and natural variability, making it suitable for investigations of climate patterns, extremes, and variability. In contrast, ensemble means of general circulation models primarily represent the forced climate signal, with much of the natural variability averaged out. Importantly, we used ERA5 reanalysis as a representation of observed atmospheric conditions, rather than relying on weather forecasts or purely modelled climate projections. For each parameter, spatial means were computed over the broader United Kingdom area (60°–50°N, 7°W–1°E) to derive regional time series for analysis. The greater grid was selected for more accurate measures of atmospheric variables (in contrast to smaller scale local parameters). As a result, the following climate metrics were used for the analysis: the wind speed 10 metres above the Earth's surface in m/s (10 metre wind speed), the 2-metre temperature in Kelvin, 24-hour surface downward ultraviolet (UV) radiation in $J*m^{-2}$, the surface pressure in Pascal, and the total precipitation in $kg\ m^{-2}\ s^{-1}$.

268 **Table 2. Atmospheric characteristics at baseline.**

Characteristic	N = 30,831[1]
Wind speed (m/s)	6.67 (5.71, 7.53)
Temperature in Kelvin	282.9 (280.5, 286.5)
Surface pressure in Pascal	100,673 (100,342, 100,861)
UV radiation in J*m^{-2}	1,123,889 (507,593, 2,059,785)
Total precipitation in kg m^{-2} s^{-1}	0.0031 (0.0024, 0.0036)

^{1} Median (IQR)

Statistical Analyses

First, we ran simple linear models predicting brain variables (*B,* cortical thickness, fractional

anisotropy, radial, axial, and mean diffusivity) from atmospheric patterns or weather

variables (*W*, see Weather Data section) at baseline, controlling for site, household income,

sex, age, intracranial volume (ICV), and surface holes (SH). ICV has previously been shown to

influence DTI[33] and volumetric scalars[34] and needs therefore to be controlled for.

$$B_j = b_0 + b_1 W_i + b_2 sex + b_3 income + b_4 age + b_5 site + b_6 ICV + b_7 SH + u \qquad (1)$$

Second, we added either psychological and or AD PGRS (*PsyG* in the formula) separately to

each of the estimated models.

$$B_j = b_0 + b_1 W_i + b_2 sex + b_3 income + b_4 age + b_5 site + b_6 ICV + b_7 SH + b_8 PsyG + u \qquad (2)$$

We then used likelihood ratio tests to examine whether the added psychological and

genetic factors improved model performance.

As supplemental analyses (results are reported in the Supplement), we first show additional

baseline and longitudinal associations of PGRS of common psychiatric disorder using

formula 2. Second, we report the outcomes from linear mixed effect models mirroring the

formula 1 but adding follow-up data and a random effect of participant to examine the

longitudinal effects for participants for which two scans were available. Among the brain

variables *B$_j$*, we also added brain age. Brain age models were trained on the cross-sectional

data not entailing the participants from the longitudinal set using a simple linear regression

model. This procedure, including strong model performance, has also been described

previously.[35] Third, we used the extracted tracts and brain volume average from the

Desikan-Killiany atlas to run the same models as presented in formula 1, with the region-

and tract-level metrics as the respective dependent variable.

The alpha level set at 0.05 was Benjamini-Hochberg corrected to control for the false

discovery rate. For comparability, we reported standardised regression coefficients.

Ethics approval

This study was approved by the Norwegian Ethics Commission REK 567301, PVO 17/21624

(Ole Andreassen). The study has been conducted using UKB data under Application 27412.

UKB has received ethics approval from the National Health Service National Research Ethics

Service (ref 11/NW/0382).

Data availability

Analysis code can be found at https://github.com/MaxKorbmacher/WeatherBrain.

ERA5 monthly averaged data on single levels are available from the Copernicus Climate Data

Store (https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels-monthly-

means) (accessed 07 July 2025).

Contains modified Copernicus Climate Change Service information 2025. Neither the

European Commission nor ECMWF are responsible for any use that may be made of the

Copernicus information or data it contains.

Acknowledgements

We want to thank Dennis van der Meer for estimating the polygenic risk scores for the

presented analyses. We also want to thank all UK Biobank study facilitators and participants.

References

1. Chen, S. *et al.* Long-term impacts of heatwaves on accelerated ageing. *Nat. Clim. Chang.*
15, 1000–1007 (2025).

2. Ebi, K. L. *et al.* Hot weather and heat extremes: health risks. *The Lancet* **398**, 698–708
(2021).

3. Aspvik, N. P. *et al.* Do weather changes influence physical activity level among older
adults? - The Generation 100 study. *PLoS One* **13**, e0199463 (2018).

4. Buell, J. S. *et al.* 25-Hydroxyvitamin D, dementia, and cerebrovascular pathology in elders
receiving home services. *Neurology* **74**, 18–26 (2010).

5. Dunster, G. P. *et al.* Daytime light exposure is a strong predictor of seasonal variation in
sleep and circadian timing of university students. *Journal of Pineal Research* **74**, e12843
(2023).

6. Grant, L. *et al.* Global emergence of unprecedented lifetime exposure to climate
extremes. *Nature* **641**, 374–379 (2025).

7. Chan, C. B., Ryan, D. A. & Tudor-Locke, C. Relationship between objective measures of
physical activity and weather: a longitudinal study. *Int J Behav Nutr Phys Act* **3**, 1–9 (2006).

8. Edwards, N. M. *et al.* Outdoor Temperature, Precipitation, and Wind Speed Affect Physical
Activity Levels in Children: A Longitudinal Cohort Study. *J Phys Act Health* **12**, 1074–1081
(2015).

9. Twohig-Bennett, C. & Jones, A. The health benefits of the great outdoors: A systematic
review and meta-analysis of greenspace exposure and health outcomes. *Environ Res* **166**,
628–637 (2018).

10.   Kelly, P. *et al.* Systematic review and meta-analysis of reduction in all-cause mortality

from walking and cycling and shape of dose response relationship. *International Journal*

of Behavioral Nutrition and Physical Activity **11**, 132 (2014).

11.   Zhou, M., Xie, Y., Wang, C., Shen, L. & Mauzerall, D. L. Impacts of current and climate

induced changes in atmospheric stagnation on Indian surface PM2.5 pollution. *Nat*

Commun **15**, 7448 (2024).

12.   Block, M. L. & Calderón-Garcidueñas, L. Air pollution: mechanisms of

neuroinflammation and CNS disease. *Trends Neurosci* **32**, 506–516 (2009).

13.   Power, M. C., Adar, S. D., Yanosky, J. D. & Weuve, J. Exposure to air pollution as a

potential contributor to cognitive function, cognitive decline, brain imaging, and

dementia: A systematic review of epidemiologic research. *NeuroToxicology* **56**, 235–253

(2016).

14.   Shi, L. *et al.* Long-term effects of PM2·5 on neurological disorders in the American

Medicare population: a longitudinal cohort study. *Lancet Planet Health* **4**, e557–e565

(2020).

15.   Wang, M.-Y. *et al.* The within-subject stability of cortical thickness, surface area, and

brain volumes across one year. 2024.06.01.596956 Preprint at

https://doi.org/10.1101/2024.06.01.596956 (2024).

16.   Alfaro-Almagro, F. *et al.* Image processing and Quality Control for the first 10,000

brain imaging datasets from UK Biobank. *NeuroImage* **166**, 400–424 (2018).

17.   Maximov, I. I., Alnæs, D. & Westlye, L. T. Towards an optimised processing pipeline for

diffusion magnetic resonance imaging data: Effects of artefact corrections on diffusion

metrics and their age associations in UK Biobank. *Human Brain Mapping* **40**, 4146–4162

(2019).

18.   Smith, S. M. *et al.* Advances in functional and structural MR image analysis and

implementation as FSL. *NeuroImage* **23**, S208–S219 (2004).

19.   Basser, P. J., Mattiello, J. & LeBihan, D. MR diffusion tensor spectroscopy and imaging.

Biophys J **66**, 259–267 (1994).

20.   Henriques, R. N., Jespersen, S. N., Jones, D. K. & Veraart, J. Toward more robust and

reproducible diffusion kurtosis imaging. *Magnetic Resonance in Medicine* **86**, 1600–1613

(2021).

21.   Smith, S. M. *et al.* Tract-based spatial statistics: voxelwise analysis of multi-subject

diffusion data. *NeuroImage* **31**, 1487–1505 (2006).

22.   Maximov, I. I. *et al.* Fast qualitY conTrol meThod foR derIved diffUsion Metrics

(YTTRIUM) in big data analysis: UK Biobank 18,608 example. *HBM* **42**, 3141–3155 (2021).

23.   Elyounssi, S. *et al.* Addressing artifactual bias in large, automated MRI analyses of

brain development. *Nat Neurosci* **28**, 1787–1796 (2025).

24.   Rosen, A. F. *et al.* Quantitative assessment of structural image quality. *NeuroImage*

169, 407–418 (2018).

25.   Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral

cortex on MRI scans into gyral based regions of interest. *NeuroImage* **31**, 968–980 (2006).

26.   Mori, S., Wakana, S., Nagae-Poetscher, L. & Van Zijl, P. MRI atlas of human white

matter. *American Journal of Neuroradiology* **27**, 1384 (2006).

27.   Privé, F., Arbel, J. & Vilhjálmsson, B. J. LDpred2: better, faster, stronger. *Bioinformatics*

36, 5424–5431 (2021).

28.   Wightman, D. P. *et al.* A genome-wide association study with 1,126,563 individuals

identifies new risk loci for Alzheimer's disease. *Nat Gen* **53**, 1276–1282 (2021).

29.   Tank, R. *et al.* Association between polygenic risk for Alzheimer's disease, brain

structure and cognitive abilities in UK Biobank. *Neuropsychopharmacol.* **47**, 564–569

(2022).

30.   Dutt, R. K. *et al.* Mental health in the UK Biobank: A roadmap to self-report measures

and neuroimaging correlates. *Human Brain Mapping* **43**, 816–832 (2022).

31.   ERA5 monthly averaged data on single levels from 1940 to present.

https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels-monthly-

means?tab=overview.

32.   Buontempo, C. *et al.* The Copernicus Climate Change Service: Climate Science in

Action. *Bulletin of the American Meteorological Society* **103**, E2669–E2687 (2022).

33.   Eikenes, L., Visser, E., Vangberg, T. & Håberg, A. K. Both brain size and biological sex

contribute to variation in white matter microstructure in middle-aged healthy adults. *Hum*

Brain Mapp **44**, 691–709 (2022).

34.   Sanchis-Segura, C., Ibañez-Gual, M. V., Aguirre, N., Cruz-Gómez, Á. J. & Forn, C.

Effects of different intracranial volume correction methods on univariate sex differences in

grey matter volume and multivariate sex prediction. *Sci Rep* **10**, 12953 (2020).

35.   Korbmacher, M. *et al.* Cross-Sectional Brain Age Assessments Are Limited in

Predicting Future Brain Change. *Human Brain Mapping* **46**, e70203 (2025).