<u>**Title**</u>

Benefits of hydrogen-rich jelly during constant load testing in healthy adults: A pilot

double-blind randomized crossover trial

Yuki Muramoto[1#], Genta Toshima[2#], Emi Minaguchi[1], Kazuhisa Sugai[1], Yuji Iwasawa[1], Kohsuke

Shirakawa[3], Kengo Nagashima[2], Yasunori Sato[2,4], Motoaki Sano[5], Kazuki Sato[1], Yoshinori

Katsumata[1,3*]

Yuki Muramoto & Genta Toshima are co-first author

[1] Institute for Integrated Sports Medicine, Keio University School of Medicine, Tokyo, Japan

[2] Clinical and Translational Research Center, Keio University Hospital, Tokyo, Japan

[3] Department of Cardiology, Keio University School of Medicine, Tokyo, Japan

[4] Department of Biostatistics, Keio University School of Medicine, Tokyo, Japan

[5] Department of Medicine and Clinical Science, Yamaguchi University Graduate School of

Medicine, Yamaguchi, Japan

Running Title: Effects of Hydrogen Jelly During Exercise

*Corresponding author

Yoshinori Katsumata, MD, PhD

Department of Cardiology, Keio University School of Medicine

35 Shinanomachi Shinjuku-ku, Tokyo, Japan 160-8582

Tel: +81-3-3353-1211, Fax: +81-3-5363-3875, E-mail: goodcentury21@keio.jp

Abstract

Purpose: Exercise-induced fatigue hinders activity. Molecular hydrogen may enhance mitochondrial function and efficiency, but its effects in non-athletes are unclear. We evaluated whether hydrogen-rich jelly (HRJ) enhances high-intensity exercise in non-athletes.

Methods: This double-blind, randomized, placebo-controlled crossover trial (UMIN000053818) enrolled 50 healthy adults, who were randomly assigned in a 1:1 ratio to ingest HRJ first or placebo jelly (PJ) first. Participants consumed 10 g of HRJ or PJ on three occasions: 24 h, 90 min, and immediately before exercise. An incremental test determined anaerobic threshold (AT), followed by 30 min of cycling at 120% of AT. Oxygen consumption (VO_2) and heart rate (HR) were continuously monitored; blood lactate was sampled every 5 min. Condition differences were assessed with paired t-tests and mixed-effects models.

Results: Of the 50 participants, 48 completed all measurements, and analyses were restricted to the 28 who completed the entire 30-min exercise session (13 women, 46.4%). The mean age was 41.7 ± 11.5 years; mean BMI 22.3 ± 2.6 kg/m²; and mean peak VO_2, 26.8 ± 5.9 mL/kg/min. In the PJ condition, maximum lactate reached 5.12 mmol/L. Compared with PJ, HRJ resulted in significantly lower lactate levels at 5 and 10 min, as well as a lower peak lactate (mean difference −0.75, p = 0.014). VO_2 at 10 and 30 min and peak VO_2 were also lower with HRJ (peak difference −1.17, p = 0.017). HR showed no significant difference between conditions.

Conclusion: HRJ reduced peak blood lactate and VO_2, suggesting suppressed metabolite accumulation and improved exercise efficiency.

47 **Keywords:** Exercise efficiency; Hydrogen-rich jelly; Lactate; Metabolic load; Mitochondrial

48 function; Molecular hydrogen; Non-athlete; Oxygen uptake

1. <u>Introduction</u>

A plethora of substantial benefits have been attributed to exercise, including weight reduction,[1,2] improved exercise tolerance,[1,2] and enhanced quality of life.[3,4] Exercise plays a pivotal role in both the primary and secondary prevention of disease onset and progression, thereby reducing the overall mortality risk.[1-4] To maximize these benefits, the World Health Organization (WHO) recommends approximately 75 min per week of vigorous-intensity exercise, defined as activity that results in perspiration and a sense of breathlessness.[5] Importantly, vigorous-intensity exercise elicits greater improvements in oxygen consumption compared with low-intensity exercise[6] and is consistently associated with a reduced risk of mortality.[7-9] Thus, the health benefits of exercise have been demonstrated from multiple perspectives, and physical activity has been increasingly recommended for the general population in various guidelines and position statements.[1,4,5] Consequently, there has been an observed trend of individuals increasingly participating in regular exercise as part of their daily routines.[10] However, the recommended level of vigorous-intensity exercise corresponds to an intensity that partly relies on anaerobic metabolism for energy production and is often accompanied by fatigue, which limits the ability to sustain the prescribed workload.[11] To facilitate the incorporation of relatively high-intensity exercise into daily life, strategies that enhance mitochondrial function (oxidative phosphorylation) and alleviate exercise-induced fatigue may be beneficial. Based on this rationale, we focused on molecular hydrogen.

Molecular hydrogen attenuates oxidative stress at the cellular level.[12] Its efficacy and safety have been documented in both disease treatment and general health maintenance.[13,14] Hydrogen-rich water (HRW), in which molecular hydrogen is dissolved, has been shown to reduce post-

exercise lactate accumulation and alleviate muscle fatigue, with particularly notable effects in

endurance athletes such as cyclists and marathon runners.[15-18] One proposed mechanism involves

the selective elimination of potent reactive oxygen species, such as hydroxyl radicals (•OH), via

heme proteins.[19] In addition, molecular hydrogen enhances mitochondrial membrane potential,

thereby promoting mitochondrial activity and ATP production.[20] Collectively, these processes are

thought to suppress excessive lactate production. Physiological responses to high-intensity

exercise have been extensively examined in athletes. However, much of the existing evidence

derives from athlete cohorts, and the physiological effects of molecular hydrogen during exercise

in non-athletes remain insufficiently explored. Recently, the methods of hydrogen administration

have diversified. Our laboratory developed a technique to encapsulate molecular hydrogen

within a jelly matrix composed of polysaccharides and gelatin, achieving a higher hydrogen

concentration than conventional HRW.[21] Each 10 g portion of this high-concentration hydrogen-

rich jelly (HRJ) delivers hydrogen equivalent to approximately 600 mL of HRW. HRJ therefore

represents a novel approach to daily hydrogen intake, offering greater convenience and

efficiency than traditional HRW.

In this study, we investigated the effects of HRJ on metabolic responses during high-intensity

exercise in non-athletes with moderate fitness levels. We hypothesized that HRJ intake would

reduce inefficient oxygen utilization, suppress exercise-induced lactate accumulation, and

thereby promote more efficient metabolic control from the perspective of energy production.

2. Materials and Methods

2.1. Hydrogen gas content in the hydrogen jelly

The HRJ used in this study was produced in a facility certified under the FSSC 22000 standard,

a globally recognized food safety certification scheme approved by the Global Food Safety

Initiative. The jelly contained uniformly dispersed molecular hydrogen, and each portion was

individually packaged in aluminum pouches to preserve hydrogen content. The concentration of

dissolved hydrogen gas was stable for up to 8 months (Supplemental Fig. 1, Supplemental

Digital Content, Hydrogen molecule content in hydrogen-rich jelly). Using the headspace gas

chromatography method, we confirmed that hydrogen levels remained above 30 ppm for up to

18 months after production.

The HRJ was stored at an ambient temperature of 25 °C, a safe and stable condition that

preserved the integrity of the hydrogen concentration in the product.

2.2. Sample size

This study represents a secondary analysis focusing on exercise-related outcomes of HRJ. The

sample size was determined for the primary analysis, which evaluated heart rate variability

(HRV), with emphasis on the high-frequency (HF) component as an indicator of parasympathetic

nervous system activity.[22] Based on preliminary data, the required sample size to detect

significant changes in HF was calculated using G*Power with a two-tailed α of 0.05 and a power

(1–β) of 0.90, resulting in 36 participants. However, as approximately 20% of participants in the

pilot data exhibited HF values exceeding three standard deviations above the mean (indicating

potential outliers), the target sample size was increased to 45. Accounting for an anticipated

dropout rate of 10%, the final recruitment target was set at 50 participants. For this secondary

analysis, the dataset included 28 participants who successfully completed the exercise test from

within this predefined sample.

2.3. Participants

Fifty healthy adults with varying fitness levels, independent of their exercise habits, were

recruited between May and November 2024. Participants were aged 22 to 60 years, representing

a broad demographic range. Inclusion criteria were as follows: absence of cardiovascular,

respiratory, or metabolic diseases; absence of athletic injuries; non-smoking status; and no use of

dietary supplements or medications for at least 1 week before participation. The study protocol

was approved by the Institutional Review Board of Keio University School of Medicine

(approval number: 20231194) and conducted in accordance with the principles of the Declaration

of Helsinki. All participants were informed about the study and provided written informed

consent. The study was registered in the University Hospital Medical Information Network

(UMIN000053818).

2.4. Study design

This study employed a double-blind, randomized, placebo-controlled crossover design. The

study protocol consisted of three sessions (**Visit 1, 2, and 3; Fig. 1**). On the first day (Visit 1),

body weight, body fat, and skeletal muscle mass were assessed using a bioelectrical impedance

analyzer (Inbody 470; Inbody Japan Inc., Tokyo, Japan). The anaerobic threshold (AT) was then

determined with a ramp load test for exhaustion, performed on an electromagnetically braked

ergometer (StrengthErgo8 V2; Fukuda Denshi Co., Ltd., Tokyo, Japan) equipped with an exhaled

gas analyzer (Aeromonitor®, MINATO Medical Science Co., Ltd., Osaka, Japan) and a heart

rate (HR) monitor (Polar H10, Polar Electro Japan Co., Ltd., Tokyo, Japan). After AT

determination, participants returned on two separate occasions to consume either 10 g of HRJ

(30–40 ppm; Shinryo Co., Ltd., Fukuoka, Japan) or placebo jelly (PJ), followed by a 30-min

constant-load test at 120% of AT **(Visits 2 and 3; Fig. 1).** In addition, exhaled acetone (Ex-ace)

was collected from 20 consenting participants before and after exercise using the same device

(ReCIVA Breath Sampler).

Participants were randomly assigned (1:1) to two groups using an allocation sequence generated

in Microsoft Excel, which determined whether HRJ or PJ was administered first. Administration

followed a double-blind design, with containers labeled ID_1 or ID_2. Participants consumed the

ID_1 drink before Visit 2 and the ID_2 drink before Visit 3, ensuring that neither the participants

nor the investigators could identify the hydrogen water condition. The two conditions were

measured with a washout interval of at least 7 days. The concentration of hydrogen in the body

reaches its peak within a few minutes after hydrogen gas inhalation,[16] and hydrogen in the

exhaled breath is completely lost 60 min after intake.[17] Therefore, participants were instructed to

drink 10 g of HRJ or PJ within 2 min at the following time points: 24 h, 90 min, and immediately

before the test.

2.6. Ramp test protocol

Participants were asked to fast for 3 h before the test and were prohibited from consuming

caffeine, which can enhance exercise performance by mitigating or delaying fatigue, and alcohol,

which may accelerate fatigue, for 12 h before the test. After measuring the remaining baseline

data for 2 min, the participants performed a warm-up exercise for 2 min at a 20-W load and then

exercised at increasing intensities until they could no longer maintain the pedaling rate

(volitional exhaustion). The load was increased by 15 W at 1-min intervals. The rotational speed

was maintained at 70 rpm.

2.7. Respiratory gas analysis and AT

Expired gas flow was assessed with a breath-by-breath automated system, and respiratory gas

exchange variables (minute ventilation [VE], oxygen uptake [VO_2], and carbon dioxide output

[VCO_2]) were continuously monitored and averaged over 10 s. A three-step calibration process

was performed for the flow-volume sensor, including delay-time adjustment. Peak VO_2 was

measured as the average oxygen consumption during the last 30 s of exercise. The ventilatory

efficiency slope (VE-VCO_2 slope) was calculated using linear regression of data collected from

the start of exercise until the respiratory compensation point. Initial HR was defined as the

average heart rate recorded during the 2 min preceding exercise in a seated position. AT was

assessed using the ventilatory equivalent, excess carbon dioxide, and modified V-slope

methods.[23] AT decisions were made by experienced cardiac rehabilitation doctors and physical

therapists.

2.8. Constant load test protocol

This test was conducted at 120% of the AT workload for 30 min. The rotational speed was set at

70 rpm. During the constant-load exercise, an exhaled gas analyzer and an HR monitor were

attached. Blood lactate values were obtained by auricular pricking and gentle squeezing of the

earlobe using a blood lactate analyzer (Lactate Pro 2, ARKRAY Inc., Kyoto, Japan). Blood

lactate concentrations were measured at baseline and at 5-min intervals during the 30-min

exercise bout. Exercise termination was defined as the inability to sustain pedaling at 70 rpm

because of fatigue.

2.9. Exhaled Acetone Sampling and Acetone Quantification

Exhaled acetone (Ex-ace) was collected according to a standardized protocol using the Breath

Biopsy® platform (Owlstone Medical). Sampling was performed with the ReCIVA® Breath

Sampler, coupled with the CASPER® Portable Air Supply, and breath was adsorbed onto Tenax

TA/Carbograph 5TD-packed tubes (Markes International, Llantrisant, UK). Breathing patterns

were monitored in real time using CO_2 concentration and pressure sensors, and breath sampling

was triggered when the exhaled CO_2 concentration reached approximately 0.6%.

Approximately 1.0 L of exhaled breath was collected over 8–12 min at a flow rate of 225

mL·min^{-1}, adjusted to each participant's tidal volume. Ambient contamination was minimized by

using the CASPER® Portable Air Supply. Before sampling, the adsorbent tubes were

conditioned at 280 °C under a nitrogen flow (20 psi) for 30 min. After collection, breath samples

were stored at −40 °C and analyzed the next day using a Shimadzu TD-30R thermal desorption

system (Shimadzu Corporation, Kyoto, Japan) coupled to a GCMS-QP2010 Ultra mass

spectrometer. For Ex-ace peak analysis, individual peaks were identified using a pre-established

system library based on retention indices and mass spectra. Optimized integration parameters

were applied to each peak, and the corresponding peak areas were calculated.

2.10. Statistical analysis

Continuous variables were summarized as means with standard deviations, while categorical variables were expressed as frequencies with percentages.

Differences in HR, VO_2, and blood lactate levels between conditions during constant-load exercise were analyzed using mixed-effects models for repeated measures. These differences in HR, VO_2, and lactate changes were assessed using paired t-tests and mixed-effects models for repeated measures, with 95% confidence intervals (CIs) estimated.

For VOC analysis, among the 20 participants whose VOCs were successfully measured, nine participants who completed the constant-load exercise were included for comparison. Exhaled acetone peak area changes were compared between HRJ and PJ conditions using paired t-tests and mixed-effects models, consistent with the analyses of HR, VO_2, and lactate.

Additional subgroup analyses were performed to explore participant characteristics associated with the difference in HR, VO_2, and blood lactate levels changes between the HRJ and PJ conditions. Subgroups were defined according to nine characteristics: sex (male/female), smoking status (no/yes), alcohol intake (daily alcohol consumption of < 20 g / ≥ 20 g), physical activity habits (<3 times per week / ≥ 3 times per week and <4 times per week / ≥ 4 times per week), age (< 47 / ≥ 47), body mass index (BMI) (<22.8 kg·m⁻² / ≥ 22.8 kg·m⁻²), body fat percentage ($< 24.9\%$ / $\geq 24.9\%$), and skeletal muscle mass (<40 kg / ≥ 40 kg). Cutoff values for BMI, body fat percentage, skeletal muscle mass, and age were determined from baseline sample medians.

Effect sizes were calculated using Cohen's d, based on within-subject mean differences and standard deviations. All p-values were two-tailed, and statistical significance was set at $p < 0.05$. Statistical analyses were performed using SAS software, version 9.4 (SAS Institute, Japan).

3. Results

Fifty participants were initially recruited for the study; however, two withdrew prior to commencement due to scheduling conflicts. Among the remaining participants, 48 (27 females) completed the HRJ and PJ interventions and participated in both the constant-load exercise test (Table 1). However, 20 participants were unable to complete the constant-load exercise for 30 min and were excluded from the primary statistical analysis. Thus, the analysis included data from 28 participants (Figure 2).

3.1. Effects of HRJ on Metabolic Responses During Exercise

HRJ intake significantly reduced blood lactate levels and VO_2 during exercise compared with PJ. Blood lactate concentrations were notably lower under the HRJ condition at 5 min (mean difference = –0.49, p = 0.004) and 10 min (mean difference = –0.63, p = 0.008). A trend toward lower values was also observed at 25 min (mean difference = –0.48, p = 0.076). (Table 2, Fig.3). Additionally, peak blood lactate levels during exercise were significantly lower in the HRJ condition (mean difference = –0.75, p = 0.014; Table 2, Fig.3). Similarly, VO_2 was significantly lower in the HRJ condition at 10 min (mean difference = –1.20, p = 0.016) and 30 min (mean difference = –1.40, p = 0.017). A trend toward lower VO_2 was also observed at 25 min (mean difference = –1.13, p = 0.081; Table 2, Fig. 4). Additionally, peak VO_2 during constant-load exercise was also significantly lower with HRJ than with PJ (mean difference = –1.17, p = 0.014; Table 2, Fig. 4).No significant differences were observed between the two conditions in HR at any time point or in peak HR during exercise (Table 2, Fig.5). Regarding the peak area of Ex-ace that emerged before and after exercise, a trend toward a lower response was observed with HRJ than with PJ (mean difference = –4,129,905, p = 0.090).

3.2. Participant Characteristics Associated with the Impact of HRJ on Peak Exercise Response

Participants who exhibited a reduction in peak blood lactate levels under the HRJ condition were primarily characterized by being female (mean difference = –0.90, 95% confidence interval [CI]: –1.64 to –0.15, p = 0.023), daily alcohol consumption ≥20 g (mean difference = –0.86, 95% CI: –1.53 to –0.20, p = 0.014), exercising less than three times per week (mean difference = –0.75, 95% CI: –1.27 to –0.23, p = 0.008), and body fat percentage of ≥24.9% (mean difference = –1.09, 95% CI: –1.93 to –0.25, p = 0.018). Participants younger than 47 years also showed a trend toward a greater reduction (mean difference = –0.84, 95% CI: –1.72 to 0.03, p = 0.058), although this did not reach statistical significance (Supplemental Fig. 2, Supplemental Digital Content, Characteristics of participants who exhibited a reduction in peak blood lactate under the hydrogen-rich jelly (HRJ) condition).

Participants showing a reduction in peak VO□ under the HRJ condition were characterized by a BMI of <22.8 kg·m⁻² (mean difference = –1.75, 95% CI: –2.83 to –0.68, p = 0.004), body fat percentage <24.9% (mean difference = –1.58, 95% CI: –2.70 to –0.47, p = 0.009), skeletal muscle mass ≥40 kg (mean difference = –1.45, 95% CI: –2.81 to –0.10, p = 0.037), and a habitual exercise frequency ≥4 times per week (mean difference = –1.51, 95% CI: –2.79 to –0.22, p = 0.033). Male participants (mean difference = –1.59, 95% CI: –3.24 to 0.07, p = 0.059) and participants aged ≥47 years (mean difference = –1.39, 95% CI: –2.96 to 0.18, p = 0.075) also showed a trend toward greater reductions, although these differences did not reach statistical significance (Supplemental Fig. 3, Supplemental Digital Content, Characteristics of participants

284 who exhibited a reduction in peak oxygen uptake (VO$_2$) under the hydrogen-rich jelly (HRJ)

285 condition.).

4. Discussion

This study examined the effects of HRJ intake on physiological responses to high-intensity

exercise in non-athletes. Our findings corroborate those of earlier studies, showing that HRJ

blunted the rise in blood lactate during the first 10 min of exercise and reduced peak lactate

levels during exercise. A novel observation was that peak VO$_2$ during exercise was significantly

lower with HRJ. Collectively, these findings suggest that HRJ reduces unnecessary oxygen

utilization and attenuates lactate accumulation, thereby improving exercise efficiency in non-

athletes, even at comparable intensities (Fig.6).

4.1. Effects of HRJ on Metabolic Responses During Exercise

Previous studies have reported that HRW suppressed blood lactate accumulation in athletes

during high-intensity exercise, such as that performed at 75% of peak VO$_2$ or during the Quebec

90-s test.[15,16] In the current trial, the mean blood lactate concentration under the placebo

condition was 5.12 mmol·L^{-1}, which was within the range of 5–12 mmol·L^{-1} observed in previous

studies.[15,16] These results indicated that, even in non-athletes, aerobic exercise of sufficient

intensity to elevate lactate levels to a comparable range revealed the lactate-suppressing effect of

hydrogen gas. Here, the lactate level increase within the first 10 min of exercise and the peak

lactate level were attenuated. At the onset of high-intensity exercise, oxygen supply falls short of

demand, resulting in an oxygen deficit. Consequently, glycogen is anaerobically broken down to

generate ATP, leading to lactate accumulation.[23,24] Lactate is subsequently reutilized as a

substrate for ATP resynthesis within mitochondria, thereby supporting sustained exercise

performance.[25] In addition, lactate produced within skeletal muscles is transported to the liver

and converted to glucose via the Cori cycle, which is then reutilized as an energy substrate.[26]

Thus, suppression of lactate elevation during high-intensity exercise could be attributed to both

reduced lactate production and enhanced lactate utilization in skeletal muscles, as well as

augmented gluconeogenic capacity in hepatocytes. With respect to the former, the effect was

likely mediated by improved mitochondrial function within skeletal muscles. Our findings also

showed that HRJ intake suppressed VO_2 during exercise. This suggests that mitochondrial

oxygen utilization efficiency was enhanced, thereby allowing the same exercise intensity to be

maintained with lower oxygen consumption. Skeletal muscle mitochondria generate energy

through oxygen consumption; however, during this process, superoxide (O_2^-) and hydrogen

peroxide (H_2O_2) are produced. In the presence of transition metals, such as iron (Fe), the

Fenton reaction generates highly reactive hydroxyl radicals (•OH), which subsequently trigger

lipid peroxidation.[27] More so, lipid peroxide accumulation impairs mitochondrial oxygen uptake

capacity.[28] Molecular hydrogen selectively neutralizes hydroxyl radicals (•OH) and suppresses

lipid peroxidation chain reactions.[12,19,29] This antioxidative action prevents a decline in oxygen

utilization efficiency, thereby contributing to the observed reduction in VO_2. Such

improvements in mitochondrial function were also likely to reduce skeletal muscle lactate

production and facilitate energy production via lactate reutilization. Generally, hydrogen gas

inhaled via the gastrointestinal tract is absorbed and almost completely exhaled through the

lungs.[30] However, during exercise, the shortened pulmonary capillary transit time and increased

intrapulmonary shunting reduced hydrogen exhalation, suggesting that hydrogen gas might have

remained in the arterial circulation. Future studies are required to investigate the hemodynamics

of hydrogen gas during exercise in greater detail.

HRJ intake also tended to suppress the post-exercise rise in Ex-ace. High-intensity exercise

typically results in depletion of muscle glycogen stores, leading to a metabolic shift toward fatty

acid oxidation.[31,32] Consequently, ketone bodies are generated in the liver, with acetone being

produced as a byproduct.[31,32] Therefore, the attenuated increase in Ex-ace under the HRJ

condition suggests that improved oxygen utilization efficiency promoted glucose metabolism and

lactate reutilization while limiting excessive reliance on fatty acid oxidation. Orally ingested

molecular hydrogen delivers high concentrations of hydrogen to the liver via the portal

vein.[30,33,34] The liver plays a key role in lactate reutilization through the Cori cycle, and

improvement of hepatic function induced by hydrogen intake likely contributed to the enhanced

lactate clearance. Furthermore, in the present study, HRJ was ingested 24 h before exercise, as

well as 90 min and immediately before exercise. Considering hydrogen gas rapidly disappears

from the body after ingestion, acute effects of hydrogen gas on the organs were unlikely present

24 h after ingestion. Instead, a preconditioning effect might have been induced via activation of

the PGC-1α pathway, promoting functional improvement of existing mitochondria and

stimulating mitochondrial biogenesis.[33] Thus, both skeletal muscle and liver metabolism were

primed, resulting in more pronounced suppression of lactate accumulation and improvement in

oxygen utilization efficiency.

4.2. Participant Characteristics Associated with the Impact of HRJ on Peak Exercise

Response

We examined the characteristics of participants whose peak blood lactate concentration and

peak VO□ were reduced by HRJ intake during exercise. The reduction in peak lactate was

significant primarily in women, individuals exercising less than three times per week, those with

body fat percentages above the median, and those consuming more than 20 g of alcohol per day.

Conversely, peak VO₂ reduction was significant mainly in men with greater skeletal muscle

mass (≥40 kg), below-median BMI and body fat percentages, and regular exercise four or more

times per week.

A plausible explanation for these sex- and body composition-related differences is variations in

skeletal muscle fiber composition and metabolic properties. Generally, women possess a higher

proportion of type I (slow-twitch) fibers, characterized by superior oxidative metabolic capacity,

whereas men typically have a higher proportion of type II (fast-twitch) fibers, associated with

greater reliance on anaerobic metabolism and higher oxygen consumption during high-intensity

exercise.[35,36] Beyond fiber distribution, sex differences have also been reported in lactate

production and clearance, with women exhibiting a greater capacity for lactate reutilization and

more efficient lactate removal during recovery compared with men.[37] Molecular hydrogen has

been shown to enhance mitochondrial function.[20] Accordingly, it is plausible that lactate

production during exercise was suppressed in women through mitochondrial action, thereby

reducing metabolic load. In addition, individuals with higher body fat percentages or habitual

alcohol consumption are prone to elevated oxidative stress; however, HRJ intake may have

strengthened antioxidant defense mechanisms, which in turn could have contributed to the

suppression of lactate elevation.

Conversely, reductions in peak VO₂ were primarily observed in men with greater skeletal

muscle mass and more frequent exercise habits. These individuals are generally predisposed to

higher oxygen consumption during muscle activity; however, HRJ intake may have improved the

efficiency of oxygen delivery and utilization, thereby reducing the VO₂ required to sustain the

same exercise intensity. Molecular hydrogen has also been reported to improve blood flow,[38]

suggesting that enhanced peripheral oxygen delivery through vasodilation and improved

microcirculation may have contributed to this outcome. These circulatory improvements could

further enhance muscular oxygen utilization, ultimately reducing the VO₂ required to sustain

exercise performance.

4.3. Limitation

This study has several limitations. First, it was designed to examine whether HRJ influences

physiological indices during high-intensity exercise in non-athletes. As a result, nearly half of the

participants did not complete the study protocol. It remains unclear whether HRJ provides

beneficial effects during moderate, daily life exercise in non-athletes. Second, the present study

investigated only the acute effects of HRJ intake, and its long-term impact on exercise

performance and physiology is still unknown. Further studies are warranted to elucidate the

potential long-term physiological and performance-related effects of HRJ. Third, the analysis of

Ex-ace was based on a small analyzable sample (n = 9), limiting statistical power. A power

analysis performed with G*Power 3.1 estimated that 16 participants would be necessary to

achieve 80% statistical power at a two-sided α of 0.05, highlighting the potential risk of type II

error with the current sample. Exercise performance is also influenced by daily physiological

variations, which could contribute to inter-session variability.

Future studies should address these limitations by examining the effects of HRJ during exercise

at daily life intensities, evaluating the impact of longer-term intake, and incorporating analyses

that consider participants' fitness levels. Such approaches will help to more clearly elucidate the

potential physiological effects of HRJ.

5. Conclusions

In the present study, intake of HRJ led to reductions in both peak blood lactate concentration

and peak VO$_2$. Furthermore, HRJ was associated with a suppression of the rise in Ex-ace,

suggesting that excessive metabolite production may be attenuated even at the same exercise

intensity. Collectively, these findings indicate that HRJ may enhance exercise efficiency and

reduce metabolic load, providing a potentially useful strategy even for non-athletes.

<u>Acknowledgements:</u>

This work was supported by Shinryo CO., LTD. However, this corporation was not involved in

data analysis and manuscript writing. The authors would like to thank the participants who

actively participated in this study and SOUKEN Corporation (https://www.souken-lab.co.jp/) for

their help with recruitment. The manuscript has been edited by Editage (www.editage.jp). The

authors express their sincere appreciation to Noriko Akiyama for her substantial contributions to

the measurement of VOC.

<u>Conflict of interest</u>

Yoshinori Katsumata reports that Shinryo Co., Ltd. provided both financial support and

hydrogen-rich jelly for this study. The company had no influence on the study design, data

analysis, or manuscript writing. No other competing interests are declared.

<u>Ethical Statement:</u>

The study protocol was approved by the Institutional Review Board of the Keio University

School of Medicine (approval number: 20221205) and was conducted in accordance with the

principles of the Declaration of Helsinki. All participants were informed about the study and

asked to provide their names on a consent form. The study was registered with the University

Hospital Medical Information Network (UMIN000050872).

Clinical Trial:

The clinical trial described in this paper was registered with the University Hospital Medical

Information Network under the registration number UMIN000050872.

Data availability statement

All data from this study are presented in this manuscript or are available from the corresponding

author upon reasonable request.

CRediT author statement

The author contributions are stated as follows: Y.M., G.T., and Y.K. drew the manuscript. Y.M.,

G.T., and Y.K. prepared the images. Y.M., K.S., Y.I., K.S., and Y.K. collected the patient

information. K.N. and G.T were conducted statical analysis. Y.S., K.S., M.S., and Y.K. critically

revised the manuscript to ensure intellectual content and supervision. All authors have approved

all aspects of our work and have read and approved the manuscript.

References

1. Coletta AM, Marquez G, Thomas P, et al. Clinical factors associated with adherence to aerobic and resistance physical activity guidelines among cancer prevention patients and survivors. *PLOS One*. 2019;14(8): e0220814. doi: 10.1371/journal.pone.0220814.

2. Westcott WL. Resistance training is medicine: effects of strength training on health. *Curr. Sports Med Rep*. 2012;11(4):209–6. doi: 10.1249/JSR.0b013e31825dabb8.

3. Courneya KS, Vardy JL, O'Callaghan CJ, et al. Structured exercise after adjuvant chemotherapy for colon cancer. *N Engl J Med*. 2025;393(1):13–25. doi: 10.1056/NEJMoa2502760

4. Zhao M, Veeranki SP, Magnussen CG, Xi B. Recommended physical activity and all cause and cause specific mortality in US adults: prospective cohort study. *BMJ*. 2020;370:m2031. doi: 10.1136/bmj.m2031.

5. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451–62. doi: 10.1136/bjsports-2020-102955.

6. Yin H, Zhang J, Lian M, Zhang Y. A systematic review and meta-analysis of the effectiveness of high-intensity interval training for physical fitness in university students. *BMC Public Health*. 2025;25(1):1601. doi: 10.1186/s12889-025-22829-7.

7. Stamatakis E, Ahmadi MN, Friedenreich CM, et al. Vigorous intermittent lifestyle physical

activity and cancer incidence among nonexercising adults: the UK Biobank

accelerometry study. *JAMA Oncol*. 2023;9(9):1255–9. doi:

10.1001/jamaoncol.2023.1830

8. Stamatakis E, Ahmadi MN, Gill JMR, et al. Association of wearable device-measured

vigorous intermittent lifestyle physical activity with mortality. *Nat Med*.

2022;28(12):2521–9. doi: 10.1038/s41591-022-02100-x.

9. Arem H, Moore SC, Patel A, et al. Leisure time physical activity and mortality: a detailed

pooled analysis of the dose-response relationship. JAMA Intern Med. 2015;175(6):959–

67. doi: 10.1001/jamainternmed.2015.0533

10. Yang L, Cao C, Kantor ED, et al. Trends in sedentary behavior among the US population,

2001-2016. *JAMA*. 2019;321(16):1587–97. doi: 10.1001/jama.2019.3636.

11. Bennie JA, De Cocker K, Teychenne MJ, Brown WJ, Biddle SJH. The epidemiology of

aerobic physical activity and muscle-strengthening activity guideline adherence among

383,928 U.S. adults. *Int J Behav Nutr Phys Act*. 2019;16(1):34. doi: 10.1186/s12966-

019-0797-2.

12. Ohsawa I, Ishikawa M, Takahashi K, et al. Hydrogen acts as a therapeutic antioxidant by

selectively reducing cytotoxic oxygen radicals. *Nat Med*. 2007;13(6):688–94. doi:

10.1038/nm1577.

13. Sano M, Tamura T. Hydrogen gas therapy: from preclinical studies to clinical trials. *Curr*

Pharm Des. 2021;27(5):650–8. doi: 10.2174/1381612826666201221150857.

14. Katsumata Y, Sano F, Abe T, et al. The effects of hydrogen gas inhalation on adverse left

ventricular remodeling after percutaneous coronary intervention for ST-elevated

myocardial infarction—first pilot study in humans. *Circ J*. 2017;81(7):940–7. doi:

10.1253/circj.CJ-17-0105.

15. Aoki K, Nakao A, Adachi T, Matsui Y, Miyakawa S. Pilot study: effects of drinking

hydrogen-rich water on muscle fatigue caused by acute exercise in elite athletes. *Med*

Gas Res. 2012;2:12. doi: 10.1186/2045-9912-2-12.

16. Botek M, Krejčí J, McKune AJ, Sládečková B, Naumovski N. Hydrogen rich water

improved ventilatory, perceptual and lactate responses to exercise. *Int J Sports Med*.

2019;40(14):879–85. doi: 10.1055/a-0991-0268.

17. Dobashi S, Takeuchi K, Koyama K. Hydrogen-rich water suppresses the reduction in

blood total antioxidant capacity induced by 3 consecutive days of severe exercise in

physically active males. *Med Gas Res*. 2020;10(1):21–6. doi: 10.4103/2045-

9912.279979.

18. Timón R, Olcina G, González-Custodio A, Camacho-Cardenosa M, Camacho-Cardenosa

A, Martínez Guardado I. Effects of 7-day intake of hydrogen-rich water on physical

performance of trained and untrained subjects. *Biol Sport*. 2021;38(2):269–75. doi:

10.5114/biolsport.2020.98625.

19. Jin Z, Zhao P, Gong W, Ding W, He Q. Fe-porphyrin: A redox-related biosensor of

hydrogen molecule. *Nano Res*. 2023;16(2):2020–5. doi: 10.1007/s12274-022-4860-y.

20. Murakami Y, Ito M, Ohsawa I. Molecular hydrogen protects against oxidative stress-

induced SH-SY5Y neuroblastoma cell death through the process of mitohormesis. *PLOS*

One. 2017;12(5):e0176992. doi: 10.1371/journal.pone.0176992.

21. Maruyama T, Takayama E, Tokuno S, Morita M, Ekuni D. Efficacy of oral intake of hydrogen-rich jelly intake on gingival inflammation: A double-blind, placebo-controlled and exploratory randomized clinical trial. *Healthcare (Basel)*. 2025;13(5):577. doi: 10.3390/healthcare13050577.

22. Muramoto Y, Toshima G, Minaguchi E, et al. Hydrogen-rich jelly exhibits a positive effect on heart rate variability during orthostatic loading in healthy adults: A pilot randomized trial. *medRxiv*. doi: 10.1101/2025.08.27.25334136

23. Poole DC, Jones AM. Oxygen uptake kinetics. *Compr Physiol*. 2012;2(2):933–96. doi: 10.1002/cphy.c100072.

24. di Prampero PE, Ferretti G. The energetics of anaerobic muscle metabolism: a reappraisal of older and recent concepts. *Respir Physiol*. 1999;118(2-3):103–15. doi: 10.1016/s0034-5687(99)00083-3.

25. Brooks GA. The science and translation of lactate shuttle theory. *Cell Metab*. 2018 3;27(4):757–85. doi: 10.1016/j.cmet.2018.03.008.

26. Van Hall G. Lactate kinetics in human tissues at rest and during exercise. Acta Physiol (Oxf). 2010;199(4):499–508. doi: 10.1111/j.1748-1716.2010.02122.x.

27. Zheng Y, Sun J, Luo Z, Li Y, Huang Y. Emerging mechanisms of lipid peroxidation in regulated cell death and its physiological implications. Cell Death Dis. 2024;15(11):859. doi: 10.1038/s41419-024-07244-x.

28. Aoi W, Naito Y, Takanami Y, et al. Astaxanthin improves muscle lipid metabolism in exercise via inhibitory effect of oxidative CPT I modification. *Biochem Biophys Res Commun*. 2008;366(4):892–7. doi: 10.1016/j.bbrc.2007.12.019.

29. Iketani M, Ohsawa I. Molecular hydrogen as a neuroprotective agent. *Curr Neuropharmacol*. 2017;15(2):324-331. doi: 10.2174/1570159x14666160607205417.

30. Jin J, Yue L, Du M, Geng F, Gao X, Zhou Y, Lu Q, Pan X. Molecular hydrogen therapy: Mechanisms, delivery methods, preventive, and therapeutic application. MedComm. 2025;6:e70194. doi:10.1002/mco2.70194.

31. Bovey F, Cros J, Tuzson B, et al. Breath acetone as a marker of energy balance: an exploratory study in healthy humans. *Nutr Diabetes*. 2018;8(1):50. doi: 10.1038/s41387-018-0058-5.

32. Königstein K, Abegg S, Schorn AN, et al. Breath acetone change during aerobic exercise is moderated by cardiorespiratory fitness. *J Breath Res*. 2020;15(1):016006. doi: 10.1088/1752-7163/abba6c.

33. Kamimura N, Ichimiya H, Iuchi K, Ohta S. Molecular hydrogen stimulates the gene expression of transcriptional coactivator PGC-1α to enhance fatty acid metabolism. NPJ Aging Mech Dis. 2016;2:16008. doi:10.1038/npjamd.2016.8.

34. Ichihara G, Katsumata Y, Moriyama H, Kitakata H, Hirai A, Momoi M, Ko S,Shinya Y, Kinouchi K, Kobayashi E, Sano M. Pharmacokinetics of hydrogen after ingesting a hydrogen-rich solution: A study in pigs. Heliyon. 2021;7(11):e08359. doi: 10.1016/j.heliyon.2021.e08359. PMID: 34816046; PMCID:PMC8591508.

35. Haizlip KM, Harrison BC, Leinwand LA. Sex-based differences in skeletal muscle

kinetics and fiber-type composition. *Physiology (Bethesda)*. 2015;30(1):30–9. doi:

10.1152/physiol.00024.2014.

36. Nuzzo JL. Sex differences in skeletal muscle fiber types: A meta-analysis. *Clin Anat*.

2024;37(1):81–91. doi: 10.1002/ca.24091.

37. Jones G, Cornett K, Crowell S, Carlyle J, Roland K, Jakobi J. Lactate clearance following

exhaustive exercise differs between sexes and whole body vibration does not enhance

recovery. *Ann Sports Med Res*. 2018;2(5):1031. doi: 10.47739/2379-0571/1031.

38. Nakao A, Sugimoto R, Billiar TR, McCurry KR. Therapeutic antioxidant medical gas. *J*

Clin Biochem Nutr. 2009;44(1):1–13. doi: 10.3164/jcbn.08-193R.

<u>Figure legends</u>

Fig. 1. Protocols for tests 1, 2, and 3.

Red triangle: measurement point of blood lactate; W·min^{-1}: workload per min; VO□: oxygen

uptake.

This study was conducted as a double-blind, randomized, placebo-controlled crossover trial

consisting of three sessions (Visits 1–3).

At Visit 1, body composition (body weight, body fat, and skeletal muscle mass) was measured

using bioelectrical impedance (Inbody 470; Inbody Japan Inc., Tokyo, Japan), and the anaerobic

threshold (AT) was determined using a ramp load test to exhaustion (StrengthErgo8 V2; Fukuda

Denshi Co., Ltd., Tokyo, Japan) with an exhaled gas analyzer (Aeromonitor®, MINATO Medical

Science Co., Ltd., Osaka, Japan) and a heart rate (HR) monitor (Polar H10, Polar Electro Japan

Co., Ltd., Tokyo, Japan).

At Visits 2 and 3, participants consumed either 10 g of HRJ (30–40 ppm; Shinryo Co., Ltd.,

Fukuoka, Japan) or placebo jelly (PJ) in random order, followed by a 30-min constant-load

exercise test at 120% AT.

Fig. 2. CONSORT flow diagram of the study protocol.

Fifty participants were initially recruited, of whom two withdrew before the start of the study due

to scheduling conflicts. The remaining 48 participants completed both the hydrogen-rich jelly

(HRJ) and placebo jelly (PJ) interventions and underwent the constant-load exercise test. Among

them, 20 participants were unable to complete the 30-min constant-load exercise and were

excluded from the primary statistical analysis. Consequently, data from 28 participants were

included in the final analysis.

Fig. 3. Blood lactate responses during the constant-load exercise test.

(A) Time course of blood lactate concentrations measured every 5 min during exercise under the

hydrogen-rich jelly (HRJ) and placebo jelly (PJ) conditions. Blood lactate concentrations were

significantly lower with HRJ at 5 min (mean difference = –0.49, p = 0.004) and 10 min (mean

difference = –0.63, p = 0.008), with a trend toward lower values at 25 min (mean difference = –

0.48, p = 0.076).

(B) Peak blood lactate levels during exercise. Peak values were significantly lower with HRJ

than with PJ (mean difference = –0.75, p = 0.014).

Data are presented as mean ± SD. *p < 0.05 vs. PJ.

Fig. 4. Oxygen consumption during the constant-load exercise test.

(A) Time course of oxygen uptake (VO□) measured every 5 min during exercise under the

hydrogen-rich jelly (HRJ) and placebo jelly (PJ) conditions. VO□ was significantly lower in the

HRJ condition at 10 min (mean difference = –1.20, p = 0.016) and 30 min (mean difference = –

1.40, p = 0.017), with a trend toward lower values at 25 min (mean difference = –1.13, p =

0.081).

(B) Peak VO□ during exercise. Peak values were significantly lower with HRJ than with PJ

(mean difference = –1.17, p = 0.014).

Data are presented as mean ± SD. *p < 0.05 vs. PJ.

Fig. 5. Heart rate during the constant-load exercise test.

(A) Time course of heart rate (HR) measured every 5 min during exercise under the hydrogen-

rich jelly (HRJ) and placebo jelly (PJ) conditions.

(B) Peak HR during exercise.

No significant differences were observed between the two conditions at any time point or in peak

HR.

Data are presented as mean ± SD.

Fig.6. Graphical summary of the effects of hydrogen-rich jelly (HRJ) intake on exercise
efficiency during high-intensity exercise in non-athletes.

A double-blind, randomized, placebo-controlled crossover trial was conducted with 48

participants, and data from 28 participants who completed the 30-min constant-load exercise

were analyzed. Participants ingested HRJ or placebo jelly (PJ, 10 g × 3 doses: 24 h, 90 min, and

immediately before exercise), performed an incremental cycling test to determine the anaerobic

threshold (AT), and then completed 30 min of constant-load cycling at 120% of AT. Compared

with PJ, HRJ suppressed the rise in blood lactate at 5 and 10 min and significantly reduced peak

lactate levels (-0.75 mmol·L^{-1}, $p = 0.014$). In addition, VO$\square$ was significantly lower at 10 and 30

min as well as at peak values (-1.17 mL·min^{-1}·kg^{-1}, $p = 0.017$). No significant difference was

observed in heart rate between conditions.

List of Supplemental Content

Online supplemental figures.docx

Test protocol for determining anaerobic thresholds	Test protocol for constant load tests
Exercise protocol • Ramp load

Measurement

• Heart rate (continuously)
• VO_2 (breath by breath) | Exercise protocol

• Constant load (120% of AT)

Measurement

• Blood lactate (every 5minutes in exercise)
• Heart rate (continuously)
• VO_2 (breath by breath) |

Visit 1

Visit 2, 3

Impact of Hydron-Rich Jelly on Exercise Efficiency in Non-Athletes

Double-blind, randomized, placebo-controlled crossover trial

Total participants (N = 48)

HRJ (n = 24)

PJ (n = 24)

Timing and dosage
10 g × 3 doses:
24 hours, 90 minutes, immediately before exercise

Test protocol
• Incremental cycling test for AT
• 30-minute constant-load cycling at 120% of AT

Analysis
• Blood lactate (every five minutes)
• Oxygen uptake (VO_2, continuous)
• HR

Key findings

Compared to PJ, HRJ showed:

28 participants completed a 30-minute exercise

↓ Blood lactate at 5 and 10 minutes, and ↓ peak lactate (−0.75 mmol/L; $p = 0.014$)

↓ VO_2 at 10 and 30 minutes, and ↓ peak VO_2 (−1.17 mL/kg/min; $p = 0.017$)

HR showed no significant difference

HRJ: Hydrogen rich jelly; PJ: Placebo jelly; HR: Heart rate; AT: Anaerobic threshold

HRJ suppresses lactate accumulation and oxygen consumption during high-intensity exercise, thereby enhancing efficiency in non-athletes