Untargeted longitudinal ultra deep metagenomic sequencing of wastewater provides

a comprehensive readout of expected and unexpected viral pathogens

Clayton Rushford[a], Devon Gregory[b], Emma Copen[b], Jonathan Naydenov[b], Alejandro Tovar-

Mendez[b], Po-E Li[c], Rose S. Kantor[d], Migun Shakya[c], Nelson Ruth[c], Patrick S. G. Chain[c],

Hsinyeh Hsieh[e], Torin Hunter[b], Josh Kome[b], Matt Frank[b], Terri Lyddon[b], Jeff Kaufman[f], David

H. O'Connor[g], Marc C. Johnson[b#]

Author Affiliations

[a]University of Missouri, Department of Veterinary Pathobiology

[b]University of Missouri, Department of Molecular Microbiology and Immunology

[c]B-GEN, Bioscience Division, Los Alamos National Laboratory

[d]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory

[e]University of Missouri, School of Natural Resources

[f]Nucleic Acid Observatory, SecureBio

[g]University of Wisconsin-Madison, Department of Pathology and Laboratory Medicine

[#]Corresponding Author: marcjohnson@health.missouri.edu

Abstract

Wastewater surveillance has become a powerful tool to monitor circulating viruses

at a community level. Currently, most wastewater surveillance efforts use target-based

approaches such as quantitative PCR techniques or hybrid capture. This study explores

the feasibility of using untargeted ultra-deep metagenomic sequencing as a

comprehensive approach to wastewater surveillance. To test this, composite influent

wastewater samples were collected weekly at a single site from January 2024 through June

2025. Sequencing was performed using random hexamers on all samples, with an average

depth of 1.1 billion reads per sample. Human enteric viruses such as rotaviruses,

astroviruses, and noroviruses were detected at high levels in virtually every sample. SARS-

CoV-2 was also detected in most samples and the counts per sample positively correlated

with digital PCR (dPCR) measurements. Less abundant respiratory pathogens such as

influenza A and B, rhinoviruses, parainfluenzaviruses, and human coronaviruses 229E,

OC43, NL63, and HKU1 were also regularly detected. However, those pathogens displayed

distinct and reproducible winter and spring seasonality. Several unexpected viruses were

also detected, such as several detections of highly pathogenic avian influenza H5N1 (HPAI

H5N1) genotype B3.13, a month-long surge of hepatitis A virus, and a large season-specific

surge in influenza C virus. The most abundant known virus detected was

the *Tobamovirus* tomato brown rugose fruit virus, which was present stably year-round at

high abundance. However, other tobamoviruses such as tomato mosaic virus were

detected primarily in the late growing season. This eighteen-month study highlights that

ultra deep sequencing enables detection of expected and unexpected viral pathogens

without targeted enrichment.

Importance

This study demonstrates that untargeted ultra deep metagenomic sequencing can

provide a comprehensive tool for wastewater surveillance of viral pathogens. By generating

approximately 1 billion reads per sample across 78 consecutive weeks, we captured

expected pathogens such as SARS-CoV-2, noroviruses, and influenza viruses. Additionally,

we captured unexpected viral signals such as influenza C and highly pathogenic avian

influenza H5N1. The wide range of viral taxa captured in this study also displays

epidemiologically relevant seasonality. We also observed a correlation between

metagenomic SARS-CoV-2 read counts and dPCR values to validate this method against

other wastewater surveillance methods currently in use. Our findings highlight how ultra

deep metagenomics can enhance pandemic preparedness, enable early detection of non-

standard and clinically overlooked species, and broaden the scope of One Health

monitoring by capturing human, animal, and plant viral signatures from a composite

wastewater sample.

Keywords

Wastewater surveillance, metagenomic, deep sequencing, SARS-CoV-2, influenza

H5N1, influenza C, tomato brown rugose fruit virus, *NVD*, *GOTTCHA2*, One Health, public

health

Introduction

Wastewater surveillance is a powerful tool for public health that leverages

minimally biased, population-level sampling, and supports pandemic preparedness by

providing early warnings for circulating pathogens based on sequence data [1, 2].

Compared to clinical surveillance, wastewater-based approaches reduce biases

associated with asymptomatic infections and limited access to healthcare. Several labs

have demonstrated this concept through reverse transcription PCR (RT-PCR), real time

quantitative PCR (RT-qPCR), and digital PCR (dPCR) designed to target human-infecting

pathogens such as SARS-CoV-2, influenza, and *Enterovirus* spp. [3-6]. However, PCR-

based approaches require occasional modifications due to evolution around primer

binding sites and are limited to known targets [7]. Alternatively, probe-capture approaches

facilitate detection of a wider range of pathogens in wastewater, have potential to detect

pathogens that differ by up to 40% from reference genomes, and are commercially

available [8-14]. While PCR and probe-capture approaches facilitate detection of viruses

with *a priori* knowledge of genetic makeup, they are unable to detect highly divergent,

novel viral sequences that may be circulating in a population.

Untargeted sequencing and metagenomics enable detection of a broader diversity

of known and unknown pathogens if they receive sufficient coverage. Pathogen

concentration methods, such as polyethylene glycol (PEG) precipitation and vacuum-

based direct capture, have been shown to be the most reproducible while improving depth

and breadth of viral signals [15]. However, a deep dataset is a prerequisite for untargeted

metagenomic surveillance due to a low target-to-nontarget ratio of sequences [16].

Fortunately, the rapidly decreasing cost of sequencing has facilitated data generation that

would, until recently, have been infeasible [17].

The goal of this study was to test the feasibility of viral surveillance using minimally

biased metagenomic sequencing. A variety of previous studies have utilized metagenomic

approaches for screening antimicrobial resistance genes, microbiomes, and viromes in

wastewater [2, 18-26]. Most of these take place over a short period, yet a few longitudinal

studies have surveyed wastewater for longer than a year to determine how detectable

sequences change over time [11, 18, 27]. While previous studies have demonstrated the

efficacy of metagenomics for viral surveillance, our study is unique due to our combination

of longitudinal sampling (78 consecutive weeks), use of PEG precipitation for viral

concentration, and sequencing at a depth greater than previous work [2, 11, 18-27]. To

provide a high-confidence multi-kingdom readout of present species, we assigned

sequence reads through *GOTTCHA2* (Genomic Origin Through Taxonomic CHAllenge)

classification using an expanded reference database. For the detection of human-infecting

viral sequences, we implement *NVD* (New Virus Discovery), a workflow designed to

assemble putative viral sequences from human-infecting virus families and validate

assemblies to be used as references for mapping abundance values.

Methods

Sample collection

Influent wastewater from Columbia, MO was collected weekly as a 24-hour

composite sample using an ISCO autosampler. A 35 mL aliquot was transferred to sterile

50 mL conical and stored at 4°C prior to processing. At the time of collection, water quality

parameters including temperature, pH, total suspended solids, average influent flow over

sampling period, and chemical oxygen demand were recorded and provided by the

Missouri's Department of Health and Senior Services. A 35 mL aliquot of nuclease free

water was used as a negative control for each batch of samples processed after October

8th, 2024.

Viral concentration and RNA extraction for Metagenomics

Samples were centrifuged for 2000 xg at 4°C for 5 minutes and the supernatant was

filtered through 0.22 µm filters (Millipore, SCGP00525) to remove solids and larger

microbes. A subset of samples collected between January 2nd and March 19th, 2024 were

then treated with 10,000 ng of RNAse A (ThermoFisher, R1253), 2 units of TURBO DNAse

(Invitrogen, AM1907), and a 4.75 mL of 5 mM $CaCl_2$ and 25 mM $MgCl_2$ buffer, in an attempt

to remove bacterial ribosomal RNA. Separately, another sample subset was treated with

nucleases (as above) and 30 mM EDTA to dissociate rRNA from ribosomes. For nuclease

treated samples, 400 units of RNaseOUT™ Recombinant Ribonuclease Inhibitor

(Invitrogen, 10777019) was added prior to RNA extraction. A breakdown of treatments per

sample date is summarized in Supplemental Table 1.

Untreated, nuclease treated, and nuclease and EDTA treated sample filtrates were

mixed with 12.5 mL of 50% (w/vol) PEG (Research Products International, P48080-1000.0)

and 1.2M NaCl for 1 h at 4°C. Nucleic acids were extracted from pellets using QIAamp Viral

RNA Mini Kit (Qiagen, 52906) using the manufacturer's instructions excluding the addition

of carrier RNA. Samples were not DNase treated after extraction.

RT-qPCR of pepper mild mottle virus

Viral RNA in samples was assessed prior to library preparation through comparative

RT-qPCR targeting pepper mild mottle virus (PMMoV) to determine if viruses were

successfully concentrated. Pepper mild mottle virus was selected as an endogenous

control due to its ubiquitous distribution in wastewater [28]. Forward (5'

GGCGTAGATCCATTGGTGC 3') and reverse (5' CGAACCTTCCTCCTTTGATG 3') primers

were designed for RT-qPCR to target the PMMoV genome were mixed and resuspended at

100 µM concentrations. A Taqman probe (VIC-5' GCTGTGGTTTCAAATGAGAGTGG 3'-QSY)

was prepared at 100 µM concentration. The RT-qPCR reaction was carried out using

TaqPath 1-Step RT-qPCR Master Mix, CG (4x) (Applied Biosystems, A15299), forward and

reverse primers (500 nM each), taqman probe (125 nM), and 5 µL of RNA template for a

total reaction volume of 20 µL. Reactions were carried out on an Applied Biosystems 7500

Fast System, with thermocycler settings of 48°C for 15 min, 95°C for 10 min, followed by 40

cycles of 95°C for 15 s and 60°C for 1 min.

Library preparation and deep sequencing

Total RNA concentrations were quantified through Qubit RNA High Sensitivity

Assay, although some samples were below the detection limit for this assay. Libraries were

prepped from RNA for deep sequencing using Illumina's Stranded mRNA Prep, Ligation

(Illumina, 20040534) and Illumina RNA UD Indexes Sets A-D (Illumina, 20091655,

20091657, 20091659, 20091661) following manufacturers recommendations with slight

modifications listed below. RNA inputs for library preparation were normalized based on

Qubit results to 10-15 ng in 9.5 µL. If input concentrations were <0.2 ng/µl then 9.5 µL was

used for input. The mRNA capture step was omitted and library preps began with RNA

fragmentation and cDNA synthesis. Normalized RNA inputs were evenly split into two

aliquots. One aliquot was fragmented so longer RNAs were broken into a target range

compatible with sequencing, while the other half was left unfragmented to retain

sequences already within the target range. After fragmentation, aliquots were pooled

together for cDNA synthesis. Lastly, final indexed libraries underwent size selection during

the final cleanup by mixing 30 µL of AMPure XP (Beckman Coulter, A63881) with 50 µL of

post-indexing PCR amplicons to bind to long library sequences. Magnetic beads were

incubated on a magnetic stand for 5 minutes. The supernatant was separated from the

bead pellet and moved to a new tube where 20 µL of AMPure XP beads were added to bind

to shorter library sequences. These shorter fragments were washed, eluted in 30 µL of

resuspension buffer, and used for sample pooling. Final libraries were sequenced using a

NovaSeq 6000 until April 16[th] 2024, and subsequent libraries were sequenced using a

NovaSeq X with PE150 reads (Supplementary Table 1).

Assessing sequencing depth and rRNA abundance in samples

Analysis of rRNA abundance (described below) and qPCR of PMMoV showed no

noticeable deviation in percent rRNA or Ct values for our endogenous control in sample

dates that underwent multiple treatments. Therefore, treatment use was discontinued,

and fastq files from samples with multiple treatments per date were combined prior to

taxonomic assignment. Fastq files were trimmed for adapters, homopolymers, low entropy

sequences, and short length sequences using fastp (version 0.23.4) using the following

options, "-w 16 --dont_eval_duplication -l 50 -r -c --trim_poly_g --poly_g_min_len=5 --

trim_poly_x --poly_x_min_len=5 -y --adapter_sequence=CTGTCTCTTATACACATCT --

adapter_sequence_r2=CTGTCTCTTATACACATCT". Samples were split into rRNA and non-

rRNA fractions to assess the approximate depth of potential viral sequences. Ribosomal

RNA separation was performed using bbduk.sh from BBmap (version 38.18) and a custom

rRNA database curated by Brian Bushnell [29]. The depth of rRNA fractions and non-rRNA

fractions were determined using stats.sh from BBmap. Both BBTools commands were

performed using default settings.

Taxonomic assignment with GOTTCHA2.

Overall sample composition was determined with high confidence through a

species-level *GOTTCHA2* (version 2.1.9.0) analysis using fastp trimmed sequences and

arguments "-nc -rm no" [30]. *GOTTCHA2* was selected for reporting for definitive species

presence due to its curated multi-kingdom reference database which uses unique

genomic signatures at the queried taxonomic level. Using the arguments listed above, the

detection threshold for this analysis requires sequences to have a minimum linear

coverage of 0.005, minimum of 3 reads, minimum length of 60 bases, maximum estimated

z-score of 30 for the depth of mapped region, and a minimum mapped fraction of 50% to

the signature fragment.

Taxonomic assignments were grouped by species, read counts were summed for

each species, and read counts were normalized to reads per million (RPM). Host domains

for each detected viral phylum were manually annotated by searching host range of the

detected phylum. Sources and reasoning for annotations are listed in Supplemental Table

2. To ensure that community composition derived from *GOTTCHA2* is consistent with a

more permissive classifier, *Kraken2* was also used for taxonomic assignment. *Kraken2*

(version 2.1.3) was run on trimmed R1 and R2 files using the arguments "--use-names --

paired" and the PlusPFP database (accessed June 2024). *Kraken2* report read counts were

normalized to RPM and only reads classified on the species level were used for

comparison with *GOTTCHA2*. Unclassified reads from *GOTTCHA2* and *Kraken2* were not

included in outputs.

Taxonomic assignment with NVD.

We developed a Snakemake-based pipeline to detect and classify viral sequences

from high-throughput sequencing data. The pipeline was containerized using Apptainer

(formerly Singularity) to ensure reproducibility across computing environments. Raw

sequencing reads were first interleaved using BBMap (v38.18). To reduce computational

requirements and improve specificity, reads were pre-filtered using NCBI's Sequence

Taxonomy Analysis Tool (STAT; aligns_to v3.1.1) to retain only those with k-mer matches to

known human-infecting virus families, including *Adenoviridae*, *Anelloviridae*, *Arenaviridae*,

Astroviridae, *Bornaviridae*, *Peribunyaviridae*, *Caliciviridae*, *Coronaviridae*, *Filoviridae*,

Flaviviridae, *Hepadnaviridae*, *Hepeviridae*, *Orthoherpesviridae*, *Orthomyxoviridae*,

Papillomaviridae, *Paramyxoviridae*, *Parvoviridae*, *Picobirnaviridae*, *Picornaviridae*,

Pneumoviridae, *Polyomaviridae*, *Poxviridae*, *Sedoreoviridae*, *Retroviridae*, *Rhabdoviridae*,

Togaviridae, and *Kolmioviridae*. *Arteriviridae* were also included due to their ability to infect

other primate species. Unclassified reads at this step were discarded.

The filtered reads were assembled using SPAdes (v4.0.0) in sewage mode. Full

genome reconstruction was not a goal of this study, but generation of assemblies would

create a reference database to be used for subsequent abundance mapping. The resulting

contigs underwent quality control using BBMask to remove low-complexity regions using

an entropy threshold of 0.9. Following masking, contigs were filtered to retain only

sequences with at least 200 consecutive non-N bases after trimming terminal N's, as

shorter sequences can interfere with subsequent classification steps. The quality-

controlled contigs then underwent a hierarchical classification process beginning with

analysis using a NCBI STAT's tree index database provided by Kenneth Katz (downloaded

from NCBI on 2024-08-30; tree_filter.dbss, tree_filter.dbss.annotation, tree_index.dbs).

From this first-pass classification, a focused taxa list was generated and used for a

second, more detailed classification using STAT's dbss database. This two-stage STAT

analysis used a STAT cutoff-percent value of 0.001% (the default is 0,01) to more

sensitively include matching contigs for subsequent BLAST analysis.

The BLAST-based classification employed a two-stage approach for maximum

sensitivity. Initially, contigs were analyzed using MEGABLAST against NCBI's nucleotide

database with default parameters to identify close matches to known sequences. Contigs

that remained unclassified after MEGABLAST underwent a more sensitive BLASTN search

using default parameters. The results from both BLAST stages were filtered to remove non-

viral matches and bacteriophages, then merged to create a comprehensive classification

of the viral contigs. The workflow does not enforce explicit detection thresholds for

alignment length, percent identity, coverage, or E-value, but instead relies on BLAST

defaults and returning the top 5 hits per contig and selects a single hit based on the lowest

E-value. To validate the assembled viral sequences and assess their coverage, the initially

filtered reads were mapped back to the identified viral contigs with minimap2 using the

options "-ax sr" (v2.1). Coverage calculations were performed using samtools (v1.20),

retaining only primary alignments for accurate depth estimation. A SQL script was

generated to query BLAST validated assemblies for human specific pathogen names and

group them based on tropism for use in heatmap generation. Similarly to *GOTTCHA2* read

counts, *NVD* read counts were normalized to RPM for each species detected.

NVD derived influenza H5N1 assemblies were aligned for visualization using

minimap2 on default settings with Influenza A virus

(A/Grackle/Texas/USL_047/2024(H5N1)) as a reference genome because 5 of the 9

assembled contigs were blast validated as this strain. Next, samtools was used to filter out

quality scores less than 30 and secondary alignments and index the alignment for viewing

with IGV (version 2.16.2). Influenza C assemblies were also aligned using the same method

and genome coverage over the course of the study was assessed with samtools using the

option "depth -aa". The most abundantly detected influenza C strain, influenza C virus

(C/Sri Lanka/68/2023), was used for reference.

Digital PCR targeting SARS-CoV-2

Raw samples were prepped in accordance with Missouri's SARS-CoV-2 wastewater

surveillance program. Twenty-four-hour composite samples as described above were

acquired at Missouri wastewater treatment facilities (WWTF) and stored at 4°C until

processing. After mixing the samples briefly by gently inverting the tubes, a total of 9.75 mL

of wastewater per sample was loaded into 24-deep-well plates. A total of 20 µL of the

previously described NL4-3 derived HIV with the CMV-driven puromycin-resistance gene

and lacking the *env*, *nef*, *vif*, and *vpr* genes were added to each sample as a processing

control [31]. Then, a total of 100 µL of Nanotrap Enhancement Reagent 2 (Ceres

Nanosciences, 10112-10) and 150 µL of Nanotrap Microbiome A Particles (Ceres

Nanosciences, 44202) were added to each sample to capture, concentrate, and enhance

downstream detection of viruses. A total of 5.6 µL of carrier RNA (Qiagen, 52906) was

mixed with 700 µL of Buffer AVL (Qiagen, 52906) on a separate 24-deep-well plate to lyse

viruses and help with RNA recovery from the wastewater samples. All plates were loaded

into the KingFisher Apex (ThermoFisher Scientific) and processed. After processing in the

KingFisher Apex, 700 µL of the lysate was loaded into the QIAcube Connect (Qiagen), and

RNA was extracted and eluted into a final volume of 60 µL using the QIAamp Viral RNA Mini

Kit (Qiagen, 52906), according to the manufacturer's instructions.

Forward (5' GACCCCAAAATCAGCGAAAT 3' and 5' TTACAAACATTGGCCGCAAA 3')

and reverse (5' TCTGGTTACTGCCAGTTGAATCTG 3' and 5' GCGCGACATTCCGAAGAA 3')

primers and TaqMan probes (FAM-5' ACCCCGCATTACGTTTGGTGGACC 3'-MGB-NFQ and

ABY-5' ACAATTTGCCCCCAGCGCTTCAG 3'-QSY) were designed for RT-dPCR to target the

N1 and N2 regions of N gene of SARS-CoV-2, respectively. The primers were resuspended

at 100 µM concentrations as TaqMan probe. The RT-dPCR reaction was carried out using

QIAcuity One Step Advanced Probe kits (Qiagen, 250131), forward and reverse primers

(500nM each), TaqMan probe (125 nM) and 10 µL of RNA template for a total reaction

volume of 40 µL in a QIAcuity Four digital PCR system, with thermocycler settings of 50°C

for 30 min, 95°C for 2 min, followed by 45 cycles of 95°C for 10 s and 55°C for 30 s, after

priming.

Statistical analysis.

Relative abundance of rRNA was assessed for correlation against collection

temperature, water pH, total suspended solids, average influent flow over sampling

period, and chemical oxygen demand using Spearman's Correlation Coefficient.

For *GOTTCHA2* assigned sequences, Shannon–Wiener diversity (H') at the species

level was calculated separately for prokaryotic-host and eukaryotic-host viruses in each

sample using the diversity function in the Vegan package (version 2.6.8) of R (version

4.4.2). Paired differences in H' between prokaryote-infecting and eukaryote-infecting

viruses were tested for normality by Shapiro–Wilk test, then compared using a two-tailed

paired Student's t-test. A Wilcoxon signed-rank test was also performed as a

nonparametric alternative. A 95% confidence interval for the mean H' difference was

estimated by nonparametric bootstrap (10,000 replicates) via the boot package (v1.3-31).

Statistical significance was defined as $p < 0.05$.

Metagenomic read counts for SARS-CoV-2 were compared with dPCR reports

through Missouri's SARS-CoV-2 wastewater surveillance program to determine if the

metagenomic approach showed consistent epidemiological trends and was consistent

with our current SARS-CoV-2 surveillance protocol. For each sampling week, dPCR results

were flow normalized by multiplying mean copies per liter by average influent flow during

the sampling period to calculate total SARS-CoV-2 viral load. The metagenomic SARS-

CoV-2 (NC_045512.2), PMMoV (NC_003630.1) and ToBRFV (KT383474.1) sequences were

extracted from trimmed fastq files using minimap2 (version 2.14-r883) using the argument

"-ax sr". Secondary alignments and matches with MAPQ scores <30 were removed using

samtools (version 1.20). The association between flow-normalized dPCR or with SARS-

CoV-2 counts normalized to RPM, PMMoV, or ToBRFV were quantified in R using Pearson's

correlation (two-sided).-For visualization only, the metagenomic counts were linearly

rescaled to overlay the dPCR results on a secondary y-axis.

For *NVD* assigned sequences and *GOTTCHA2* sequences assigned to the same viral

families, beta diversity was longitudinally assessed on the species level through Bray-

Curtis dissimilarity of both outputs using the vegdist function from the vegan package in R.

Permutational analysis of variance (PERMANOVA) was performed to evaluate the effects of

season on the full data set. Since samples were collected longitudinally from a single site,

results were interpreted as indicative of overall compositional differences rather than as

strictly independent statistical tests, recognizing the potential for temporal autocorrelation

among samples. Seasons were assigned based on solstice and equinox dates.

PERMANOVA was also used to evaluate season, year, and their interaction on a subset of

data focusing on winter and spring. Non-Metric Multidimensional Scaling (NMDS) was

performed with k = 2 dimensions and ordination was repeated with multiple random starts

(up to 200 iterations) to ensure stable solutions. The first two NMDS axes were plotted to

visualize community dissimilarity.

Results

Sequencing depth and sample quality

Raw 24-hour composite Influent wastewater samples from Columbia, MO were

collected weekly from January 2024 through June 2025. Samples were filtered, PEG

concentrated, and extracted nucleic acids were sequenced using untargeted ultra-deep

sequencing. While this study targeted RNA viruses due to the method used for library

preparation, purified nucleic acids contained both RNA and DNA, and DNA viruses were

included in the output. Over the study period, 85.8 billion read pairs were obtained with an

average depth of 1.1 billion read pairs per sample (**Figure 1**). On average, ribosomal RNA

(rRNA) accounted for 66.2 ± 20.7% of reads per sample (range 14.2% - 95.5%). No strong

correlations were observed ($|\rho| \geq 0.5$) between rRNA abundance and temperature, water

pH, total suspended solids, average influent flow rate over the sampling period, or

chemical oxygen demand associated with samples. However, high rRNA abundances are

expected in nonselective metagenomic sample preps [21, 22]. The average RT-qPCR Ct

value for the wastewater-ubiquitous plant virus PMMoV was 24.2 ± 1.1, providing evidence

of viral concentration, even when overall RNA concentrations were low or below the

detection limit. Negative control samples were prepped on a per-batch basis starting on

October 8th 2024 and used to rule out contamination of suspicious hits when assessing

taxonomic profiles.

Definitive taxonomic profile

We used *GOTTCHA2* to identify reads that matched unique genomic regions

specific to each taxonomic level. The objective of this classification method was not to

provide a readout of total sample composition, but provide a baseline for species that are

unambiguously present in each sample. The settings used for this analysis enforce a

stringent mapping threshold, ensuring that a read must match at least 30 bp of a signature

fragment. The average percentage of sequences that could be assigned at the species

level using *GOTTCHA2* was 3.79% ± 1.78% representing approximately 40 million reads per

sample. In nearly all samples, eukaryotic viruses were more abundant than prokaryotic

viruses (**Figure 2A**). The most abundant viral family, *Virgaviridae*, consists mostly of tomato

brown rugose fruit virus (ToBRFV), with PMMoV also consistently present throughout the

year but at lower levels (**Figures 2B and 2D**). Notably, other plant viruses such as tomato

mosaic virus (TMV) and cucumber green mottle mosaic virus (CGMMV) were detected

predominantly during US growing seasons (**Figure 2D**). A similar breakdown of abundant

prokaryotic virus families is shown in **Supplemental Figure 1**. Taxonomic assignment

using *Kraken2* generated community profiles nearly identical taxonomic distribution and

the same highly abundant viral families as *GOTTCHA2* (Supplemental Figure 2 and 3).

Although prokaryotic viruses were less abundant in most samples, the prokaryotic
virus fraction was much more diverse compared to the eukaryotic virus fraction
(**Supplemental Figure 4**). Mean Shannon diversity was 3.15 ± 0.42 for prokaryotic viruses
and 1.10 ± 0.24 for eukaryotic viruses with a paired mean difference of 2.05 (95% CI 1.94-
2.16) that was statistically significant (paired t-test $p < 2.2 \times 10^{-16}$, Wilcoxon $p = 2.51 \times 10^{-14}$).
If the distribution of species were even, then these values would roughly equate to
prokaryotic virus fractions having 8-fold more detectable species than the eukaryotic virus
fraction. However, this difference is primarily due to the strong ToBRFV signal skewing the
diversity of the eukaryotic virus fraction, causing subsequent underreporting of eukaryotic
viral diversity.

Comparison between dPCR and metagenomic read counts

Many metagenomic studies normalize read counts by an endogenous control, such
as PMMoV, so counts are more reflective of true viral signal trends and not skewed by
changes in wastewater flow rate or variability in sample preparation. Since ToBRFV was the
most abundant virus while also displaying the smallest coefficient of variance throughout
the study of the highly abundant viruses such as PMMoV, we compared SARS-CoV-2
metagenomic counts normalized to RPM, PMMoV, and ToBRFV to dPCR findings from
Missouri's wastewater surveillance program. SARS-CoV-2 counts normalized to ToBRFV
showed the best correlation with flow normalized dPCR measurements ($p = 4.2e^{-12}$, $R^2 =$
0.471) compared to RPM ($p = 1.8e^{-10}$, $R^2 = 0.41$) and PMMoV ($p = 4.5e^{-6}$, $R^2 = 24$; **Figure 3**).

Human virus profiles

To better investigate human infecting viruses, we developed and applied a custom
pipeline to extract putative human infecting virus reads, assemble them into contigs for
BLAST validation, and derive abundance values by mapping raw reads to assembled
contigs. This pipeline facilitates detection of a wide breadth of pathogens that are non-
standard in probe panels for wastewater or overlooked in clinical settings. Many viruses
and viral groups were detected across the study period, such as SARS-CoV-2,
enteroviruses, rhinoviruses, and parechovirus A (**Figure 4**) while other viruses and viral

serotypes were only present at distinct seasons or timeframes (**Figure 5**). For example,

during the winter and early spring months of both years, the classic respiratory seasonal

viruses such as influenza A virus, and human coronaviruses NL63 and OC43 were readily

detected, while during the summer months these viruses were virtually absent (**Figure 5**).

Some respiratory viruses appeared to surge during specific seasons. For instance, in both

2024 and 2025 influenza A virus was predominantly detected from January to March, while

parainfluenza virus 3 detection was primarily from March to June, which is consistent with

what is observed clinically [32, 33]. Enterovirus D68, on the other hand, was detected

primarily in the fall, consistent with clinical observations [34]. Some of the viruses

detected had a more complex distribution when the specific serotypes were considered.

For instance, rhinoviruses were detected year-round, with higher detection levels in the

winter and spring (**Figure 4**). However, when individual serotypes were analyzed, there

were distinct waves (**Figure 5**). While most rhinovirus serotypes were abundant in the late

spring months (e.g. A58, A18, and A68 in the spring of 2024), some serotypes surged in the

fall and early winter, such as rhinovirus C42 in late 2024. Notably, while rhinoviruses were

most pronounced in the spring, the season-specific serotypes were not the same from

year to year. For instance, rhinoviruses A61 and A30 were both prevalent in the spring of

2025, but were completely absent in the spring of 2024. Rhinovirus infections are rarely

typed, so it is difficult to say if these observations are consistent with what was seen

clinically.

The parechoviruses also displayed a year-to-year serotype change. In 2024,

parechovirus 5 was the main serotype detected in 2024, while parechovirus 3 was the

primary serotype in 2025. This is consistent with a report that demonstrated that

parechovirus 5 emerged as the dominant serotype in nearby Kansas City in 2024 [35].

Some of the human-infecting viruses detected were not expected. One example

was detection of the respiratory pathogen influenza C virus, an influenza type that

generally causes milder disease and is typically not included in respiratory panels (**Figures

4 and 5**). Influenza C virus was the most frequently detected species of influenza in the

winter of 2023/24, but did not return the following winter. Coverage ranged from 31-100%

across coding sequences, with all seven Influenza C genome segments represented

(Supplemental Figure 5). Another unexpected finding from wastewater was the detection

of influenza A virus subtype H5N1 from early March-May of 2024 (Supplemental Figure 6).

The assembled sequence matched the influenza H5N1 strain B3.13 found in the mammary

tissue of dairy cattle and reported by the USDA in late March 2024, followed by

dissemination to 16 states by the end of the year [36]. Additionally, our H5N1 detections

align to the same timeframe as detections throughout Texas wastewater [37]. While

Missouri did not have any reported H5N1 dairy herds, the signal was believed to be derived

from a local dairy that received milk for processing from a state with H5N1-infected herds.

This finding highlights the fact that this metagenomic approach can detect pathogen

surges that are likely to be missed by standard patient surveillance.

Viruses associated with gastrointestinal disease, such as norovirus genogroup II,

norovirus genogroup I, rotavirus A, astroviruses, Sapporo virus, human mastadenovirus F,

and human picobirnavirus were detected throughout the study (**Figure 6**). Some viruses,

such as rotavirus A and astrovirus MLB, displayed slight increases during colder months. A

month-long surge of hepatitis A virus was detected the summer of 2024, but the signal did

not return for the remainder of this study. As expected for wastewater, read counts for

enteric viruses were higher than detected respiratory viruses (**Figures 4-6**). Despite the

lower counts associated with respiratory viruses, all reported human-infecting viruses

were captured with sufficient coverage for raw reads to assemble with SPAdes, undergo

BLAST-based classification, and read-mapping–based coverage assessment. When

analyzed in terms of the relative abundance of viruses at the species level using only viral

families listed in *NVD* analysis, season explained 18.8% of the variance of sample

groupings for *NVD*, and 27.3% of the variance for *GOTTCHA2* ($p < 0.001$ for both methods;

Figure 7). Additionally, both year and the interaction between season and year were

significant ($p < 0.01$) for spring and winter suggesting that seasonal trends are mostly

consistent between years with slight shifts in community composition from year to year.

The seasonality associated with this data reflects known epidemiological patterns and

reinforces previous studies that show untargeted metagenomic sequencing captures

expected biological signals in wastewater samples.

Discussion

This single-site longitudinal feasibility study demonstrates that an untargeted

sequencing approach to detect human infecting viral sequences is possible from

wastewater samples. By generating approximately one billion reads per week, we were

able to detect a wide array of viruses, across a range of hosts, with sufficient sensitivity to

track temporal trends and detect low-abundant or clinically overlooked pathogens. A

major challenge in generating this type of data is the high proportion of rRNA sequences in

the final output. While these ribosomal signals dilute target signals, they also provide

important contextual information about the microbial communities, potential hosts, and

other species signatures in wastewater. Additionally, removing rRNA during sample

preparation introduces risks of removing target sequences. Given the low rate of

classification of non-rRNA sequences using standard databases, there is a demand to

generate a database of wastewater associated sequences for classifying future samples.

While low classification rates in this study are consistent with previous work, Guajardo-

Leiva, S. et al have recently shown that family level resolution is currently possible [24, 26,

38].

The detection of influenza C virus demonstrate that minimally biased sequencing

approaches can capture clinically overlooked pathogens in a time-sensitive manner. While

newly emerging HPAI strains such as H5N1 are available on commercially available

panels, such as the Twist Comprehensive Viral Research, this minimally biased

sequencing method recovered enough H5N1 sequences to facilitate partial assembly of

five genomic segments with sufficient sequence information to resolve the virus to the

B3.13 lineage. Additionally, the correlation between metagenomic read counts, and dPCR

suggest that this method provides similar trends to current methods already being used for

surveillance. The consistent detection of clinically relevant viruses such as SARS-CoV-2,

parechovirus A, rhinovirus A, norovirus genogroup II, norovirus genogroup I, rotavirus A,

astroviruses, Sapporo virus, human mastadenovirus F, and human picobirnavirus across

all seasons strengthens the value of metagenomics for tracking endemic viral transmission

in wastewater. Seasonal increases in virus abundance were observed for several of these

pathogens, such as influenza virus, enterovirus A, and rotavirus A during colder months,

which is consistent with established clinical trends. The similar human virus trends

observed by multiple different taxonomic assignment methods between seasons

reinforces the observations made in this study that are reflective of a wastewater dataset

that follows traditionally expected temporal trends [2]

Beyond human health, the wide breadth of other detectable species emphasizes

the expanded utility of metagenomic surveillance. We selected *GOTTCHA2* for definitive

unambiguous characterization because this method is well suited for highly

heterogeneous samples such as wastewater that contain a wide range of organisms such

as viruses, bacteria, archaea, fungi, plants, and animals. The low classification rates using

this method are a direct result of the fact that the reference database contains only unique

kmers, and therefore the percentage classified is an underestimation of the total

percentage of classifiable sequences. Other classifiers, such as *Kraken2* using the

PlusPFP database, have potential to classify 93-99% of rRNA sequences or 14-55% of non-

rRNA sequences across all taxonomy levels (data not shown). However, because that

method contains non-unique kmers in the reference database, classification is less

accurate. This ambiguity led to the decision to use *GOTTCHA2* for baseline

characterization in this study due to its low false positive rate.

The detection of highly abundant plant viruses such as ToBRFV and TMV highlights

this expanded utility in detecting unexpected sequences. While these viruses have

geographic distributions that overlap with Missouri, it is likely that these viruses are

introduced to wastewater through consumption of crops within *Solanaceae*, such as

tomato and peppers, or suggest disease outbreaks in nearby plant communities, and

further highlights the utility of an expanded breadth of detectable outbreaks when using

metagenomic approaches to detect and identify viruses. While this study provides

evidence that ToBRFV an optimal virus for use as an endogenous control in some

locations, since ToBRFV detection is dependent on food sources, it may not be appropriate

in all parts of the world with different diets and supply chains [26]. In addition to ToBRFV,

the detection of plant viruses in this study is consistent with previous work showing high

abundance of plant viruses in wastewater [23]. These results, in combination with human

viral detection, demonstrate that deep metagenomic sequencing can support timely,

minimally biased, and ecologically broad surveillance that supports the initiatives of

CDC's One Health. Heatmaps, similar to **Figures 4-6**, are publicly available so that trends

can be followed in real time during the continuation of this study (https://lungfish-

science.github.io/wastewater-dashboard/). The dataset produced here provides a starting

point for other research efforts to begin development of new bioinformatic workflows to

further improve the detection of impactful nucleic acid signatures in deep data.

As sequencing costs continue to decline and analytical workflows improve, these

findings could support broader implementation of this approach as a scalable solution for

public health preparedness and environmental monitoring. However, at current prices this

sequencing approach may not be cost effective for all surveillance programs. This study is

also limited due to single-site geographic representation. Additional sampling at new

locations is required to determine if these findings can be generalized, and to fully evaluate

scalability of this ultra deep sequencing method. Another limitation of this study is the

evaluation of only non-solid fractions of wastewater. While it is known that some viruses

are enriched in the solid fraction, we have not found an efficient method of separating viral

material from solids that is sufficiently pure for untargeted metagenomics. Evaluation of

the supernatant allowed us to utilize a PEG concentration method that adequately

concentrated viruses without the use of rRNA depletion from samples. While these

limitations restrict generalization of our findings, this study is a first look at how ultra deep

sequencing at this depth can be applied to wastewater surveillance programs. Lastly, the

data generated in this study can be analyzed many different ways, and analyses presented

here should be viewed as one of several possible perspectives rather than a complete

characterization.

Acknowledgments

Authors would like to thank the University of Missouri Genomics Core for technical support in weekly sequencing efforts and optimization of library preparation protocols. Additionally, authors would like to thank the Research Support Solutions in the Division of IT at the University of Missouri and the Center for High Throughput Computing at the University of Wisconsin-Madison for access to high performance computing infrastructure (Columbia MO DOI: https://doi.org/10.32469/10355/97710). We would also like to thank Tami Hansen and the wastewater treatment plant in Columbia, MO for weekly sample collection. This research was funded by Inkfish, and by Coefficient Giving through a grant to SecureBio.

Author Contributions

Clayton Rushford, Data curation, Formal analysis, Investigation, Methodology, Project Administration, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review and editing | Devon Gregory, Data curation, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – review and editing | Emma Copen, Data Curation, Investigation, Methodology, Resources, Writing – review and editing | Jonathan Naydenov, Methodology | Alejandro Tovar-Mendez, Methodology, Writing – review and editing | Po-E Li, Data curation, Formal analysis, Software | Rose S. Kantor, Data curation, Methodology, Validation, Writing – review and editing | Migun Shakya, Data curation, Methodology, Validation, Writing – review and editing | Nelson Ruth, Software | Patrick S. G. Chain, Software | Hsinyeh Hsieh, Investigation, Methodology, Resources, Writing – Review & Editing | Torin Hunter, Investigation, Methodology, Writing – Review & Editing | Josh Kome, Software | Matt Frank, Investigation | Terri Lyddon, Resources | Jeff Kaufman, Conceptualization, Funding Acquisition, Methodology, Project administration, Writing – Review & Editing | David H. O'Connor, Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Methodology, Resources, Software, Validation, Visualization, Writing – Original Draft, Writing – Review & Editing | Marc C. Johnson, Conceptualization, Funding Acquisition,

Methodology, Project Administration, Resources, Supervision, Validation, Writing –

Original Draft, Writing – Review & Editing.

Data Availability

Sequence data has been uploaded to the NCBI Sequence Read Archive under

accession number PRJNA1247874. The workflow for identifying human-infecting virus

sequences with high confidence is available at https://github.com/dhoconno/nvd/. The

workflow for *GOTTCHA2* analysis, and subsequent visualization of *NVD* and *GOTTCHA2*

results are available at

https://github.com/ClaytonRushford/Columbia_MO_viral_surveillance_2024-2025/.

Supplemental material

Table S1: Breakdown of treatments in each sample and sequencing platform used.

Table S2: Host annotation for *GOTTCHA2* detected phyla.

Conflict of Interest

Authors declare no conflict of interest.

Funding

569   **Figures and Descriptions**

572   **Figure 1)** Sequencing depth and rRNA abundance of Columbia MO wastewater samples

573   from January 2024 through June 2025. Blue bars represent total read counts for each

574   sample date. The x-axis indicates the first sample date of the corresponding month

575   labeled.

Figure 2) Normalized abundance (RPM) of *GOTTCHA2* classified viral reads out of total

classified viral reads, colored by host domain based on manual annotation (A). Normalized

abundance (RPM) of the seven most abundant eukaryotic infecting virus families (B).

Normalized abundance (RPM) of the seven most abundant families within eukaryotic

viruses after removal of the family *Virgaviridae* (C). Breakdown of the most abundant

species within the family *Virgaviridae* (D).

Figure 3) Comparison of SARS-CoV-2 read counts derived from metagenomics using

minimap2 against reference genomes and dPCR results from Missouri's Wastewater

Surveillance Program from January 2024 through June 2025. SARS-CoV-2 read counts

normalized to RPM (A), PMMoV (B), and ToBRFV (C) were scaled prior to plotting by dividing

the max value of clinical cases or total viral load by the max value of SARS-CoV-2 read

counts.

Figure 4) Normalized abundance (RPM) of human respiratory viruses in Columbia MO
wastewater from January 2024 to June 2025. Reads counts were derived by mapping raw
sequences to NVD assembled contigs and counts were normalized to mapped reads per
million (RPM) prior to plotting as opposed to mapped reads per kilobase per million (RPKM)
due to groupings of viruses with different genome lengths in some rows. A $\log_{10}$ color scale
was used for visualization to enhance dynamic range and improve interpretability of
variation across samples. Vertical dashed lines signify solstice and equinox dates to
highlight season timeframes.

Figure 5) Normalized abundance (RPM) of individual *NVD* assigned viruses associated with
human respiratory disease that exhibit a seasonal trend. Reads counts were derived by
mapping raw sequences to NVD assembled contigs and counts were normalized to
mapped reads per million prior to plotting to keep normalization consistent across
heatmaps. A $\log_{10}$ color scale was used for visualization to enhance dynamic range and
improve interpretability of variation across samples. Vertical dashed lines signify solstice
and equinox dates to highlight season timeframes.

Figure 6) Normalized abundance (RPM) of *NVD* assigned virus reads associated with
human gastrointestinal disease in Columbia MO wastewater from January 2024 to June
2025. Reads counts were derived by mapping raw sequences to NVD assembled contigs
and counts were normalized to mapped reads per million prior to plotting as opposed to
mapped reads per kilobase per million due to high level groupings of viruses with different
genome lengths. A $\log_{10}$ color scale was used for visualization to enhance dynamic range
and improve interpretability of variation across samples. Vertical dashed lines signify
solstice and equinox dates to highlight season timeframes.

Figure 7) Sample clustering using non-metric multidimensional scaling of Bray-Curtis
dissimilarity using human infecting virus counts derived from *NVD* and *GOTTCHA2*.
Seasonal groupings were significant based on PERMANOVA results (p < 0.001 for both
analysis).

Supplemental Figure 1) Normalized abundance (RPM) of *GOTTCHA2* classified viral

families with prokaryotic hosts.

Supplemental Figure 2) Normalized abundance (RPM) of *Kraken2* classified viral reads out

of total classified viral reads, colored by host domain based on manual annotation (A).

Normalized abundance (RPM) of the seven most abundant eukaryotic infecting virus

families (B). Normalized abundance (RPM) of the seven most abundant families within

eukaryotic viruses after removal of the family *Virgaviridae* (C). Breakdown of the most

abundant species within the family *Virgaviridae* (D).

678　**Supplemental Figure 3)** Normalized abundance (RPM) of *Kraken2* classified viral families

679　with prokaryotic hosts.

684 **Supplemental Figure 4**) Shannon-Wiener diversity index values for *GOTTCHA2* assigned

685 prokaryotic infecting (Blue) and eukaryotic infecting (Orange) virus groups. H' was analyzed

686 on the species level.

Supplemental Figure 5) Alignment of NVD derived Influenza C assemblies mapped to

coding sequences of Influenza C virus (C/Sri Lanka/68/2023) throughout the study period.

Alignments were visualized on IGV, and images were cropped to only display the reference

and coverage of the assembled contigs.

Supplemental Figure 6) Alignment of NVD derived H5N1 assemblies mapped to Influenza

A virus (A/Grackle/Texas/USL_047/2024(H5N1)). Alignments were visualized on IGV, and

images were cropped to only display the reference and coverage of the assembled contigs.

Sample dates associated with each contig are annotated in mm-dd-yyyy format. One

contig H5N1 contig (218bp, BLAST validated as Influenza A virus (A/cat/ID/24-015315-001-

original/2024(H5N1)) segment 3 polymerase) was filtered out from this alignment.

References

1. Grassly, N.C., A.G. Shaw, and M. Owusu, *Global wastewater surveillance for pathogens with pandemic potential: opportunities and challenges*. The Lancet Microbe, 2024.

2. Smith, M.F., et al., *Seasonality of respiratory, enteric, and urinary viruses revealed by wastewater genomic surveillance*. Msphere, 2024. **9**(5): p. e00105-24.

3. Kilaru, P., et al., *Wastewater surveillance for infectious disease: a systematic review*. American journal of epidemiology, 2023. **192**(2): p. 305-322.

4. Majumdar, M., et al., *Detection of Enterovirus D68 in wastewater samples from the United Kingdom during outbreaks reported globally between 2015 and 2018*. BioRxiv, 2019: p. 738948.

5. Jahn, K., et al., *Early detection and surveillance of SARS-CoV-2 genomic variants in wastewater using COJAC*. Nature Microbiology, 2022. **7**(8): p. 1151-1160.

6. Kim, N.-Y., et al., *Wastewater Knows Pathogen Spread: Analysis of Residential Wastewater for Infectious Microorganisms including SARS-CoV-2*. Infection & Chemotherapy, 2023. **55**(2): p. 214.

7. Kantor, R.S. and M. Jiang, *Considerations and Opportunities for Probe Capture Enrichment Sequencing of Emerging Viruses from Wastewater*. Environmental Science & Technology, 2024. **58**(19): p. 8161-8168.

8. Santos, A.F., et al., *Wastewater Metavirome Diversity: Exploring Replicate Inconsistencies and Bioinformatic Tool Disparities*. International Journal of Environmental Research and Public Health, 2025. **22**(5): p. 707.

9. Pogka, V., et al., *Targeted virome sequencing enhances unbiased detection and genome assembly of known and emerging viruses—the example of SARS-CoV-2*. Viruses, 2022. **14**(6): p. 1272.

10. McCall, C., et al., *Targeted metagenomic sequencing for detection of vertebrate viruses in wastewater for public health surveillance*. ACS ES&T Water, 2023. **3**(9): p. 2955-2965.

11. Tisza, M., et al., *Wastewater sequencing reveals community and variant dynamics of the collective human virome*. Nature Communications, 2023. **14**(1): p. 6878.

12. Zulli, A., et al., *Probe-Based Enrichment Sequencing Applied to Wastewater Surveillance Accurately Tracks Multiple Viral Respiratory Outbreaks*. ACS ES&T Water, 2025.

13. Javornik Cregeen, S., et al., *Sequencing-Based Detection of Measles in Wastewater: Texas, January 2025*. American Journal of Public Health, 2025(0): p. e1-e5.

14. Clark, J.R., et al., *Wastewater pandemic preparedness: Toward an end-to-end pathogen monitoring program*. Frontiers in Public Health, 2023. **11**: p. 1137881.

15. Thornton, M., et al., *Comparative wastewater virome analysis with different enrichment methods*. Water Research, 2025: p. 123985.

16. Liu, S., et al., *Analysis of metagenomic data*. Nature Reviews Methods Primers, 2025. **5**(1): p. 5.

17. Wetterstrand, K. *DNA Sequencing Costs: Data from the NHGRI Genome Sequencing Program*.

18. Rajput, V., et al., *Metagenomics based longitudinal monitoring of antibiotic resistome and microbiome in the inlets of wastewater treatment plants in an Indian megacity*. Science of The Total Environment, 2025. **986**: p. 179691.

19. Becsei, Á., et al., *Time-series sewage metagenomics distinguishes seasonal, human-derived and environmental microbial communities potentially allowing source-attributed surveillance*. Nature communications, 2024. **15**(1): p. 7551.

20. Child, H.T., et al., *Comparison of metagenomic and targeted methods for sequencing human pathogenic viruses from wastewater*. MBio, 2023. **14**(6): p. e01468-23.

21. Li, Y., et al., *A broad wastewater screening and clinical data surveillance for virus-related diseases in the metropolitan Detroit area in Michigan*. Human Genomics, 2024. **18**(1): p. 14.

22. Cantalupo, P.G., et al., *Raw sewage harbors diverse viral populations*. MBio, 2011. **2**(5): p. 10.1128/mbio. 00180-11.

23. Nieuwenhuijse, D.F., et al., *Setting a baseline for global urban virome surveillance in sewage*. Scientific reports, 2020. **10**(1): p. 13748.

24. Guajardo-Leiva, S., J. Chnaiderman, A. Gaggero, and B. Díez, *Metagenomic insights into the sewage RNA virosphere of a large city*. Viruses, 2020. **12**(9): p. 1050.

25. Gulino, K., et al., *Initial mapping of the New York City wastewater virome*. Msystems, 2020. **5**(3): p. 10.1128/msystems. 00876-19.

26. Guajardo-Leiva, S., et al., *From sewage to genomes: Expanding our understanding of the urban and semi-urban wastewater RNA virome*. Environmental Research, 2025. **276**: p. 121509.

27. Wyler, E., et al., *Pathogen dynamics and discovery of novel viruses and enzymes by deep nucleic acid sequencing of wastewater*. Environment International, 2024. **190**: p. 108875.

28. Kitajima, M., H.P. Sassi, and J.R. Torrey, *Pepper mild mottle virus as a water quality indicator*. NPJ Clean Water, 2018. **1**(1): p. 19.

29. Bushnell, B. *BBmap*. 2025 2025-06-05 [cited 2025 6-20]; Available from: https://sourceforge.net/projects/bbmap/.

30. Freitas, T.A.K., P.-E. Li, M.B. Scholz, and P.S. Chain, *Accurate read-based metagenome characterization using a hierarchical suite of unique signatures*. Nucleic acids research, 2015. **43**(10): p. e69-e69.

31. Robinson, C.A., et al., *Defining biological and biophysical properties of SARS-CoV-2 genetic material in wastewater*. Science of The Total Environment, 2022. **807**: p. 150786.

32. Russell, E., A. Yang, S. Tardrew, and M.G. Ison, *Parainfluenza virus in hospitalized adults: a 7-year retrospective study*. Clinical Infectious Diseases, 2019. **68**(2): p. 298-305.

33. Control, C.f.D. and Prevention, *Influenza activity in the United States during the 2023–2024 season and composition of the 2024–2025 influenza vaccine*. 2024.

34. Grizer, C.S., K. Messacar, and J.J. Mattapallil, *Enterovirus-D68–a reemerging non-polio enterovirus that causes severe respiratory and neurological disease in children*. Frontiers in Virology, 2024. **4**: p. 1328457.

797   35.   Dhar, D., et al., *Emergence of Parechovirus-A5 central nervous system infections in*
798   *children from Kansas City, Missouri, USA.* Journal of Clinical Virology, 2025: p.
799   105835.
800   36.   Campbell, A., K. Brizuela, and S.S. Lakdawala, *mGem: Transmission and exposure*
801   *risks of dairy cow H5N1 influenza virus.* MBio, 2025. **16**(3): p. e02944-24.
802   37.   Tisza, M.J., et al., *Virome Sequencing Detects H5N1 Avian Influenza in Wastewater*
803   *in Ten Cities.* The New England journal of medicine, 2024. **391**(12): p. 1157.
804   38.   Tamaki, H., et al., *Metagenomic analysis of DNA viruses in a wastewater treatment*
805   *plant in tropical climate.* Environmental microbiology, 2012. **14**(2): p. 441-452.

806