Untargeted longitudinal ultra deep metagenomic sequencing of wastewater provides

a comprehensive readout of expected and unexpected viral pathogens

Clayton Rushford[a], Devon Gregory[b], Emma Copen[b], Jonathan Naydenov[b], Alejandro Tovar-

Mendez[b], Po-E Li[c], Rose S. Kantor[d], Migun Shakya[c], Nelson Ruth[c], Patrick S. G. Chain[c],

Hsinyeh Hsieh[e], Torin Hunter[b], Josh Kome[b], Matt Frank[b], Terri Lyddon[b], Jeff Kaufman[f], David

H. O'Connor[g], Marc C. Johnson[b#]

Author Affiliations

[a]University of Missouri, Department of Veterinary Pathobiology

[b]University of Missouri, Department of Molecular Microbiology and Immunology

[c]B-GEN, Bioscience Division, Los Alamos National Laboratory

[d]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory

[e]University of Missouri, School of Natural Resources

[f]Nucleic Acid Observatory, SecureBio

[g]University of Wisconsin-Madison, Department of Pathology and Laboratory Medicine

[#]Corresponding Author: marcjohnson@health.missouri.edu

Abstract

Wastewater surveillance has become a powerful tool to monitor circulating viruses

at a community level. Currently, most wastewater surveillance efforts use target-based

approaches such as quantitative PCR techniques or hybrid capture. This study explores

the feasibility of using unbiased ultra-deep metagenomic sequencing as a comprehensive

approach to wastewater surveillance. To test this, composite influent wastewater samples

were collected weekly from January 2024 through June 2025. Unbiased sequencing was

performed on all samples with an average depth of 1.1 billion reads per sample. Human

enteric viruses such as rotaviruses, astroviruses, and noroviruses were detected at high

levels in virtually every sample. SARS-CoV-2 was also detected in most samples and the

counts per sample positively correlated with digital PCR (dPCR) measurements. Less

abundant respiratory pathogens such as influenza A and B, rhinoviruses,

parainfluenzaviruses, and human coronaviruses 229E, OC43, NL63, and HKU1 were also

regularly detected. However, those pathogens displayed distinct and reproducible winter

and spring seasonality. Several unexpected viruses were also detected, such as several

detections of highly pathogenic avian influenza H5N1 (HPAI H5N1) genotype B3.13, a

month-long surge of hepatitis A virus, and a large season-specific surge in influenza C

virus. The most abundant known virus detected was the *Tobamovirus* tomato brown

rugose fruit virus, which was present year-round at high abundance. However, other

tobamoviruses such as tomato mosaic virus were detected primarily in the late growing

season. This eighteen-month study highlights that unbiased ultra deep sequencing

enables detection of expected and unexpected viral pathogens without targeted

enrichment.

Importance

This study demonstrates that untargeted ultra deep metagenomic sequencing can

provide a comprehensive and scalable tool for wastewater surveillance of viral pathogens.

By generating approximately 1 billion reads per sample across 78 consecutive weeks, we

captured expected pathogens such as SARS-CoV-2, noroviruses, and influenza viruses.

Additionally, we captured unexpected viral signals such as influenza C and highly

pathogenic avian influenza H5N1. The wide range of viral taxa captured in this study also

displays epidemiologically relevant seasonality. We also observed a correlation between

metagenomic SARS-CoV-2 read counts and dPCR values to validate this method against

other wastewater surveillance methods currently in use. Our findings highlight how ultra

deep metagenomics can enhance pandemic preparedness, enable early detection of

novel or re-emerging viruses, and broaden the scope of One Health monitoring by

capturing human, animal, and plant viral signatures from a composite wastewater sample.

Keywords

Wastewater surveillance, metagenomic, deep sequencing, SARS-CoV-2, influenza

H5N1, influenza C, tomato brown rugose fruit virus, *NVD*, *GOTTCHA2*, OneHealth, public

health

Introduction

Wastewater surveillance is a powerful tool for public health that leverages

minimally biased sampling on a population level to facilitate pandemic preparedness by

providing early warnings for circulating pathogens in sampled communities based on

sequence data [1, 2]. This form of community-level sampling reduces biases inherent to

clinical surveillance, such as underreporting due to asymptomatic infections or limited

access to healthcare. Several labs have demonstrated this concept through reverse

transcription PCR (RT-PCR), real time quantitative PCR (RT-qPCR), and digital PCR (dPCR)

designed to target human-infecting pathogens such as SARS-CoV-2, influenza, and

Enterovirus spp. [3-5]. Rapid multiplex PCR panels are also available for clinically relevant

viruses and have been used for tracking multiple viruses in wastewater [6]. However, PCR-

based approaches require frequent modifications due to evolution around primer binding

sites and are limited to known targets [7]. Alternatively, probe-capture approaches

facilitate detection of a wider range of pathogens, have been utilized in wastewater

systems, and have potential to detect pathogens that differ by up to 40% from reference

genomes [8, 9]. Additionally, commercially available probe panels have been developed

and implemented for wastewater surveillance [10-14]. While PCR and probe-capture

approaches facilitate detection of viruses with *a priori* knowledge of genetic makeup, they

are unable to detect highly divergent, novel viral sequences that may be circulating in a

population.

Unbiased sequencing and metagenomics can detect a wider range of known and

unknown pathogens compared to targeted and hybrid capture methods if they receive

sufficient coverage. Additionally, pathogen concentration methods, such as polyethylene

glycol (PEG) precipitation and vacuum-based direct capture, have been shown to be the

most reproducible while improving depth and breadth of viral signals [15]. However, a deep

dataset is a prerequisite for unbiased metagenomic surveillance due to a low target-to-

nontarget ratio of sequences [16]. Fortunately, the rapidly decreasing cost of sequencing

has facilitated data generation at scales that would, until recently, have been infeasible

[17].

The goal of this study was to test the feasibility of viral surveillance using unbiased

metagenomic sequencing. A variety of previous studies have utilized metagenomic

approaches for screening antimicrobial resistance genes, microbiomes, and viromes in

wastewater [2, 18-26]. Most of these take place over a short period, yet a few longitudinal

studies have surveyed wastewater for longer than a year to determine how detectable

sequences change over time [11, 18, 27]. While previous studies have demonstrated the

efficacy of metagenomics for viral surveillance, our study is unique due to our combination

of longitudinal sampling (78 consecutive weeks), use of PEG precipitation for viral

concentration, and sequencing at a depth greater than previous work. To provide a high-

confidence multi-kingdom breakdown of sample composition, we assigned sequence

reads through *GOTTCHA2* (Genomic Origin Through Taxonomic CHAllenge) classification

using an expanded reference database. For the detection of human-infecting viral

sequences, we implement *NVD* (New Virus Discovery), a workflow designed to assemble

putative viral sequences from human-infecting virus families and validate assemblies.

Methods

Sample collection

Influent wastewater from Columbia, MO was collected weekly as a 24-hour composite sample using an ISCO autosampler. A 35 mL aliquot was transferred to sterile 50 mL conical and stored at 4°C prior to processing. At the time of collection, water quality parameters including temperature, pH, total suspended solids, average influent flow over sampling period, and chemical oxygen demand were recorded and provided by the Missouri's Department of Health and Senior Services.

Viral concentration and RNA extraction for Metagenomics

Samples were centrifuged for 2000 xg at 4°C for 5 minutes and the supernatant was filtered through 0.22 µm filters (Millipore, SCGP00525) to remove solids and larger microbes. A subset of samples collected between January 2nd and March 19th, 2024 were then treated with 10,000 ng of RNAse A (ThermoFisher, R1253), 2 units of TURBO DNAse (Invitrogen, AM1907), and a 4.75 mL of 5 mM $CaCl_2$ and 25 mM $MgCl_2$ buffer, in an attempt to remove bacterial ribosomal RNA. Separately, another sample subset was treated with nucleases (as above) and 30 mM EDTA to dissociate rRNA from ribosomes. For nuclease treated samples, 400 units of RNaseOUT™ Recombinant Ribonuclease Inhibitor (Invitrogen, 10777019) was added prior to extraction. A breakdown of treatments per sample date is summarized in Supplementary Table 1.

Untreated, nuclease treated, and nuclease and EDTA treated sample filtrates were mixed with 12.5 mL of 50% (w/vol) PEG (Research Products International, P48080-1000.0) and 1.2M NaCl for 1 h at 4°C. Nucleic acids were extracted from pellets using QIAamp Viral RNA Mini Kit (Qiagen, 52906) using the manufacturer's instructions excluding the addition of carrier RNA. Samples were not DNase treated after extraction.

RT-qPCR of pepper mild mottle virus

Viral RNA in samples was assessed prior to library preparation through comparative RT-qPCR targeting pepper mild mottle virus (PMMoV) to determine if viruses were

successfully concentrated. Pepper mild mottle virus was selected as an endogenous

control due to its ubiquitous distribution in wastewater [28]. Forward (5'

GGCGTAGATCCATTGGTGC 3') and reverse (5' CGAACCTTCCTCCTTTGATG 3') primers

were designed for RT-qPCR to target the PMMoV genome were mixed and resuspended at

100 µM concentrations. A Taqman probe (VIC-5' GCTGTGGTTTCAAATGAGAGTGG 3'-QSY)

was prepared at 100 µM concentration. The RT-qPCR reaction was carried out using

TaqPath 1-Step RT-qPCR Master Mix, CG (4x) (Applied Biosystems, A15299), forward and

reverse primers (500 nM each), taqman probe (125 nM), and 5 µL of RNA template for a

total reaction volume of 20 µL. Reactions were carried out on an Applied Biosystems 7500

Fast System, with thermocycler settings of 48°C for 15 min, 95°C for 10 min, followed by 40

cycles of 95°C for 15 s and 60°C for 1 min.

Library preparation and deep sequencing

Total RNA concentrations were quantified through Qubit RNA High Sensitivity

Assay, although some samples were below the detection limit for this assay. Libraries were

prepped from RNA for deep sequencing using manufacturers recommendations with slight

modifications. RNA inputs for library preparation were normalized based on Qubit results

to 10-15 ng in 9.5 µL. If input concentrations were <0.2 ng/µl then 9.5 µL was used for

input. The mRNA capture step was removed with library preps beginning with RNA

fragmentation and cDNA synthesis. Normalized RNA inputs were evenly split into two

aliquots. One aliquot was fragmented so longer RNAs were broken into a target range

compatible with sequencing, while the other half was left unfragmented to retain

sequences already within the target range. After fragmentation, aliquots were pooled

together for cDNA synthesis. Lastly, final indexed libraries underwent size selection during

the final cleanup by mixing 30 µL of AMPure XP (Beckman Coulter, A63881) with 50 µL of

post-indexing PCR amplicons to bind to long library sequences. Magnetic beads were

incubated for 5 minutes, and the supernatant was moved to a new tube where 20 µL of

AMPure XP beads were added to bind to shorter library sequences. These shorter

fragments were washed, eluted in 30 µL of resuspension buffer, and used for sample

pooling. Final libraries were sequenced using a NovaSeq 6000 until April 16[th] 2024, and

subsequent libraries were sequenced using a NovaSeq X with PE150 reads

(Supplementary Table 1).

Assessing sequencing depth and rRNA abundance in samples

Analysis by qPCR of PMMoV showed no noticeable deviation in Ct values for sample

dates that underwent multiple treatments. However upon analysis of rRNA abundance

(described below), treatments did not show any reliable reduction of rRNA in samples.

Therefore, treatment use was discontinued, and fastq files from samples with multiple

treatments per date were combined for analysis. Fastq files were trimmed for adapters,

homopolymers, low entropy sequences, and short length sequences using fastp (version

0.23.4) using the following options, "-w 16 --dont_eval_duplication -l 50 -r -c --trim_poly_g -

-poly_g_min_len=5 --trim_poly_x --poly_x_min_len=5 -y --

adapter_sequence=CTGTCTCTTATACACATCT --

adapter_sequence_r2=CTGTCTCTTATACACATCT". Samples were split into rRNA and non-

rRNA fractions to assess the approximate depth of potential viral sequences. Ribosomal

RNA separation was performed using bbduk.sh from BBmap (version 38.18) and a custom

rRNA database curated by Brian Bushnell [29]. The depth of rRNA fractions and non-rRNA

fractions were determined using stats.sh from BBmap. Both BBTools commands were

performed using default settings.

Taxonomic assignment with GOTTCHA2.

Overall sample composition was determined with high confidence through a

species-level *GOTTCHA2* (version 2.1.9.0) analysis using fastp trimmed sequences and

arguments "-nc -rm no" [30]. *GOTTCHA2* was selected for reporting for definitive sample

composition due to its curated multi-kingdom reference database which uses unique

genomic signatures at the queried taxonomic level. Taxonomic assignments were grouped

by species, read counts were summed for each species, and read counts were normalized

to reads per million (RPM). Host domains for each detected viral phylum were manually

annotated (Supplemental Table 2).

Taxonomic assignment with NVD.

We developed a Snakemake-based pipeline to detect and classify viral sequences

from high-throughput sequencing data. The pipeline was containerized using Apptainer

(formerly Singularity) to ensure reproducibility across computing environments. Raw

sequencing reads were first interleaved using BBMap (v38.18). To reduce computational

requirements and improve specificity, reads were pre-filtered using NCBI's Sequence

Taxonomy Analysis Tool (STAT; aligns_to v3.1.1) to retain only those with k-mer matches to

known human-infecting virus families, including *Adenoviridae*, *Anelloviridae*, *Arenaviridae*,

Astroviridae, *Bornaviridae*, *Peribunyaviridae*, *Caliciviridae*, *Coronaviridae*, *Filoviridae*,

Flaviviridae, *Hepadnaviridae*, *Hepeviridae*, *Orthoherpesviridae*, *Orthomyxoviridae*,

Papillomaviridae, *Paramyxoviridae*, *Parvoviridae*, *Picobirnaviridae*, *Picornaviridae*,

Pneumoviridae, *Polyomaviridae*, *Poxviridae*, *Sedoreoviridae*, *Retroviridae*, *Rhabdoviridae*,

Togaviridae, and *Kolmioviridae*. *Arteriviridae* were also included due to their ability to infect

other primate species.

The filtered reads were assembled using SPAdes (v4.0.0) in sewage mode. The

resulting contigs underwent quality control using BBMask to remove low-complexity

regions using an entropy threshold of 0.9. Following masking, contigs were filtered to retain

only sequences with at least 200 consecutive non-N bases after trimming terminal N's, as

shorter sequences can interfere with subsequent classification steps. The quality-

controlled contigs then underwent a hierarchical classification process beginning with

analysis using a NCBI STAT's tree index database provided by Kenneth Katz (downloaded

from NCBI on 2024-08-30; tree_filter.dbss, tree_filter.dbss.annotation, tree_index.dbs).

From this first-pass classification, a focused taxa list was generated and used for a

second, more detailed classification using STAT's dbss database. This two-stage STAT

analysis retained matches with greater than 0.001% confidence and formed the basis for

subsequent BLAST analysis.

The BLAST-based classification employed a two-stage approach for maximum

sensitivity. Initially, contigs were analyzed using MEGABLAST against NCBI's nucleotide

database with default parameters to identify close matches to known sequences. Contigs

that remained unclassified after MEGABLAST underwent a more sensitive BLASTN search

using default parameters. The results from both BLAST stages were filtered to remove non-

viral matches and bacteriophages, then merged to create a comprehensive classification

of the viral contigs. To validate the assembled viral sequences and assess their coverage,

the initially filtered reads were mapped back to the identified viral contigs using minimap2

(v2.1). Coverage calculations were performed using samtools (v1.20), retaining only

primary alignments for accurate depth estimation. A SQL script was generated to query

BLAST validated assemblies for human specific pathogen names and group them based on

tropism for use in heatmap generation. Similarly to *GOTTCHA2* read counts, *NVD* read

counts were normalized to RPM for each species detected.

Digital PCR targeting SARS-CoV-2

Raw samples were prepped in accordance with Missouri's SARS-CoV-2 wastewater

surveillance program. Twenty-four-hour composite samples as described above were

acquired at Missouri wastewater treatment facilities (WWTF) and stored at 4°C until

processing. After mixing the samples briefly by gently inverting the tubes, a total of 9.75 mL

of wastewater per sample was loaded into 24-deep-well plates. A total of 20 µL of the

previously described NL4-3 derived HIV with the CMV-driven puromycin-resistance gene

and lacking the *env*, *nef*, *vif*, and *vpr* genes were added to each sample as a processing

control [31]. Then, a total of 100 µL of Nanotrap Enhancement Reagent 2 (Ceres

Nanosciences, 10112-10) and 150 µL of Nanotrap Microbiome A Particles (Ceres

Nanosciences, 44202) were added to each sample to capture, concentrate, and enhance

downstream detection of viruses. A total of 5.6 µL of carrier RNA (Qiagen, 52906) was

mixed with 700 µL of Buffer AVL (Qiagen, 52906) on a separate 24-deep-well plate to lyse

viruses and help with RNA recovery from the wastewater samples. All plates were loaded

into the KingFisher Apex (ThermoFisher Scientific) and processed. After processing in the

KingFisher Apex, 700 µL of the lysate was loaded into the QIAcube Connect (Qiagen), and

RNA was extracted and eluted into a final volume of 60 µL using the QIAamp Viral RNA Mini

Kit (Qiagen, 52906), according to the manufacturer's instructions.

Forward (5' GACCCCAAAATCAGCGAAAT 3' and 5' TTACAAACATTGGCCGCAAA 3')

and reverse (5' TCTGGTTACTGCCAGTTGAATCTG 3' and 5' GCGCGACATTCCGAAGAA 3')

primers and TaqMan probes (FAM-5' ACCCCGCATTACGTTTGGTGGACC 3'-MGB-NFQ and

ABY-5' ACAATTTGCCCCCAGCGCTTCAG 3'-QSY) were designed for RT-dPCR to target the

N1 and N2 regions of N gene of SARS-CoV-2, respectively. The primers were resuspended

at 100 µM concentrations as TaqMan probe. The RT-dPCR reaction was carried out using

QIAcuity One Step Advanced Probe kits (Qiagen, 250131), forward and reverse primers

(500nM each), TaqMan probe (125 nM) and 10 µL of RNA template for a total reaction

volume of 40 µL in a QIAcuity Four digital PCR system, with thermocycler settings of 50°C

for 30 min, 95°C for 2 min, followed by 45 cycles of 95°C for 10 s and 55°C for 30 s, after

priming.

Statistical analysis.

Relative abundance of rRNA was assessed for correlation against collection

temperature, water pH, total suspended solids, average influent flow over sampling

period, and chemical oxygen demand using Spearman's Correlation Coefficient.

For *GOTTCHA2* assigned sequences, Shannon–Wiener diversity (H') at the species

level was calculated separately for prokaryotic-host and eukaryotic-host viruses in each

sample using the diversity function in the Vegan package (version 2.6.8) of R (version

4.4.2). Paired differences in H' between prokaryote-infecting and eukaryote-infecting

viruses were tested for normality by Shapiro–Wilk test, then compared using a two-tailed

paired Student's t-test. A Wilcoxon signed-rank test was also performed as a

nonparametric alternative. A 95 % confidence interval for the mean H' difference was

estimated by nonparametric bootstrap (10 000 replicates) via the boot package (v1.3-31).

Statistical significance was defined as $p < 0.05$.

Metagenomic read counts for SARS-CoV-2 were compared with dPCR reports
through Missouri's SARS-CoV-2 wastewater surveillance program to determine if the
metagenomic approach showed consistent epidemiological trends and was consistent
with our current SARS-CoV-2 surveillance protocol. For each sampling week, dPCR results
were flow normalized by multiplying mean copies per liter by average influent flow during
the sampling period to calculate total SARS-CoV-2 viral load. The metagenomic SARS-
CoV-2 (NC_045512.2) and ToBRFV (KT383474.1) sequences were extracted from trimmed
fastq files using minimap2 (version 2.14-r883) using the argument "-ax sr". Secondary
alignments and matches with MAPQ scores <30 were removed using samtools (version
1.20). The association between flow-normalized dPCR or with SARS-CoV-2 counts
normalized to RPM or with ToBRFV were quantified in R using Pearson's correlation (two-
sided). For visualization only, the metagenomic counts were linearly rescaled to overlay
the dPCR results on a secondary y-axis.

For *NVD* assigned sequences and *GOTTCHA2* sequences assigned to the same viral
families, beta diversity was longitudinally assessed on the species level through Bray-
Curtis dissimilarity of both outputs using the vegdist function from the vegan package in R.
Permutational analysis of variance (PERMANOVA) was performed to evaluate the effects of
season on the full data set. Since samples were collected longitudinally from a single site,
results were interpreted as indicative of overall compositional differences rather than as
strictly independent statistical tests, recognizing the potential for temporal autocorrelation
among samples. Seasons were assigned based on solstice and equinox dates.
PERMANOVA was also used to evaluate season, year, and their interaction on a subset of
data focusing on winter and spring. Non-Metric Multidimensional Scaling (NMDS) was
performed with k = 2 dimensions and ordination was repeated with multiple random starts
(up to 200 iterations) to ensure stable solutions. The first two NMDS axes were plotted to
visualize community dissimilarity.

Results

Sequencing depth and sample quality

Raw 24-hour composite Influent wastewater samples from Columbia, MO were

collected weekly from January 2024 through June 2025. Samples were filtered, PEG

concentrated, and extracted nucleic acids were sequenced using unbiased ultra-deep

sequencing. While this study targeted RNA viruses due to the method used for library

preparation, purified nucleic acids contained both RNA and DNA, and DNA viruses were

included in the output. Over the study period, 85.8 billion read pairs were obtained with an

average depth of 1.1 billion read pairs per sample (**Figure 1**). On average, ribosomal RNA

(rRNA) accounted for 66.2 ± 20.7% of reads per sample (range 14.2% - 95.5%). No strong

correlations were observed ($|\rho| \geq 0.5$) between rRNA abundance and temperature, water

pH, total suspended solids, average influent flow rate over the sampling period, or

chemical oxygen demand associated with samples. However, high rRNA abundances are

expected in nonselective metagenomic sample preps [21, 22]. The average RT-qPCR Ct

value for the wastewater-ubiquitous plant virus PMMoV was 24.2 ± 1.1, providing evidence

of viral concentration, even when overall RNA concentrations were low or below the

detection limit.

Definitive taxonomic profile

We used *GOTTCHA2* to identify reads that matched unique genomic regions

specific to each taxonomic level to provide a baseline for species that are unambiguously

present in each sample. The average percentage of sequences that could be assigned at

the species level using *GOTTCHA2* was 3.79% ± 1.78%. This represents approximately 40

million reads per sample, even though this excludes sequences that map to more than one

species. In nearly all samples, eukaryotic viruses were more abundant than prokaryotic

viruses (**Figure 2A**). The most abundant viral family, *Virgaviridae*, consists mostly of tomato

brown rugose fruit virus (ToBRFV), with PMMoV also consistently present throughout the

year but at lower levels (**Figures 2B and 2D**). Notably, other plant viruses such as tomato

mosaic virus (TMV) and cucumber green mottle mosaic virus (CGMMV) were detected

predominantly during US growing seasons (**Figure 2D**). A similar breakdown of abundant

prokaryotic virus families is shown in **Supplemental Figure 1**.

319   Although prokaryotic viruses were less abundant in most samples, the prokaryotic

virus fraction was much more diverse compared to the eukaryotic virus fraction

(**Supplemental Figure 2**). Mean Shannon diversity was 3.15 ± 0.42 for prokaryotic viruses

and 1.10 ± 0.24 for eukaryotic viruses with a paired mean difference of 2.05 (95% CI 1.94-

2.16) that was statistically significant (paired t-test $p < 2.2 \times 10^{-16}$, Wilcoxon $p = 2.51 \times 10^{-14}$).

If the distribution of species were even, then these values would roughly equate to

prokaryotic virus fractions having 8-fold more detectable species than the eukaryotic virus

fraction. However, this difference is primarily due to the strong ToBRFV signal skewing the

diversity of the eukaryotic virus fraction, causing subsequent underreporting of eukaryotic

viral diversity.

Comparison between dPCR and metagenomic read counts

330   Many metagenomic studies normalize read counts by an endogenous control, such

as PMMoV, so counts are more reflective of true viral signal trends and not skewed by

changes in wastewater flow rate or variability in sample preparation. Since ToBRFV was the

most abundant and consistent viral signal in our data, we compared SARS-CoV-2

metagenomic counts normalized to RPM and SARS-CoV-2 counts normalized against

ToBRFV to dPCR findings from Missouri's wastewater surveillance program. Both SARS-

CoV-2 counts in RPM and ToBRFV normalized counts were positively correlated with flow

normalized dPCR measurements ($R^2 = 0.416$ and $R^2 = 0.471$ respectively), showing

normalization slightly improved correlations (**Figure 3**).

Human virus profiles

340   To better investigate the viruses that infect humans, we developed and applied a

custom pipeline. Many viruses and viral groups were detected across the study period,

such as SARS-CoV-2, enteroviruses, rhinoviruses, and parechovirus A (**Figure 4**) while

other viruses and viral serotypes were only present at distinct seasons or timeframes

(**Figure 5**). For example, during the winter and early spring months of both years, the

classic respiratory seasonal viruses such as influenza A virus, and human coronaviruses

NL63 and OC43 were readily detected, while during the summer months these viruses

were virtually absent (**Figure 5**). Some respiratory viruses appeared to surge during specific

seasons. For instance, in both 2024 and 2025 influenza A virus was predominantly

detected from January to March, while parainfluenza virus 3 detection was primarily from

March to June, which is consistent with what is observed clinically [32, 33]. Enterovirus

D68, on the other hand, was detected primarily in the fall, consistent with clinical

observations [34]. Some of the viruses detected had a more complex distribution when the

specific serotypes were considered. For instance, rhinoviruses were detected year-round,

with higher detection levels in the winter and spring (**Figure 4**). However, when individual

serotypes were analyzed, there were distinct waves (**Figure 5**). While most rhinovirus

serotypes were abundant in the late spring months (e.g. A58, A18, and A68 in the spring of

2024), some serotypes surged in the fall and early winter, such as rhinovirus C42 in late

2024. Notably, while rhinoviruses were most pronounced in the spring, the season-specific

serotypes were not the same from year to year. For instance, rhinoviruses A61 and A30

were both prevalent in the spring of 2025, but were completely absent in the spring of

2024. Rhinovirus infections are rarely typed, so it is difficult to say if these observations are

consistent with what was seen clinically.

The parechoviruses also displayed a year-to-year serotype change. In 2024,

parechovirus 5 was the main serotype detected in 2024, while parechovirus 3 was the

primary serotype in 2025. This is consistent with a report that demonstrated that

parechovirus 5 emerged as the dominant serotype in nearby Kansas City in 2024 [35].

Some of the human-infecting viruses detected were not expected. One example

was detection of the respiratory pathogen influenza C virus, an influenza type that

generally causes milder disease and is typically not included in respiratory panels (**Figures**

4 and 5). Influenza C virus was the most frequently detected species of influenza in the

winter of 2023/24, but did not return the following winter. Another unexpected finding from

wastewater was the detection of influenza A virus subtype H5N1 from early March-May of

2024. The assembled sequence matched the influenza H5N1 strain B3.13 found in the

mammary tissue of dairy cattle and reported by the USDA in late March 2024, followed by

dissemination to 16 states by the end of the year [36]. Additionally, our H5N1 detections

align to the same timeframe as detections throughout Texas wastewater [37]. While

Missouri did not have any reported H5N1 dairy herds, the signal was believed to be derived

from a local dairy that received milk for processing from a state with H5N1-infected herds.

This finding highlights the fact that this metagenomic approach can detect pathogen

surges that are likely to be missed by standard patient surveillance.

Viruses associated with gastrointestinal disease, such as norovirus genogroup II,

norovirus genogroup I, rotavirus A, astroviruses, Sapporo virus, human mastadenovirus F,

and human picobirnavirus were detected throughout the study (**Figure 6**). Some viruses,

such as rotavirus A and astrovirus MLB, displayed slight increases during colder months. A

month-long surge of hepatitis A virus was detected the summer of 2024, but the signal did

not return for the remainder of this study. As expected for wastewater, read counts for

enteric viruses were higher than detected respiratory viruses (**Figures 4-6**). Despite the

lower counts associated with respiratory viruses, all reported human-infecting viruses

were captured with sufficient coverage to assemble enough reads to confidently assign

taxonomy.

When analyzed in terms of the relative abundance of viruses at the species level

using only viral families listed in *NVD* analysis, season explained 18.8% of the variance of

sample groupings for *NVD*, and 27.3% of the variance for *GOTTCHA2* ($p < 0.001$ for both

methods; **Figure 7**). Additionally, both year and the interaction between season and year

were significant ($p < 0.01$) for spring and winter suggesting that seasonal trends are mostly

consistent between years with slight shifts in community composition from year to year.

The seasonality associated with this data reflects known epidemiological patterns and

reinforces previous studies that show unbiased metagenomic sequencing captures

expected biological signals in wastewater samples.

Discussion

This study demonstrates the feasibility of an untargeted sequencing approach to

detect human infecting viral sequences in deep wastewater data. By generating

approximately one billion reads per week, we were able to detect a wide array of viruses,

across a range of hosts, with sufficient sensitivity to track temporal trends and detect low-

abundant or clinically overlooked pathogens. A major challenge in generating this type of

data is the high proportion of rRNA sequences in the final output. While these ribosomal

signals dilute target signals, they also provide important contextual information about the

microbial communities, potential hosts, and other species signatures in wastewater.

Additionally, removing rRNA during sample preparation introduces risks of removing target

sequences. Given the low rate of classification of non-rRNA sequences using standard

databases, there is a demand to generate a database of wastewater associated

sequences for classifying future samples. While low classification rates in this study are

consistent with previous work, Guajardo-Leiva, S. et al have recently shown that family

level resolution is currently possible [24, 26, 38]

The detection of influenza C virus, and H5N1 influenza demonstrate that minimally

biased sequencing approaches can capture novel and clinically overlooked pathogens in a

time-sensitive manner. Additionally, the correlation between metagenomic read counts,

dPCR, and clinical cases suggests that this method provides similar trends to current

methods already being used for surveillance. The consistent detection of clinically relevant

viruses such as SARS-CoV-2, parechovirus A, rhinovirus A, norovirus genogroup II,

norovirus genogroup I, rotavirus A, astroviruses, Sapporo virus, human mastadenovirus F,

and human picobirnavirus across all seasons strengthens the value of metagenomics for

tracking endemic viral transmission in wastewater. Seasonal increases in virus abundance

were observed for several of these pathogens, such as influenza virus, enterovirus A, and

rotavirus A during colder months, which is consistent with established clinical trends. The

similar human virus trends observed by two different taxonomic assignment methods

between seasons reinforces the observations made in this study that are reflective of a

wastewater dataset that follows traditionally expected temporal trends [2]

Beyond human health, the wide breath of other detectable species emphasizes the

expanded utility of metagenomic surveillance. We selected *GOTTCHA2* for definitive

baseline characterization because this method is well suited for highly heterogeneous

samples such as wastewater that contain a wide range of organisms such as viruses,

bacteria, archaea, fungi, plants, and animals. The low classification rates using this

method are a direct result of the reference database only containing unique kmers, and

our classified percentages underrepresent the total amount of classifiable sequences.

Other classifiers, such as Kraken2 using the PlusPFP database, have potential to classify

93-99% of rRNA sequences or 14-55% of non-rRNA sequences (data not shown). However,

because this method contains non-unique kmers in the reference database, classified

reads are not always accurate. This ambiguity led us to our decision to use *GOTTCHA2* for

baseline characterization in this study due to its low false positive rate.

The detection of highly abundant plant viruses such as ToBRFV and TMV highlights

this expanded utility in detecting unexpected sequences. While these viruses have

geographic distributions that overlap with Missouri, it is likely that these viruses are

introduced to wastewater through consumption of crops within *Solanaceae*, such as

tomato and peppers, or suggest disease outbreaks in nearby plant communities, and

further highlights the utility of an expanded breath of detectable outbreaks when using

metagenomic approaches to detect and identify viruses. While this study provides

evidence that ToBRFV may be prevalent enough to be used as an endogenous control,

similar to PMMoV, the presence of ToBRFV is not consistent across all studies [26]. In

addition to ToBRFV, the detection of plant viruses in this study is consistent with previous

work showing high abundance of plant viruses in wastewater [23]. These results, in

combination with human viral detection, demonstrate that deep metagenomic sequencing

can support timely, minimally biased, and ecologically broad surveillance that supports

the initiatives of CDC's OneHealth.

The dataset generated here has already proven valuable for the development of

foundation large language models that reduce computational burden for downstream

users, particularly those lacking high-performance computing resources, to detect low

abundant human viruses and putative chimeric viruses [39]. Additionally, similar

heatmaps, such as **Figures 4-6**, are publicly available so that trends can be followed in real

time during the continuation of this study. The dataset produced here provides a starting

point for other research efforts to begin development of new bioinformatic workflows to

further improve the detection of impactful nucleic acid signatures in deep data. As

sequencing costs continue to decline and analytical workflows improve, our findings

support broader implementation of this approach as a scalable solution for public health

preparedness and environmental monitoring.

Acknowledgments

Authors would like to thank the University of Missouri Genomics Core for technical

support in weekly sequencing efforts and optimization of library preparation protocols.

Additionally, authors would like to thank the Research Support Solutions in the Division of

IT at the University of Missouri and the Center for High Throughput Computing at the

University of Wisconsin-Madison for access to high performance computing infrastructure

(Columbia MO DOI: https://doi.org/10.32469/10355/97710). We would also like to thank

Tami Hansen and the wastewater treatment plant in Columbia, MO for weekly sample

collection.

Author Contributions

Clayton Rushford, Data curation, Formal analysis, Investigation, Methodology,

Project Administration, Software, Supervision, Validation, Visualization, Writing – original

draft, Writing – review and editing | Devon Gregory, Data curation, Formal analysis,

Investigation, Methodology, Software, Validation, Visualization, Writing – review and

editing | Emma Copen, Data Curation, Investigation, Methodology, Resources, Writing –

review and editing | Jonathan Naydenov, Methodology | Alejandro Tovar-Mendez,

Methodology, Writing – review and editing | Po-E Li, Data curation, Formal analysis,

Software | Rose S. Kantor, Data curation, Methodology, Validation, Writing – review and

editing | Migun Shakya, Data curation, Methodology, Validation, Writing – review and

editing | Nelson Ruth, Software | Patrick S. G. Chain, Software | Hsinyeh Hsieh,

Investigation, Methodology, Resources, Writing – Review & Editing | Torin Hunter,

Investigation, Methodology, Writing – Review & Editing | Josh Kome, Software | Matt Frank,

Investigation | Terri Lyddon, Resources | Jeff Kaufman, Conceptualization, Funding

Acquisition, Methodology, Project administration, Writing – Review & Editing | David H.

O'Connor, Conceptualization, Data Curation, Formal Analysis, Funding Acquisition,

Methodology, Resources, Software, Validation, Visualization, Writing – Original Draft,

Writing – Review & Editing | Marc C. Johnson, Conceptualization, Funding Acquisition,

Methodology, Project Administration, Resources, Supervision, Validation, Writing –

Original Draft, Writing – Review & Editing.

Data Availability

Sequence data has been uploaded to the NCBI Sequence Read Archive under

accession number PRJNA1247874. The workflow for identifying human-infecting virus

sequences with high confidence is available at https://github.com/dhoconno/nvd/. The

workflow for *GOTTCHA2* analysis, and subsequent visualization of *NVD* and *GOTTCHA2*

results are available at

https://github.com/ClaytonRushford/Columbia_MO_viral_surveillance_2024-2025/.

Supplemental material

Table S1: Breakdown of treatments in each sample and sequencing platform used.

Table S2: Host annotation for *GOTTCHA2* detected phyla.

Conflict of Interest

Authors declare no conflict of interest.

Funding

Funder	Grant	Author
Open Philanthropy		Jeff Kaufman
Inkfish		Marc Johnson
		David H. O'Connor

Figure 1) Sequencing depth and rRNA abundance of Columbia MO wastewater samples from January 2024 through June 2025. Blue bars represent total read counts for each sample date. The x-axis indicates the first sample date of the corresponding month labeled.

Figure 2) Relative abundance of *GOTTCHA2* classified viral reads out of total classified

viral reads, colored by host domain based on manual annotation (A). Relative abundance

of the seven most abundant eukaryotic infecting virus families (B). Relative abundance of

the seven most abundant families within eukaryotic viruses after removal of the family

Virgaviridae (C). Breakdown of the most abundant species within the family *Virgaviridae*

(D).

Supplemental Figure 1) Relative abundance of *GOTTCHA2* classified viral families with

prokaryotic hosts.

Supplemental Figure 2) Shannon-Wiener diversity index values for *GOTTCHA2* assigned

prokaryotic infecting (Blue) and eukaryotic infecting (Orange) virus groups. H' was analyzed

on the species level.

540   **Figure 3)** Comparison of SARS-CoV-2 read counts derived from metagenomics and dPCR

541   results from Missouri's Wastewater Surveillance Program from January 2024 through June

542   2025. SARS-CoV-2 read counts normalized to ToBRFV (A) or RPM (B) were scaled prior to

543   plotting by dividing the max value of clinical cases or total viral load by the max value of

544   SARS-CoV-2 read counts.

Figure 4) Relative abundance of human respiratory viruses in Columbia MO wastewater
from January 2024 to June 2025. Reads counts were collected by *NVD* and normalized to
mapped reads per million (RPM) prior to plotting as opposed to mapped reads per kilobase
per million (RPKM) due to groupings of viruses with different genome lengths in some rows.
A $\log_{10}$ color scale was used for visualization to enhance dynamic range and improve
interpretability of variation across samples. Vertical dashed lines signify solstice and
equinox dates to highlight season timeframes.

Figure 5) Distribution of individual *NVD* assigned viruses associated with human respiratory disease that exhibit a seasonal trend. Reads counts were normalized to mapped reads per million prior to plotting to keep normalization consistent across heatmaps. A $\log_{10}$ color scale was used for visualization to enhance dynamic range and improve interpretability of variation across samples. Vertical dashed lines signify solstice and equinox dates to highlight season timeframes.

Figure 6) Distribution of *NVD* assigned virus reads associated with human gastrointestinal disease in Columbia MO wastewater from January 2024 to June 2025. Reads counts were normalized to mapped reads per million prior to plotting as opposed to mapped reads per kilobase per million due to high level groupings of viruses with different genome lengths. A $\log_{10}$ color scale was used for visualization to enhance dynamic range and improve interpretability of variation across samples. Vertical dashed lines signify solstice and equinox dates to highlight season timeframes.

589 **Figure 7)** Sample clustering using non-metric multidimensional scaling of Bray-Curtis

590 dissimilarity using human infecting virus counts derived from *NVD* and *GOTTCHA2*.

591 Seasonal groupings were significant based on PERMANOVA results (p < 0.001 for both

592 analysis).

References

1. Grassly, N.C., A.G. Shaw, and M. Owusu, *Global wastewater surveillance for pathogens with pandemic potential: opportunities and challenges*. The Lancet Microbe, 2024.

2. Smith, M.F., et al., *Seasonality of respiratory, enteric, and urinary viruses revealed by wastewater genomic surveillance*. Msphere, 2024. **9**(5): p. e00105-24.

3. Kilaru, P., et al., *Wastewater surveillance for infectious disease: a systematic review*. American journal of epidemiology, 2023. **192**(2): p. 305-322.

4. Majumdar, M., et al., *Detection of Enterovirus D68 in wastewater samples from the United Kingdom during outbreaks reported globally between 2015 and 2018*. BioRxiv, 2019: p. 738948.

5. Jahn, K., et al., *Early detection and surveillance of SARS-CoV-2 genomic variants in wastewater using COJAC*. Nature Microbiology, 2022. **7**(8): p. 1151-1160.

6. Kim, N.-Y., et al., *Wastewater Knows Pathogen Spread: Analysis of Residential Wastewater for Infectious Microorganisms including SARS-CoV-2*. Infection & Chemotherapy, 2023. **55**(2): p. 214.

7. Kantor, R.S. and M. Jiang, *Considerations and Opportunities for Probe Capture Enrichment Sequencing of Emerging Viruses from Wastewater*. Environmental Science & Technology, 2024. **58**(19): p. 8161-8168.

8. Santos, A.F., et al., *Wastewater Metavirome Diversity: Exploring Replicate Inconsistencies and Bioinformatic Tool Disparities*. International Journal of Environmental Research and Public Health, 2025. **22**(5): p. 707.

9. Pogka, V., et al., *Targeted virome sequencing enhances unbiased detection and genome assembly of known and emerging viruses—the example of SARS-CoV-2*. Viruses, 2022. **14**(6): p. 1272.

10. McCall, C., et al., *Targeted metagenomic sequencing for detection of vertebrate viruses in wastewater for public health surveillance*. ACS ES&T Water, 2023. **3**(9): p. 2955-2965.

11. Tisza, M., et al., *Wastewater sequencing reveals community and variant dynamics of the collective human virome*. Nature Communications, 2023. **14**(1): p. 6878.

12. Zulli, A., et al., *Probe-Based Enrichment Sequencing Applied to Wastewater Surveillance Accurately Tracks Multiple Viral Respiratory Outbreaks*. ACS ES&T Water, 2025.

13. Javornik Cregeen, S., et al., *Sequencing-Based Detection of Measles in Wastewater: Texas, January 2025*. American Journal of Public Health, 2025(0): p. e1-e5.

14. Clark, J.R., et al., *Wastewater pandemic preparedness: Toward an end-to-end pathogen monitoring program*. Frontiers in Public Health, 2023. **11**: p. 1137881.

15. Thornton, M., et al., *Comparative wastewater virome analysis with different enrichment methods*. Water Research, 2025: p. 123985.

16. Liu, S., et al., *Analysis of metagenomic data*. Nature Reviews Methods Primers, 2025. **5**(1): p. 5.

17. Wetterstrand, K. *DNA Sequencing Costs: Data from the NHGRI Genome Sequencing Program*.

18. Rajput, V., et al., *Metagenomics based longitudinal monitoring of antibiotic*
resistome and microbiome in the inlets of wastewater treatment plants in an Indian
megacity. Science of The Total Environment, 2025. **986**: p. 179691.

19. Becsei, Á., et al., *Time-series sewage metagenomics distinguishes seasonal,*
human-derived and environmental microbial communities potentially allowing
source-attributed surveillance. Nature communications, 2024. **15**(1): p. 7551.

20. Child, H.T., et al., *Comparison of metagenomic and targeted methods for*
sequencing human pathogenic viruses from wastewater. MBio, 2023. **14**(6): p.
e01468-23.

21. Li, Y., et al., *A broad wastewater screening and clinical data surveillance for virus-*
related diseases in the metropolitan Detroit area in Michigan. Human Genomics,
2024. **18**(1): p. 14.

22. Cantalupo, P.G., et al., *Raw sewage harbors diverse viral populations*. MBio, 2011.
2(5): p. 10.1128/mbio. 00180-11.

23. Nieuwenhuijse, D.F., et al., *Setting a baseline for global urban virome surveillance in*
sewage. Scientific reports, 2020. **10**(1): p. 13748.

24. Guajardo-Leiva, S., J. Chnaiderman, A. Gaggero, and B. Díez, *Metagenomic insights*
into the sewage RNA virosphere of a large city. Viruses, 2020. **12**(9): p. 1050.

25. Gulino, K., et al., *Initial mapping of the New York City wastewater virome.*
Msystems, 2020. **5**(3): p. 10.1128/msystems. 00876-19.

26. Guajardo-Leiva, S., et al., *From sewage to genomes: Expanding our understanding*
of the urban and semi-urban wastewater RNA virome. Environmental Research,
2025. **276**: p. 121509.

27. Wyler, E., et al., *Pathogen dynamics and discovery of novel viruses and enzymes by*
deep nucleic acid sequencing of wastewater. Environment International, 2024. **190**:
p. 108875.

28. Kitajima, M., H.P. Sassi, and J.R. Torrey, *Pepper mild mottle virus as a water quality*
indicator. NPJ Clean Water, 2018. **1**(1): p. 19.

29. Bushnell, B. *BBmap*. 2025 2025-06-05 [cited 2025 6-20]; Available from:
https://sourceforge.net/projects/bbmap/.

30. Freitas, T.A.K., P.-E. Li, M.B. Scholz, and P.S. Chain, *Accurate read-based*
metagenome characterization using a hierarchical suite of unique signatures.
Nucleic acids research, 2015. **43**(10): p. e69-e69.

31. Robinson, C.A., et al., *Defining biological and biophysical properties of SARS-CoV-2*
genetic material in wastewater. Science of The Total Environment, 2022. **807**: p.
150786.

32. Russell, E., A. Yang, S. Tardrew, and M.G. Ison, *Parainfluenza virus in hospitalized*
adults: a 7-year retrospective study. Clinical Infectious Diseases, 2019. **68**(2): p.
298-305.

33. Control, C.f.D. and Prevention, *Influenza activity in the United States during the*
2023–2024 season and composition of the 2024–2025 influenza vaccine. 2024.

34. Grizer, C.S., K. Messacar, and J.J. Mattapallil, *Enterovirus-D68–a reemerging non-*
polio enterovirus that causes severe respiratory and neurological disease in
children. Frontiers in Virology, 2024. **4**: p. 1328457.

35.   Dhar, D., et al., *Emergence of Parechovirus-A5 central nervous system infections in children from Kansas City, Missouri, USA*. Journal of Clinical Virology, 2025: p. 105835.

36.   Campbell, A., K. Brizuela, and S.S. Lakdawala, *mGem: Transmission and exposure risks of dairy cow H5N1 influenza virus*. MBio, 2025. **16**(3): p. e02944-24.

37.   Tisza, M.J., et al., *Virome Sequencing Detects H5N1 Avian Influenza in Wastewater in Ten Cities*. The New England journal of medicine, 2024. **391**(12): p. 1157.

38.   Tamaki, H., et al., *Metagenomic analysis of DNA viruses in a wastewater treatment plant in tropical climate*. Environmental microbiology, 2012. **14**(2): p. 441-452.

39.   Liu, O., et al., *METAGENE-1: Metagenomic Foundation Model for Pandemic Monitoring*. arXiv preprint arXiv:2501.02045, 2025.

697