Title

Reduced brain entropy in migraine with partial restoration during attacks: A resting-state fMRI study

Authors

Saberi Majid[1], Dajung J Kim[1], Xiao-Su Hu[1], Alexandre F DaSilva[1*]

   1- Headache and Orofacial Pain Effort (H.O.P.E.), Biologic and Materials Science & Prosthodontics, University of Michigan School of Dentistry, Ann Arbor, MI, USA

Corresponding Author:

adasilva@umich.edu

Abstract

Migraine is a prevalent and disabling neurological disorder, characterized by difficulties in regulating headache activity, sensory processing, and cognitive-emotional states. Brain entropy quantifies the complexity of neural dynamics, where reduced entropy may reflect diminished neural adaptability, but its assessment with fMRI in migraine remains limited. Here, we examined alterations in brain entropy and their associations with clinical burden, attack timing, and symptomatology. Resting-state fMRI data were acquired from adults with episodic migraine, chronic migraine, and healthy controls. Following standard preprocessing, voxel-wise sample entropy was computed, and group differences were assessed using ANCOVA with age and sex as covariates. Associations with clinical burden and symptom measures were examined within affected regions. In chronic migraine, attack timing-related changes in entropy were further explored, and the largest Lyapunov exponent (LLE) was estimated to characterize chaotic dynamics underlying attack-related complexity changes. Migraine patients showed reduced entropy in sensory, attentional, and default mode regions compared to controls, most pronounced in chronic migraine. Lower entropy correlated with greater headache frequency and longer illness duration. In chronic migraine, entropy relatively increased during attacks in multisensory integration regions and was associated with positive and elevated LLEs, indicating partially restored complexity with weakly chaotic dynamics. Patients experiencing phonophobia and nausea also exhibited increased entropy in multisensory and default mode regions. Our findings demonstrate widespread reductions in brain entropy in migraine, reflecting impaired neural adaptability, whereas attacks may transiently restore complexity partially through chaotic dynamics. These results advance understanding of migraine pathophysiology and highlight potential targets for therapeutic intervention.

Keywords

Migraine; resting-state fMRI; brain entropy; neural complexity; Lyapunov exponent; brain dynamics

Highlights

- Migraine is marked by reduced brain entropy across sensory, attentional and default mode regions, which correlates with disease burden.
- Reduced entropy reflects constrained neural adaptability within affected regions.
- Migraine attacks transiently restore entropy, suggesting partial recovery of neural adaptability.
- Positive and elevated Lyapunov exponents indicate a shift toward weakly chaotic dynamics during attacks.
- Symptoms such as phonophobia and nausea are linked to increased entropy in multisensory integration and default mode regions.

Introduction

Migraine is a prevalent and disabling neurological disorder affecting over one billion individuals worldwide. Between 1990 and 2021, its prevalence rose by 58%, incidence by 42%, and disability-adjusted life years (DALYs) by 58%, underscoring its growing public health burden (Dang et al., 2025). Clinically, migraine is classified as a primary headache disorder characterized by recurrent attacks of moderate to severe headache, often unilateral and pulsating, lasting 4-72 hours, and commonly accompanied by symptoms such as nausea, photophobia, and phonophobia (Headache Classification Committee of the International Headache Society (IHS), 2018). Chronic migraine is defined as ≥15 headache days per month, with ≥8 days meeting migraine criteria, while episodic migraine involves fewer than 15 headache days monthly.

Recent research increasingly conceptualizes migraine not merely as a vascular condition but as a disorder of brain network dysfunction involving widespread disruptions in sensory, affective, and cognitive regulation (Ashina et al., 2021). Functional MRI studies have revealed altered connectivity and neural activations in several large-scale networks, including the default mode network, dorsal attention network, salience network, visual network, and limbic regions. Each of these networks contributes to core migraine symptoms such as sensory hypersensitivity, impaired attention, and emotional and mood disturbances (Zhou et al., 2023; Zhou et al., 2024). Atypical neural functionality in multisensory integration regions such as the superior parietal lobule and superior temporal gyrus further suggests increased sensitivity to negative emotional input and is associated with migraine frequency (Dong et al. 2023; Klamer et al., 2025). Additionally, alterations in functional connectivity across the preictal, ictal, and postictal phases support the characterization of migraine as a cyclic disorder (Esteves et al., 2025).

Despite substantial progress in characterizing large-scale network abnormalities in migraine, most functional MRI research has focused on static measures of brain activity. Recently, a growing body of research has begun to explore the dynamic properties of brain function in migraine, such as time-varying functional connectivity (Lee et al., 2019; Chen et al., 2023) and variability in BOLD signal fluctuations (Lim et al., 2021). However, sample entropy (Richman & Moorman, 2000), a measure of signal complexity reflecting adaptability and regional information processing, has been increasingly applied in fMRI (Wang et al., 2014) but remains rarely investigated in the context of migraine (Wang et al., 2022). This measure has been applied in resting-state fMRI studies of psychiatric disorders such as depression (Liu et al., 2022), anxiety (Fan et al., 2023), schizophrenia (Xue et al., 2019), and ADHD (Sokunbi et al., 2013), as well as neurological disorders including epilepsy (Zhou et al., 2019), multiple sclerosis (Zhou et al., 2016), chronic pain (Del Mauro et al., 2025), and Alzheimer's disease (Xue et al., 2018), revealing clinically relevant alterations across key networks. Brain entropy analysis could advance

understanding of migraine pathophysiology by characterizing the complexity of brain
dynamics across regions and phases and may help identify neural markers that bridge
mechanistic understanding with potential treatment targets.

While entropy effectively captures the degree of signal complexity, it cannot distinguish
whether complexity arises from structured, deterministic (chaotic) dynamics or
unstructured, stochastic sources. Chaos theory provides tools that address this distinction
(Sprott, 2003). One such measure is the largest Lyapunov exponent (LLE), which
describes how rapidly two nearly identical system states diverge over time in a
reconstructed phase space (wolf et al., 1985). In simpler terms, it indicates how sensitive
a system is to its initial conditions: even tiny differences at the start can lead to
increasingly different trajectories. This growing separation reflects dynamical instability,
small perturbations are amplified rather than dampened, thereby limiting how far into the
future the system's behavior can be predicted. When such instability occurs in a
deterministic system, it is a hallmark of chaotic dynamics.

Taken together, entropy and LLE provide complementary perspectives: entropy captures
the overall complexity of a signal, while LLE characterizes the nature of its underlying
dynamics (Stam, 2005). For example, simultaneous increases in entropy and LLE
suggest enhanced deterministic chaos, whereas high entropy with low LLE indicates that
the complexity is primarily stochastic. Although LLE has been widely applied in EEG
research, particularly in studies of epilepsy and seizure activity (Brari & Belghith, 2022;
Yakovleva et al., 2020), its application in fMRI remains limited despite its high potential
(Xie et al., 2008; Keiholz et al., 2020). In migraine, where abnormal sensory integration
may underlie susceptibility to attacks (Kernick, 2005), LLE could help reveal whether
altered brain dynamics reflect deterministic chaotic processes or stochastic variability,
providing mechanistic insights beyond what entropy alone can offer.

In this study, we investigated the complexity of resting-state brain activity in individuals
with episodic migraine, chronic migraine, and healthy controls using voxel-wise entropy
analysis of fMRI data. Our primary aim was to identify brain regions exhibiting altered
signal entropy in migraine, expecting these alterations to be amplified with increasing
disease severity. We hypothesized that the diagnosis of migraine would be associated
with reduced entropy, indicating constrained neural processing and diminished neural
adaptability, in sensory integration and cognitive-emotional cortical systems where
flexible functionality is critical.

To capture dynamic changes associated with migraine attacks, we further examined how
brain signal complexity varied across the ictal (during headache), post-ictal (soon after
headache), and interictal (outside attack) phases by relating entropy of affected regions
to the timing of the most recent attack in chronic migraine patients. We also computed
the LLE to complement the entropy analysis and further characterize the nonlinear

dynamics underlying migraine attacks. This dual-measure approach allows us to determine whether entropy changes reflect a shift toward enhanced deterministic chaos or a rise in stochastic, noise-dominated activity. We hypothesized that migraine attacks would be associated with increased entropy and elevated Lyapunov exponents, consistent with a state of greater dynamical instability and deterministic chaos. This transient shift may disrupt overly rigid neural states and partially restore adaptive processing in affected systems. Finally, we explored the association between entropy in affected regions and specific migraine-related symptoms, providing further insight into how signal complexity may relate to symptom expression in migraine.

Collectively, This work aims to advance understanding of migraine pathophysiology and highlight novel neural targets for therapeutic intervention.

Methods

Participants

Participants included adults aged 18-65 years diagnosed with episodic migraine, chronic migraine, or classified as healthy controls. All participants provided written informed consent, and the study received approval from the Institutional Review Board of the University of Michigan Medical School.

The chronic migraine group included individuals meeting ICHD-3β criteria for chronic migraine (Headache Classification Committee of the International Headache Society (IHS), 2018), defined as experiencing headaches on ≥15 days per month for more than 3 months, with ≥8 days per month having migraine features. Additional inclusion criteria included migraine onset before age 50, and a stable medication regimen for at least 3 months, if applicable. The episodic migraine group consisted of individuals experiencing between 1 and 14 migraine days per month for at least 3 months. These participants were recruited as part of an earlier study cohort and were scanned during the interictal period. All participants were required to have no history of opioid use in the past 6 months and no overuse of analgesics (defined as use on ≥15 days per month). Those with any other chronic pain condition, major psychiatric or neurological disorder, substance use, or MRI contraindications were excluded.

The healthy control group included individuals without a history of migraine, chronic pain, or systemic medical or psychiatric conditions. They were not taking any regular medications affecting the central nervous system and had no MRI contraindications.

Clinical Assessments

To assess migraine burden, participants with a migraine diagnosis reported the average
number of headache days per month and the duration of their illness, defined as the
number of years since migraine onset, during the baseline clinical intake visit.

At the time of the MRI scan, participants completed a structured Recent Migraine
Questionnaire, a study-specific instrument designed to capture the temporal and
symptomatic context of their most recent migraine episode. This questionnaire was
developed for this study to characterize migraine phase and symptoms in proximity to
scanning and included items on the time since last migraine attack (e.g., current, within
the past 12/24/48 hours, or more than 48 hours ago) and presence of symptoms such as
photophobia, phonophobia, and nausea/vomiting. This instrument was administered only
to the chronic migraine cohort; episodic migraine participants were recruited in an earlier
cohort in which recent migraine timing and ictal/interictal state were not systematically
collected.

MRI Data Acquisition

All imaging data were collected on a GE Discovery MR750 3T scanner using a Nova 32-
channel head coil. The protocol included resting-state fMRI (rs-fMRI), high-resolution T1-
weighted structural scans, and dual-echo fieldmaps for B0 distortion correction.

Resting-state fMRI data were acquired using a multiband EPI sequence (TR = 800 ms,
TE = 30 ms, flip angle = 52°, voxel size = 2.4 mm³, 90 × 90 matrix, multiband factor = 6,
duration ≈ 6 min). Participants kept their eyes open, fixated on a cross, and minimized
movement. T1-weighted images were acquired using a 3D BRAVO sequence (TR = 12.2
ms, TE = 5.2 ms, TI = 500 ms, flip angle = 15°, voxel size = 1 mm³) for anatomical
normalization and coregistration. Fieldmaps were collected using dual-echo gradient-
echo scans with opposing phase encoding (AP/PA), TR = 7.4 ms, TE = 80 ms, flip angle
= 90°, voxel size = 1.687 × 1.687 × 2.4 mm³. These were used to compute voxel
displacement maps for EPI distortion correction.

Preprocessing of fMRI Data

Resting-state fMRI data were preprocessed using FSL (Jenkinson et al., 2012), fMRIPrep
(Esteban et al., 2019), and AFNI (Cox, 1996). First, all images were converted to BIDS
format (Gorgolewski et al., 2016). To correct for susceptibility-induced distortions, B0
distortion correction was performed using the FSL TOPUP algorithm, based on dual-echo
fieldmaps acquired with opposing phase-encoding directions.

Following distortion correction, data were preprocessed using fMRIPrep, which included
head motion correction, co-registration to the T1-weighted structural image, spatial
normalization to MNI152 standard space, and brain segmentation. As part of this pipeline,

nuisance components, including principal components of white matter and cerebrospinal
fluid (aCompCor (Behzadi et al., 2007)) and six motion parameters, were automatically
extracted for denoising.

Next, additional denoising was performed using FSL commands to remove the six motion
parameters and aCompCor components. The denoised data were then temporally filtered
using AFNI commands, applying a bandpass filter from 0.01 to 0.1 Hz to retain biologically
relevant low-frequency resting-state fluctuations. Finally, spatial smoothing was applied
using FSL, with a Gaussian kernel of 6 mm full width at half maximum (FWHM).

All preprocessing steps were visually checked to ensure data quality and alignment.

Entropy Computation

To quantify the complexity of resting-state brain dynamics, we computed voxel-wise
sample entropy maps from the final smoothed 4D fMRI data (Richman & Moorman, 2000;
Wang et al, 2014). For each participant, this process yielded a single 3D entropy map
representing local signal irregularity across the brain. Sample entropy was computed
using a custom implementation in R with embedded C++ code for efficient processing.

For each voxel, the time series was extracted and entropy was calculated using an
embedding dimension (m) of 2 and a similarity threshold (r) equal to 20% of the standard
deviation of the voxel's time series. Entropy computation was restricted to voxels within
a binarized MNI152 brain mask to reduce computational load and exclude non-brain
regions. These individual entropy maps were subsequently used for voxel-wise group-
level statistical analyses to identify regional differences in signal complexity across study
groups. In this framework, lower entropy indicates more constrained neural dynamics,
while higher entropy reflects more adaptable neural dynamics.

Lyapunov Analysis

To further investigate the nonlinear dynamical characteristics underlying changes in
signal complexity in affected regions, we examined the LLE of their mean time series.
The LLE quantifies the average rate at which nearby trajectories in a reconstructed phase
space diverge over time, serving as a marker of deterministic chaos by capturing
sensitivity to initial conditions, as opposed to stochastic variability (Wolf et al., 1985;
Rosenstein et al., 1993).

The estimation of the LLE follows a previously published method (Xu et al., 2011). Time-
delay embedding was used to reconstruct phase space dynamics from the fMRI signal.
The optimal time delay and embedding dimension were determined using established
methods: average mutual information for delay selection and the false nearest neighbors

algorithm to identify the appropriate embedding dimension. To minimize the influence of
autocorrelation, a Theiler window was applied based on the signal's dominant frequency
component.

LLE values were then computed using the algorithm which estimates divergence over
time between neighboring trajectories (Rosenstein et al., 1993). This approach is robust
to noise and particularly well-suited for short physiological time series like those in fMRI.
In this framework, positive LLE values indicate divergence and dynamical instability
(consistent with chaos), values near zero indicate neutral or marginally stable dynamics,
and negative values indicate convergence toward stable attractors. Given the technical
limitations of short and noisy fMRI time series, the magnitude of LLE values is typically
small; therefore, in this study we interpret positive but low values as indicative of weakly
chaotic behavior rather than strong deterministic chaos (Bunimovich & Su, 2025;
Heiligenthal et al., 2011).

Statistical Analysis

Voxel-wise group comparisons of entropy were conducted using analysis of covariance
(ANCOVA), with group (chronic migraine, episodic migraine, healthy controls) as the main
factor and age and sex as covariates. To correct for multiple comparisons, a cluster-wise
threshold was applied to voxels with $p < 0.05$. Residuals from the voxel-wise models were
used with AFNI's tools to estimate a minimum cluster size for statistical significance.
Clusters were defined using a 26-voxel connectivity criterion, where voxels were
considered contiguous if they shared a face, edge, or corner, to control the family-wise
error rate (FWE) at $p < 0.05$. Four clusters exceeded the threshold (187 contiguous
voxels) and were identified using FSL tools for subsequent analyses.

Mean entropy values were extracted from significant clusters and used in between-group
post hoc analyses and correlations with clinical measures such as headache days per
month and illness duration.

To examine transient changes in signal complexity related to migraine attacks,
participants with chronic migraine were categorized based on the time since their most
recent attack: ictal, <12 hours, 12-24 hours, 24-48 hours, and >48 hours (interictal).
Spearman's rank correlations were performed between entropy values and time since
attack, and between LLEs and time since attack in the identified clusters.

To determine whether altered entropy during migraine attacks reflects deterministic chaos
or stochastic dominance, we evaluated the relationship between LLE and entropy values
in identified clusters using Spearman correlation. A strong positive correlation would
support a state of deterministic chaos, whereas a weak or negative correlation would
suggest entropy is primarily driven by stochastic processes.

For symptom-based analyses, independent-samples t-tests were conducted to compare
entropy values between participants with chronic migraine who did versus did not report
phonophobia, nausea, or vomiting during their most recent attack.

Statistical analyses were performed using R (R Core Team, 2025) and relevant packages,
including Rcpp (Eddelbuettel & François, 2011) for computational efficiency in entropy
computation, and oro.nifti (Whitcher et al., 2011) for neuroimaging data handling.
Visualizations were generated with ggplot2 (Wickham, 2016) for statistical plots and
MRIcronGL (Rorden, 2025) for brain maps.

Results

Demographic and Clinical Characteristics

The final sample included 66 participants: 25 with episodic migraine (EM), 15 with chronic
migraine (CM), and 24 healthy controls (HC). Mean age was 41.7 ± 14.4 years in CM,
30.2 ± 11.5 in EM, and 36.1 ± 16.1 in HC. The proportion of females was 93% in CM,
88% in EM, and 87% in HC. CM participants reported 18.3 ± 4.6 headache days per
month and an illness duration of 27.4 ± 15.1 years; EM participants reported 5.5 ± 3
headache days and 12.9 ± 8.6 years of illness. Symptom data were only collected
systematically in the CM group during the scanning session; among CM participants, 64%
reported photophobia, 21% phonophobia, and 43% nausea or vomiting. In total, 43% of
CM participants were scanned during an ictal period and all EM participants were
scanned during the interictal period.

Entropy decreases in Sensory, Attentional, and Default Mode Regions in Migraine.

Four distinct clusters exhibited altered entropy (all surviving cluster-level FWE correction,
Figure 1 and Table 1): the occipital cortex (OC), right supramarginal gyrus and superior
parietal lobule (rSMG+rSPL), precuneus and posterior cingulate cortex (PCu+PCC), and
medial prefrontal cortex (mPFC). In each of these regions, entropy was significantly
reduced in the migraine groups compared to healthy controls, with the most pronounced
reductions observed in the chronic migraine group (peak voxel statistics are reported in
Table 1).

Specifically, in the PCu+PCC regions, individuals with chronic migraine showed
significantly lower entropy than healthy controls, while those with episodic migraine
showed a similar trend that did not reach significance. Chronic migraine patients also had
significantly lower entropy than episodic migraine patients in both the PCu+PCC and
rSMG+rSPL regions, with a similar trend observed in the mPFC. These graded entropy
reductions are consistent with greater disease burden and may reflect increasing
constraint in neural dynamics in key sensory, attentional, and default mode regions.

Complementary group-level comparisons of regional entropy values corresponding to the
lower panel of Figure 1 are presented in Supplementary Table 1.

Figure 1. Group-level differences in regional brain entropy. Colored voxels in brain maps indicate group-level significant differences in signal entropy among chronic migraine, episodic migraine, and healthy control subjects. The color bar represents F-statistics from the analysis of covariance (ANCOVA). Voxels have passed correction for false positives and are categorized into four distinct regions, displayed in three viewing orientations. For each region, group comparisons are shown in the lower row. Horizontal dashed lines represent group-level entropy averages, while box plots indicate medians, interquartile ranges and confidence intervals. Post hoc significant differences are denoted by connecting lines and asterisks in box plots. (Abbreviations: OC - Occipital Cortex, rSMG+rSPL - right Supramarginal Gyrus and right Superior Parietal Lobule, PCu+PCC - Precuneus and Posterior Cingulate Cortex, mPFC - medial Prefrontal Cortex, CM - Chronic Migraine, EM - Episodic Migraine, HC - Healthy Control)

Region	Brain Area	Function	Brain Network	Cluster Size (Voxel)	Peak MNI (x, y, z)	Peak F	Cluster-level pFWE
OC	Occipital cortex	Visual processing	Visual network	675	(7, -99, 16)	13.06	1.9e-05
rSMG+rSPL	Right supramarginal gyrus and right superior parietal lobule	Sensory integration and attentional control	Dorsal attention network	471	(42, -63, 47)	13.29	1.62e-05

PCu+PCC	Precuneus and posterior cingulate cortex	Self-referential processing, episodic memory, and pain perception	Default mode network	335	(10, -56, 15)	10.58	1.13e-04
mPFC	Medial prefrontal cortex	Attentional modulation and emotional regulation of pain	Default mode network and limbic network	270	(-1, 56, -11)	11.89	4.38e-05

Table 1. Brain regions showing significantly reduced entropy in migraine patients compared to healthy controls.

Reduced Brain Entropy Correlates with Migraine Burden and Illness Duration.

Reduced entropy was significantly associated with migraine-related clinical measures across the four regions identified in Figure 1. As shown in Figure 2A, lower entropy values were significantly associated with greater number of headache days per month, particularly in the rSMG+rSPL (r = -0.46, p = 0.0026) and PCu+PCC (r = -0.36, p = 0.021). Additionally, longer illness duration (years with migraine) was significantly associated with lower entropy in most regions, including the rSMG+rSPL (r = -0.45, p = 0.0048), PCu+PCC (r = -0.49, p = 0.0015), and mPFC (r = -0.5, p = 0.0013) (Figure 2B). These findings suggest that chronic exposure to migraine is associated with more constrained regional neural dynamics.

Figure 2. Correlation analysis between brain entropy and general migraine measures across four brain regions with reduced entropy. Each column corresponds to a specific brain region, while the rows represent correlations between brain entropy and headache days per month **(A)**, and brain entropy and years of migraine **(B)**. Each panel displays a scatter plot with a fitted regression line based on data from individuals with migraine. Pearson's correlation coefficient and the corresponding p-value are reported, with significant correlations marked by an asterisk.

Migraine Attacks Transiently Elevate Brain Signal Entropy in Key Regions.

In chronic migraine, brain entropy varied by time since the most recent attack. Entropy was relatively higher during and shortly after migraine attacks, particularly in rSMG+rSPL (r = -0.55, p = 0.043) and PCu+PCC (r = -0.57, p = 0.034) (Figure 3). Although not significant, similar trends were noted in the OC (r = -0.37, p = 0.19) and mPFC (r = -0.20, p = 0.48). This pattern suggests that although the entropy levels are generally lower in individuals with migraine (inside or outside attacks), migraine attacks temporarily increase brain signal complexity, particularly in parietal and default mode regions involved in multisensory integration, self-referential thought and pain perception, which may reflect a temporary release from the constrained neural state of migraine.

Figure 3. Correlation analysis between brain entropy and the time since the last migraine attack in individuals with chronic migraine. Each column corresponds to a specific brain region that showed altered entropy compared to healthy controls. The x-axis shows categorized time intervals since the last migraine attack (migraine period), while the y-axis represents regional brain entropy. Each panel displays individual data points, a fitted regression line, and the 95% confidence interval. Spearman's correlation coefficient (r) and the corresponding p-value are reported, with significant correlations (p < 0.05) marked by an asterisk.

Weakly Chaotic Dynamics Underlie Transient Increases in Brain Entropy During Migraine Attacks.

In chronic migraine, LLE values in the rSMG+rSPL were negatively correlated with time since last attack (r = -0.66, p = 0.01), indicating greater chaotic dynamics near the time of attack (Figure 4A). A marginal positive correlation was observed between LLE and entropy in this region (r = 0.50, p = 0.057), suggesting that entropy increases may reflect underlying chaos rather than noise (Figure 4B). Critically, LLE values in this cluster were significantly greater than zero (one-sample t-test vs. 0: t = 0.012, p = 5.75e-05), confirming the presence of diverging trajectories consistent with deterministic chaos. However, the small magnitude of these exponents (<0.1), indicates this is a state of weak chaos or dynamical instability. No such correlation between entropy and LLE was found in controls or episodic migraine participants scanned during the interictal period (Figure 4C),

 supporting the specificity of this weakly chaotic state to the attack phase in chronic
 migraine. These findings suggest that transient disruption of rigid neural states during
 migraine attacks is accompanied by weakly chaotic nonlinear dynamics.

Figure 4. Association between largest Lyapunov exponent (LLE), migraine period, and brain entropy in rSMG+rSPL region. (A) Correlation between LLE and time since the last migraine attack in chronic migraine. The x-axis shows categorized time intervals since the most recent attack, while the y-axis represents LLE values. **(B)** Correlation between brain entropy and LLE in chronic migraine. Each dot represents an individual, with red dots indicating patients who were in the ictal period of migraine at the time of MRI assessment. **(C)** Correlation between brain entropy and LLE across individuals with episodic migraine and health controls, discriminated by colors. Each panel includes a fitted regression line with 95% confidence intervals. Spearman's correlation coefficient (r) and the corresponding p-value are reported, with significant correlations ($p < 0.05$) marked by an asterisk.

330 *Migraine-Related Symptoms Are Associated with Elevated Entropy in Sensory and*
331 *Default Mode Regions.*

332 Although chronic migraine patients generally showed reduced entropy, those who
333 experienced phonophobia in recent attack showed relatively higher entropy in

Figure 5. Group differences in brain entropy based on migraine-associated symptoms across individuals with chronic migraine. (A) Comparison of brain entropy in the rSMG+rSPL region between individuals with and without phonophobia. (B) Comparison of brain entropy in the PCu+PCC region between individuals with and without nausea/vomiting. Entropy values are represented by notched box plots showing individual data points, group means (dashed lines), medians, interquartile ranges and confidence intervals. P-values from group comparisons are shown above each panel, with significant differences ($p < 0.05$) marked by an asterisk.

rSMG+rSPL (t = 2.51, p = 0.034) (Figure 5A), while those reporting nausea or vomiting
showed relatively higher entropy in the PCu+PCC (t = 2.67, p = 0.026) (Figure 5B). The
rSMG+rSPL is a key hub for multisensory integration and attentional reorienting, and
increased entropy in this region may reflect enhanced neural reactivity in response to
sound sensitivity during migraine. The PCu+PCC is part of the default mode network,
involved in self-referential thought and interoception. Altered internal state processing or
disrupted homeostatic regulation associated with visceral symptoms such as nausea may
contribute to elevated entropy in this region.

Discussion

Summary of Findings

In this study, we investigated local brain signal complexity in individuals with episodic
migraine, chronic migraine, and healthy controls using voxel-wise entropy and Lyapunov
exponent analyses of resting-state fMRI data. Our findings reveal that migraine is
associated with widespread reductions in brain entropy, particularly in regions involved in
sensory integration, visual processing, attentional and self-referential functions, and
affective regulation. These reductions were most pronounced in the chronic migraine
group and were related to clinical burden, suggesting a progressive constraint in neural
adaptability with increasing disease severity. Importantly, we found that entropy values
were transiently elevated closer to the time of a migraine attack, the ictal phase, especially
in the multisensory integration region, indicating a short-term and partial restoration of the
reduced entropy triggered by the attack. The transient ictal increase was further
characterized by elevated Lyapunov exponents, providing evidence that the observed
complexity changes reflect underlying weakly chaotic dynamics rather than purely
stochastic noise. This suggests that by inducing weakly chaotic dynamics, migraine
attacks destabilize rigid neural states and potentially reset adaptive neural processing.
Additionally, patients who experienced phonophobia showed increased entropy in
multisensory integration areas, while those reporting nausea or vomiting exhibited greater
entropy in default mode regions, supporting a link between symptom expression and
disruption of migraineurs rigid neural processing. These findings collectively highlight the
association between regional brain complexity and migraine burden, symptom
expression, and migraine phase.

Cross-Modal Support from EEG Studies

Our findings align with an EEG study that reported reduced entropy during the interictal
phase in migraine compared to controls (Cao et al., 2017). Their reductions were limited
to the prefrontal cortex, with no changes in occipital regions. In partial agreement, our
fMRI results showed reduced entropy in the medial prefrontal cortex but also in occipital
regions, contrasting with their findings. We further identified reductions in areas linked to

sensory integration, attention, self-referential, and pain-related processing. While EEG
captures rapid scalp-level fluctuations, our voxel-wise fMRI analysis offers higher spatial
resolution. These cross-modal findings underscore the value of signal complexity
measures in tracking the spatial and temporal dynamics of migraine.

Comparisons with fMRI-Based Entropy Studies in Migraine

Although we did not differentiate migraine with and without aura, our findings both align
with and diverge from a recent fMRI study on patients without aura (Wang et al., 2022).
That study used multiscale entropy and reported increased complexity across several
networks, including the default mode, frontoparietal, and sensorimotor, depending on
temporal scale, with reduced entropy in the posterior DMN at the slowest scale. In
contrast, we observed broadly reduced entropy, especially in association with migraine
chronicity. Notably, their analysis lacked correction for multiple comparisons and did not
account for covariate effects of age and sex, despite known effects of these factors on
entropy (Wang, 2021). These methodological and sample differences may explain the
discrepancies. Nonetheless, both studies support altered brain signal entropy in networks
linked to self-referential processing, sensory integration, and attentional control in
migraine.

Parallels with Epilepsy and Shared Entropy Mechanisms

Because both migraine and epilepsy are characterized by episodic alterations in cortical
excitability and synchronization, comparisons between them may reveal shared
mechanisms of disrupted neural dynamics. Our observation of reduced brain entropy in
migraine aligns with epilepsy studies showing decreased neural complexity during
interictal periods. Intracranial EEG has consistently demonstrated lower entropy in
epileptogenic versus non-epileptogenic regions (Protzner & Valiante, 2010), and gamma-
band analyses suggest this reflects excessive neural regularity due to synchronized
inhibitory activity (Sato et al., 2019). A resting-state fMRI study in right temporal lobe
epilepsy also reported interictal entropy reductions, particularly in attention and sensory
integration areas (Zhou et al., 2019). These cross-condition findings suggest that reduced
brain entropy may reflect region-specific disruptions in neural adaptability that are
common to episodic neurological disorders.

Seizure studies further show that ictal periods involve transient increases in entropy and
complexity, with elevated values observed across EEG measures such as distribution
entropy, sample entropy, and Lempel-Ziv complexity (Li et al., 2015; Jouny & Bergey,
2012). Similarly, we found increased entropy during migraine attacks, especially in
sensory integration regions, suggesting that such paroxysmal events may temporarily
disrupt rigid neural states. This supports the idea that both seizures and migraine attacks

can transiently restore brain complexity as a compensatory mechanism (Zhou et al.,
2016; Zhou et al., 2019).

Chaotic Dynamics During Migraine Attacks

In addition to entropy, our analysis of Lyapunov exponents offers deeper insight into the
nature of increased signal complexity during migraine attacks. LLE values rose in sensory
integration regions near the time of attack, indicating nonlinear dynamical instability. This
contrasts with seizure studies, where LLE typically decreases during ictal phases (Khoa
et al., 2012). Because fMRI BOLD signals are short and noisy, and the LLE values we
obtained were small and close to zero, we interpret these findings cautiously as evidence
for weakly chaotic instability rather than definitive proof of strong chaos. These results
suggest that migraine attacks may release the brain from rigid states through weakly
chaotic destabilization, enabling escape from pathological attractors.

Symptom-Related Brain Entropy Patterns

Our results demonstrate that individuals with chronic migraine who experienced
phonophobia during their most recent attack showed higher entropy in the right
supramarginal gyrus and superior parietal lobule, key multisensory integration regions.
This aligns with evidence that overlapping sensory hypersensitivities contribute to
migraine-related disability and highlight the role of multisensory processing in migraine
(Suzuki et al., 2021; Schwedt, 2013). Structural and functional changes in these regions
have also been linked to hypersensitivity and migraine attacks (Chong et al., 2016; Zhang
et al., 2022). Our observation of increased entropy associated with phonophobia in these
multisensory hubs is consistent with the notion that the migraine brain exhibits abnormal
multisensory processing.

We also observed increased entropy in the precuneus and posterior cingulate cortex
among individuals reporting nausea or vomiting. These regions are involved in
interoception and self-referential processing, and their function has been linked to nausea
(Farmer et al., 2015; Ruffle et al., 2019). Our findings suggest that increased signal
entropy in default mode regions may reflect disrupted homeostatic regulation during
migraine-associated nausea.

Limitations and Future Directions

This study has some limitations that should be considered. First, its cross-sectional
design limits inference about dynamic changes across the migraine cycle, including the
preictal phase. Second, the Recent Migraine Questionnaire was administered only in the
chronic migraine cohort, which restricts direct comparison of attack-related effects

between groups. Third, migraine with and without aura were analyzed together, meaning
potential subtype-specific differences could not be addressed in the present dataset.

Future research should employ longitudinal or time-locked designs to track preictal, ictal,
and postictal transitions in larger and more balanced migraine samples, enabling a clearer
understanding of dynamic changes across the migraine cycle. Neuromodulation studies
targeting regions with altered complexity may help determine whether restoring neural
regulation can reduce migraine burden. Finally, stratifying patients by aura subtype,
hormonal status, or sensory features may reveal distinct neurophysiological patterns and
guide more personalized interventions.

Conclusions

This study provides compelling evidence that migraine is characterized by altered brain
signal complexity, with widespread reductions in entropy during resting state, especially
in chronic cases. These reductions reflect constrained neural processing and less
adaptability in regions supporting sensory processing, attentional regulation, and self-
referential thought. Importantly, we observed that migraine attacks are associated with
transient increases in entropy that were characterized by elevated Lyapunov exponents,
confirming their origin in weakly chaotic neural dynamics rather than random noise. This
suggests a potential compensatory release from rigid neural states through a
deterministic chaotic destabilization. Symptom-specific variations in entropy further
emphasize the relevance of subjective experiences in shaping brain signal complexity in
migraine. Together, our findings highlight the utility of complexity analysis for capturing
dynamic brain alterations in migraine and point toward future directions for personalized
interventions.

Declarations

Acknowledgment

We gratefully acknowledge Jacqueline Dobson and Ayesha Kousar for their invaluable
contributions in project coordination, patient recruitment, and imaging support.

Data Availability

The datasets generated and analyzed during the current study are not publicly available
due to participant confidentiality and institutional regulations but are available from the
corresponding author on reasonable request.

Declaration of Conflicting Interests

The authors declare no potential conflicts of interest with respect to the research,
authorship, and/or publication of this article.

Ethics Approval and Consent to Participate

Funding

This research was supported by the National Institute of Neurological Disorders and
Stroke (NINDS) under grant numbers K23 NS062946 and R01 NS094413, awarded to
Dr. Alexandre F. DaSilva.

Authors' Contributions

Majid Saberi contributed to the conceptualization, data curation, formal analysis,
investigation, methodology, software development, validation, visualization, and writing
of both the original draft and subsequent review and editing of the manuscript. Dajung J.
Kim contributed to the investigation and to the review and editing of the manuscript. Xiao-
Su Hu contributed to software development and to the review and editing of the
manuscript. Alexandre F. DaSilva contributed to the conceptualization, methodology,
investigation, supervision, project administration, funding acquisition, resources,
validation, and to the review and editing of the manuscript.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT (OpenAI) in order to
improve the readability and language of the manuscript. After using this tool, the authors
reviewed and edited the content as needed and take full responsibility for the content of
the published article.

References

Ashina, S., Bentivegna, E., Martelletti, P., & Eikermann-Haerter, K. (2021). Structural and functional brain changes in migraine. *Pain and Therapy, 10*(1), 211-223.

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage, 37*(1), 90-101.

Brari, Z., & Belghith, S. (2022). A new algorithm for Largest Lyapunov Exponent determination for noisy chaotic signal studies with application to Electroencephalographic signals analysis for epilepsy and epileptic seizures detection. *Chaos, Solitons & Fractals, 165,* 112757.

Bunimovich, L. A., & Su, Y. (2025). Statistics of Weakly Chaotic Systems. arXiv preprint arXiv:2507.03530.

Cao, Z., Lai, K. L., Lin, C. T., Chuang, C. H., Chou, C. C., & Wang, S. J. (2018). Exploring resting-state EEG complexity before migraine attacks. *Cephalalgia, 38*(7), 1296-1306.

Chen, Z., Liu, H., Wei, X. E., Wang, Q., Liu, Y., Hao, L., ... & Rong, L. (2023). Aberrant dynamic functional network connectivity in vestibular migraine patients without peripheral vestibular lesion. *European Archives of Oto-Rhino-Laryngology, 280*(6), 2993-3003.

Chong, C. D., Starling, A. J., & Schwedt, T. J. (2016). Interictal photosensitivity associates with altered brain structure in patients with episodic migraine. *Cephalalgia, 36*(6), 526-533.

Cox, R. W. (1996). AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research, 29*(3), 162-173.

Del Mauro, G., Li, Y., Yu, J., Kochunov, P., Sevel, L. S., Boissoneault, J., ... & Wang, Z. (2025). Chronic pain is associated with greater brain entropy in the prefrontal cortex. *The Journal of Pain,* 105421.

Dong, L., Dong, W., Jin, Y., Jiang, Y., Li, Z., & Yu, D. (2025). The global burden of migraine: A 30-Year trend review and future projections by age, sex, country, and region. *Pain and Therapy, 14*(1), 297-315.

Dong, L., Fan, X., Fan, Y., Li, X., Li, H., & Zhou, J. (2023). Impairments to the multisensory integration brain regions during migraine chronification: correlation with the vestibular dysfunction. *Frontiers in Molecular Neuroscience, 16,* 1153641.

Eddelbuettel, D., & François, R. (2011). Rcpp: Seamless R and C++ integration. *Journal of Statistical Software, 40,* 1-18.

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., ... & Gorgolewski, K. J. (2019). fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature Methods, 16*(1), 111-116.

Esteves, I., Fouto, A. R., Ruiz-Tagle, A., Caetano, G., Nunes, R. G., da Silva, N.
A., ... & Figueiredo, P. (2025). Uncovering longitudinal changes in the brain
functional connectome along the migraine cycle: a multilevel clinical connectome
fingerprinting framework. *The Journal of Headache and Pain, 26*(1), 29.

Fan, S., Yu, Y., Wu, Y., Kai, Y., Wang, H., Chen, Y., ... & Tian, Y. (2023). Altered
brain entropy and functional connectivity patterns in generalized anxiety disorder
patients. *Journal of Affective Disorders, 332,* 168-175.

Farmer, A. D., Ban, V. F., Coen, S. J., Sanger, G. J., Barker, G. J., Gresty, M. A.,
... & Aziz, Q. (2015). Visually induced nausea causes characteristic changes in
cerebral, autonomic and endocrine function in humans. *The Journal of Physiology,*
593(5), 1183-1196.

Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P.,
... & Poldrack, R. A. (2016). The brain imaging data structure, a format for
organizing and describing outputs of neuroimaging experiments. *Scientific Data,*
3(1), 1-9.

Headache Classification Committee of the International Headache Society (IHS).
(2018). The international classification of headache disorders, 3rd edition.
Cephalalgia, 38(1), 1-211.

Heiligenthal, S., Dahms, T., Yanchuk, S., Jüngling, T., Flunkert, V., Kanter, I., ... &
Kinzel, W. (2011). Strong and weak chaos in nonlinear networks with time-delayed
couplings. Physical review letters, 107(23), 234102.

Hu, X. S., Hong, K. S., & Ge, S. S. (2011). Recognition of stimulus-evoked
neuronal optical response by identifying chaos levels of near-infrared
spectroscopy time series. *Neuroscience Letters, 504*(2), 115-120.

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M.
(2012). Fsl. *Neuroimage, 62*(2), 782-790.

Jouny, C. C., & Bergey, G. K. (2012). Characterization of early partial seizure
onset: Frequency, complexity and entropy. *Clinical Neurophysiology, 123*(4), 658-
669.

Keilholz, S., Maltbie, E., Zhang, X., Yousefi, B., Pan, W. J., Xu, N., ... & Guo, Y.
(2020). Relationship between basic properties of BOLD fluctuations and calculated
metrics of complexity in the human connectome project. *Frontiers in Neuroscience,*
14, 550923.

Kernick, D. (2005). Migraine—new perspectives from chaos theory. *Cephalalgia,*
25(8), 561-566.

Khoa, T. Q. D., Thi Minh Huong, N., & Toi, V. V. (2012). Detecting epileptic seizure
from scalp EEG using Lyapunov spectrum. *Computational and Mathematical*
Methods in Medicine, 2012(1), 847686.

Klamer, K., Craig, J., Haines, C., Sullivan, K., Seres, P., & Ekstrand, C. (2025).
Differential fMRI neural synchrony associated with migraine during naturalistic

stimuli with negative emotional valence. *The Journal of Headache and Pain, 26*(1), 62.

Lee, M. J., Park, B. Y., Cho, S., Park, H., Kim, S. T., & Chung, C. S. (2019). Dynamic functional connectivity of the migraine brain: a resting-state functional magnetic resonance imaging study. *Pain, 160*(12), 2776-2786.

Li, P., Yan, C., Karmakar, C., & Liu, C. (2015, August). Distribution entropy analysis of epileptic EEG signals. In *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)* (pp. 4170-4173). IEEE.

Lim, M., Jassar, H., Kim, D. J., Nascimento, T. D., & DaSilva, A. F. (2021). Differential alteration of fMRI signal variability in the ascending trigeminal somatosensory and pain modulatory pathways in migraine. *The Journal of Headache and Pain, 22*(1), 4.

Liu, X., Song, D., Yin, Y., Xie, C., Zhang, H., Zhang, H., ... & Yuan, Y. (2020). Altered Brain Entropy as a predictor of antidepressant response in major depressive disorder. *Journal of Affective Disorders, 260,* 716-721.

Protzner, A. B., Valiante, T., Kovacevic, N., McCormick, C., & McAndrews, M. P. (2010). Hippocampal signal complexity in mesial temporal lobe epilepsy: a noisy brain is a healthy brain. *Archives Italiennes de Biologie, 148*(3), 289-297.

R Core Team. (2025). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/

Richman, J. S., & Moorman, J. R. (2000). Physiological time-series analysis using approximate entropy and sample entropy. *American Journal of Physiology-Heart and Circulatory Physiology, 278*(6), H2039-H2049.

Rorden, C. (2025). MRIcroGL: voxel-based visualization for neuroimaging. *Nature Methods,* 1-2.

Rosenstein, M. T., Collins, J. J., & De Luca, C. J. (1993). A practical method for calculating largest Lyapunov exponents from small data sets. *Physica D: Nonlinear Phenomena, 65*(1-2), 117-134.

Ruffle, J. K., Patel, A., Giampietro, V., Howard, M. A., Sanger, G. J., Andrews, P. L., ... & Farmer, A. D. (2019). Functional brain networks and neuroanatomy underpinning nausea severity can predict nausea susceptibility using machine learning. *The Journal of Physiology, 597*(6), 1517-1529.

Sato, Y., Ochi, A., Mizutani, T., & Otsubo, H. (2019). Low entropy of interictal gamma oscillations is a biomarker of the seizure onset zone in focal cortical dysplasia type II. *Epilepsy & Behavior, 96,* 155-159.

Schwedt, T. J. (2013). Multisensory integration in migraine. *Current Opinion in Neurology, 26*(3), 248-253.

Sokunbi, M. O., Fung, W., Sawlani, V., Choppin, S., Linden, D. E., & Thome, J.
(2013). Resting state fMRI entropy probes complexity of brain activity in adults with
ADHD. *Psychiatry Research: Neuroimaging, 214*(3), 341-348.
Sprott, J. C. (2003). Chaos and time-series analysis. Oxford University Press.
Stam, C. J. (2005). Nonlinear dynamical analysis of EEG and MEG: review of an
emerging field. *Clinical Neurophysiology, 116*(10), 2266-2301.
Suzuki, K., Suzuki, S., Shiina, T., Okamura, M., Haruyama, Y., Tatsumoto, M., &
Hirata, K. (2021). Investigating the relationships between the burden of multiple
sensory hypersensitivity symptoms and headache-related disability in patents with
migraine. *The Journal of Headache and Pain, 22*(1), 77.
Wang, X., Zhang, Y., Qi, W., Xu, T., Wang, Z., Liao, H., ... & Zhao, L. (2022).
Alteration in functional magnetic resonance imaging signal complexity across
multiple time scales in patients with migraine without aura. *Frontiers in*
Neuroscience, 16, 825172.
Wang, Z. (2021). The neurocognitive correlates of brain entropy estimated by
resting state fMRI. *NeuroImage, 232,* 117893.
Wang, Z., Li, Y., Childress, A. R., & Detre, J. A. (2014). Brain entropy mapping
using fMRI. *PloS One, 9*(3), e89948.
Whitcher, B., Schmid, V. J., & Thorton, A. (2011). Working with the DICOM and
NIfTI Data Standards in R. *Journal of Statistical Software, 44,* 1-29.
Wickham, H. (2016). Data analysis. In *ggplot2: elegant graphics for data analysis*
(pp. 189-201). Cham: Springer International Publishing.
Wolf, A., Swift, J. B., Swinney, H. L., & Vastano, J. A. (1985). Determining
Lyapunov exponents from a time series. *Physica D: Nonlinear Phenomena, 16*(3),
285-317.
Xie, X., Cao, Z., & Weng, X. (2008). Spatiotemporal nonlinearity in resting-state
fMRI of the human brain. *Neuroimage, 40*(4), 1672-1685.
Xue, S. W., Guo, Y., & Alzheimer's Disease Neuroimaging Initiative. (2018).
Increased resting-state brain entropy in Alzheimer's disease. *Neuroreport, 29*(4),
286-290.
Xue, S. W., Yu, Q., Guo, Y., Song, D., & Wang, Z. (2019). Resting-state brain
entropy in schizophrenia. *Comprehensive Psychiatry, 89,* 16-21.
Yakovleva, T. V., Kutepov, I. E., Karas, A. Y., Yakovlev, N. M., Dobriyan, V. V.,
Papkova, I. V., ... & Krysko, V. A. (2020). EEG analysis in structural focal epilepsy
using the methods of nonlinear dynamics (Lyapunov exponents, Lempel-Ziv
complexity, and multiscale entropy). *The Scientific World Journal, 2020*(1),
8407872.
Zhang, L., Yu, W., Xu, M., Cui, F., Song, W., Yan, M., ... & Zhang, Z. (2022). The
hypothalamus may mediate migraine and ictal photophobia: evidence from
Granger causality analysis. *Neurological Sciences, 43*(10), 6021-6030.

Zhou, F., Zhuang, Y., Gong, H., Zhan, J., Grossman, M., & Wang, Z. (2016).
Resting state brain entropy alterations in relapsing remitting multiple sclerosis.
PloS One, 11(1), e0146080.

Zhou, M., Jiang, W., Zhong, D., & Zheng, J. (2019). Resting-state brain entropy in
right temporal lobe epilepsy and its relationship with alertness. *Brain and Behavior,*
9(11), e01446.

Zhou, Q., Zhao, R., Qin, Z., Qi, Y., Tang, W., Liu, L., ... & Du, X. (2024). Altered
intra-and inter-network functional activity among migraine, chronic migraine, and
trigeminal neuralgia. *Molecular Pain, 20,* 17448069241300939.

Zhou, Y., Gong, L., Yang, Y., Tan, L., Ruan, L., Chen, X., ... & Ruan, J. (2023).
Spatio-temporal dynamics of resting-state brain networks are associated with
migraine disability. *The Journal of Headache and Pain, 24*(1), 13.