Establishment and Quality Assessment of a Hospital-Associated Disability Database in Japan

Shinsuke Hori[a], Hidetaka Wakabayashi[b], Shinta Nishioka[c], Takako Nagai[b], Eiji Kose[d], Nao Hashida[e],

Kenta Ushida[a], Ryo Momosaki[a, f, g]

[a]Department of Rehabilitation, Mie University Hospital, Mie, Japan

[b]Department of Rehabilitation Medicine, Tokyo Women's Medical University Hospital, Tokyo, Japan

[c]Graduate School of Regional Design and Creation, University of Nagasaki, Nagasaki, Japan

[d]Department of Pharmacy, Juntendo University Hospital, Tokyo, Japan

[e]Swallowing Center, The University of Osaka Hospital, Osaka, Japan.

[f]Department of Rehabilitation Medicine, Mie University Graduate School of Medicine, Mie, Japan

[g]Scientific Research WorkS Peer Support Group (SRWS-PSG)

*Corresponding Author:

Ryo Momosaki, MD, MPH, PhD

Department of Rehabilitation Medicine, Mie University Graduate School of Medicine, 2-174 Edobashi

Tsu, Mie, 514-8507, Japan

Tel: +81-59-232-1111, Fax: +81-59-231-5012

E-mail: momosak@med.mie-u.ac.jp

Running title: Development of HAD Database in Japan

Keywords: database, hospital-associated disability, Japan, rehabilitation

Abstract

Objectives:

Hospital-associated disability (HAD) refers to a decline in activities of daily living (ADL) during

hospitalization and is associated with prolonged hospital stay, higher mortality, and increased readmission

rates among older adults. This study aimed to establish and assess the quality of a multicenter HAD

registry database that includes detailed data on nutrition, medication, swallowing, and walking ability.

Methods:

This prospective multicenter study was conducted at nine hospitals in Japan. Patients aged ≥70 years who

were independent in ADL (Katz Index = 6) before admission to a general ward for rehabilitation were

enrolled. Data were collected at admission and discharge using REDCap. Assessment items included

demographics, Katz Index, Functional Comorbidity Index, Clinical Frailty Scale, Global Leadership

Initiative on Malnutrition (GLIM) criteria, Food Intake LEVEL Scale (FILS), and Functional Ambulation

Categories. HAD was defined as a Katz Index score of less than 6 on discharge.

Results:

A total of 209 patients (median age = 79.0 [75–85] years; 60.8% male) were enrolled. The prevalence of

HAD, hospital-associated dysphagia, and weight loss during hospitalization was 29.1%, 16.7%, and

75.4%, respectively. Median length of stay was 19 days (IQR: 13–34), and 82.3% were discharged home.

While most variables had low levels of missing data, variables involving body weight demonstrated

comparatively higher missingness.

Conclusions:

A multicenter HAD registry database was successfully established, demonstrating the feasibility of prospective data collection across multiple institutions. This registry may facilitate future studies on risk factor identification, predictive model development, and early intervention strategies for HAD prevention.

Introduction

Hospital-Associated Disability (HAD) refers to decline in Activities of Daily Living (ADL) that occurs during hospitalization.[1] Once patients develop HAD, they often fail to regain their pre-hospitalization level of function after discharge.[2] Decreases in ADL associated with hospitalization include mobility impairments and limitations in ADL, such as bathing, dressing, eating, and toileting.[3] Another functional impairment associated with hospitalization is hospital-associated dysphagia.[4,5] Hospital-associated dysphagia occurs as the patient's general condition deteriorates during hospitalization, leading to a decline in swallowing ability and an increased risk of aspiration pneumonia.[6] HAD is particularly prevalent in older and frail patients and is associated with longer hospital stays, higher readmission prevalence and increased short- and long-term mortality.[7,8] In addition, the onset of HAD increases medical costs and the burden of care, posing a significant challenge to patients and to society as a whole.[9] Therefore, the need for HAD prevention and appropriate intervention is heightened.

Major risk factors for HAD include age, low activity during hospitalization, frailty, delirium, multiple medications, malnutrition, and prolonged hospitalization.[10,11] Reduced activity during hospitalization results in loss of muscle strength and physical fitness, making it challenging to perform ADL. Delirium is linked to reduced cognitive function and ADL, and it is known to negatively impact long-term prognosis. Moreover, polypharmacy increases the risk of adverse drug interactions and may contribute to the development of HAD. Furthermore, malnutrition after hospitalization can cause muscle weakness and inhibit recovery after discharge. Consequently, a variety of risks associated with HAD can be expected, and further research is necessary.

HAD, in a broad sense, refers to a decline in ADL related to hospitalization regardless of ADL status before hospitalization. On the other hand, narrowly defined HAD refers to the loss of independence in patients who were independent in ADL before hospitalization. Patients who were independent in ADL before hospitalization may require long-term care after losing their independence, which is a significant challenge in healthcare policy. The most critical point in conducting strict HAD research is accurately assessing the ADL status before hospitalization. However, previous databases often lack sufficient information on preadmission ADL, making it difficult to accurately identify HAD. Even prospectively collected databases have not accumulated detailed data on nutrition and medication use, ambulation, cognitive function, and oral intake ability.[12] The purpose of this study was to develop and evaluate the quality of a multicenter registry database on HAD, including detailed information on nutrition and medications, ambulation, and oral intake ability.

Materials and Methods

Mie University Hospital served as the leading research institute, and partner facilities were recruited from the Japanese Society of Rehabilitation Nutrition. A total of nine facilities participated and prospectively constructed the database. The database employs a web-based registration system and was built using Research Electronic Data Capture (REDCap), software developed by Vanderbilt University.[13] The data registration period was from February 1, 2023, to March 31, 2025.

Patients included in the database were ≥70 years old, admitted to a general ward, and underwent rehabilitation. Eligible patients were required to have a diagnosis of a respiratory, cardiovascular, gastrointestinal, infectious, or malignant disease and a hospital stay of at least 48 hours. In addition, independence in ADL before admission was required, as assessed using the 6-item Katz Index.[14] The

Katz Index is a tool used to evaluate the degree of independence in ADL among older adults or persons
with disabilities, assessing six basic activities as either "independent (1 point)" or "dependent (0 points)."
A total score of 6 points was considered indicative of independence in ADL. Preadmission Katz Index
scores were determined by therapists based on information obtained from medical records, as well as
from patients and/or their family members. Patients for whom preadmission ADL data were unavailable
were not enrolled in the study.
To ensure data accuracy, manuals were provided to data entry personnel at each facility for entering
patient data on admission and discharge. If the Katz Index on discharge was less than six points, HAD
was considered to have developed. The study was approved by the Ethics Committee of Mie University
Hospital (No. H2023-015), and informed consent was obtained from the participants in written form
before registration.
102

Assessment Items on Admission
Assessment items on admission include age, sex, type of the disease that led to admission, type of
admission (emergency admission or scheduled admission), whether admitted by ambulance, stage of
cancer modified Early Warning Score (MEWS),[15] use of ventilator support on admission, use of
vasopressor drugs, consciousness status (Japan Coma Scale: JCS),[16] surgeries requiring general
anesthesia, history of falls, and smoking status, as well as the Clinical Frailty Scale.[17] The MEWS is
evaluated based on respiratory rate, heart rate, systolic blood pressure, level of consciousness, and
temperature, with higher scores indicating greater severity. The JCS is a scoring system used to evaluate
arousal status across four categories. Level 0 indicates full arousal, level 1 signifies arousal without
stimulation, level 2 represents arousal with some stimulation, and level 3 denotes an inability to achieve
arousal even with strong stimulation. The Clinical Frailty Scale is an index for evaluating frailty in older
adults, consisting of nine levels ranging from 1 (very healthy) to 9 (terminal stage). It is based on a
comprehensive evaluation of physical and cognitive functions. Additionally, Functional Comorbidity
Index,[18] cognitive function score,[19] place of residence before admission (facility, hospital, home, other),
medications used on admission (antipsychotics, anxiolytics, antidepressants, hypnotic–sedatives,
antidementia medications, and number of medications), and laboratory values (albumin [Alb], C-reactive
protein [CRP], hemoglobin [Hb]) on admission were recorded. The Functional Comorbidity Index is an
index developed to assess comorbidities that affect physical function. It scores the presence of 13 chronic
conditions, such as arthritis, osteoporosis, and depression, with higher total scores indicating a higher risk
of physical decline. The cognitive function score assesses the level of independence among older adults
with dementia. It features a ranking system from 0 to 5, where 0 indicates complete independence and 5
signifies significant mental symptoms and problematic behaviors that require specialized medical care.
This index helps to reflect the need for assistance in daily living activities and the stability of the
individual's behavior. Additionally, the following information was recorded on admission: the Katz Index,
the Food Intake LEVEL Scale (FILS),[20] Functional Ambulation Categories (FAC),[21] and Global
Leadership Initiative on Malnutrition (GLIM) criteria.[22] Other recorded factors included height, weight,
and weight from 3 to 6 months prior as well as the need for infection control in rehabilitation
implementation, bed rest orders, urinary catheter placement, and the use of physical restraints. The FILS
is a rating scale for eating and swallowing abilities, consisting of 10 steps from 1 (unable to eat food
orally) to 10 (able to eat normal food), which indicates the current status of oral intake. The FAC is a
walking ability rating scale that categorizes walking capability on a 6-point scale from 0 (unable to walk)
to 5 (independent walking), based on the level of assistance required for walking. The GLIM criteria are
internationally standardized criteria for diagnosing malnutrition. The diagnosis using the GLIM criteria is
a two-step process. First, patients are identified using a validated screening tool. To be diagnosed as

malnourished, they must meet at least one criterion from the phenotypic category, which includes weight
loss, low body mass index (BMI), and muscle mass loss, and the etiologic category, which includes
factors such as inflammation, disease burden, and insufficient intake. Cases diagnosed as malnourished
were considered severely malnourished if any one of the three phenotypic criteria exceeds the threshold
for severe malnutrition; otherwise, they are categorized as moderately malnourished. Laboratory values
were obtained within 3 days of admission using the values closest to the admission date.
143
Discharge Assessment Items
Items evaluated on discharge include the length of stay and the discharge destination, which can be a
general ward, convalescent ward, long-term care ward, home, facility, or death. Additionally, the
assessment covers any complications experienced during hospitalization, such as pneumonia, falls,
depression, or delirium. It also includes whether physical therapy, occupational therapy, or speech therapy
was provided, along with the total number of therapy units administered (1 unit = 20 minutes).
Additionally, the FILS score on discharge and the rehabilitation start location (bedside or training room),
as well as the Katz Index and FAC on discharge, were also recorded. The presence of dental treatments,
whether the patient is edentulous, denture use, weight on discharge, length of nothing by mouth period
after admission, infectious diseases during hospitalization, presence of parenteral nutrition, presence of
enteral nutrition, medications and number of medication types administered on discharge were also
recorded. HAD was defined as a Katz Index score < 6 on discharge. Hospital-associated dysphagia was
defined as a decrease in FILS after admission (Food Intake LEVEL Scale on discharge < Food Intake
LEVEL Scale on admission).
Additionally, facility Information: the number of general hospital beds, hospital type, total number of
therapists, number of patients receiving rehabilitation, whether weekend rehabilitation was offered, and
the availability of dentists and dental hygienists, number of full-time dietitians and ward dietitians at each
facility was also recorded. Facility information was collected from each participating institution at the
time of enrollment.
163
Data Quality Assurance
The data dictionary is provided as a supplementary table. The data capture system was developed by the
Osaka Metropolitan University REDCap Group. For ordinal variables, values outside the defined range
cannot be selected. For continuous variables, outliers cannot be entered. A brief data entry manual was
distributed to all data registrants. Each facility's database manager was asked to monitor eligibility and
case registration status within REDCap on an ongoing basis. Data entry was scheduled at two time points:
admission and discharge. Clinical variables were entered based on physicians' assessments;
rehabilitation-related variables based on evaluations by rehabilitation professionals; and nutrition-related
variables based on the judgment of ward staff or registered dietitians. The diagnosis of the condition
leading to admission and comorbidities was based on physicians' clinical judgment. For items that might
be prone to ambiguity, explanatory notes were included in the data entry sheet. In particular, visual
reference materials outlining the assessment criteria were provided for the Clinical Frailty Scale, the
cognitive function score, the Katz Index, FILS, FAC, and the GLIM criteria. For patients who died during
hospitalization, we requested that their clinical status immediately prior to death be recorded. In addition,
the corresponding author reviewed the REDCap database monthly to identify outliers or missing values in
required fields and requested clarification or data completion from participating sites when necessary.
180
Statistical Analysis

Descriptive statistics were used to analyze the data. Frequencies and percentages were reported for
categorical variables. As this study was descriptive in nature, no statistical hypothesis testing was
conducted.　IBM SPSS ver. 26 (IBM Corporation; Armonk, New York, US) was used for statistical
analysis. Nominal data were expressed as numbers and percentages (%), whereas continuous data were
reported as mean ± standard deviation (SD) for parametric values and as median (interquartile range
[IQR]: 25th–75th percentile) for nonparametric values assessed via histogram. The number of missing
values was also recorded.
We conducted an additional facility-level comparative analysis limited to the facilities that enrolled ≥10
patients. Facility-level analyses were performed for the main outcome, HAD, and for missingness in
GLIM-defined malnutrition.
This article was previously posted as a preprint on medRxiv (DOI:
https://doi.org/10.1101/2025.11.01.25339287) on Nov 01, 2025. This work is licensed under a CC BY 4.0
License.
195

Results
Among the nine hospitals involved in the HAD database, the total number of registered patients was 209.
Table 1 displays the data recorded on admission. Variables with missing values exceeding 5% included
cancer stage (13.3%), BMI on admission (5.0%), malnutrition on admission (20.6%), weight 3–6 months
before admission (41.6%), and average energy intake within 48 hours after admission (5.0%). The median
age of patients in the study was 79.0 [75–85] years, with 60.8% of participants being men. Cancers
accounted for 39.7% of hospitalizations, followed by cardiovascular diseases at 29.7% and respiratory
diseases at 15.3%. In terms of hospitalization type, 52.2% were scheduled admissions, 47.8% were
emergency admissions, and 28% involved ambulance transports. Among patients with cancer, 28.9%
were Stage 3 and 20.5% were Stage 4.
On admission, the clinical status included a median MEWS of 0.0 [0.0–2.0], a JCS score of 0 for the
majority (94.3%), use of a ventilator in 9.6% of cases, and administration of vasopressors in 2.4% of
patients. Moreover, 30.6% had received surgery under general anesthesia, 9.1% had a history of falls, and
15.8% of the patients had a history of smoking. Regarding preadmission frailty indicators, 70.6% of
patients had a Clinical Frailty Scale score of 3 or higher. The Functional Comorbidity Index had the
highest score of 1 in 35.9% of patients. On admission, 12.9% of patients used hypnotic–sedatives, 3.8%
used antidementia drugs, and <3% used antipsychotics, anxiolytics, or antidepressants. The median Katz
Index on admission was 6 [5–6]. Among the patients, 65.2% had normal oral intake, whereas 12.1% were
considered difficult to feed orally (FILS 1–6). The largest proportion of patients (47.1%) had a FAC score
of 5, indicating independent walking, whereas 12.5% of patients were classified as FAC 0, meaning they
were unable to walk. On admission, nutrition-related indices indicated a median BMI of 21.9 [19.7–24.5]
and a median weight of 55.2 [47.9–63.7] kg over the previous 3–6 months. The average energy intake
during the 48 hours following admission was 1400 [800–1600] kcal. According to the GLIM criteria for
diagnosing malnutrition, 59.6% of patients were not malnourished, 22.3% had moderate malnutrition, and
18.1% had severe malnutrition. Additionally, 16.9% were on bed rest, 17.5% had urinary catheters, 3.4%
underwent physical restraint, and 1.9% required infection control during rehabilitation.
Table 2 presents the data recorded on discharge. The three variables with missing values that exceeded
5% were BMI on discharge (15.3%), the length of nothing by mouth period (9.1%), and weight loss after
admission (14.3%). The prevalence of HAD was 29.1%, whereas the prevalence of hospital-associated
dysphagia was 16.7%. During hospitalization, 75.4% of patients experienced weight loss, and 15.9%
decreased FAC score. The median length of hospital stay was 19.0 [13–34] days. The most common
discharge destination was home (82.3%), followed by general wards (6.2%) and convalescent wards

(5.3%). During hospitalization, adverse events included pneumonia (8.6%), falls (6.3%), depression
(1.0%), and delirium (9.6%). Physical therapist intervention was present in 97.6% of cases, whereas
occupational therapists were involved in 25.8%, and speech therapists in 14.4%. The median total number
of rehabilitation units was 16.5 [8.0–31.0]. Swallowing ability on discharge based on FILS was as
follows: 6.7% without oral intake, 4.3% with oral intake plus alternative nutrition, 28.2% with oral intake
only, and 59.8% normal. Rehabilitation was initiated at the bedside in 72.7% of cases and in the training
room in 27.3%. The median Katz Index on discharge was 5.0 [5.0–6.0]. For walking ability (FAC),
47.6% scored 5 and 26.9% scored 4. Cognitive function score was 63.8% for rank 0, 22.7% for rank 1,
and 8.7% for rank 2, indicating that most patients maintained relatively high levels of independence.
During hospitalization, 17.9% of patients received dental interventions. Of the patients, 8.8% were
edentulous, 46.1% used dentures. The median body weight on discharge was 53.1 [45.4–61.3], with a
missing data rate of 13.9%. The median length of nothing by mouth period after admission was 0 days
[0–1]. Dietitians were involved in 50.7% of cases. Total parenteral nutrition was administered to 16.3% of
patients, whereas enteral nutrition was given to 12.0%. The median number of drug types on discharge
was 6 [3–8]. Usage of medications were as follows: antipsychotics 2.4%, anxiolytics 3.8%,
antidepressants 2.4%, hypnotic–sedatives 20.2%, and antidementia drugs 2.9%.
Table 3 contains information about the participating facilities. University hospitals made up 44.4% of
the total. The average number of general beds was 472.3 (±218.5), and the median number of therapists
was 41 [24–61]. All facilities had systems for holiday rehabilitation. The median number of ward
dietitians was 1.0 [0.5–7.5]. The median number of registered patients per facility was 17 [11–26], with a
substantial proportion of cases enrolled from general hospitals, including university hospitals. The
minimum and maximum numbers of registered patients per facility were 1 and 88, respectively.
Table 4 shows an additional facility-level comparative analysis limited to the seven facilities that
enrolled ≥10 patients. Facility-level analyses were performed for the main outcome, HAD, and for
GLIM-defined malnutrition, which had a high proportion of missing data. The incidence of HAD varied
across facilities. However, given that some facilities had relatively small sample sizes, direct comparison
of incidence rates between facilities is limited by statistical instability. Missingness in GLIM-defined
malnutrition was concentrated in specific facilities.
256

Discussion

We established a multicenter prospective database on HAD and collected detailed clinical and functional
data on 209 hospitalized older adults. In this database, HAD occurred in 29.1% of patients.
Hospital-associated dysphagia occurred in 16.7% of patients, and 75.4% of hospitalized patients
experienced weight loss during hospitalization. Additionally, the FAC score decreased in 15.9% of
patients during hospitalization. While most variables had low levels of missing data, variables involving
body weight—such as GLIM-defined malnutrition, body weight before admission, weight on discharge,
and weight loss during hospitalization —demonstrated comparatively higher missingness.
The incidence of HAD in this study was 29.1%, which was consistent with that reported in previous
studies. The reported incidence of HAD varied among studies, likely due to differences in patients'
baseline functional levels, timing of assessment, and HAD definitions. Zaslavsky et al.[23] and Brown et
al.[24] reported higher incidence in older adults hospitalized in acute care settings. They assessed ADL at
multiple time points, from before admission to 1 month after discharge. Using the Modified Barthel Index,
Zaslavsky et al. defined HAD as failure to regain pre-hospital functional status after discharge, reporting
an incidence of approximately 44%. In a randomized controlled trial of cognitively intact older adults,
Brown et al. reported a 43% incidence of HAD in the usual care group. However, unlike our study, these
studies did not exclusively include patients who were independent in ADL before hospitalization, which

may have influenced the observed incidence of HAD. Differences in assessment timing, whether pre- and
post-hospitalization data were available; baseline ADL; age and cognitive function; length of stay; and
whether rehabilitation was provided could all contribute to variations in the reported incidence of HAD.
In this study, hospital-associated dysphagia was observed in 16.7% of the patients. In the previous study
by Maeda et al.,[25] hospital-associated dysphagia was reported in 4.3% of patients, which is lower than the
incidence observed in the present study. Nagano et al.[26] reported that 5.6% of patients who already had
dysphagia on admission experienced a further decline in their swallowing ability during hospitalization.
In that study, the group with decreased swallowing ability showed significantly higher rates of
malnutrition, insufficient energy intake, and longer nothing per mouth durations. The difference in
definition of hospital-associated dysphagia may account for the discrepancy in incidence between our
study and previous research. In our study, we defined hospital-associated dysphagia as a decrease in the
FILS, whereas the previous study defined it based on a change in diet form. Our study also differed from
those by Maeda et al. and Nagano et al. in patient characteristics, including age and reasons for
hospitalization, which may have influenced the outcomes. In the report by Nagano et al., the Functional
Oral Intake Scale (FOIS) was used to assess swallowing function, and decreased swallowing ability was
defined as FOIS ≤5. Although the FOIS and FILS are assessment tools that evaluate actual feeding
conditions, the FILS provides a more detailed assessment. This difference may also explain the variation
in the incidence of hospital-associated dysphagia between our study and previous research. Swallowing
ability is closely associated with overall muscle strength, as represented by the concept manuscript   of
sarcopenic dysphagia.[27] Reduced physical activity and malnutrition during hospitalization can lead to
deterioration of general health and impaired swallowing function. In our study, a high incidence of weight
loss during hospitalization (75.4%) was also observed, suggesting that the presence or progression of
sarcopenia may have contributed to the decline in swallowing ability. Unlike previous studies, our study
included patients who were undergoing rehabilitation. Patients requiring rehabilitation may be more
susceptible to sarcopenic effects, making them more prone to developing hospital-associated dysphagia.
In this study, 75.4% of the patients experienced weight loss during hospitalization. In previous studies,
41.2%–45.5% of patients were reported to have lost weight during hospitalization,[28,29] which was lower
than in the present study. Zannidi et al.[30] reported that 19% of patients lost 10% or more of their weight
during their hospital stay, and the longer the stay, the higher the risk of weight loss. The average age of
the patients in this study was higher, and the average length of hospital stay was longer than that of
previous studies. Additionally, many patients in this study had cardiovascular disease (29.7%) and may
have experienced edema on admission, which could have contributed to weight loss as the edema
improved during hospitalization.
In this study, 15.9% of patients experienced a decline in walking ability during hospitalization. Valiani
et al.[31] evaluated activity using the Braden Scale and reported that 20.6% of patients showed a decline in
walking ability. Compared with the previous study, the patients in our study differed in several
characteristics, including being originally independent in ADL and undergoing rehabilitation during
hospitalization. The previous study reported a median hospital stay of 4 days, whereas in the present
study it was 19 days. These differences may have contributed to the variation in the incidence of
hospital-associated walking ability decline.
While most variables had low levels of missing data, variables involving body weight—such as
GLIM-defined malnutrition, body weight before admission, weight on discharge, and weight loss during
hospitalization—demonstrated comparatively higher missingness. These variables require accurate body
weight measurement or reliable information on previous body weight; however, in routine clinical
practice, obtaining such data is not always feasible. Furthermore, the need for multiple parameters to
diagnose malnutrition according to the GLIM criteria may have increased the assessment burden and

contributed to incomplete documentation in clinical practice. Missingness in GLIM-defined malnutrition
was concentrated in specific facilities. This pattern may reflect differences in assessment systems or data
entry processes across institutions. Although it may be challenging to mandate a full GLIM assessment, in
Japan's Diagnosis Procedure Combination system, body weight at both admission and discharge is
already a required data item. Therefore, we are considering making body weight on discharge a
mandatory variable in future database construction. To improve data completeness in future studies,
standardization of required data elements and strengthening of centralized monitoring procedures will be
necessary.
This study has several limitations. First, the number of participating facilities was limited, which may
affect the representativeness of the patient population. Second, some evaluation indicators were based on
simple screening tools and did not include detailed clinical diagnoses. Third, we did not systematically
record the specific screening tools used prior to GLIM-based diagnosis. In addition, we did not establish a
detailed protocol for body weight assessment, nor did we standardize the method for confirming previous
body weight (3–6 months prior to admission). Therefore, concerns remain regarding the validity of the
nutritional diagnosis and body weight–related data. Fourth, all patients included in this study were
followed up until discharge. However, the total number of potentially eligible patients, the number
screened, and the number excluded were not available. In addition, there was substantial variability in the
number of registered patients across facilities, which may limit the generalizability of the findings. Some
facilities had relatively small sample sizes, and direct comparison of incidence rates between facilities is
limited by statistical instability. Therefore, it is difficult to draw definitive conclusions regarding
interfacility differences in HAD incidence based on the present data, and the findings should be
interpreted with caution. Fifth, it was difficult to obtain data on patients not referred for rehabilitation;
therefore, the study population was limited to patients receiving rehabilitation, which may restrict
generalizability. Although each participating facility was requested to enroll consecutive cases, adherence
to this approach could not be monitored and may not have been fully ensured. In addition, patients
without available preadmission ADL data were excluded, which may have reduced the sample size and
introduced selection bias.
347
Nevertheless, this study, which gathered data prospectively from multiple centers, is significant for
understanding the current status of HAD in clinical practice and for laying the groundwork for future
analytical research.
This study elucidates the realities of HAD and its associated functional decline in an older adult
inpatient population, providing a descriptive analysis of its characteristics, incidence, and background
factors. In this study, 29.1% of patients experienced HAD, whereas 16.7% developed hospital-associated
dysphagia. These findings are important for understanding the functional decline and risks experienced by
older patients during hospitalization. The dataset obtained in this study may be utilized in future analyses
to identify predictors of HAD and to develop early intervention programs for high-risk patients. The
registry provides a structured foundation for future analytical studies.
358

359 **Acknowledgments:**
360 This work was supported by JSPS KAKENHI (grant number 22K19669).
361

362 **Conflict of Interest:**
363 The authors declare no conflict of interest.
364
References

1.   Covinsky KE, Pierluissi E, Johnston CB. Hospitalization-associated disability: "She was probably
able to ambulate, but I'm not sure." JAMA. 2011;306(16):1782–1793.
http://dx.doi.org/10.1001/jama.2011.1556

2.   Covinsky KE, Palmer RM, Fortinsky RH, Counsell SR, Stewart AL, Kresevic D, Burant CJ,
Landefeld CS. Loss of independence in activities of daily living in older adults hospitalized with
medical illnesses: increased vulnerability with age. J Am Geriatr Soc. 2003;51(4):451–458.
https://pubmed.ncbi.nlm.nih.gov/12657063/

3.   Jo SJ, Lee SH, Min HJ, Kim HJ, Kong HH. Mortality outcomes and contributing risk factors in
patients with hospital-associated disability. J Clin Med. 2024;13(16):4798.
https://pubmed.ncbi.nlm.nih.gov/39200940/

4.   Chen Y, Almirall-Sánchez A, Mockler D, Adrion E, Domínguez-Vivero C, Romero-Ortuño R.
Hospital-associated deconditioning: Not only physical, but also cognitive. Int J Geriatr Psychiatry.
2022;37(3). https://pubmed.ncbi.nlm.nih.gov/35142397/

5.   Mathews SB, Arnold SE, Epperson CN. Hospitalization and cognitive decline: Can the nature of the
relationship be deciphered? Am J Geriatr Psychiatry. 2014;22(5):465–480.
https://pubmed.ncbi.nlm.nih.gov/23567430/

6.   Matsuo H, Yoshimura Y, Maeno Y, Tanaka S. Possible sarcopenia and its association with
hospital-associated dysphagia and decline in physical function: Findings from a heart failure patients
prospective cohort study. Clin Nutr ESPEN. 2024;63:364–370.
https://pubmed.ncbi.nlm.nih.gov/38971404/

7.   Brown CJ, Friedkin RJ, Inouye SK. Prevalence and outcomes of low mobility in hospitalized older
patients: Prevalence and outcomes of low mobility. J Am Geriatr Soc. 2004;52(8):1263–1270.
https://pubmed.ncbi.nlm.nih.gov/15271112/

8.   Fortinsky RH, Covinsky KE, Palmer RM, Landefeld CS. Effects of functional status changes before
and during hospitalization on nursing home admission of older adults. J Gerontol A Biol Sci Med
Sci. 1999;54(10):M521-6. https://pubmed.ncbi.nlm.nih.gov/10568535/

9.   Loyd C, Markland AD, Zhang Y, Fowler M, Harper S, Wright NC, Carter CS, Buford TW, Smith
CH, Kennedy R, Brown CJ. Prevalence of Hospital-Associated Disability in Older Adults: A
Meta-analysis. J Am Med Dir Assoc. 2020;21(4):455-461.e5.
http://dx.doi.org/10.1016/j.jamda.2019.09.015

10. Hao X, Zhang H, Zhao X, Peng X, Li K. Risk factors for hospitalization-associated disability among
older patients: A systematic review and meta-analysis. Ageing Res Rev. 2024;101(102516):102516.
http://dx.doi.org/10.1016/j.arr.2024.102516

11. Gillick MR, Serrell NA, Gillick LS. Adverse consequences of hospitalization in the elderly. Soc Sci
Med. 1982;16(10):1033–1038. https://pubmed.ncbi.nlm.nih.gov/6955965/

12. Reichardt LA, Aarden JJ, van Seben R, van der Schaaf M, Engelbert RHH, Bosch JA, Buurman BM,
Hospital-ADL study group. Unravelling the potential mechanisms behind hospitalization-associated
disability in older patients; the Hospital-Associated Disability and impact on daily Life

404  (Hospital-ADL) cohort study protocol. BMC Geriatr. 2016;16:59.
405  http://dx.doi.org/10.1186/s12877-016-0232-3

406 13. Van Bulck L, Wampers M, Moons P. Research Electronic Data Capture (REDCap): tackling data
407  collection, management, storage, and privacy challenges. Eur J Cardiovasc Nurs. 2022;21(1):85–91.
408  http://dx.doi.org/10.1093/eurjcn/zvab104

409 14. Laan W, Zuithoff NPA, Drubbel I, Bleijenberg N, Numans ME, De Wit NJ, Schuurmans MJ.
410  Validity and reliability of the Katz-15 scale to measure unfavorable health outcomes in
411  community-dwelling older people. J Nutr Health Aging. 2014;18(9):848–854.
412  http://dx.doi.org/10.1007/s12603-014-0558-5

413 15. Kaho Umeki Megumi Ishibashi Misako Takahashi and Megumi Kamitani. Utilize MEWS and
414  underlying diseases of emergency inpatients for prediction of sudden change. Medical journal of
415  Shimane Central Hospital. 2022;46:15–18.

416 16. Okada Y, Kiguchi T, Iiduka R, Ishii W, Iwami T, Koike K. Association between the Japan Coma
417  Scale scores at the scene of injury and in-hospital outcomes in trauma patients: an analysis from the
418  nationwide trauma database in Japan. BMJ Open. 2019;9(7):e029706.
419  http://dx.doi.org/10.1136/bmjopen-2019-029706

420 17. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, Mitnitski A. A global
421  clinical measure of fitness and frailty in elderly people. CMAJ. 2005;173(5):489–495.
422  http://dx.doi.org/10.1503/cmaj.050051

423 18. Groll DL, To T, Bombardier C, Wright JG. The development of a comorbidity index with physical
424  function as the outcome. J Clin Epidemiol. 2005;58(6):595–602.
425  http://dx.doi.org/10.1016/j.jclinepi.2004.10.018

426 19. Tago M, Katsuki NE, Yaita S, Nakatani E, Yamashita S, Oda Y, Yamashita SI. High inter-rater
427  reliability of Japanese bedriddenness ranks and cognitive function scores: a hospital-based
428  prospective observational study. BMC Geriatr. 2021;21(1):168.
429  http://dx.doi.org/10.1186/s12877-021-02108-x

430 20. Kunieda K, Ohno T, Fujishima I, Hojo K, Morita T. Reliability and validity of a tool to measure the
431  severity of dysphagia: the Food Intake LEVEL Scale. J Pain Symptom Manage. 2013;46(2):201–206.
432  http://dx.doi.org/10.1016/j.jpainsymman.2012.07.020

433 21. Holden MK, Gill KM, Magliozzi MR, Nathan J, Piehl-Baker L. Clinical gait assessment in the
434  neurologically impaired. Reliability and meaningfulness. Phys Ther. 1984;64(1):35–40.
435  http://dx.doi.org/10.1093/ptj/64.1.35

436 22. Cederholm T, Jensen GL, Correia MITD, Gonzalez MC, Fukushima R, Higashiguchi T, Baptista G,
437  Barazzoni R, Blaauw R, Coats A, Crivelli A, Evans DC, Gramlich L, Fuchs-Tarlovsky V, Keller H,
438  Llido L, Malone A, Mogensen KM, Morley JE, Muscaritoli M, Nyulasi I, Pirlich M, Pisprasert V, de
439  van der Schueren MAE, Siltharm S, Singer P, Tappenden K, Velasco N, Waitzberg D, Yamwong P,
440  Yu J, Van Gossum A, Compher C, GLIM Core Leadership Committee, GLIM Working Group.
441  GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical
442  nutrition community. Clin Nutr. 2019;38(1):1–9. http://dx.doi.org/10.1016/j.clnu.2018.08.002

23.   Zaslavsky O, Zisberg A, Shadmi E. Impact of functional change before and during hospitalization on
functional recovery 1 month following hospitalization. J Gerontol A Biol Sci Med Sci.
2015;70(3):381–386. http://dx.doi.org/10.1093/gerona/glu168

24.   Brown CJ, Foley KT, Lowman JD Jr, MacLennan PA, Razjouyan J, Najafi B, Locher J, Allman RM.
Comparison of posthospitalization function and community mobility in hospital mobility program
and usual care patients: A randomized clinical trial. JAMA Intern Med. 2016;176(7):921–927.
http://dx.doi.org/10.1001/jamainternmed.2016.1870

25.   Maeda K, Ishida Y, Nonogaki T, Shimizu A, Yamanaka Y, Matsuyama R, Kato R, Mori N.
Development and predictors of sarcopenic dysphagia during hospitalization of older adults. Nutrients.
2019;12(1):70. http://dx.doi.org/10.3390/nu12010070

26.   Nagano A, Onaka M, Maeda K, Ueshima J, Shimizu A, Ishida Y, Nagami S, Miyahara S, Nishihara
K, Yasuda A, Satake S, Mori N. Prevalence and characteristics of the course of dysphagia in
hospitalized older adults. Nutrients. 2023;15(20):4371. http://dx.doi.org/10.3390/nu15204371

27.   Nagai T, Wakabayashi H, Nishioka S, Momosaki R. Functional prognosis in patients with
sarcopenic dysphagia: An observational cohort study from the Japanese sarcopenic dysphagia
database. Geriatr Gerontol Int. 2022;22(10):839–845. http://dx.doi.org/10.1111/ggi.14466

28.   Leandro-Merhi VA, Srebernich SM, Gonçalves GMS, de Aquino JLB. In-hospital weight loss,
prescribed diet and food acceptance. Arq Bras Cir Dig. 2015;28(1):8–12.
http://dx.doi.org/10.1590/S0102-67202015000100003

29.   Merhi VNAL, Teles MV, Aquino JL, Mendon JA, Mendes ET. Do more overweight patients at
admission lose weight during hospitalization. Authorea, Inc.; 2020.
http://dx.doi.org/10.22541/au.159355933.38613731

30.   Zannidi D, Patel PS, Leventea E, Paciepnik J, Dobson F, Heyes C, Goudie RJB, Griep LMO, Preller
J, Spillman LN. Factors associated with significant weight loss in hospitalised patients with
COVID-19: A retrospective cohort study in a large teaching hospital. Nutrients. 2022;14(19):4195.
http://dx.doi.org/10.3390/nu14194195

31.   Valiani V, Gao S, Chen Z, Swami S, Harle CA, Lipori G, Sourdet S, Wu S, Nayfield SG, Sabbá C,
Pahor M, Manini TM. In-hospital mobility variations across primary diagnoses among older adults. J
Am Med Dir Assoc. 2016;17(5):465.e1-465.e8. http://dx.doi.org/10.1016/j.jamda.2016.02.003

Table. 1 Admission data (n=209)

	Number of missing (%)	n (%), mean ± SD, median [IQR]
Age, years	0	79.0 [75–85]
Sex, Male	0	127 (60.8)
Primary disease leading to hospitalization	0	
Malignant tumor		83 (39.7)
Circulatory disease		62 (29.7)
Gastrointestinal disease		13 (6.2)
Infectious disease		19 (9.1)
Respiratory disease		32 (15.3)
Hospitalization type	0	
Scheduled hospitalization		109 (52.2)
Emergency hospitalization		100 (47.8)
Place of residence before admission	0	
General ward		3 (1.4)
Home		199 (95.2)
Facility		7 (3.3)
Stage of Cancer (among 83 patients with cancer)	11 (13.3)	
0		3 (3.6)
1		14 (16.9)
2		14 (16.9)
3		24 (28.9)
4		17 (20.5)
Admission via ambulance	2 (1)	58 (28)
Modified early warning score on admission	0	0.0 [0.0–2.0]
Japan Coma Scale on admission	0	
0		197 (94.3)
1		8 (3.8)
2		3 (1.4)
3		1 (0.5)
Vasopressors use on admission	2 (1)	5 (2.4)
Ventilator use on admission	0	20 (9.6)
Surgery with general anesthesia during hospitalization	0	64 (30.6)
History of falls in the year before admission	0	19 (9.1)
Smoking habits in the year before admission	1 (0.5)	33 (15.8)
Cognitive function score before hospitalization	0	
0		148 (70.8)

1		42 (20.1)
2		15 (7.2)
3		3 (1.4)
4		1 (0.5)
Preadmission Clinical Frailty Scale	1 (0.5)	
1		27 (13)
2		34 (16.3)
3		61 (29.3)
4		40 (19.2)
5		46 (22.1)
Functional comorbidity index on admission	0	
0		51 (24.4)
1		75 (35.9)
2		50 (23.9)
3		23 (11.0)
4		6 (2.9)
5		4 (1.9)
Number of drug types on admission	5 (2.4)	5.5 [3.0–8.0]
Antipsychotics on admission	1 (0.5)	5 (2.4)
Antianxiety medications on admission	1 (0.5)	7 (3.3)
Antidepressants on admission	1 (0.5)	5 (2.4)
Hypnotic–sedatives on admission	1 (0.5)	27 (12.9)
Antidementia medications on admission	1 (0.5)	8 (3.8)
Alb on admission, g/dL	5 (2.4)	3.5 [3.1–3.9]
CRP on admission, mg/dL	8 (3.8)	0.9 [0.1–6.1]
Hb on admission, g/dL	3 (1.4)	11.6 [10.2–12.9]
Katz Index on admission	0	6.0 [5.0–6.0]
Food Intake LEVEL Scale on admission	2 (1)	
No oral intake (1–3)		17 (8.2)
Oral intake and alternative nutrition (4–6)		8 (3.9)
Oral intake only (7–9)		47 (22.7)
Normal (10)		135 (65.2)
Functional Ambulation Categories on admission	1 (0.5)	
0		26 (12.5)
1		7 (3.4)
2		6 (2.9)
3		22 (10.6)

4		49 (23.6)
5		98 (47.1)
Diagnosis of malnutrition on admission (GLIM criteria)	43 (20.6)	
No malnutrition		99 (59.6)
Moderate malnutrition		37 (22.3)
Severe malnutrition		30 (18.1)
Weight on admission, kg	7 (3.3)	55.2 [47.9–63.7]
Body mass index on admission	10 (5)	21.9 [19.7–24.5]
Weight 3–6 months before admission, kg	87 (41.6)	57.0 [48.0–65.0]
Average energy intake within 48 hours of admission, kcal	10 (5)	1400 [800–1600]
Bed rest order on admission	2 (1)	35 (16.9)
Need for infection control in rehabilitation implementation	1 (0.5)	4 (1.9)
Urinary catheter placement on admission	3 (1.4)	36 (17.5)
Physical restraint on admission	2 (1)	7 (3.4)

473
474

Table. 2 Discharge Data (n=209)

	Number of missing (%)	n (%), mean ± SD, median [IQR]
Hospital-associated disability (Katz Index < 6 on discharge)	3 (1.4)	60 (29.1)
Hospital-associated dysphagia (Food Intake LEVEL Scale on discharge < Food Intake LEVEL Scale on admission)	4 (1.9)	35 (16.7)
Weight loss during hospitalization (weight on discharge < weight on admission)	30 (14.3)	135 (75.4)
Decrease in Functional Ambulation Categories during hospitalization	2 (1)	33 (15.9)
Length of hospital stay, days	0	19 [13–34]
Place of discharge	0	
General ward		13 (6.2)
Convalescent ward		11 (5.3)
Long-term care ward		2 (1)
Home		172 (82.3)
Facility		5 (2.4)
Death		6 (2.9)
Pneumonia (in hospital)	0	18 (8.6)
Falls (in hospital)	1 (0.5)	13 (6.3)
Depression (in hospital)	1 (0.5)	2 (1.0)
Delirium (in hospital)	1 (0.5)	20 (9.6)
Physical Therapist involvement	0	204 (97.6)
Occupational Therapist involvement	0	54 (25.8)
Speech Therapist involvement	1 (0.5)	30 (14.4)
Total rehabilitation units	3 (1.4)	16.5 [8.0–31.0]
Food Intake LEVEL Scale on discharge	2 (1)	
No oral intake (1–3)		14 (6.7)
Oral intake and alternative nutrition (4–6)		9 (4.3)
Oral intake only (7–9)		59 (28.2)
Normal (10)		125 (59.8)
Rehabilitation start location	4 (1.9)	
Bedside		149 (72.7)
Training room		56 (27.3)
Katz Index on discharge	3 (1.4)	5.0 [5.0–6.0]
Functional Ambulation Categories on discharge	1 (0.5)	
0		12 (5.8)
1		10 (4.8)
2		14 (6.7)

3		17 (8.2)
4		56 (26.9)
5		99 (47.6)
Cognitive function score on discharge	2 (1)	
0		132 (63.8)
1		47 (22.7)
2		18 (8.7)
3		4 (1.9)
4		2 (1)
5		4 (1.9)
Dental professionals' intervention during hospitalization	2 (1)	37 (17.9)
Edentulous	5 (2.4)	18 (8.8)
Denture use	3 (1.4)	95 (46.1)
Weight on discharge	29 (13.9)	53.1 [45.4–61.3]
Length of nothing by mouth period after admission	19 (9.1)	0 [0–1]
Dietitian involvement	4 (1.9)	104 (50.7)
Total parenteral nutrition during hospitalization	0	34 (16.3)
Enteral nutrition during hospitalization	0	25 (12.0)
Number of drug types on discharge	1 (0.5)	6 [3–8]
Antipsychotics on discharge	0	5 (2.4)
Antianxiety medications on discharge	0	8 (3.8)
Antidepressants on discharge	1 (0.5)	5 (2.4)
Hypnotic–sedatives on discharge	1 (0.5)	42 (20.2)
Antidementia medications on discharge	2 (1)	6 (2.9)

475
476

Table. 3 Facility Information (n=9)

	Number of missing (%)	n (%), mean ± SD, median [IQR]
University hospital	0	3 (33.3)
Number of general hospital beds	0	472.3 ± 218.5
Total number of therapists	0	41 [24–61]
Weekend rehabilitation	0	9 (100)
Dentists	0	5 (55.5)
Dental hygienists	0	5 (55.5)
Number of ward dietitians	0	1 [0.5–7.5]

477
478

479 Table 4. Facility-level comparative analysis

Hospital ID	1	2	3	4	5	6	7
Number of eligible patients	11	11	17	88	30	20	26
Hospital-associated disability, n (%)	4 (36.4)	0 (0)	12 (70.6)	13 (14.8)	8 (26.7)	2 (10.0)	18 (69.2)
Missingness in GLIM-defined malnutrition, n (%)	0 (0)	0 (0)	0 (0)	29 (33)	0 (0)	2 (10.0)	12 (46.2)