Deep Radiomics in 18F-FDG-PET/CT for Survival Prediction in Multiple Myeloma: Novel Embeddings Using A Foundational Model's Memory

Javier Guinea-Pérez[1*], Silvia Uribe[1], Sara Peluso[2,3],
Gastone Castellani[2,3], Cristina Nanni[4], Federico Álvarez[1]

[1]Universidad Politécnica de Madrid, Madrid,Spain.
[2]Department of Medical and Surgical Sciences, Alma Mater
Studiorum-University of Bologna, Bologna, Italy.
[3]IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy.
[4]Nuclear Medicine, IRCCS Azienda Ospedaliero-Universitaria di
Bologna, Bologna, Italy.

*Corresponding author(s). E-mail(s): javier.guinea.perez@upm.es;

Abstract

Purpose: To test whether internal memory states from a medical foundational segmentation model can serve as compact, mask-aware embeddings for predicting progression-free survival (PFS) in multiple myeloma (MM) from whole-body $[^{18}F]$FDG PET/CT.

Methods: We analyzed 227 newly diagnosed MM patients with PET/CT and clinical data. For two regions of interest (spine-dilated and full skeleton), we prompted MedSAM2 slice-wise using mask-derived bounding boxes and cached the final spatio-temporal memory tensor per modality. We compared two downsampling strategy to obtain per-study embeddings: channel×memory averaging with a small CNN head, and a squeeze-and-excitation–style depth-attention pooling. PET and CT embeddings were combined by late fusion (concatenation or scalar-gated mixing) and passed to a DeepSurv head. We evaluated image-only and multimodal (image+clinical) models with stratified 5-fold cross-validation. The primary endpoint was Harrell's c-index (mean $\pm$ SE across folds).

Results: Image-only models using the averaging downsampler achieved up to **0.659 $\pm$ 0.015** c-index (PET, spine-dilated), comparable to baseline radiomics results. Multimodal models improved discrimination to **0.710 $\pm$ 0.032** (CT,

spine-dilated), with similar performance for other PET/CT+clinical variants (**0.703–0.710**), improving clinical-only baselines $\sim$ **6.5%**. Averaging consistently outperformed depth-attention; concatenation and gated fusion performed comparably. PET outperformed CT within the same mask in image-only settings.
Conclusion: Mask-aware memory embeddings extracted from a foundational segmentation model provide effective, data-efficient imaging biomarkers for MM PFS and, when fused with routine clinical covariates, significantly improve risk stratification over clinical-only or radiomics baselines. This offers a practical path to prognostic modeling on small medical cohorts without feature design.

Keywords: Multiple Myeloma, Survival Analysis, Radiomics, Foundational models, Deep Radiomics

1 Introduction

Multiple myeloma (MM) is a common cancer of bone marrow (BM) plasma cells that accounts for almost 10% of all hematologic diseases, having had more than 180000 diagnoses and 12000 deaths in 2022 alone [1, 2]. Few cases are diagnosed before metastasis (3% [3]), when disease markers are common and subtle symptoms [4]. Accurate risk stratification at diagnosis is central to tailoring therapy and surveillance, which can significantly improve 5-year survival rates [3].

Beyond clinical staging systems, imaging adds complementary information on tumor burden and disease biology. Whole-body [18F] fluorodeoxyglucose positron emission tomography-computed tomography (FDG PET/CT) has been included in the updated criteria for the diagnosis of MM of the International Myeloma Working Group [4] as it has proven to be useful for assessing the state of the disease and treatment strategies [5].

Risk stratification of patients is crucial for treatment selection [6], but it has not yet been delegated to imaging techniques. Standardized criteria for PET/CT interpretation in MM have been proposed [7], but visual interpretation can be difficult and remains limited in clinical practice [8]. Because of this, patient stratification is being carried out after invasive BM studies that identify chromosomal abnormalities [9]. FDG PET has the potential to improve risk stratification by detecting medullary and extramedullary focal lesions [10].

Recent years have seen a growing interest in the application of machine learning (ML) for FDG PET/CT interpretation in MM. Currently, images are processed using classical radiomic approaches, where handcrafted image features extract intensity, texture, and shape descriptors from segmented lesions. These features are then fed to ML algorithms with a clinical objective, be it risk stratification, biomarker discovery [11, 12], prognosis [13, 14], patient classification, diagnosis, and therapy response assessment [8]. These studies have shown that radiomic features can correlate with tumor burden and predict outcomes, but these features rely on manually predefined descriptors that can fail to capture high order interactions or spatial context across the skeleton.

In parallel, survival analysis has taken a step forward as deep learning-based survival models have extended traditional Cox proportional-hazards to model non linear survival data [15, 16]. These methods have been successfully integrated with radiomic feature extraction for survival analysis and biomarker discovery in MM and other ailments [17].

Foundational models are large-scale deep learning models trained on vast, heterogeneous datasets to acquire transferable representations that can be adapted to a wide range of downstream tasks with relatively little task-specific supervision [18]. In computer vision, examples include CLIP [19] and SAM [20], which have shown that training on diverse, high-volume datasets produces general-purpose embeddings that capture both semantic and structural image properties. These embeddings can then be fine-tuned or prompted to solve specialized tasks with far less annotated data than would otherwise be required. In the medical imaging domain, foundational models are being developed by pretraining on millions of scans spanning multiple modalities (CT,

MRI, PET, ultrasound) and anatomical regions [21]. Adaptations of SAM, for example, have been fine-tuned to segment medical structures robustly across organ systems without retraining from scratch [22] [23].

In this work, we propose a survival analysis framework that overcomes the limitations of radiomics feature design by creating mask-aware data-efficient representations of FDG PET/CT volumes by using the internal memory state of a state of the art foundational segmentation model. These memory embeddings are generated during mask-guided volume propagation, where the model integrates anatomical prompts with imaging context. We extract and compress the final memory state into compact embeddings that capture information that radiomics descriptors cannot. We further explore late fusion strategies for PET and CT and evaluate our method in contrast to traditional radiomics features as well as measuring the improvement on a clinical feature only model.

In doing so, we address the gap between handcrafted radiomics and deep learning approaches: radiomics pipelines embed anatomical priors but are constrained by feature engineering, while deep survival networks can model complex non-linear risks but struggle to converge on small medical cohorts. Foundational medical imaging models provide a middle ground, offering robust, transferable representations that can be harnessed without retraining on small datasets. By leveraging the memory state embeddings of MedSAM2, we show that it is possible to build clinically relevant, mask-aware survival predictors from FDG PET/CT in multiple myeloma. Related work has begun to explore SAM-derived embeddings for survival in 2D medical images in multi-omics datasets [24].

In summary, the contributions of this study are threefold: (i) we propose the use of foundational segmentation model memory states as a novel embedding for survival analysis; (ii) we benchmark PET, CT, and PET+CT late fusion embeddings against clinical and radiomics baselines; and (iii) we demonstrate that combining clinical variables with imaging embeddings yields superior progression-free survival (PFS) stratification. This positions foundational model embeddings as a promising bridge between radiomics and end-to-end deep learning in imaging-based prognostication for MM.

2 Materials and methods

2.1 Dataset

This study is based on a cohort of 262 newly diagnosed MM patients collected since 2007 by the Italian IRCCS Azienda Ospedaliero-Universitaria Sant'Orsola-Malpighi in Bologna. For each patient, whole-body [^{18}F] FDG PET/CT imaging was available along with clinical annotations including demographic variables, laboratory data, disease progression information, and survival outcomes.

For the purpose of this study, we focused on PFS as the clinical endpoint. PFS was defined as the time from diagnosis until relapse, progression, or death, with censoring applied to patients who did not experience an event during follow-up. To ensure consistency of the endpoint, patients without PFS annotations were excluded, leaving a final dataset of 227 patients with both imaging and PFS information.

Among this population, 160 patients (70.5%) experienced the event of interest, while 67 (29.5%) were censored. The cohort included 132 men (58.1%) and 95 women (41.9%), with a mean age of 62.5 years (standard deviation 9.7 years). Baseline staging according to the Revised International Staging System (R-ISS) [25] was available for most patients: 49 (21.6%) were stage I, 95 (41.9%) stage II, 25 (11.0%) stage III, while staging information was not reported for 58 (25.6%) of the cases.

The dataset included clinical covariates with established relevance to MM prognosis: age, sex, creatinine, hemoglobin, platelets, antibody isotypes light chain ratio, and calcium. Additionally, the R-ISS stage (I/II/III) was present, which condensates information about clinical variables, some not present: $\beta 2$ microglobulin, albumin, creatine, lactate dehydrogenase and chromosomal abnormalities. Variables with extensive missingness in this cohort (e.g., creatinine clearance) were excluded. Categorical variables were encoded and continuous variables standardized within each training fold prior to model fitting.

2.2 Preprocessing

PET and CT volumes were first loaded as NIfTI files and inspected. Because the two modalities differed in voxel origin and spacing despite identical qform/sform matrices, PET volumes were rigidly resampled onto the CT grid (same origin, voxel size, and dimensions) using linear interpolation with SimpleITK. No deformable registration or artifact correction was applied.

Mask generation followed the same automatic segmentation procedure described in our previous work [11]: CT scans were segmented using the Multi-Organ Objective Segmentation (MOOSE 2.0) model [26], which provides robust bone segmentation (median Dice coefficient 0.90 for most bones). From this segmentation, multiple regions of interest (ROIs) were derived through morphological operations aligned with the IMPeTUs protocol, which emphasizes bone marrow, spine, and paramedullary regions.

For the present study, we restricted the analysis to the two masks that demonstrated the best prognostic performance:

- **Spine dilated**: including vertebrae, internal spinal canal and surrounding paramedullary regions, designed to capture both intramedullary and paramedullary disease.
- **Skeleton**: a comprehensive mask encompassing the entire segmented skeleton and the dilated spine mask, designed to include total body bone disease burden.

Both masks were automatically generated on the CT images and then resampled to PET resolution, ensuring one-to-one alignment across modalities.

2.3 Architecture

Our pipeline converts FDG PET/CT volumes (paired with the automatically generated masks) into compact, mask-aware embeddings using the internal memory state of a foundational segmentation model, and then estimates the log-risk of progression with a DeepSurv head. PET and CT are processed in parallel with modality-specific

Fig. 1: Overview of the proposed pipeline. The workflow is divided into four stages: (1) Memory Extraction: For each ROI (spine dilated or skeleton) and modality (PET or CT), axial slices are prompted with mask-derived bounding boxes and propagated through MedSAM2 [27]; the final memory state is cached. (2) Downsampler: The large memory tensor is compressed into a compact embedding (either with channel/memory averaging and a small CNN head and or a lightweight depth-attention block). (3) Fusion Head: Modality-specific embeddings are late-fused (with optional clinical covariates) into a single representation either using a learned gated sum or concatenation. (4) DeepSurv: A fully connected survival head outputs the linear predictor. The top progress bar indicates trainability. Memory Extraction is frozen while Downsampler, Fusion Head and DeepSurv are trained together end-to-end.

embedding towers and are optionally combined with clinical data through late fusion before a survival head. Figure 1 summarizes the overall architecture.

2.3.1 Embedding generation

We extract image representations from **MedSAM2** [27], a medical adaptation of the SAM2 [28] segmentation foundation model. The model allows for segmentation of medical images with prompts by processing them as videos. In our case, for each study, we first select a ROI mask (either the *spine dilated* or the *full skeleton*, see Section 2.2). We generate prompts by computing the tight 2D bounding box of the ROI on each axial slice containing mask pixels and propagate the segmentation top-to-bottom through the volume. To alleviate memory requirements, only one of every two mask containing slices are propagated. MedSAM2 maintains a spatio-temporal memory during this slice-wise propagation; we cache the final memory state after the last slice and use it as our representation.

(a) CNN embedding head used to convert the $2 \times 32 \times 64 \times 64$ ROI map into a 128-D vector. The pipeline first aplies a global average to the channel and memory dimensions, followed by *Conv–ReLU–MaxPool* twice, continued by *Conv–ReLU*, global average pooling (AdaptiveAvgPool$(1,1)$), and flatten. Intermediate feature-map sizes are annotated below operations.

(b) **Light attention embedding module**. The pipeline shrinks the in-plane resolution, mixes channels to *base*, applies normalization and GELU, performs depth attention to collapse D into a 2D map of intermediate size (S), then applies two-stage 2D pooling and a linear projection to *out_dim*. The inset details the depth-attention steps, where FC stands for fully connected layers.

Fig. 2: Downsampler architectures: (a) average embedding head; (b) light attention head.

Concretely, for each modality (PET and CT) and ROI, MedSAM2 outputs a tensor of shape $\mathbf{M} \in \mathbb{R}^{C \times D \times H \times W}$ (memory layers $\times$ channels $\times$ spatial height $\times$ width). In order to have all embeddings have the same memory layer size, they were cropped/padded in that dimension, ending with a tensor of shape: $\mathbb{R}^{2 \times 32 \times 64 \times 64}$.

Given the cohort size, we consider two strategies to transform $\mathbf{M}$ into a compact embedding:

1. **Average embedding.** We perform global averaging across the memory and channel dimensions, reducing $\mathbf{M}$ to a 64×64 map. This 64×64 sized tensor was then processed through a small 2D convolutional neural network (CNN) head that maps an input with to a 128×1 sized embedding. Starting from $(1, 64, 64)$, the two 2×2 max-pool layers reduce the resolution to $(64, 16, 16)$. A final 3×3 convolution produces $(128, 16, 16)$, which is collapsed by global average pooling (GAP, implemented as AdaptiveAvgPool2d$(1, 1)$) to $(128, 1, 1)$ and flattened to a vector in

$\mathbb{R}^{128}$. This head is intentionally small to limit capacity and overfitting given cohort size, while GAP provides translation-robust, size-agnostic summarization of local patterns learned by the convolutions. Figure 2a details the process.

2. **Light attention embedding.** Given a memory tensor $x \in \mathbb{R}^{C \times D \times H \times W}$, we first reduce the in–plane resolution with average 3D pooling over $(1, 2, 2)$ to obtain $(C, D, 32, 32)$. A $1 \times 1 \times 1$ pointwise 3D convolution mixes the C input channels into a compact width *base*, followed by normalization and GELU, yielding $(base, D, 32, 32)$.

   Our deph attention block follows the squeeze-and-excitation operator [29]. We squeeze spatial information by global average pooling over (H, W) to produce depth descriptors $g \in \mathbb{R}^{base \times D}$. A two-layer MLP with GELU then excites these descriptors to obtain slice-wise logits, and a softmax across D provides normalized attention weights $w \in \mathbb{R}^{1 \times D}$. The weighted sum collapses depth:

$$\tilde{x}_{c,h,w} = \sum_{d=1}^{D} w_d \, x_{c,d,h,w}, \qquad \tilde{x} \in \mathbb{R}^{base \times 32 \times 32}.$$

   The resulting 2D map is resized to $S \times S$ via AdaptiveAvgPool2D, globally averaged to $(base, 1, 1)$, flattened to $\mathbb{R}^{base}$, and linearly projected to the final embedding $\in \mathbb{R}^{out\_dim}$. Figure 2b details the attention pipeline.

Discarded configurations

We also evaluated three additional downsampling heads: (i) a 3D CNN head operating on $((C \times D \times) 64 \times 64)$, (ii) direct flattening of the 64×64 maps, and (iii) PCA-based flattening, all under the same outer splits and training protocol. All three variants performed at chance (mean c-index ≈ 0.50, high p-values) and were therefore excluded from the final pipeline.

2.3.2 Fusion head

We combine modality embeddings (and optionally clinical covariates) via late fusion before the survival head.

(i) Concatenation (1D stacking).

Each modality-specific tower outputs a fixed-length vector $\mathbf{z}_{\mathrm{PET}}, \mathbf{z}_{\mathrm{CT}} \in \mathbb{R}^d$. Concatenation is performed as simple 1D stacking:

$$\mathbf{z}_{\mathrm{img}} = [\, \mathbf{z}_{\mathrm{PET}} \parallel \mathbf{z}_{\mathrm{CT}} \,] \in \mathbb{R}^{2d}.$$

When including clinical variables, a vector $\mathbf{z}_{\mathrm{clin}} \in \mathbb{R}^{d_{\mathrm{clin}}}$, the final fused input to the survival head is:

$$\mathbf{z}_{\mathrm{fused}} = [\, \mathbf{z}_{\mathrm{img}} \parallel \mathbf{z}_{\mathrm{clin}} \,] \in \mathbb{R}^{2d + d_{\mathrm{clin}}}.$$

This scheme is parameter-efficient, preserves modality-specific information, and naturally supports single-modality inputs.

(ii) Lightweight gated fusion (scalar mixing).

Alternatively, we add modality vectors $\mathbf{z}_{\text{PET}}, \mathbf{z}_{\text{CT}} \in \mathbb{R}^d$ using a scalar gate $\alpha \in (0, 1)$ learned via logistic regression on their 1D concatenation:

$$\alpha \;=\; \sigma\!\big(\mathbf{w}^\top [\,\mathbf{z}_{\text{PET}} \,\|\, \mathbf{z}_{\text{CT}}\,] + b\big), \qquad \mathbf{z}_{\text{img}} \;=\; \alpha\,\mathbf{z}_{\text{PET}} + (1 - \alpha)\,\mathbf{z}_{\text{CT}} \in \mathbb{R}^d.$$

This gate adds only $2d + 1$ parameters and yields an interpretable per-case modality weighting.

If clinical covariates are used, we again stack 1D:

$$\mathbf{z}_{\text{fused}} \;=\; [\,\mathbf{z}_{\text{img}} \,\|\, \mathbf{z}_{\text{clin}}\,] \in \mathbb{R}^{d + d_{\text{clin}}}$$

2.3.3 DeepSurv Head

The survival head maps the fused vector $\mathbf{z}_{\text{fused}}$ to a scalar linear predictor $\hat{h}(x_i)$ (higher $\hat{h}(x_i) =$ higher risk) using a deep feed-forward neural network. We update the standard DeepSurv model [15] to include weight decay, and early stopping on validation partial likelihood. The model is trained with the average negative log partial likelihood with regularization as the loss function:

$$l(\theta) := -\frac{1}{N_{E=1}} \sum_{i: E_i = 1} \left(\hat{h}_\theta(x_i) - \log \sum_{j \in \mathrm{R}(T_i)} \exp \hat{h}_\theta(x_i) \right) + \lambda\,||\theta||^2$$

where $N_{E=1}$ is the number of observed events, θ the parameters of the survival head, $\mathcal{R}(T_i)$ is the risk set just prior to T_i, and λ controls weight decay. Dropout and batch normalization are applied to hidden layers.

2.4 Experiments

2.4.1 Experimental setup

We evaluated three families of models on PFS: (i) image-only embeddings from PET/CT, (ii) *clinical-only* baselines, and (iii) *multimodal* models combining imaging embeddings with clinical covariates. MedSAM2 memory extraction was frozen, while the downsampler, fusion, and DeepSurv head were trained end-to-end per training fold. We used stratified 5-fold cross-validation preserving event proportion and approximate time quantiles. Optimization used AdamW cosine/step decay, dropout, BN, gradient clipping, and early stopping on validation partial likelihood. Hyperparameters were tuned by maximizing 5-fold cross-validated concordance using Sobol sequences in Optunity [30] to search the parameter space. Detailed search ranges and results are provided in the supplementary spreadsheet.

2.4.2 Models setup

Image-only

We ablated (a) modality: PET, CT, PET+CT; (b) ROI: Spine-dilated vs Skeleton; and (c) downsampler: Average embedding vs Light attention. PET+CT used late fusion (Section 2.3.2) and additionally tested fusion strategies: concatenation vs gated mixing.

Clinical-only

Two clinical baselines were trained using the dataset's clinical variables (section 2.1): CoxPH (ridge-penalized) and DeepSurv. Categorical variables were one-hot encoded, continuous variables were z-scored within each training fold and missing values were imputed from training-fold statistics.

Multimodal

We extend the best image-only configuration (downsampler + fusion) by adding clinical covariates. We report PET+CT+Clinical and single-modality+Clinical variants to assess complementarity with imaging.

2.4.3 Metrics and statistical analysis

Our primary endpoint is the mean Harrell's c-index [31] on outer test folds, reported as mean $\pm$ standard error (SE). For the best performing model we include the Kaplan-Mayer curve of the least and most at risk patient subsets. Significance is assessed with a one-sample t-test against a random model (c-index of 0.5); model comparisons use paired t-tests on fold-wise differences (as we used identical splits).

3 Results

We evaluated image-only ablations, multimodal variants, and baselines. Summary metrics are in Tables 1–3.

Ablations on downsampling and fusion were conducted using image-only models (Table 1). Across origins and masks, averaging consistently outperformed attention. The top results were obtained with (i) PET + Spine Dilated + Averaging (no image fusion; 0.659 ± 0.015, $p = 2.08 \times 10^{-4}$) and (ii) Combined + Full Skeleton + Averaging with Gated fusion (0.659 ± 0.018, $p = 4.84 \times 10^{-4}$).In contrast, Attention-based variants underperformed across all configurations (0.548–0.639 c-index).

Multimodal configurations were trained using the best downsampling choice from the ablation, that is, averaging, and are summarized in Table 2. The best multimodal configuration reached 0.710 ± 0.032 (CT + Spine Dilated), closely followed by Combined + Spine Dilated (0.710 ± 0.034) and PET + Spine Dilated (0.710 ± 0.040). Compared with the best image-only result, this corresponds to a relative improvement of $\sim 7.8\%$; compared with the best clinical baseline (Table 3), the improvement is $\sim 6.5\%$.

Table 1: Image-only configurations (sorted by c-index). Reported c-indexes are mean $\pm$ SE. An en dash (—) denotes not applicable.

Origin	Mask	Downsampling	Fusion	Test c-index	p-value
PET	Spine Dilated	Averaging	—	0.659 ± 0.015	2.081e-04
Combined	Full Skeleton	Averaging	Gated	0.659 ± 0.018	4.836e-04
Combined	Full Skeleton	Averaging	Concatenation	0.655 ± 0.022	9.975e-04
PET	Full Skeleton	Averaging	—	0.654 ± 0.022	1.128e-03
CT	Spine Dilated	Averaging	—	0.654 ± 0.008	1.983e-05
CT	Full Skeleton	Averaging	—	0.651 ± 0.021	9.451e-04
Combined	Spine Dilated	Averaging	Gated	0.651 ± 0.013	1.382e-04
Combined	Spine Dilated	Averaging	Concatenation	0.647 ± 0.008	3.085e-05
Combined	Full Skeleton	Attention	Concatenation	0.639 ± 0.013	2.497e-04
Combined	Spine Dilated	Attention	Gated	0.638 ± 0.024	2.344e-03
Combined	Spine Dilated	Attention	Concatenation	0.618 ± 0.012	3.136e-04
Combined	Full Skeleton	Attention	Gated	0.598 ± 0.007	9.568e-05
CT	Spine Dilated	Attention	—	0.586 ± 0.009	2.990e-04
CT	Full Skeleton	Attention	—	0.571 ± 0.015	4.956e-03
PET	Full Skeleton	Attention	—	0.560 ± 0.011	3.039e-03
PET	Spine Dilated	Attention	—	0.548 ± 0.012	8.132e-03

Table 2: Multimodal configurations (sorted by c-index).

Origin	Mask	Test c-index	p-value
CT	Spine Dilated	0.710 ± 0.032	1.308e-03
Combined	Spine Dilated	0.710 ± 0.034	1.697e-03
PET	Spine Dilated	0.710 ± 0.040	3.104e-03
Combined	Full Skeleton	0.709 ± 0.030	1.057e-03
PET	Full Skeleton	0.706 ± 0.031	1.244e-03
CT	Full Skeleton	0.703 ± 0.029	1.144e-03

Figure 3 shows Kaplan–Meier curves for the first and fourth risk quartiles produced by the best overall model (clinical + CT with the spine mask). Quartiles were computed from out-of-fold risk estimates standardized using in-fold statistics. The high- vs low-risk separation is statistically significant (log-rank $p = 3.14 \times 10^{-3}$).

4 Discussion

Our novel embedding strategy proves the viability of the usage internal states of foundational medical segmentation models as information rich embeddings for survival analysis in MM. Not only are the results comparable with radiomics models (without defining the features), but results show that our embedding added to clinical models shows an improvement to clinical only models (see Table 3).

In all settings (modality and mask), simple averaging across memory layers and channels outperformed the light attention downsampling alternative (Figure 5 shows

Table 3: Best models c-index (5-fold cross validation mean $\pm$ SE).

Model	c-index
Multimodal Model - Our Embedding + Clinical	0.710 ± 0.030
Clinical Only - DeepSurv	0.667 ± 0.035
Clinical Only - Cox	0.661 ± 0.025
Image Only - Our Embedding	0.659 ± 0.014
Image Only - Radiomics Baseline	0.657 ± 0.016

Fig. 3: Kaplan-Meyer curves of the first and fourth quartile risk patients from the best performing model (clinical variables + CT with the spine mask). Risk quartiles calculated from out-of-fold risk estimations standardized with in-fold estimations. Log-rank test p-value of 3.14e-03.

the difference between downsampling strategies). Memory states are highly redundant and smooth across slices, so averaging behaves as a variance-reducing low-pass filter that suppresses slice-specific noise and propagation artifacts. Additionally, the attention head introduces additional degrees of freedom that might not be reliably constrained, increasing the risk of overfitting. Consistent with this interpretation, averaging yielded lower standard errors and p-values.

With respect to multi-image setups, fusion of PET and CT, Fig. 4 shows that 1D concatenation and scalar-gated mixing deliver statistically indistinguishable discrimination (paired tests, $p > 0.05$). Both strategies access the same modality-specific vectors, and the DeepSurv head can effectively reweight concatenated features. Nevertheless, the gate remains attractive from an interpretability standpoint, as it provides a per-patient estimate of modality importance.

(a) Model performances by fusion strategy (hue) and mask (position) for multi-image model configurations with attention downsampling.

(b) Model performances by fusion strategy (hue) and mask (position) for multi-image model configurations with average downsampling.

Fig. 4: Comparison between both fusion strategies in imaging only models.

Within image-only models, PET consistently outperformed CT under the same mask, in line with prior work and clinical research [11, 32].

Finally, mask differences were statistically insignificant both in unimodal and multimodal models (p-values > 0.05).

Fig. 5: Aggregated model performances by downsampling strategy (hue) and origin (position) for multi-image model configurations. Reported c-index values correspond to the mean and terror across folds, obtained by first averaging the c-index of each model within each fold and then aggregating these fold-wise means across the cross-validation splits.

5 Limitations and future work

Limitations of this work include a small dataset with limited clinical information. The number of events restricts statistical power, increases fold-to-fold variability, and constrains the capacity of the downsampler and fusion heads. Additionally, external, multi-center validation is needed to assess generalization under scanner/reconstruction variability.

Additionally, the clinical covariates used when building the multimodal model were limited. We only used available variables with acceptable completeness. Using richer multi-omics variables would likely improve performance and allow for a deeper explainability analysis.

Our mask prompts introduce a prior over "where to look": masks focus the model on regions deemed anatomically informative. Although prompts were derived from an automatic CT segmentation trained independently of the folds, residual concerns about prompt-induced bias remain, as well as how much of the signal comes from the memory versus the prompt prior. Future work could compare our mask-derived prompts to coarser boxes and random prompts, test a "no-prompt" variant using only the encoder of the foundational model (e.g., Hiera [33]) to quantify the incremental value of memory states; and evaluate whether prompting on anatomically uninformative regions degrades performance to chance, clarifying the contribution of mask priors.

The scalar-gated fusion provides per-patient modality weighting that may aid case review; projecting embedding saliency back to the masks (e.g., Grad-CAM-style analysis over the downsampler) could further expose image patterns associated with risk, a deeper explainability study could explore these ideas.

6 Conclusion

This study demonstrates that mask-aware memory embeddings derived from the internal states of a foundational segmentation model can serve as effective imaging biomarkers for PFS in multiple myeloma.

Without handcrafting features, our memory-based embeddings achieved performance comparable to a strong radiomics baseline and improved a clinical-only model when fused with routine covariates (Table 3). These results support the central premise that foundational medical models encode rich, transferable information that can be repurposed—via lightweight heads and rigorous evaluation—to address prognostic tasks in small, censored cohorts.

Simple averaging across memory×channel dimensions consistently outperforms a lightweight attention alternative, suggesting a favorable bias–variance trade-off in this data regime. PET outperforms CT under identical masks, consistent with metabolic burden driving risk, while late-fusion choices (concatenation vs. scalar gating) deliver comparable discrimination.

Overall, foundational-model memory embeddings offer a bridge between handcrafted radiomics and end-to-end deep survival networks: anatomically grounded, computationally light, and readily extensible to multimodal fusion.

Supplementary information. Spreadsheet with optimization range and best model hyperparameters available.

Acknowledgements. Thanks to Matteo Bastico for the idea and to Pablo Perdomo-Quinteiro, Alessandro Ceresi, Anaida Fernández García and Francisco Moreno García for their comments and support.

Declarations

- Funding. This work was supported by the European Project: SYNTHIA (Synthetic Data Generation framework for integrated validation of use cases and AI healthcare applications) Grant No. 101172872 within the Innovative Health Initiative (IHI).
- Ethics approval and consent to participate: The data used in this work was gathered by IRCCS Azienda Ospedaliero-Universitaria Sant'Orsola-Malpighi in Bologna in compliance with Italian law.
  Data was accessed by UPM researchers with ethics compliance following Genomed (Grant no. 101017549) and SynthIA's (Grant No. 101172872) data sharing agreements.
- Consent for publication. Not applicable.
- Data availability: Data is not publicly available.
- Materials availability: Not applicable.
- Code availability: Upon request.

- Author contribution: **Javier Guinea-Pérez**: Conceptualization, Methodology, Software, Formal Analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization. **Silvia Uribe**: Writing - Review & Editing, Funding acquisition. **Sara Peluso**: Writing - Review & Editing, Data Curation, Resources. **Gastone Castelani**: Resources, Funding acquisition, Project administration. **Cristina Nanni**: Resources, Data Curation. **Federico Álvarez**: Supervision, Writing - Review & Editing, Methodology, Conceptualization, Funding acquisition, Project administration.

References

[1] Mafra, A. *et al.* The global multiple myeloma incidence and mortality burden in 2022 and predictions for 2045. *JNCI: Journal of the National Cancer Institute* djae321 (2024). URL https://doi.org/10.1093/jnci/djae321.

[2] Ludwig, H., Novis Durie, S., Meckl, A., Hinke, A. & Durie, B. Multiple Myeloma Incidence and Mortality Around the Globe; Interrelations Between Health Access and Quality, Economic Resources, and Patient Empowerment. *The Oncologist* **25**, e1406–e1413 (2020). URL https://doi.org/10.1634/theoncologist.2020-0141.

[3] SEER*Explorer Application. URL https://seer.cancer.gov/statistics-network/explorer/application.html?site=1&data_type=1&graph_type=2&compareBy=sex&chk_sex_3=3&chk_sex_2=2&rate_type=2&race=1&age_range=1&hdn_stage=101&advopt_precision=1&advopt_show_ci=on&hdn_view=0&advopt_show_apc=on&advopt_display=2#resultsRegion0.

[4] Rajkumar, S. V. *et al.* International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *The Lancet Oncology* **15**, e538–e548 (2014). URL https://www.thelancet.com/article/S1470-2045(14)70442-5/fulltext. Publisher: Elsevier.

[5] Agarwal, A., Chirindel, A., Shah, B. A. & Subramaniam, R. M. Evolving role of FDG PET/CT in multiple myeloma imaging and management. *AJR. American journal of roentgenology* **200**, 884–890 (2013).

[6] Rajkumar, S. V. & Kumar, S. Multiple myeloma current treatment algorithms. *Blood Cancer Journal* **10**, 94 (2020). URL https://www.nature.com/articles/s41408-020-00359-2. Publisher: Nature Publishing Group.

[7] Nanni, C. *et al.* Interpretation criteria for FDG PET/CT in multiple myeloma (IMPeTUs): final results. IMPeTUs (Italian myeloma criteria for PET USe). *European Journal of Nuclear Medicine and Molecular Imaging* **45**, 712–719 (2018). URL https://doi.org/10.1007/s00259-017-3909-8.

[8] Manco, L. *et al.* Positron Emission Tomography-Derived Radiomics and Artificial Intelligence in Multiple Myeloma: State-of-the-Art. *Journal of Clinical Medicine*

12, 7669 (2023). URL https://www.mdpi.com/2077-0383/12/24/7669. Publisher: Multidisciplinary Digital Publishing Institute.

[9] Kumar, S. K. & Rajkumar, S. V. The multiple myelomas — current concepts in cytogenetic classification and therapy. *Nature Reviews Clinical Oncology* **15**, 409–421 (2018). URL https://www.nature.com/articles/s41571-018-0018-y. Publisher: Nature Publishing Group.

[10] Terpos, E., Moulopoulos, L. A. & Dimopoulos, M. A. Advances in imaging and the management of myeloma bone disease. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* **29**, 1907–1915 (2011).

[11] Guinea-Pérez, J. *et al.* Radiomics feature analysis for survival prediction in multiple myeloma: An automated PET/CT approach. *Computer Methods and Programs in Biomedicine* **271**, 109019 (2025). URL https://www.sciencedirect.com/science/article/pii/S0169260725004365.

[12] Shreve, J. *et al.* P921: ARTIFICIAL INTELLIGENCE BASED FDG PET/CT RADIOMICS FOR RISK STRATIFICATION IN NEWLY DIAGNOSED MULTIPLE MYELOMA. *HemaSphere* **7**, e156011e (2023). URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10431033/.

[13] Zhong, H. *et al.* 18F-FDG PET/CT based radiomics features improve prediction of prognosis: multiple machine learning algorithms and multimodality applications for multiple myeloma. *BMC Medical Imaging* **23**, 87 (2023). URL https://doi.org/10.1186/s12880-023-01033-2.

[14] Ni, B. *et al.* Machine Learning Model Based on Optimized Radiomics Feature from 18F-FDG-PET/CT and Clinical Characteristics Predicts Prognosis of Multiple Myeloma: A Preliminary Study. *Journal of Clinical Medicine* **12**, 2280 (2023).

[15] Katzman, J. L. *et al.* DeepSurv: personalized treatment recommender system using a Cox proportional hazards deep neural network. *BMC Medical Research Methodology* **18**, 24 (2018). URL https://doi.org/10.1186/s12874-018-0482-1.

[16] Lee, C., Zame, W., Yoon, J. & Schaar, M. v. d. DeepHit: A Deep Learning Approach to Survival Analysis With Competing Risks. *Proceedings of the AAAI Conference on Artificial Intelligence* **32** (2018). URL https://ojs.aaai.org/index.php/AAAI/article/view/11842. Number: 1.

[17] Wiegrebe, S., Kopper, P., Sonabend, R., Bischl, B. & Bender, A. Deep learning for survival analysis: a review. *Artificial Intelligence Review* **57**, 65 (2024). URL https://doi.org/10.1007/s10462-023-10681-3.

[18] Bommasani, R. *et al.* On the Opportunities and Risks of Foundation Models (2022). URL http://arxiv.org/abs/2108.07258. ArXiv:2108.07258 [cs].

[19] Radford, A. *et al.* Learning Transferable Visual Models From Natural Language Supervision (2021). URL http://arxiv.org/abs/2103.00020. ArXiv:2103.00020 [cs].

[20] Kirillov, A. *et al.* Segment Anything (2023). URL http://arxiv.org/abs/2304.02643. ArXiv:2304.02643 [cs].

[21] Khan, W. *et al.* A Comprehensive Survey of Foundation Models in Medicine (2025). URL http://arxiv.org/abs/2406.10729. ArXiv:2406.10729 [cs].

[22] Ma, J. *et al.* Segment anything in medical images. *Nature Communications* **15**, 654 (2024). URL https://www.nature.com/articles/s41467-024-44824-z. Publisher: Nature Publishing Group.

[23] Ma, J. *et al.* Segment Anything in Medical Images and Videos: Benchmark and Deployment (2024). URL http://arxiv.org/abs/2408.03322. ArXiv:2408.03322 [eess].

[24] Zhu, W., Chen, Y., Nie, S. & Yang, H. *SAMMS: Multi-modality Deep Learning with the Foundation Model for the Prediction of Cancer Patient Survival*, 3662–3668 (2023). URL https://ieeexplore.ieee.org/document/10385661. ISSN: 2156-1133.

[25] Palumbo, A. *et al.* Revised International Staging System for Multiple Myeloma: A Report From International Myeloma Working Group. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* **33**, 2863–2869 (2015).

[26] Sundar, L. K. S. *et al.* Fully Automated, Semantic Segmentation of Whole-Body ^{18}F-FDG PET/CT Images Based on Data-Centric Artificial Intelligence. *Journal of Nuclear Medicine* **63**, 1941 (2022). URL http://jnm.snmjournals.org/content/63/12/1941.abstract.

[27] Ma, J. *et al.* MedSAM2: Segment Anything in 3D Medical Images and Videos (2025). URL http://arxiv.org/abs/2504.03600. ArXiv:2504.03600 [eess].

[28] Ravi, N. *et al.* SAM 2: Segment Anything in Images and Videos (2024). URL http://arxiv.org/abs/2408.00714. ArXiv:2408.00714 [cs].

[29] Hu, J., Shen, L., Albanie, S., Sun, G. & Wu, E. Squeeze-and-Excitation Networks (2019). URL http://arxiv.org/abs/1709.01507. ArXiv:1709.01507 [cs].

[30] Claesen, M., Simm, J., Popovic, D., Moreau, Y. & Moor, B. D. Easy Hyperparameter Search Using Optunity (2014). URL http://arxiv.org/abs/1412.1114. ArXiv:1412.1114.

[31] Harrell, F. E., Califf, R. M., Pryor, D. B., Lee, K. L. & Rosati, R. A. Evaluating the yield of medical tests. *JAMA* **247**, 2543–2546 (1982).

[32] Zamagni, E. *et al.* 18F-FDG PET/CT focal, but not osteolytic, lesions predict the progression of smoldering myeloma to active disease. *Leukemia* **30**, 417–422 (2016). URL https://www.nature.com/articles/leu2015291. Publisher: Nature Publishing Group.

[33] Ryali, C. *et al.* *Hiera: a hierarchical vision transformer without the bells-and-whistles*, Vol. 202 of *ICML '23*, 29441–29454 (JMLR.org, Honolulu, Hawaii, USA, 2023).