Coverage and Utilization of Digital Healthcare Services: A Registry-Based Observational Study

Alexandra Dahlberg[1,2*], Taavi Kaartinen[2,3], Sakari Jukarainen[2,3], Petja Orre[1,3]

[1]Harjun terveys, Lahti, Finland

[2]Faculty of Medicine, University of Helsinki, Helsinki, Finland

[3]Mehiläinen, Helsinki, Finland

*Lead author and corresponding author, alexandra.dahlberg@harjunterveys.fi

November 6th, 2025

Abstract

A retrospective, registry-based analysis describing the uptake, user characteristics, and clinical scope of a digital primary care clinic within the Finnish public health system. The study utilized data from 2,635,708 primary care encounters recorded between 2019 and 2024 to examine patterns of care following the digital clinic's introduction. By 2024, digital visits accounted for 12.7% of all primary care encounters. Users of the digital clinic were significantly younger, more likely female, and had fewer comorbidities compared to those using traditional care, with notably low uptake among adults over 65. The most common conditions managed digitally were acute respiratory infections, dermatologic issues, and urinary tract infections. Follow-up analysis showed that 21% of patients had a new encounter within 30 days, with in-person visits being the most common subsequent modality. This study demonstrates that a digital clinic can expand access and resolve nearly half of cases without even a digital physician consultation, while also functioning as a triage system for cases requiring further care. However, targeted strategies are needed to increase adoption among older adults and individuals with chronic illnesses.

Introduction

Digital health platforms are transforming how patients access primary care. Asynchronous messaging and live chat, video visits, remote monitoring, patient portals, and digital care pathways create new routes to improve accessibility, timeliness, and operational efficiency[1-3]. In many countries, the COVID-19 pandemic accelerated the adoption of digital care models[4,5], though interest in digital-first approaches predated the crisis[6]. For publicly funded health systems like Finland's, where digital transformation is a strategic priority and equitable access remains a central policy objective, the implementation of these technologies requires thorough assessment to ensure they support accessibility, equity, and overall system efficiency.

Economic evaluations suggest that digital-first pathways can be cost-effective in real-clinical settings[7,8]. However, understanding the demographic and clinical characteristics of digital care users is essential for designing inclusive and scalable digital health systems. Identifying patterns in utilization – such as differences by age, sex, or diagnosis – and examining changes in overall coverage over time can shed light on both the potential and limitations of digital modalities. For instance, studies in European contexts have shown that digital care users tend to be younger, more likely female, and more socioeconomically advantaged than the general patient population[9,10].

This study evaluates population coverage and real-world utilization. Using registry data from Harjun terveys in Finland, we examine the integration of a 24/7 digital primary care clinic within the publicly funded service model. Specifically, we assessed: (1) trends in overall primary care coverage; (2) the volume and proportion of digital visits

and their follow-ups; (3) user characteristics across modalities, including age, sex, and

comorbidity; and (4) the diagnostic spectrum of the digital encounters.

Methodology

Study design and setting

We conducted a retrospective cohort study using routinely collected health service

data from Harjun terveys in Päijät-Häme, Finland. The observation window spanned

January 1, 2019, to December 31, 2024, providing two pre-implementation years

(2019-2020) and up to four post-implementation years (2021-2024) following the

Harjun terveys joint-venture model. System-wide digital services were introduced in

January 2021, followed by the launch of a centralized digital clinic with a new operating

model in January 2023 and the implementation of new software in April 2023.

Finland provides universal healthcare coverage primarily through publicly funded

services, with private providers in a complementary role. Three components operate

in parallel: public, private, and occupational healthcare[11]. In some municipalities, parts

of public service delivery are outsourced to private healthcare providers to address

local capacity constraints, enhance service quality, and improve cost efficiency.

Harjun terveys is a joint venture that delivers publicly funded primary health care,

including outpatient medical and dental services, in the Päijät-Häme region of Finland,

serving approximately 130,000 residents. Established in 2021 as a partnership

between the regional wellbeing authority and a private healthcare company, the

venture was created to address insufficient local capacity to meet patient demand. In

addition to expanding service coverage, Harjun terveys implemented a

multidisciplinary team-based model and a "digi-physical" service pathway. All residents

can access a 24/7 digital clinic via the Päijät-Sote application, which provides chat-

based consultations with nurses and physicians alongside traditional telephone and in-

person appointments. The model is intended to reduce appointment delays and

improve access.

All primary care encounters – whether a physical visit at a health center, a telephone

triage, or an asynchronous chat consultation via the app – are documented in the

national unified electronic health record (EHR) system. The digital clinic is staffed by

nurses and general practitioners (GPs) who provide consultation, prescriptions, and

referrals through chat messaging, with the ability to escalate patients to physical

appointments when needed. Harjun terveys serves an urban-rural mixed population,

with a slightly older age distribution than the national average[12].

Data Extraction

We obtained pseudonymized encounter-level data. For each primary care episode of

care, we extracted: patient demographics (year of birth, sex, and home municipality),

encounter modality, dates and times of encounters, provider type (nurse or physician),

and diagnosis codes. Chronic conditions (e.g., diabetes, ischemic heart disease,

asthma/COPD) were identified using prior diagnostic history for the studied period.

Each patient's Charlson Comorbidity Index (CCI) was calculated using an adapted

algorithm for ICD-10 code mapping[13].

Primary outcome measures included service coverage and utilization patterns. Service

coverage was defined as the proportion of the population (Päijät-Häme residents

served by Harjun terveys) with at least one primary care encounter during a year, and

utilization as encounters per 1,000 residents. A primary care encounter was defined

as any encounter with a nurse or a general practitioner at a primary care facility, for

example the digital clinic, or a GP office, excluding encounters related to health

promotion such as vaccinations and maternity or child preventative health clinic visits.

Coverage was stratified by modality to assess engagement via digital versus traditional

channels. Population denominators were drawn from national population data[12].

Digital service utilization was analyzed by calculating the number and share of unique

digital users annually, alongside the total number of digital encounters. Digital users

were defined as individuals with at least one digital clinic encounter during the studied

year; non-digital users were those who did not use the digital clinic.

To characterize the clinical scope of digital consultations, we examined the most

frequent diagnoses recorded in digital clinic encounters during 2023-2024. Diagnoses

were analyzed using both ICPC-2 codes, which reflect presenting symptoms and

primary care-level assessments, and ICD-10 codes, which were used when patients

were escalated to a physician for diagnosis or prescription.

Statistical Analysis

We present proportions, means (with standard deviations), or medians (with

interquartile ranges) where appropriate. Differences in demographic characteristics

between digital users and non-users were tested using χ^2 tests for categorical variables

and t-tests for continuous variables. All statistical tests were two-tailed with a

significance level of $P<.05$. Data analyses were performed using Python[14-18].

The study was approved by the Päijät-Häme Wellbeing Services County for secondary

use of health data (decision number HA/187/07.01.04.05/2025). This study was based

on secondary use of pseudonymized administrative data. In accordance with the

Finnish Act on the Secondary Use of Health and Social Data (552/2019)[19] and the

research permit, individual patient consent or separate ethical review was not required.

Results

The initial extract comprised 2,706,347 primary care encounters from 1 January 2019 through 31 December 2024 (inclusive). After deduplication, exclusion of encounters with unknown profession codes or out-of-scope professional categories, and removal of records with missing date of birth, 2,635,708 encounters remained for analysis. This represents a net reduction of 70,639 records, or 2.61%.

Primary Care Coverage and Service Utilization

We examined trends in primary care coverage and service utilization in the Päijät-Häme region from 2019 to 2024, a period that encompasses the introduction of digital services and the subsequent implementation of a new centralized operational model for the digital clinic. For a comprehensive interpretation of the findings, it is crucial to contextualize the state of healthcare services prior to the intervention. Upon the initiation of the joint venture, the services provided by the local municipal authorities had been considered suboptimal for a prolonged period, particularly in terms of access to care and availability. Consequently, a significant portion of the subsequent increase in service utilization may be attributed to broad enhancements aimed at better meeting the population's healthcare needs, rather than being a direct outcome of the digital clinic's introduction.

Annual coverage increased gradually from 25.3% in 2019 to 28.3% in 2024 (Table 1). Over the same period, total utilization rose from 1,473 to 1,886 encounters per 1,000 residents. The uptick in utilization accelerated after 2021 and continued through 2024. Because the centralized digital clinic began on 1 January 2023, changes observed in 2021-2022 are partly interpreted as part of an ongoing system transition rather than effects of the centralized clinic.

Table 1. Annual primary care coverage (%) and encounters per 1,000 population, 2019–2024. *Digital share of the encounters. Coverage based on national population data [12].*

Year	Coverage (%)	Encounters per 1,000	Digital Share (%)
2019	25.3	1473	0
2020	24.4	1382	0
2021	25.9	1548	2.9
2022	27.9	1771	4.0
2023	28.3	1888	8.7
2024	28.3	1886	12.7

Digital encounters expanded quickly, accounting for 12.7% of all recorded interactions

in 2024. In-person visits remained the predominant modality but declined in relative

terms, while phone contacts were stable.

Conversely, among older adults, particularly those over 75, digital care did not

substantially expand reach as seen in Figure 1. Coverage in this group remained near

pre-pandemic levels, highlighting a potential digital divide in access or preferences for

traditional modes of care.

Figure 1 Health service coverage in the area by age distribution. *Proportion of residents in each age group with*
at least one healthcare encounter during the year. The gray line indicates the difference in coverage between areas
with and without access to digital health services in 2024.

Table 1 and Figure 1 together highlight a consistent pattern: the introduction of digital

primary care improved service reach, particularly among digitally literate and younger

populations.

Demographic Profile of Digital Service Users

By 2024, digital health services had been adopted across a broad range of age groups,

though uptake remained uneven. Younger adults were most likely to engage with the

digital clinic, particularly those aged 20 to 39 years. Uptake declined progressively in

older age brackets, with notably lower participation among those aged 65 and above.

As shown in Table 2, digital service users were substantially younger than those using

traditional modalities. This was reflected in both mean and median age comparisons,

with statistically significant differences observed across years. Women were more

likely than men to access digital services.

Table 2 Demographic characteristics of digital and traditional service users in 2023 and 2024. *Comparison of age and sex distribution between users of digital clinic services and those utilizing traditional encounter types. CCI stands for the Charlson comorbidity index. P-values for age are calculated using a two-sided t-test; sex distribution and users with CCI scores are compared using the chi-square test.*

	2023			2024		
	Digital user	Traditional user	*P*-value	Digital user	Traditional user	*P*-value
Healthcare users (n)	523	10,558	-	7,334	10,146	-
Healthcare users (% of total)	33.1%	66.9%	-	42.0%	58.0%	-
Mean age (SD)	22.1 (18.9)	36.0 (24.8)	< .001	22.8 (18.8)	38.1 (25.1)	< .001
Median age (IQR)	22.0 (29.0)	32.0 (41.0)	< .001	22.0 (29.0)	35.0 (42.0)	< .001
Female (%)	61.2%	55.4%	< .001	60.8%	54.7%	< .001
Male (%)	38.8%	44.6%	< .001	39.2%	45.3%	< .001
CCI ≥ 1 (%)	7.1%	12.2%	< .001	6.0%	12.2%	< .001

Users of the digital clinic also differed clinically. A smaller proportion of digital users

documented chronic conditions compared to those using traditional encounters. This

suggests a preference or suitability of digital modalities for relatively healthier patients

or those with less complex care needs. These patterns were consistent with the

previous year's comparison and were statistically significant based on chi-square

analysis.

Figure 2 illustrates the distribution of encounter types by age group. For each

encounter modality – digital, phone, and in-person – only a single encounter per person

was counted. This allows for a clearer interpretation of service penetration within each

age band. Physical visits and phone calls were most common among the oldest age

groups, whereas digital usage peaked in the groups for early adulthood and declined

thereafter.

Figure 2 Age distribution of encounter types in 2024. *Proportion of the population with at least one healthcare*
encounter in 2024, by age group and encounter type. Encounter types include digital clinics, physical visits, and
phone calls. Proportions are calculated using unique individuals per age group and encounter type, relative to the
age-matched population. Only one encounter per person per type is counted.

Digital Consultation Characteristics: Diagnoses and Conditions Managed

Figure 3 displays the 20 most common ICPC-2 codes recorded in the digital clinic,

collectively representing 48.2% of all interactions. The most common reason for

contact was respiratory infections (R74 and R83, 13.0%). Other frequent codes

included general symptom consultations such as health maintenance or preventive

advice (-45 and A98, 8.1%), various skin symptoms (S29 and S21, 5.0%), and throat

or eye-related complaints (R21, F03, and F29, 6.8%). Notably, administrative

interactions related to prescription renewal also featured (-50, 2.7%), indicating that

digital services were used not only for acute care but also for preventive and

administrative functions.

Figure 3. Most common ICPC-2 diagnosis categories in the digital clinic (2023–2024). *The 20 most frequent*
ICPC-2 codes recorded in digital clinic consultations, shown as the percentage of all digital clinic encounters.

Figure 4 summarizes the 20 most frequent ICD-10 diagnoses among patients

escalated to the digital clinics physician-level care. The most common conditions were

prescription renewal requests (Z76.0, 1.3%), acute cystitis (N30.0, 1.2%), and

conjunctival infections (H10.x series, 1.7%). Additional diagnoses reflected a typical

profile of low-complexity primary care issues, including cellulitis, dermatitis, and

streptococcal pharyngitis. These findings suggest that patients referred to digital

consultations with physicians typically require evaluation for infections or

dermatological conditions, often with a prescription.

Figure 4. Most common ICD-10 diagnosis categories in the digital clinic (2023-2024). *Top 10 ICD-10 codes*
recorded during digital consultations with a physician, following escalation from the nurse encounter at the digital
clinic. Percentages represent the share of all digital clinic encounters.

Together, these distributions highlight the dual clinical role of the digital clinic:

managing common acute complaints directly, and efficiently triaging cases that require

further care. The high frequency of minor infections and symptom-based codes further

supports the interpretation that digital consultations are primarily used for low-acuity,

high-frequency presentations that benefit from rapid, accessible care.

Follow-up Analysis

To evaluate care continuity following digital nurse consultations, we analyzed

subsequent patient encounters over a 30-day period. The cohort included 86,258

digital nurse encounters. On the same day as the initial digital encounter, 24.7% (n =

21,336) involved a digital consultation with a physician, while 14.4% (n = 12,397)

resulted in a new appointment being booked.

To isolate unplanned or deferred care needs as best as possible, we excluded follow-

up encounters that had already been booked at the time of the digital interaction. The

analysis therefore focused exclusively on the first new encounter occurring more than

24 hours after the digital consultation.

Across all follow-up modalities, 5.0% of patients had a new encounter within 48 hours,

14.9% within 14 days, and 21.0% within 30 days. Figure 5 illustrates the cumulative

incidence of follow-up activity over the 30-day period, stratified by encounter type.

Physical visits represented the most common form of follow-up, with over 10% of

patients receiving in-person care within 30 days. Phone-based interactions, digital

revisits, and other administrative or triage-related encounters accounted for additional

follow-up activity.

Figure 5 Cumulative follow-up by encounter type over 30 days following an encounter with the digital clinic.
Follow-ups that had been booked prior to the encounter with the digital clinic are excluded. Dashed lines indicate
follow-up encounters that were scheduled on the same day as the digital clinic visit.

To examine planned follow-up, we evaluated encounters that were booked on the

same day as the digital nurse consultation but scheduled for a later date. The likelihood

of such pre-booking varied substantially by encounter type. Among patients who had

a follow-up within 30 days, 39% of phone calls were booked during the initial digital

consultation, compared to 17% of physical visits and 11% of other encounter types.

For these pre-booked follow-ups, the most common presenting reasons included

naevus/mole, laceration/cut, acute upper respiratory infections, diabetes, sleep

disturbances and more general preventive or health maintenance codes. These are

conditions for which a subsequent follow-up call or in-person evaluation is often

clinically warranted.

To illustrate follow-up care utilization, Figure 6 presents a Sankey diagram of

healthcare utilization after a digital clinic visit. Flow widths correspond to the share of

patients at each step. For readability, paths are limited to three steps, and the horizon

is 14 days. The diagram takes into account all healthcare utilization, planned and un-

planned. Nearly half of trajectories terminate at the nurse-led digital clinic visit, 14%

proceed to a physician consultation, and 14% culminate in an in-person visit, with

smaller fractions continuing via phone contacts or administrative interactions.

258
259
260
Figure 6 Sankey diagram of healthcare utilization after a digital clinic visit. *Flow widths correspond to the proportion of patients at each step. The diagram covers the first 14 days after the visit, and sequences are truncated at three steps to aid interpretation.*

Discussion

This study assessed a digital-first primary care service embedded in a Finnish public

system and additionally reported primary healthcare coverage to inform equity

considerations. Following the implementation of the digital clinic, overall healthcare

coverage expanded, with the digital clinic contributing to 12.7% of all annual primary

care encounters in the area in 2024. Because implementation coincided with other

joint-venture changes, the relative contribution of the digital channel to the coverage

increase cannot be isolated with the present design. Interpretation should therefore be

conservative with respect to attribution.

The demographic analysis showed higher adoption among younger adults and women,

consistent with findings from other settings[9,10]. These groups were more likely to

engage with the chat-based platform, while older adults and individuals with multiple

chronic conditions continued to rely primarily on traditional care. While lower

engagement among older patients could reflect usability barriers[20], preferences[21,22] or

differences in clinical appropriateness, it raises important equity concerns. Strategies

such as assisted digital onboarding or targeting digital support to specific pathways for

chronic conditions may improve inclusion.

Most digital encounters were managed entirely online, with approximately 85.6%

resolved without a need to book a new appointment. Across all follow-up modalities,

5.0% of patients had a new encounter within 48 hours, 14.9% within 14 days, and

21.0% within 30 days, excluding those booked prior to the digital encounter. These

resolution rates reflect the appropriateness of the digital modality for managing acute,

low-complexity conditions, which constituted the bulk of digital encounters. Our

findings are consistent with previous studies such as Glock et al.[10], who found that

text-based consultations were suited for uncomplicated, low-complexity presentations.

They found that about one-quarter of Swedish eVisit patients required in-person follow-

up within 14 days (for the same diagnostic group), which was higher than the 15%

observed in our cohort.

Usage patterns of the digital clinic also revealed that most interactions were episodic

rather than part of chronic care management. Few high-frequency or chronic patients

used the digital clinic for their routine follow-up needs, although a small number used

it for incidental concerns. Expanding the scope of digital services to include selected

chronic care tasks – such as remote monitoring or asynchronous prescription follow-

up – may represent an area for future innovation[23]. Similarly, digital utilization for

mental health-related concerns was also modest, despite the potential of digital

platforms to lower access barriers. Prior research suggests this may reflect limitations

in perceived personal connection, lack of trust, and concerns about privacy and

confidentiality in digital encounters[24,25]. Finland's Finnish Therapy Navigator, created

through a government-funded program and implemented in the digital clinic, offers an

example of emerging tools that could enhance digital pathway guidance and uptake[26].

Ensuring equitable access to digital health requires deliberate design and

implementation. Evidence indicates that digital services can improve access for

specific populations[27,28]. The corresponding policy question is national in scope: what

minimum level of care every public provider should be required to deliver, and through

which modalities. In line with previous recommendations[29-32], hybrid models that

preserve physical care options, improve users' digital skills, and offer personalized

support (including remote digital assistance) are crucial. Equally important is the

development of user-centered platforms with intuitive design and clear guidance for

patients and clinicans[20,32].

Finally, while we did not conduct a formal cost analysis in this study, prior work in the

same setting has demonstrated significant cost reductions for digitally initiated

episodes when compared to matched traditional care episodes[8]. Taken together, these

findings suggest that digital-first primary care, when carefully implemented and fully

integrated into the broader health system, can help to improve access and streamline

service use. Future research should continue to explore long-term outcomes, equity

impacts, and patient experiences to ensure that digital transformation enhances both

efficiency and inclusiveness in healthcare delivery.

Limitations

This study relied on registry healthcare data, which has inherent limitations. Some

outcomes like clinical resolution or patient satisfaction were not directly measured; we

inferred resolution from follow-up rates, which could miss cases where patients sought

care outside the system under study. There is also potential misclassification in

diagnostic coding. Since this was an observational study within one wellbeing services

county, generalizability may be limited; factors unique to Päijät-Häme (such as the

specific app used or population characteristics) might affect results. Additionally, our

comparisons between digital and traditional users are subject to confounding – the

groups differ by age and health status by nature.

Attribution should be cautious, concurrent system improvements under the joint

venture mean that attributing all observed changes to the digital channel risks

overstatement. As a descriptive study, we cannot establish causality.

Conclusion

The integration of a digital primary care clinic in the Harjun terveys service network has

significantly influenced healthcare coverage and utilization patterns in Päijät-Häme,

Finland. Digital health services via the Päijät-Sote app have expanded access to care,

enabling a notable portion of the population – especially younger adults – to receive

timely care for acute issues without needing physical visits. The digital-first model

achieved high resolution rates for uncomplicated conditions, and most patients did not

require additional in-person follow-up, indicating effective management. We observed

distinct user demographics, with digital services attracting younger and relatively

healthier patients, while older and chronically ill patients continued to use traditional

care channels. Common conditions like skin ailments, respiratory infections, and minor

urinary issues were handled efficiently online, whereas more complex concerns were

not as frequent in the digital clinic.

Our findings suggest that a well-implemented digital clinic can serve as a

complementary extension of primary care. However, to ensure that the benefits are

broadly shared, strategies to engage under-represented groups in digital care are

needed, as is ongoing monitoring of quality and outcomes. As health systems

worldwide continue to adopt digital health innovations, the Harjun terveys experience

provides encouraging evidence that access can be widened, and care streamlined.

Future research and quality improvement efforts will focus on refining the model – for

example, through patient education, interface improvements, and possibly expanding

digital offerings to chronic care – and rigorously evaluating long-term health outcomes

in this digital-enhanced primary care paradigm.

Declaration statements

Data Availability

All data generated or analyzed during this study are presented within the published

article. Individual-level patient data are not publicly available due to privacy and data

protection regulations.

Code Availability

The   underlying   code   for   this   study   is   available   at

https://github.com/achdahlberg/coverage-utilization-digital-clinic-2025.

Acknowledgements

The authors gratefully acknowledge the support provided by Mehiläinen Oy, Harjun terveys oy, and the Päijät-Häme Wellbeing Services County. ChatGPT [33] was used to assist in improving the clarity and coherence of the manuscript and to support routine programming tasks; it was not used for conceptual input, interpretation of findings, or generation of original content. All content was thoroughly reviewed, edited, and approved by the authors, who take full responsibility for the accuracy and integrity of the final work.

Authors' Contributions

A.D. was responsible for the study's conceptual framework and design, performed the data analysis, and prepared the first draft of the manuscript. S.J. oversaw the research, offered methodological support, and provided critical revisions to the manuscript. T.K. supported the development of the digital health context, advised on data organization, and contributed to manuscript refinement. P.O. offered strategic guidance during the study design phase, enabled access to key data sources, and assisted in manuscript development. All authors participated in the interpretation of the results and approved the final manuscript for submission.

Competing Interests

The authors disclose the following potential conflicts of interest: All authors are employees of Mehiläinen, the healthcare organization that developed and operates the digital clinic assessed in this study. Some authors also hold personal financial interests in the company. The study received funding from Mehiläinen, and institutional support was provided by the leadership of Harjun terveys throughout the research process.

References

Dorsey, E. R. & Topol Eric, J. State of Telehealth. *New England Journal of Medicine* **375**, 154-161 (2016). https://doi.org/10.1056/NEJMra1601705

De Guzman, K. R., Snoswell, C. L., Caffery, L. J. & Smith, A. C. Economic evaluations of videoconference and telephone consultations in primary care: A systematic review. *Journal of Telemedicine and Telecare* **30**, 3-17 (2021). https://doi.org/10.1177/1357633X211043380

Campbell, K. *et al.* The Impact of Virtual Consultations on the Quality of Primary Care: Systematic Review. *J Med Internet Res* **25**, e48920 (2023). https://doi.org/10.2196/48920

Neves, A. L. & Burgers, J. Digital technologies in primary care: Implications for patient care and future research. *Eur J Gen Pract* **28**, 203-208 (2022). https://doi.org/10.1080/13814788.2022.2052041

Ward, K., Vagholkar, S., Sakur, F., Khatri, N. N. & Lau, A. Y. S. Visit Types in Primary Care With Telehealth Use During the COVID-19 Pandemic: Systematic Review. *JMIR Med Inform* **10**, e40469 (2022). https://doi.org/10.2196/40469

Ministry of Social Affairs and Health.   (2023).

Gentili, A. *et al.* The cost-effectiveness of digital health interventions: A systematic review of the literature. *Frontiers in Public Health* **10** (2022). https://doi.org/10.3389/fpubh.2022.787135

Dahlberg, A., Jukarainen, S., Kaartinen, T. & Orre, P. Cost minimization analysis of digital-first healthcare pathways in primary care. *npj Digital Medicine* **8**, 546 (2025). https://doi.org/10.1038/s41746-025-01937-z

Darley, S. *et al.* Understanding How the Design and Implementation of Online Consultations Affect Primary Care Quality: Systematic Review of Evidence With Recommendations for Designers, Providers, and Researchers. *J Med Internet Res* **24**, e37436 (2022). https://doi.org/10.2196/37436

Glock, H. *et al.* eVisits to primary care and subsequent health care contacts: a register-based study. *BMC Primary Care* **25**, 297 (2024). https://doi.org/10.1186/s12875-024-02541-y

Eskola, P., Tuompo, W., Riekki, M., Timonen, M., & Auvinen, J. .   (ed Finnish Ministry of Social Affairs and Health) (Institutional Repository for the Government, 2022).

Official Statistics of Finland. *Population structure*, <https://stat.fi/tilasto/vaerak> (2025).

Brusselaers, N. & Lagergren, J. The Charlson Comorbidity Index in Registry-based Research. *Methods Inf Med* **56**, 401-406 (2017). https://doi.org/10.3414/me17-01-0051

Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357-362 (2020). https://doi.org/10.1038/s41586-020-2649-2

pandas-dev/pandas: Pandas v. latest (Zenodo, 2020).

Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering* **9**, 90-95 (2007). https://doi.org/10.1109/MCSE.2007.55

Waskom, M. L. seaborn: statistical data visualization. *Journal of Open Source Software* **6**, 3021 (2021). https://doi.org/10.21105/joss.03021

Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods* **17**, 261-272 (2020). https://doi.org/10.1038/s41592-019-0686-2

Act on the Secondary Use of Health and Social Data, <https://www.finlex.fi/fi/laki/ajantasa/2019/20190552> (552/2019).

Kaihlanen, A.-M. *et al.* Towards digital health equity - a qualitative study of the challenges experienced by vulnerable groups in using digital health services in the COVID-19 era. *BMC Health Services Research* **22**, 188 (2022). https://doi.org/10.1186/s12913-022-07584-4

Mangin, D. *et al.* Multimorbidity, eHealth and implications for equity: a cross-sectional survey of patient perspectives on eHealth. *BMJ Open* **9**, e023731 (2019). https://doi.org/10.1136/bmjopen-2018-023731

439	22	Frishammar, J., Essén, A., Bergström, F. & Ekman, T. Digital health platforms for the elderly? Key adoption and usage barriers and ways to address them. *Technological Forecasting and Social Change* **189**, 122319 (2023). https://doi.org/https://doi.org/10.1016/j.techfore.2023.122319
443	23	Melchiorre, M. G. *et al.* eHealth in integrated care programs for people with multimorbidity in Europe: Insights from the ICARE4EU project. *Health Policy* **122**, 53-63 (2018). https://doi.org/https://doi.org/10.1016/j.healthpol.2017.08.006
446	24	Berry, N., Lobban, F. & Bucci, S. A qualitative exploration of service user views about using digital health interventions for self-management in severe mental health problems. *BMC Psychiatry* **19**, 35 (2019). https://doi.org/10.1186/s12888-018-1979-1
449	25	Liberati, E. *et al.* Remote care for mental health: qualitative study with service users, carers and staff during the COVID-19 pandemic. *BMJ Open* **11**, e049210 (2021). https://doi.org/10.1136/bmjopen-2021-049210
452	26	Saarni, S. *et al.* The Finnish Therapy Navigator – Digital support system for introducing stepped care in Finland. *Psychiatria Fennica* **53**, 120-137 (2022).
454	27	Erku, D. *et al.* Digital Health Interventions to Improve Access to and Quality of Primary Health Care Services: A Scoping Review. *Int J Environ Res Public Health* **20** (2023). https://doi.org/10.3390/ijerph20196854
457	28	Talwar, G. & Sinha, S. How are digital platforms affecting the accessibility of healthcare services? *Health Services Management Research*, 09514848251358328 (2025). https://doi.org/10.1177/09514848251358328
460	29	Beaunoyer, E., Dupéré, S. & Guitton, M. J. COVID-19 and digital inequalities: Reciprocal impacts and mitigation strategies. *Computers in Human Behavior* **111**, 106424 (2020). https://doi.org/https://doi.org/10.1016/j.chb.2020.106424
463	30	Nguyen, M. H., Hargittai, E. & Marler, W. Digital inequality in communication during a time of physical distancing: The case of COVID-19. *Computers in Human Behavior* **120**, 106717 (2021). https://doi.org/https://doi.org/10.1016/j.chb.2021.106717
466	31	Xie, B. *et al.* When Going Digital Becomes a Necessity: Ensuring Older Adults' Needs for Information, Services, and Social Inclusion During COVID-19. *Journal of Aging & Social Policy* **32**, 460-470 (2020). https://doi.org/10.1080/08959420.2020.1771237
469	32	Wilson, S. *et al.* Recommendations to advance digital health equity: a systematic review of qualitative studies. *npj Digital Medicine* **7**, 173 (2024). https://doi.org/10.1038/s41746-024-01177-7
472	33	OpenAI. ChatGPT [Large language model]. (2025).