Informal social support for chronic pain and distress in people with HIV: identifying

targets for intervention

Short title: Informal social support for chronic pain and distress in people with HIV

Authors: Victoria J Madden[1,2*], Lara Court[1,3], Andiswa Gidana[1], Lucia Knight[3,4], Antonia

Wadley[5], Luyanduthando Mqadi[1], Stephan Rabie[2], Robert R Edwards[6], John A Joska[2],

Romy Parker[1]

[1] African Pain Research Initiative, Department of Anaesthesia and Perioperative Medicine,
Neuroscience Institute, University of Cape Town, Cape Town, South Africa
[2] HIV Mental Health Research Unit, Department of Psychiatry and Mental Health,
Neuroscience Institute, University of Cape Town, Cape Town, South Africa
[3] Division of Social and Behavioural Sciences, School of Public Health, University of Cape
Town, South Africa
[4] School of Public Health, University of the Western Cape, Bellville, South Africa
[5] Brain Function Research Group, Department of Physiology, Faculty of Health Sciences,
University of the Witwatersrand, Johannesburg, South Africa
[6] Department of Anesthesiology, Perioperative, and Pain Medicine, Harvard Medical School,
Boston, Massachusetts, USA
18

19 ***Corresponding author**

20 Email: torymadden [at] gmail [dot] com

Abstract

Social factors are consequences and determinants of chronic pain, yet little is known about

how informal social support operates for people with chronic pain and HIV. The overlapping

burdens of emotional distress and chronic pain in people living with HIV raise the need to

identify targets for culturally resonant treatments. This qualitative study explored how people

living with chronic or recurrent pain and HIV seek informal social support for pain and

distress from their social networks, how these networks operate, and how social support is

influenced by socio-ecological factors. Adults with virally suppressed HIV who endorsed pain

over repeated weeks were purposively sampled (n=18). Semi-structured, individual in-depth

interviews explored individuals' experiences of living with HIV and chronic pain, distress and

its relationship to pain, and help-seeking behaviours for chronic pain and distress. Codebook

thematic analysis was used within an interpretivist epistemological approach. Support for

pain involved more tangible actions by fewer people than support for distress. Disruptions in

social support were more likely when the person with pain was irritable, socially withdrawn,

or unable to fulfil social roles. Interpersonal conflict could also cause or increase pain and

distress. Current pain assessments rarely identify the social interference of pain identified

here. This work identified potential targets for individual-level interventions to sustain

informal social support: individuals' abilities to cope with daily stressors, pain self-

management strategies to facilitate socialisation and engagement in valued life activities,

and strategies to prevent support fatigue in informal social networks.

Keywords: pain, social support, hiv, emotional distress, psychological distress, qualitative

research, thematic analysis

Introduction

Chronic pain is the world's most burdensome health condition in terms of years lived with disability (1, 2). People with chronic pain incur significant difficulties, such as comorbid health problems, limited ability to work, and high healthcare costs (3).

Chronic pain has far-reaching impacts, extending beyond the individual with pain to their families, communities, and societies. It adds financial and emotional strain to the family unit as a whole (4). It increases the risk of chronic pain in younger generations, especially if a parent has chronic pain and comorbid depression (5-7). Community members with disabling pain make smaller monetary and practical contributions, while drawing on more resources than their pain-free peers, thus compromising reciprocated support (4, 8). A high prevalence of health problems in a single community could thus deplete both tangible and emotional community resources, particularly in contexts where social security systems are inadequate or absent—leaving people with pain, and their families, relatively isolated.

Social factors are recognised as both consequences and determinants of chronic pain. For example, chronic pain can introduce conflict and relational dissatisfaction, along with guilt for the person with pain and their family and friends – and lower quality of relationships in the long term (9-11). The interference of pain in work and community activities progressively isolates those with pain, contributing to loss of roles and sense of identity (12). Isolation or loneliness can compromise immune function (13, 14), increase risk of mental illness (15), and constrain cognitive function that is needed to live well with chronic health conditions (16).

Conversely, social support can improve health in people with chronic pain, partly by buffering pain-related emotional distress and facilitating adaptive coping, and possibly also by improving regulation of physiological responses to painful events (17, 18). Pain-related emotional distress is known to exacerbate pain and mental concerns, which gives the distress-relieving impact of social support relevance to both chronic pain and mental health

(19). The health benefits of social support are highest when individuals perceive their social

network to be available and supportive (20, 21), which is shaped by their experiences

receiving support when they need it, and by the impact of the support (22). However, it is

crucial to recognise that the interpersonal relationships and social networks that underpin

social support are inherently dynamic, so the social support available to an individual may

change over time.

Despite the ripple effects of chronic pain and the centrality of social support, little is known

about how social support influences, or is influenced by, chronic pain in the context of HIV

(23-25). This is surprising in light of the high burden of chronic pain in people with HIV (26-

28). It also contrasts starkly with both the HIV treatment adherence literature—where social

support interventions are already being used to improve treatment adherence—and a

consensus-based agenda for research on HIV-associated pain that included social support

as a prioritised study outcome (29, 30).

In South Africa, where HIV and chronic pain each affects approximately 1 in 5 people, the

dual burden of HIV and chronic pain is well positioned to compromise population health (31).

However, little is known about how social networks function to address pain and emotional

distress in the context of chronic pain and HIV in South Africa. Recent qualitative research

noted that rural South Africans with HIV varied in how they disclosed and sought support for

pain (32). Given that social factors may be modifiable determinants of health that could be

targeted by scalable, culturally relevant interventions, this study aimed to explore (1) how

people living with chronic or recurrent pain and HIV seek informal social support to alleviate

pain and emotional distress, (2) how informal social support networks function, and (3) how

socio-ecological factors influence informal social support, in the context of chronic or

recurrent pain and HIV. Here, informal social support is differentiated from support derived

from formal professional, or public services (i.e. health clinic, social welfare services) (33),

and understood as being received through informal relationships within an interactive field of

people – a social network (34, 35). Social network *functions* relate to the *received* and

perceived social support provided by social network members, which can include emotional,

tangible, and informational support, as well as companionship (Table 1) (36). In this work,

emotional distress refers to negative affective states such as worry, anxiety, depression (19),

and includes feelings attributed to pain or to other difficult circumstances and experiences.

The philosophical perspective for this work was an interpretivist epistemological framework;

we sought to understand the knowledge and meaning that our participants had attached to

their experiences of seeking and receiving informal social support for distress and pain,

informed by their social, cultural, and broader context.

Types of social support	Definition
Emotional support	Provision of listening support, provision of emotional comfort and security, increasing the person's perceived sense of competence.
Tangible support (also known as instrumental support)	Provision of concrete assistance, such as giving time, providing transportation, or giving financial assistance.
Informational support	Giving of advice, suggestions and guidance; sharing of information.
Companionship	Involves the sharing of leisure and other activities motivated by the internal goal of enjoyment and wellbeing; understood as both a relationship quality and a form of social support.

105   *Table 1: Social support theory constructs relevant to this study (37-39).*

106   Methods

107   Study setting

This exploratory qualitative study is a part of a larger longitudinal study of people living with

virally suppressed HIV and chronic pain or no clinical pain (40) accessing standard HIV care

(including free antiretroviral treatment, ART) at a primary health facility in a peri-urban

community near Cape Town, South Africa. The parent and current studies were approved by

the Human Research Ethics Committees at the University of Cape Town and City of Cape

Town (approvals 764/2019 and 24699). The larger study focused on relationships between

distress, inflammatory responses, and pain, by using questionnaires, blood tests, and

psychophysical testing, over six months. A subset of participants in the larger study also

responded remotely to weekly questions on their pain and distress for up to six months; the

current study purposively sampled participants from this subset who repeatedly endorsed

pain to these weekly questions. A list of possible participants was compiled, ordered by the

consistency with which each participant endorsed pain. The main data collector for the study

(AG) then worked down the list to telephonically invite participants to an in-person, individual

interview. All participants gave additional written informed consent to be interviewed. In this

peri-urban community, approximately half the people live in informal housing, one in four is

unemployed (41), and the density of HIV-positive people is thought to be the highest in the

province, with HIV prevalence between 10 and 22%(42, 43).

Data collection

The data presented here were drawn from these individual, semi-structured in-depth

interviews conducted between 20 June and 26 August 2022, in a private room adjacent to

the health facility, to capture rich descriptions of inter-individual differences and intra-

individual variations in behaviour and experiences in a confidential manner. Topics explored

individuals' experiences of living with HIV and chronic pain and perceptions of how these

affected their lives; experiences of distress and its relationship to their chronic pain; coping

strategies and help-seeking; formal healthcare and formal and informal social support, and

behaviours used for chronic pain and distress. The interview guide was consultatively

developed, based on the topic of the larger study, existing literature, and identified gaps

relevant to the local setting according to key expert and local knowledge (JAJ, AG, AW).

Interviews were conducted in participants' preferred language (i.e. either English or

IsiXhosa), by a trained and experienced female bilingual isiXhosa-English interviewer (AG),

who was resident in a neighbouring community. The interviewer already had established,

confidential relationships with all participants and had previously asked personal questions

of them (i.e. about mental health, income, social support, stressful life events and traumatic

childhood events). Participants provided written informed consent. Interviews took 56-122

minutes, during which participants were offered refreshments. The interview period

coincided with the COVID-19 pandemic restrictions and major electricity cuts in South Africa,

resulting in minimal light or static torchlight for some interviews. Participants were

compensated for their time with R150 (~US$9). Interviews were audio-recorded, verbally

translated into English audio files by AG, and then transcribed using the online, HIPPAA-

compliant OTTER software (https://otter.ai), followed by manual quality checks.

Data analysis

Codebook thematic analysis was used (led by LC) (44, 45). Following familiarization,

transcripts were coded in qualitative data analysis software, NVivo 15 (Lumivero, London,

United Kingdom). The codebook was shaped by inductive coding of transcripts (n=4),

primary study objectives, and interview guide sub-sections (i.e. lived experiences of HIV and

chronic pain; functional impacts of HIV, chronic pain and distress; help-seeking strategies for

pain and distress) (46). Consistent with an interpretivist epistemological approach, results

and processes were iteratively discussed (LC, LK, VJM) to facilitate meaning-making and

reflexivity and an audit trail was kept, using journals, memos, and notes (47, 48). To remain

consistent with the relevant body of knowledge (34, 49, 50), themes and subthemes were

refined, recategorized, and named according to social support theory constructs (51). AG

reviewed data interpretations. The Consolidated Criteria for Reporting Qualitative Research

(52) was used to guide reporting, omitting identifying data. The lead and support analysts

are all South African women. LC (Occupational Therapist, MPH) and LK (PhD) have led

qualitative research on health experiences of South Africans and had no contact with

participants; VJM (Physiotherapist, PhD) is a clinician-scientist focused on pain who led the

parent and current study and exchanged transient greetings with two parent study

participants; AG (PGDip) is a Xhosa woman research assistant, was the primary in-person

data collector for most of the parent study, conducted the current interviews, and reviewed

our data interpretations for congruity with the interviews to counteract any potential

misinterpretation by non-Xhosa women LC, VJM, and LK.

Results

1. Participant characteristics

171 Interviews were completed with 18 of 24 eligible participants (details in supplement); 17 in

172 isiXhosa and 1 in English. Four participants lived alone and mean (SD) monthly household

173 income was R2649 (2332) (~US$160 (141)).

	Sex	Age range	Employment*	Housing*	Lives with*
1	Female	40s	Full-time	Family	Family, children
2	Female	40s	Unemployed	Family	Children
3	Female	30s	Unemployed, seeking	Shack	Family
4	Female	30s	Seeking	Shack	Partner
5	Female	40s	Unemployed, seeking	Shack	Children
6	Female	50s	Unemployed	Shack	Family, children
7	Female	20s	Unemployed	Shack	Partner, children
8	Female	30s	Full-time	Backyard	Alone, children[a]
9	Female	50s	Casual	Family house	Partner, children
10	Female	30s	Casual	Shack	Partner, children
11	Female	50s	Casual	Shack	Alone
12	Female	50s	Casual	Family house	Children
13	Female	30s	Seeking	Family house	Family, children

14	Female	40s	Unemployed	Shack	Family, children[b]
15	Male	50s	Unemployed	Backyard	Alone
16	Male	30s	Full-time	Shack	Partner, children
17	Male	40s	Unemployed, seeking	Shack	Alone
18	Male	40s	Unemployed, seeking	Backyard	Partner

175   *Table 2: Demographic data as reported by interview participants at enrolment to the parent study.*

176   *Multiple selections allowed. [a]Reported living alone at the time of interview. [b]Reported living with*

177   *children at the time of interview.*

2. Findings

First, we describe the *structure* of social networks that provided informal received social

support to participants to alleviate pain and distress. Second, we describe the *functions* of

social networks, specifically how members provided emotional, informational and tangible

social support and companionship for a) chronic pain and b) emotional distress, together

with the perceived impact of this support. Third, we explore the socio-ecological factors

influencing the structure and function of social networks in the context of chronic pain,

presented under themes of how 1) chronic pain causes emotional distress, resulting in social

isolation, 2) chronic pain affects one's ability to perform social roles, and 3) inter-personal

relationships can cause emotional distress and pain.

2.1.   Structure of social support networks

Participants described seeking and receiving informal social support for pain and emotional

distress from a range of family and community members (Fig 1). Support for chronic pain

came from a smaller social network, predominantly partners and immediate and extended

family members (i.e., children, parents, sisters, aunts, and cousins). Others included people

in their ART adherence club at their local clinic (into which people are assigned when they

first achieve viral suppression), and close confidants, such as friends and neighbours. When

in distress, participants reached out to these members as well as an additional, wider group

of social network members, including church leaders and fellow churchgoers, work

colleagues, and people they sat next to at their local clinic or on public transport.

Fig 1: Members of social network accessed for social support for both chronic pain and

distress (blue & orange) or distress only (orange)

2.2. Functions of social support networks

Participants received a range of informal social support from social network members,

discussed below under categories of emotional, informational and tangible support functions,

together with the perceived impact of this support. Fig 2 shows the overlapping and distinct

support functions for pain and emotional distress.

Fig 2: Informal social support received from social network members for pain and emotional

distress, indicating unique and overlapping features.

a. Received informal social support for chronic pain

Participants framed disclosure of their chronic pain to social network members as a way of

seeking emotional support. Disclosing their chronic pain was challenging in instances when

they did not want others to know about their pain, as it might worry or burden them, or if they

were not talkative in nature. Participants preferred to disclose their pain to people they lived

with or saw often, and close confidants, who were "always there for them" and willing to

listen. Some of these trusted social network members were people they had previously

disclosed their HIV status to, or others they knew were HIV positive themselves, as they felt

they could "tell them everything":

There is someone who I would tell when I've got the pains… so I'll go to this a

neighbour of mine, the one that I told you that is also HIV positive, I will go to her and

then I will talk with her… about the things that I'm feeling. (female in 50s, living with

partner and children)

Conversely, members who were perceived as responding negatively to hearing about their

initial HIV diagnosis were not sought out for support for pain. Participants felt cared for and

understood when social network members noticed, asked about, or initiated discussions

about their pain. This solicitous emotional support typically came from those who lived with

or near them and saw them in pain, or from close confidants who phoned them. Participants

who lived on their own described feeling supported when social network members provided

solicitous support in conjunction with companionship, by visiting them, or by inviting them to

visit:

I would tell my aunt [about pain], I'll just call her and tell her that I've got the pains…

She [aunt] will come and visit me and see how am I doing … [and] I become fine, …

so, if she comes, at least that makes a difference. (female in 30s, living alone)

Provision of encouragement also eased pain-related emotional distress. This was

experienced when members motivated participants to seek formal medical care for their

pain, reassured them, or encouraged them not worry about their pain:

You know, my daughter is always there for me, and she'll tell me that, "No, Mama,

you are going to be alright you know". (female in 30s, living with partner and children)

Although demonstrative support in response to disclosure of their pain was appreciated, the

process of "sharing" their pain was what helped some participants feel "happy" and hopeful.

Talking to someone about their pain was often enough to relieve pain that the participant

attributed to emotional distress (a common attribution for headaches), or to give them

confidence about managing their pain:

I do tell my cousin, my family, they know when I've got the pains, I will tell them.

Everyone. Even with my mother, I'll phone her and tell her that I've got these pains…

I'm just happy that I have talked... (female in 30s, living with family)

In addition to motivating participants to seek formal medical care, members would often

provide informational support. Pain-focused informational support was often sought from

members who also had chronic pain or friends who were health professionals, as they were

felt to be able to provide "good" advice. Members would advise on what could be causing the

pain, based on their own experiences or personal knowledge:

I will tell my neighbour that I've got this pain, … they're the one that advised me to go

buy pills for arthritis… (female in 40s, living with children)

Members also advised on which health provider to see and recommended different

medicinal or physical treatments for pain. Participants explained that they appreciated the

advice, even though it didn't always relieve the pain:

… then they will advise me that, okay, I must use this must use [or] that. … I tried it,

but it didn't help me. But I'm happy with the support that I'm getting from them.

(female in 40s, living with children)

Tangible support received was aimed at supporting treatment of pain or assistance with

tasks that participants found challenging due to pain. Participants described how family

members, both nuclear and extended, sent money or paid for their direct or indirect medical

costs, drove them to and arranged care for them at health facilities, fetched their medication

from the health facility, or shared their own medication with them. Family members living with

them commonly provided massages using rubbing ointments or wrapping materials:

It's only my kids that knows that I have got pains… The one…will rub me, massage

my legs, wrap my neck… they go buy pills… or massage me. (female in 40s, living

with family)

Although some participants with less disabling chronic pain described not needing help with

their tasks and preferred to "force" themselves to do tasks, assistance with tasks that were

challenging due to pain was described by many as being supportive. This was predominantly

provided by members living in their household or those who lived nearby and were able to

provide physical support:

He [partner] will get me water and do the washing because now I can't go to the tap,

the tap is far and I can't carry heavy stuff... I have to get the kids to go and get water

for me… (female in 50s, living alone)

Support included assistance with getting out of bed or moving around the house, bringing

their medication to them, or performing their domestic tasks, such as helping participants to

cook and prepare meals, wash clothes or dishes, or clean their house.

b. Received informal social support for emotional distress

Participants described a range of socio-ecological factors, particularly unemployment and a

lack of financial resources, as contributing to stress, anger, depression, and anxiety. Some

who were employed described challenges with their employment, such as conflicts with

employers (i.e. who did not appreciate their work, treat them well or follow fair labour

practices), living far from their workplace, or being unsatisfied with their type of work and

salary. Although many received social grants, the grants did not cover the family's basic

needs, such as food, electricity or cooking gas and transportation, which caused significant

daily emotional distress. Insufficient finances also meant that they could not realise personal

goals and lived in environmental conditions that also worsened their pain:

When it's raining, the water will get into the house... and there is no electricity

there…I feel bad…at night the cold really affects me… I'll have these chest pains…

But I'm staying there because I don't have any other options. (female in 50s, living

alone)

Participants described how talking to someone about their emotional distress helped them

find relief and resulted in them feeling "lighter", as they had shared their problems with

someone else. This could be true even if a member did nothing more than listen. Others

described how they felt emotionally supported when members encouraged them not to

worry, especially if the reasons for their distress were out of their control, and when a

member gave them hope by telling them that "things will be alright", or expressed a belief

that they had the capacity to cope with their problem. They also felt supported when others

checked in on them and provided space to talk about their stressors:

The person that I talk to when I'm stressing is my partner… she's the one who will

notice that I'm not my usual self. She'll ask me [about] what stresses me, then she

will tell me that I need not stress a lot because there's nothing that I can change

about this situation. (male in 40s, living with partner)

Emotional support for distress was sought from a variety of social network members

depending on their availability and whether it was appropriate to seek their support. Available

members were those who were seen frequently at work or on public transport, or who made

themselves available to talk to when needed. A few participants specifically chose to talk to

"strangers" regarding commonly experienced stressors or due to their being external to the

distressing situation:

I'll talk about my stuff with strangers. I know those people don't know me. But… if I

can just speak to a stranger, someone who doesn't care about me - I feel relieved

about that. (female in 30s, living with family)

In contrast, others expressed that it was not appropriate to share their emotional distress

with "just anyone" as these people might reveal their problems to others or "make fun" of

them. These participants preferred to choose members they felt comfortable with and had an

open and trusting relationship with, who listened, would not judge them. Being empathetic to

their distress was a key consideration:

I do talk to my sister that I'm staying with, but not everything with her. Because she is

not that kind of a person who listens or who is interested. You could see, even when

you're telling her something, she doesn't respond, you know? (female in 30s, living

with family)

Participants also described valuing advice on how to deal with their stressors. Informational

support was provided by members sharing their own experiences with a similar problem and

offering advice born of these experiences, often including strategies to solve the problem or

how to cope with the resulting emotional distress. This was described as relieving

participants' distress to a degree:

I will speak to the ladies that I go with… there [at the clinic]… I will normally speak to

them, like try to find out… how do they cope with certain things in life, then I'll get

advice from them… we'll share our experiences and problems… it is good to share

your problems with other people. (female in 40s, living with family)

This support was often sought from members who knew about the participant's stressors or

if they had perceived experience with the same stressors. For example, family members who

knew their family dynamics were sought out for advice in times of sibling conflict, or

members with similar-age children were engaged with for informational support on how to

manage their own children's distressing behaviours:

I do get the support from my partner because she's the one that I always talk to when

I've got these things that are stressing me…her child is [a young adult], so she's

been in the situation that I'm in. She knows how to deal with the teenagers and then

she'll advise me. (male in 40s, living with partner)

Tangible support was provided in the forms of financial support, spiritual support, or helping

to solve the problem that was causing the emotional distress. Participants described how,

after talking about their emotional distress, members offered to do something to solve the

problem, such as brainstorming business ideas with them, talking to their "troublesome"

child, or handing in the participant's CV to employers. Spiritual support was also valued.

Participants described discussing their stressors with church members who would pray with

them, and ceremonies facilitated by spiritual leaders also helped to relieve their distress:

It has given me strength, ever since I started to go into that church, in terms of the

way that I think, it has changed me...So when I prayed to those candles there, so it's

helping me spiritually. So, they're [spiritual leaders] using candles and coarse salt,

you know, salt water. It's normal water that we're using, you just have to bring water

[to church] and then they have to pray over that water, and then they will maybe do

what we call 'isiwasho' [salt and water], and then you drink that 'isiwasho'… bath

yourself [with it]… you can spray [it] around your house for bad spirits. (female in 50s,

living with family and children)

Economic support included giving or loaning money, sometimes for specific purposes, such

as to support a particular household member or pay for basic needs.

2.3.   Factors influencing the structure and function of social networks in the context

of chronic pain

Participants described how their social support structure and functions were influenced by

their chronic pain. They described a) how chronic pain caused emotional distress, resulting

in isolation; b) how pain affected their ability to perform social roles, and c) how interpersonal

relationships could be both a prominent source of informal social support and a cause of

emotional distress and pain.

a. Chronic pain results in emotional distress, leading to social isolation

Chronic pain impacted on participants' mood and emotions and made them want to isolate

themselves from others. Notably, although some mild emotional distress was experienced

due to HIV (i.e. due to stigma experienced from health care workers, when a partner died

from HIV, or due to the need to take medication daily), many of our participants described

how their HIV did not cause emotional distress beyond the initial period of diagnosis and

initiating treatment, yet their chronic pain often resulted in ongoing distress:

The other things that are stressing me about my health, are the pains that I have.

They are the most thing that are stressing me. HIV doesn't stress me, no. The fact

that I'm on ARVs [antiretroviral treatment] doesn't really stress me. (female in 50s,

living with partner and children)

Participants described experiencing negative emotional responses to their pain, which

continued even in periods when they experienced no or mild pain, as they felt anxious that

the pain could recur at any time. Overarching low moods were described, such as feeling

"irritable," "bad" and "not happy", as well as frustrated, angry and stressed. These were due

to the pain's chronic nature, a lack of diagnostic certainty, few pain-relieving strategies, and

the current and possible future impacts of their chronic pain, specifically on their mobility,

independence and ability to work:

Sometimes I'll get afraid of the pains, when I don't know what is happening and I'll be

wondering why it doesn't go away… now I don't know which pain tablets should I

take, because they don't necessarily work…I'm afraid that I might not be able to walk

… I don't know what is going to happen in future… I feel bad. (female in 50s, living

with family)

Experiencing chronic pain also brought with it additional fears. For example, those with

diagnostic uncertainty expressed fear of death, as they were unsure if their chronic pain was

related to something terminal. Chronic pain was also related to aging, leading some

participants to describe a fear of getting old:

Now you've got that fear that oh, now I'm getting old and growing, I am going to have

pains like my mother because my mother is always like, crying about having pains.

(male in 30s, living with partner and children)

Emotional distress resulting from pain influenced participants' interpersonal relationships

with those in their social network. Feeling irritable due to their pain was described as

affecting their relationships with those they lived with, who grew tired of their complaints

about pain:

I'll be complaining that I've got the pain here, I've got the pain there, you know… I'm

getting irritable all the time. Now [my husband]'s been saying that… I'm always

complaining about the pains. So… it was interfering with the relationship…with my

kids and my husband. (female in 20s, living with partner and children)

Participants commonly described wanting to be alone when feeling emotional distress due to

their pain or when experiencing pain. They described intentionally isolating themselves at

home by withdrawing to their rooms or avoiding seeing friends and other family by staying at

home more often, particularly when experiencing pain. They also described ignoring phone

calls, not speaking to others at social gatherings, shortening their social visits with others, or

cancelling them altogether:

Sometimes you find that it [the pain] will make me not be able to go, … I'll feel like I

don't want to go anywhere… just want to stay at home. …I'll end up not going or

cancelling the things that I was supposed to do. (female in 30s, living with family and

children)

Participants described varied motivations for isolating themselves, typically feeling

disinclined to talk to others when in pain, wanting their "own space", feeling tired or "lazy" to

go out, or wanting to be somewhere comfortable when in pain. Participants also avoided

social gatherings because they did not want others to worry about—or in other cases, not

know about—their pain:

When that [pain] happens, I just want to be alone. I don't want to be seen by other

people because… I don't want them to know about my problem. (male in 40s, living

alone)

Participants described how isolating themselves affected their social relationships. Although

some friends and family continued to visit them at their home when they stayed at home

more often, they often started "meeting" people less often and neighbours and friends were

described as complaining that the participant had "been scarce". This meant that fewer

people knew about their pain and could offer support.

b. Chronic pain affects one's ability to perform social roles

Participants described how chronic pain affected their ability to perform role-specific tasks,

influencing the functions of their social network. This was contrasted to the daily impact of

HIV. Limited challenges of HIV were noted, such as negotiating intimate relationships with an

HIV positive status or engaging with family or friends while anticipating stigma. The impact of

HIV on social roles, such as parenting, working, and community roles, was felt to be minimal

and some participants even noted improvements in support:

My [social] responsibilities have never changed as much [compared to pain] … So,

when I told them [my family] about my status, the only thing that we discussed was

the way forward… So ever since then, my support system has been alright...

Otherwise, I don't see any difference with my life now… even with my [intimate]

relationship, I'm aware of everything that I have to do and I'm learning new things.

(male in 40s, living with partner)

In contrast, participants described how chronic pain reduced their ability to provide certain

tangible support for their children, perceived as crucial to the role of being a parent. Such

activities included attending sports matches, cooking meals, and washing children or their

clothes. Chronic pain also reduced the types of physical tasks participants could perform at

work, or prevented them from working at all, limiting their ability to provide financial support

for dependents. Participants described not being able to perform work-related tasks properly

due to chronic pain and needing to adapt their task performance or, in some cases, take

leave or resign from their job. Job losses occurred in rare instances, when employers felt

participants were no longer fulfilling required duties or excess leave was being taken to

manage pain or seek healthcare. Participants described how, when possible, they forced

themselves to work despite their pain to financially support younger children:

I do my responsibilities as a mother even though I have got the pains. If I need to go

to work, I do go to work even if I've got the pains because I know that I won't be able

to buy groceries or to buy food for my children if I do not go to work, so even if I've

got pains, I'll go to work. I need to go to work. (female in 40s, living with children)

Some tried to ignore their pain and continue with these necessary tasks, whereas others had

to limit their provision of tangible support to their children, which was felt to affect the quality

of their relationships with their children or result in their children feeling distressed:

I used to play with my kids .. but now after I have the pains, I'm not able to play with

them all the time….. they didn't understand what was happening, … they were saying

"Haibo!" [an expression of surprise and disapproval] I don't want to play at all

because of the pains. (female in 20s, living with partner and children)

Some participants described how their children also became distressed when seeing them in

pain or being cared for by others, which in turn caused the participants emotional distress.

Others described how older children responded by providing tangible support, taking over

the cooking or washing or getting a job to provide financial support. However, some parents

preferred not to rely on their children in this way because they perceived task assistance as

falling outside a child's role, made them feel like a burden or too dependent:

You know I don't get happy or satisfied when someone is doing something for me. I

just wish I can do it on my own, but I can't. And even with my children, I don't get

satisfied when they're doing chores in my house (female in 50s, living with family and

children)

Participants also described how chronic pain influenced their ability to perform domestic and

sexual tasks specific to that of being an intimate partner, and how partners had to take over

their domestic tasks. Although some described adapting domestic tasks to make them easier

or forcing themselves to do them, others described how they often felt too fatigued or in pain,

especially in instances where the participant was employed, because work tasks were

prioritised rather than domestic chores. At times, this adjustment could cause conflict with

partners and complaints:

You find that the house is not properly cleaned because I've got the pains. Like on

Thursday you know, my partner had to make supper. When he was back, he said

that, "Haibo, you didn't do the dishes". And I told him that I can't do anything. So, he

was complaining that the house is still dirty, the dishes have not been washed, and

he didn't carry any food with to work on that day (female in 30s, living with partner)

*C*hronic pain also made some participants disinterested in sex, limited how they engaged in

sex, or unexpectedly interrupted sex. This further complicated some intimate relationships

that were already complicated by participants' HIV status (i.e. non-disclosure of status,

negotiation of condom use, reduced sexual drive perceived to be due to ART use). Although

some partners attributed these limitations to pain, others interpreted reluctance to have sex

with suspicion, and some participants expressed concerns about how their partner

interpreted their reduced sexual activity:

There are times where I feel like, okay, and I am supposed to do something [sexual].

Then you find that I've got this problem now, I've got the pains, I can't do anything.

The thing is maybe if you've got someone, and she might not understand you and

think that you are you are lazy now, you don't want to have sex. (male in 40s, living

alone)

Roles external to the household were also described as being affected. For example,

participants explained how they could no longer engage in community activities such as

sports, helping build friends' houses, carrying heavy things, or fixing appliances. Some

ensured that where they could no longer provide tangible support, they continued to provide

companionship and emotional support by visiting their friends and attending community

events:

Sometimes there is something that is happening in my neighbourhood, maybe

they've got a traditional work or maybe a funeral or something, I'll go there, but I'm

not able to assist them... I'll just sit there... I just support them emotionally. (female in

50s, staying alone)

c. Interpersonal relationships can cause emotional distress and pain

Although, as described above, social network members play a crucial role in supporting

participants in managing their chronic pain and emotional distress, interpersonal

relationships with children, partners and siblings were also a primary source of emotional

distress in some cases. Some participants attributed their chronic pain to these interpersonal

challenges, specifically chronic headaches, muscle tension in their neck, shoulder and/or

back, or severe pain in their sides. Some participants who had lost children (from a motor

vehicle accident, fatal assaults and HIV), described their traumatic loss as a sustained

source of significant distress:

The other thing that is stressing me a lot is the death of my daughter…She had a

[detail of unanticipated death], and that is something that will come to my mind most

of the time and it made me unhappy…So, that is also causing me to have the pains,

that back pain, and you'll will find that when I think about her, I will have these signs

of pain. (female in 50s, living with partner and children)

Others expressed that their pain was exacerbated or caused by conflict and worrying about

their children's errant behaviours such as smoking, staying out late at night, and not

listening:

She [my daughter] is stressing me a lot, I will have the pain, like, if I'm talking too

much… she will go out and come home very late… I have to speak with her and that

was stressing me you know and when I've got that stress it will go to my back then I

will have pains you know. (female in 40s, living with children)

Relational challenges with intimate partners caused significant emotional distress and

reduced the amount of tangible and emotional social support that could be provided in the

relationship. Some participants also described having partners who abused substances,

were emotionally unresponsive to their needs, or did not financially support their family or

children. One participant also said that she started experiencing her chronic pain when she

found out that her partner had married another woman:

I was stressing a lot [last year when I started feeling this pain]… I saw a picture of

him with another girl and he told me that they got married… now he doesn't even

support the child… He doesn't support me. (female in 30s, living with family and

children)

A perceived lack of support and open communication from siblings was another source of

emotional distress. A participant described moving in with her sister as a form of financial

support, but conflict with her sister due to this living arrangement was perceived as the

underlying cause for her chronic pain:

The most pain that I normally have is the headache, sometimes I will think that I've

got high blood [pressure], so I went for high blood [pressure] test but I didn't have it.

So, I thought… maybe it's because I'm stressing a lot, maybe it's me because I'm so

used to having my own space, and now I have to have people that I'm sharing my

space with… so, I think stress that is causing this. (female in 40s, living with children

and extended family)

Participants also expressed that their existing chronic pain, and/or other types of pain, were

exacerbated or brought on during periods of emotional distress about interpersonal

relationships. The result was more intense pain, other locations of pain, or greater difficulties

in managing their pain, for example as emotional distress interfered with sleep, as described

by a participant experiencing distress over her child's behaviour below:

It is only this thing about my child, I don't have anything else… The neck and my

headache will be intense, will be worse when I'm stressing [about her]…the back and

the neck becomes too much... So, if maybe she has stressed me and I will tell

myself, I'm not going to speak for days, let me just relax... I don't respond to anything,

then the pains will be better. (female in 40s, living with children)

554   Discussion

This exploratory qualitative study focused on the social network structure and informal social

support received by people living with chronic pain and HIV, to address pain and emotional

distress. Participants sought support from various people spanning nuclear and extended

family members, neighbours, colleagues, friends, and acquaintances within other social

networks. Participants sought support from a wider range of people for distress than for pain.

Although the nature of support crossed emotional, tangible, informational, and

companionship support, many participants perceived benefit from merely speaking to

someone who was attentive and empathetic. The analysis revealed that chronic pain can

compromise a person's ability to fulfil their social roles, which can disrupt interpersonal

relationships with key providers of informal social support for pain, such as children and

intimate partners. This inability to fulfil social roles, together with social withdrawal (a

common strategy to cope with pain or pain-related emotional distress), may render informal

social support vulnerable to disruption. Previous work has suggested that people with HIV

may socially self-isolate in fear that any 'display' of suddenly worsening pain could prompt

others to speculate about their overall health and indirectly reveal their HIV status (24, 53).

In contrast, our participants attributed their social self-isolation to low mood and functional

limitations. Interestingly, in times of inter-personal conflict, the close relationships that usually

provided meaningful social support could switch to causing emotional distress and

exacerbating, or even causing, pain.

Participants described slightly different support-seeking strategies for pain and emotional

distress. To choose a member for distress-focused support, participants considered whether

the member would be attentive, their proximity to the problem, the nature and source of the

distress, and whether they were well positioned to help. The opportunity to disclose distress

was prioritised, and participants seemed less concerned about privacy for distress than for

pain. Given the diverse and everyday sources of distress reported, it is possible that

normalisation of stressors and worry makes it easier for people to disclose emotional

distress in a more public environment. For pain-focused support, participants described a

more careful, challenging process of deciding who to disclose to, although availability and

access was also a key consideration. For pain, participants also considered the social

network member's general behaviour, discretion, the impact of the knowledge of pain on

their roles and responsibilities, and the current health of their relationship with the person.

This selectivity echoes that reported in another sample of South Africans with chronic pain

(32) and may reflect previous negative interpersonal responses to pain disclosure (54) as

well as overlapping HIV and chronic pain stigma, which is thought to contribute to social

isolation (55). Notably, in our data, members previously chosen for HIV disclosure were also

commonly chosen for pain disclosure. People diagnosed with HIV before chronic pain may

benefit from this pre-'vetted' circle of trusted others with whom they feel safe to discuss their

chronic pain.

This study revealed a potential vulnerability arising from informal social support-seeking

behaviours and responses for pain. Participants sought support for pain from a smaller

social network than for distress, with the support sought for pain often being more practical

and hands-on in nature for household members. High intensity tangible support may be

unsustainable over time and result in support fatigue in these smaller support networks (10,

56). As in other literature, participants expressed concerns about shifts in roles, especially

between parents and children, and between life partners, describing how this could lead to

interpersonal distress and conflict (10, 25). Distrust and additional conflict due to role shifts

in relationships—especially in intimate relationships—may reduce social support available

for people with HIV and pain if these intimate relationships are disrupted. Analysis of the

social networks of people with chronic pain in other contexts suggests that larger social

networks are associated with reduced levels of interpersonal conflict, and that inter-personal

conflict has a stronger exacerbating effect on distress when the conflict occurs in denser

support networks (57). In our participants, if the few people chosen for pain support are

closely related to one another, this increase in network density with decreasing network size

may render the person with pain particularly vulnerable when conflict does arise.

The current findings stand juxtaposed with previous qualitative studies of people with chronic

pain or chronic pain and HIV, most of which were conducted in high-income countries (e.g.

25, 58, 59). In our sample, distress arose largely from concerns that were not specific to HIV

or chronic pain, such as difficult living circumstances (poverty, risky housing, income

vulnerability, employment problems), interpersonal conflict (sometimes linked to material

dependence on others), continued grief, and relational worries. These background stressors

may be common to others in the same geographical community, but their diverse and

unrelenting nature may represent a higher 'load' than in previously described samples,

rendering the current sample more vulnerable to pain and negative health outcomes (60).

That shifts in functional role performance were a trigger for interpersonal conflict in the

current study underscores the limited capacity to absorb further resource deficits. In this

context, the recursive, adverse influence of interpersonal conflict on both distress and pain

has the potential to contribute to a negative spiral of worsening pain, diminishing function,

and continued degradation of social support for the person living with chronic pain and HIV

(61).

Interestingly, the impact of chronic pain that was revealed in this qualitative study would be

missed by conventional measures of the functional interference of chronic pain. Participants

described continuing with their daily roles despite considerable pain, adapting tasks and

prioritising employment over home tasks, but rarely completely withdrawing from these. This

survival-driven stoicism has been described in resource-strapped contexts and may contrast

with pain interference in contexts with higher formal support for people with pain disability

(32, 62, 63). For this sample, the true impact of chronic pain is observed in its influence on

activities that contribute to relationships and overall wellbeing, rather than just on activities

that support survival. Participants described withdrawing from valued activities such as

caring for, supporting and playing with their children and expressing physical affection. As in

previous South African research (4), participants also described the mental and emotional

impact of chronic pain that resulted in worry, low mood, stress and irritability that potentially

strained relationships. Such consequences of chronic pain are rarely assessed, but may

best capture its true impact on the overall health and wellbeing for the person and their

networks, including inter-generational transmission of chronic pain and disability (19, 64). In

line with current evidence, the daily impact of living with HIV was almost eclipsed by the

impact of living with chronic pain; all participants who drew a comparison declared HIV to be

far more easily managed than chronic pain (25, 32, 65).

These qualitative findings present opportunities to either improve or provide social support

for people living with HIV and chronic pain and prevent worsening health. Individuals who

have existing informal social support networks may benefit from skills to maintain the health

of these interpersonal relationships. Training people in pain self-management or equipping

people with HIV to better navigate emotionally distressing situations (i.e. through training in

emotional self-regulation, workplace negotiation, parenting skills, and conflict resolution) may

improve coping skills and reduce the demand on their social network for pain and distress.

Similarly, encouraging people with HIV to reciprocate care to members of their informal

social support networks could also sustain capacity within these support networks (66).

Existing interventions could be adapted to train these skills. As examples, the ImpACT+

intervention addresses emotional coping and problem-solving among women with HIV and

trauma (67), while the Friendship Bench model supports skill-building using a culturally

resonant approach with attractive scalability (68). In the setting of the current research,

people with HIV attend monthly 'adherence club' meetings to sustain ART adherence. These

existing meetings provide an alternative opportunity for brief but regular injections of skills

training.

In contrast, for individuals who tend to socially self-isolate, more intentional facilitation of

social interactions may be beneficial. One approach could be to use the ART adherence

clubs to connect these individuals to a group-based intervention. Group interventions for

pain management have been tested for people with HIV, including in a remotely delivered

format that may feel more socially 'safe' and physically accessible, and some have led to

sustained engagements and support between participants (69, 70). Using a trained peer to

facilitate such groups is an established practice that is supported by our participants' reports

that they prefer to seek support from people they recognise as having skills or experience.

The current participants emphasised the importance of confidentiality, attentive listening, and

a sense of common struggles; an intervention that emphasises these features may have

better uptake. With respect to matching an intervention to an individual, other studies

suggest that the individual's usual preferences about social contact (e.g. intro/extraversion)

should inform the type of intervention that is suggested (71).

Limitations

The timing of this study, the sampling strategy, and the context of the current study inform

interpretation of our results. The interviews were conducted in 2021. The COVID-19

pandemic in 2020 had rendered 2.24 million people unemployed; by mid-2021, 1.44 million

people had not yet been re-employed (72). Job losses had disproportionately affected Black

South Africans and women, particularly those in face-to-face service jobs like some of our

participants (73, 74), and job loss was associated with more depressive symptoms (75). It is

likely that this situation influenced participants' perspectives, and the precarious nature of

employment and health-related fears may have exacerbated distress in our sample. The

biographical data reported in Table 2 should be interpreted with caution, given that they were

collected several months before the interview and some circumstances may have changed –

in particular, one participant was no longer living with her children at the time of interview.

We drew participants from a larger study of pain and distress that included in-person

assessments conducted by the interviewer for the current study. Participants may have been

more aware of symptoms after frequently being questioned about distress and pain. On the

other hand, the interviewer's existing relationship with the participants likely contributed to

the richness of the data obtained during the interviews.

Conclusion

Informal social support networks for pain are smaller and bear a higher load than those for

distress, due to more stringent selection of network members. Informal social support is

particularly vulnerable to support fatigue or interpersonal conflict when the person with pain

is unable to fulfil their functional social roles, or when they use social isolation to cope with

pain and pain-related distress. Conflict in interpersonal relationships can cause and increase

pain, emphasising the importance of sustaining good social relationships and support.

Individual-level interventions to sustain informal social support should target individuals'

abilities to cope with daily stressors, their knowledge of pain self-management strategies to

facilitate socialisation and engagement in valued life activities, and skills to reduce support

fatigue in their informal support social network.

Funding details

This work was funded by NIH award K43TW011442 to VJM. SR was supported by NIMH

and FIC (1K43TW012840); RRE was supported by NIH award K24 NS126570.

Conflicts of interest

VJM is an unpaid associate director of the not-for-profit organisation, Train Pain Academy.

RP receives speakers' fees for talks on pain and rehabilitation, is a director of the not-for-

profit organisation, Train Pain Academy, and serves as a councillor for the International

Association for the Study of Pain. All authors declare no other conflicts of interest.

References

References

1.   Blyth FM, Briggs AM, Schneider CH, Hoy DG, March LM. The Global Burden of

Musculoskeletal Pain-Where to From Here? American journal of public health.

2019;109(1):35-40.

2.   James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global,

regional, and national incidence, prevalence, and years lived with disability for 354 diseases

and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the

Global Burden of Disease Study 2017. The Lancet. 2018;392(10159):1789-858.

3.   Webster F, Connoy L, Sud A, Rice K, Katz J, Pinto AD, et al. Chronic Struggle: An

Institutional Ethnography of Chronic Pain and Marginalization. J Pain. 2023;24(3):437-48.

4.   de Villiers M, Maree JE, van Belkum C. The influence of chronic pain on the daily

lives of underprivileged South Africans. Pain Manag Nurs. 2015;16(2):96-104.

5.   Ojeda B, Salazar A, Dueñas M, Torres LM, Micó JA, Failde I. The impact of chronic

pain: the perspective of patients, relatives, and caregivers. Fam Syst Health.

2014;32(4):399-407.

6.   Higgins KS, Birnie KA, Chambers CT, Wilson AC, Caes L, Clark AJ, et al. Offspring of

parents with chronic pain: a systematic review and meta-analysis of pain, health,

psychological, and family outcomes. Pain. 2015;156(11):2256-66.

7.   Brown D, Rosenthal N, Könning A, Wager J. Intergenerational transmission of

chronic pain-related disability: the explanatory effects of depressive symptoms. Pain.

2021;162(2):653-62.

8.   Ampiah JA, Moffatt F, Diver C, Ampiah PK. A Qualitative Investigation of the

Psychosocial Impact of Chronic Low Back Pain in Ghana. BMJ open. 2023;13(7):e073538.

9.   Fullen B, Morlion B, Linton SJ, Roomes D, van Griensven J, Abraham L, et al.

Management of chronic low back pain and the impact on patients' personal and professional

lives: Results from an international patient survey. Pain Pract. 2022;22(4):463-77.

10.   Borthwick C, Penlington C, Robinson L. Partners' Experiences of Chronic Pain: A

Qualitative Evidence Synthesis. J Clin Psychol Med Settings. 2024;31(3):585-606.

11.   Murray CB, Groenewald CB, de la Vega R, Palermo TM. Long-term impact of

adolescent chronic pain on young adult educational, vocational, and social outcomes. Pain.

2020;161(2):439-45.

12.   Toye F, Belton J, Hannink E, Seers K, Barker K. A Healing Journey with Chronic Pain:

A Meta-Ethnography Synthesizing 195 Qualitative Studies. Pain Med. 2021;22(6):1333-44.

13.   Balter LJT, Raymond JE, Aldred S, Drayson MT, Veldhuijzen van Zanten J, Higgs S,

et al. Loneliness in healthy young adults predicts inflammatory responsiveness to a mild

immune challenge in vivo. Brain, behavior, and immunity. 2019;82:298-301.

14.   Zhao X, Li F, Cheng C, Bi M, Li J, Cong J, et al. Social isolation promotes tumor

immune evasion via β2-adrenergic receptor. Brain, behavior, and immunity. 2024;123:607-

18.

15.   Jaremka LM, Andridge RR, Fagundes CP, Alfano CM, Povoski SP, Lipari AM, et al.

Pain, depression, and fatigue: loneliness as a longitudinal risk factor. Health Psychol.

2014;33(9):948-57.

16.   Kong X, Han F, Li J, Wang W, Sun Y, Wu Y. Serial multiple mediation of loneliness

and depressive symptoms in the relationship between pain and cognitive function among

older people. Aging Ment Health. 2023;27(11):2102-10.

17.   Che X, Cash R, Ng SK, Fitzgerald P, Fitzgibbon BM. A Systematic Review of the

Processes Underlying the Main and the Buffering Effect of Social Support on the Experience

of Pain. The Clinical journal of pain. 2018;34(11):1061-76.

18.   López-Martínez AE, Esteve-Zarazaga R, Ramírez-Maestre C. Perceived social

support and coping responses are independent variables explaining pain adjustment among

chronic pain patients. J Pain. 2008;9(4):373-9.

19.   Edwards RR, Dworkin RH, Sullivan MD, Turk DC, Wasan AD. The Role of

Psychosocial Processes in the Development and Maintenance of Chronic Pain. The Journal

of Pain. 2016;17(9):T70-T92.

20.   Barrera M. Distinctions between social support concepts, measures, and models.

American Journal of Community Psychology. 1986;14(4):413-45.

21.   Wethington E, Kessler RC. Perceived support, received support, and adjustment to

stressful life events. Journal of health and social behavior. 1986;27(1):78-89.

22.   Melrose KL, Brown GDA, Wood AM. When is received social support related to

perceived support and well-being? When it is needed. Personality and Individual

Differences. 2015;77:97-105.

23.   Scott W, Arkuter C, Kioskli K, Kemp H, McCracken LM, Rice ASC, et al. Psychosocial

factors associated with persistent pain in people with HIV: a systematic review with meta-

analysis. Pain. 2018;159(12):2461-76.

24.   Merlin JS, Walcott M, Kerns R, Bair MJ, Burgio KL, Turan JM. Pain self-management

in HIV-infected individuals with chronic pain: a qualitative study. Pain Med. 2015;16(4):706-

14.

25.   Scott W, Garcia Calderon Mendoza Del Solar M, Kemp H, McCracken LM, A CdCW,

Rice ASC. A qualitative study of the experience and impact of neuropathic pain in people

living with HIV. Pain. 2020;161(5):970-8.

26.   Parker R, Stein DJ, Jelsma J. Pain in people living with HIV/AIDS: a systematic

review. J Int AIDS Soc. 2014;17(1):18719.

27. Povshedna T, Swann SA, Levy SLA, Campbell AR, Choinière M, Durand M, et al.

Global Prevalence of Chronic Pain in Women with HIV: A Systematic Review and Meta-

analysis. Open Forum Infect Dis. 2023;10(8):ofad350.

28. Madden VJ, Parker R, Goodin BR. Chronic pain in people with HIV: a common

comorbidity and threat to quality of life. Pain Manag. 2020;10(4):253-60.

29. Shushtari ZJ, Salimi Y, Sajjadi H, Paykani T. Effect of Social Support Interventions on

Adherence to Antiretroviral Therapy Among People Living with HIV: A Systematic Review

and Meta-Analysis. AIDS and behavior. 2023;27(5):1619-35.

30. Merlin JS, Hamm M, de Abril Cameron F, Baker V, Brown DA, Cherry CL, et al. The

Global Task Force for Chronic Pain in People with HIV (PWH): Developing a research

agenda in an emerging field. AIDS care. 2023;35(8):1215-23.

31. Kamerman PR, Bradshaw D, Laubscher R, Pillay-van Wyk V, Gray GE, Mitchell D, et

al. Almost 1 in 5 South African adults have chronic pain: a prevalence study conducted in a

large nationally representative sample. Pain. 2020;161(7):1629-35.

32. Mulaudzi M, Ratshinanga A, Mohale J, Ramoshai T, Evangeli M, Pincus T, et al.

Experiences and impact of chronic pain in South Africans living in a rural area: a qualitative

study. medRxiv. 2025:2025.07.22.25331712.

33. Shiba K, Kondo N, Kondo K. Informal and Formal Social Support and Caregiver

Burden: The AGES Caregiver Survey. J Epidemiol. 2016;26(12):622-8.

34. Berkman LF, Glass T, Brissette I, Seeman TE. From social integration to health:

Durkheim in the new millennium. Soc Sci Med. 2000;51(6):843-57.

35. Schwarzer R, Knoll N, Rieckmann N. Social support. In: Kaptein A, Weinman J,

editors. Introduction to health psychology. 158. Oxford, England: Blackwell; 2003. p. 847.

36. Rodriguez MS, Cohen S. Social support. Encyclopedia of mental health.

1998;3(2):535-44.

37.   Mabachi NM, Brown M, Sandbulte M, Wexler C, Goggin K, Maloba M, et al. Using a

Social Support Framework to Understand How HIV Positive Kenyan Men Engage in

PMTCT/EID Care: Qualitative Insights From Male Partners. AIDS and behavior.

2020;24(1):18-28.

38.   Drageset J. Social Support. In: Haugan G, Eriksson M, editors. Health Promotion in

Health Care – Vital Theories and Research. Cham: Springer; 2021. p. 137-44.

39.   Langford CP, Bowsher J, Maloney JP, Lillis PP. Social support: a conceptual analysis.

J Adv Nurs. 1997;25(1):95-100.

40.   Madden VJ, Msolo N, Mqadi L, Lesosky M, Bedwell GJ, Hutchinson MR, et al. Study

protocol: an observational study of distress, immune function and persistent pain in HIV.

BMJ open. 2022;12(6).

41.   Statistics South Africa. South African National Census of 2011. Pretoria, South Africa:

Stats SA; 2012.

42.   Woldesenbet SA, Lombard C, Manda S, Kufa T, Ayalew K, M. C, et al. The 2019

National Antenatal HIV Sentinel Survey. South Africa: National Department of Health; 2021.

43.   Naomi model. South Africa District HIV Estimates, December 2024 [Available from:

https://www.hivdata.org.za.

44.   Braun V, Clarke V. Conceptual and Design Thinking for Thematic Analysis. Qual

Psychol. 2022;9(1):3-26.

45.   Braun V, Clarke V. Toward good practice in thematic analysis: Avoiding common

problems and be(com)ing a knowing researcher. Int J Transgend Health. 2023;24(1):1-6.

46.   Brooks J, King N. Doing Template Analysis: Evaluating an End of Life Care Service.

Sage Research Methods Cases Part 1: SAGE Publications Ltd.; 2014.

47.   Nowell LS, Norris JM, White DE, Moules NJ. Thematic Analysis: Striving to Meet the

Trustworthiness Criteria. International Journal of Qualitative Methods.

2017;16(1):1609406917733847.

48.   Green J, Thorogood N. Analysing Qualitative Data. Qualitative Methods for Health

Research: Sage.

49.   Kelly ME, Duff H, Kelly S, McHugh Power JE, Brennan S, Lawlor BA, et al. The

impact of social activities, social networks, social support and social relationships on the

cognitive functioning of healthy older adults: a systematic review. Syst Rev. 2017;6(1):259.

50.   Langford CP, Bowsher J Fau - Maloney JP, Maloney Jp Fau - Lillis PP, Lillis PP.

Social support: a conceptual analysis. (0309-2402 (Print)).

51.   Braun V, Clarke V. Thematic Analysis: A Practical Guide. London, UK: Sage

Publications; 2021.

52.   Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research

(COREQ): a 32-item checklist for interviews and focus groups. Int J Qual Health Care.

2007;19(6):349-57.

53.   Baker V, Nkhoma K, Trevelion R, Roach A, Winston A, Sabin C, et al. "I have failed to

separate my HIV from this pain": the challenge of managing chronic pain among people with

HIV. AIDS care. 2023;35(8):1164-72.

54.   Mitchell MM, Maragh-Bass AC, Nguyen TQ, Isenberg S, Knowlton AR. The role of

chronic pain and current substance use in predicting negative social support among

disadvantaged persons living with HIV/AIDS. AIDS care. 2016;28(10):1280-6.

55.   Scott W, Buchman DZ, Vasiliou VS. The multi-dimensional stigma of chronic pain:

A narrative review. Curr Opin Psychol. 2025;62:101980.

56.   Davies RL, Rice K, Rock AJ. The Potential Burden of Informal Support: Exploring

Wellbeing and Self-Care of Social Network Members of Intimate Partner Violence Survivors.

Journal of Family Violence. 2025.

57.   Van Alboom M, Baert F, Bernardes SF, Verhofstadt L, Bracke P, Jia M, et al.

Examining the Role of Structural and Functional Social Network Characteristics in the

Context of Chronic Pain: An Ego-centered Network Design. J Pain. 2024;25(9):104525.

58.   Merlin JS, Westfall AO, Raper JL, Zinski A, Norton WE, Willig JH, et al. Pain, mood,

and substance abuse in HIV: implications for clinic visit utilization, antiretroviral therapy

adherence, and virologic failure. J Acquir Immune Defic Syndr. 2012;61(2):164-70.

59.   Sabin CA, Okhai H, Dhairyawan R, Haag K, Burns F, Gilson R, et al. Prevalence of

pain in women living with HIV aged 45-60: associated factors and impact on patient-reported

outcomes. AIDS care. 2023;35(8):1181-90.

60.   Rios R, Zautra AJ. Socioeconomic disparities in pain: the role of economic hardship

and daily financial worry. Health Psychol. 2011;30(1):58-66.

61.   Signs TL, Woods SB. Linking family and intimate partner relationships to chronic

pain: An application of the biobehavioral family model. Fam Syst Health. 2020;38(1):38-50.

62.   Mphahlele NR, Mitchell D, Kamerman PR. Pain in ambulatory HIV-positive South

Africans. Eur J Pain. 2012;16(3):447-58.

63.   Wadley AL, Mitchell D, Kamerman PR. Resilience does not explain the dissociation

between chronic pain and physical activity in South Africans living with HIV. PeerJ.

2016;4:e2464.

64.   Stone AL, Wilson AC. Transmission of risk from parents with chronic pain to offspring:

an integrative conceptual model. Pain. 2016;157(12):2628-39.

65.   Merlin JS, Walcott M, Ritchie C, Herbey I, Kertesz SG, Chamot E, et al. 'Two Pains

Together': Patient Perspectives on Psychological Aspects of Chronic Pain while Living with

HIV. PLOS ONE. 2014;9(11):e111765.

66.   Rothausen CS, Clausen AM, Voltelen B, Dieperink KB. Protective Buffering: Nurses

Facilitating Communication Between Adults With Cancer and Their Adult Family Caregivers

Who Overprotect One Another-An Integrative Review. J Fam Nurs. 2023;29(4):417-36.

67.   Sikkema KJ, Hansen NB, Tarakeshwar N, Kochman A, Tate DC, Lee RS. The clinical

significance of change in trauma-related symptoms following a pilot group intervention for

coping with HIV-AIDS and childhood sexual trauma. AIDS and behavior. 2004;8(3):277-91.

68.   Chibanda D, Weiss HA, Verhey R, Simms V, Munjoma R, Rusakaniko S, et al. Effect

of a Primary Care-Based Psychological Intervention on Symptoms of Common Mental

Disorders in Zimbabwe: A Randomized Clinical Trial. Jama. 2016;316(24):2618-26.

69.   Parker R, Jelsma J, Stein DJ. Managing Pain in Women Living With HIV/AIDS: A

Randomized Controlled Trial Testing the Effect of a Six-Week Peer-Led Exercise and

Education Intervention. J Nerv Ment Dis. 2016;204(9):665-72.

70.   Ernstzen D, Keet J, Louw K-A, Park-Ross J, Pask L, Reardon C, et al. "So, you must

understand that that group changed everything": perspectives on a telehealth group

intervention for individuals with chronic pain. BMC Musculoskeletal Disorders.

2022;23(1):538.

71.   Flowers KM, Colebaugh CA, Hruschak V, Azizoddin DR, Meints SM, Jamison RN, et

al. Introversion, Extraversion, and Worsening of Chronic Pain Impact during Social Isolation:

A Mediation Analysis. J Clin Psychol Med Settings. 2023;30(3):531-42.

72.   Altman M. Trajectories for South African employment after COVID-19. South African

Journal of Science. 2022;118(5-6):1-9.

73.   Nwosu C, Kollamparambil U, Oyenubi A. Socioeconomic inequalities in ability to work

from home during the coronavirus pandemic: The case of South Africa. National Income

Dynamics Study; 2021.

74.   Casale D, and Shepherd D. The gendered effects of the Covid-19 crisis in South

Africa: Evidence from NIDS-CRAM waves 1–5. Development Southern Africa.

2022;39(5):644-63.

75.   Posel D, Oyenubi A, Kollamparambil U. Job loss and mental health during the

COVID-19 lockdown: Evidence from South Africa. PLoS One. 2021;16(3):e0249352.