1 # Evidence for a protein leverage effect on food intake in a
2 # Norwegian population
3

Rikke Eriksen[1], Kamilla Rognmo[1], Laila A Hopstock[2], James E McCutcheon[1]

1 Dept. of Psychology, UiT The Arctic University of Norway, Tromsø, Norway

2 Dept. of Health and Care Sciences, UiT The Arctic University of Norway, Tromsø,
Norway

Keywords: dietary protein; energy intake; food frequency questionnaire; ultra-
processed food; obesity; protein leverage hypothesis

Running title: Protein leverage effect on food intake but not BMI

Contact information: James E McCutcheon, Dept. of Psychology, UiT The Arctic
University of Norway, Huginbakken 32, 9019 Tromsø, Norway

Word count: 4764

Funding: Funding was provided by Tromsø Research Foundation to JEM (19-SG-
JMcC).

Disclosure: The authors declare no conflicts of interest.

Author contributions: **Rikke Eriksen:** Conceptualization, Formal Analysis, Writing -
Original Draft; **Kamilla Rognmo:** Conceptualization, Writing – Review and Editing;
Laila A Hopstock: Conceptualization, Writing – Review and Editing, Investigation;
James E McCutcheon: Conceptualization, Writing – Review and Editing,
Visualization.

Abstract

In this pre-registered study, we aimed to explore the protein leverage hypothesis in a

general population, by studying the relationship between habitual dietary protein

intake, total energy intake, and body mass index (BMI), and whether proportion of

ultra-processed food (UPF) was associated with any of these variables. We used

regression-based analyses to investigate these phenomena in cross-sectional data

from a Norwegian population-based study, the seventh survey of Tromsø Study

2015-2016, (n = 11,152; 40-99 years). Total energy intake was negatively associated

with proportion of dietary protein (L = -0.36, p < .001) and positively associated with

dietary fat (L = 0.33, p < .001). Although we planned to test the relationship between

BMI and dietary protein, there was no positive association between total energy

intake and BMI meaning that these data were not suitable for testing an effect of

protein leverage on BMI. Proportion of UPFs was positively associated with total

energy intake (b = 554, p < .001), negatively associated with proportion of dietary

protein (b = -2.0, p < .001), and positively associated with BMI (b = 0.011, p = .026).

In summary, our study of middle-to-older aged Norwegians provides strong support

for a protein leverage effect on energy intake.

1. Introduction

The obesity epidemic is a global problem that is estimated to contribute to a huge number of deaths each year as well as posing a significant economic burden (World Health Organization, 2021). Although its causes are multifactorial, reduced intake of dietary protein has been implicated via the protein leverage hypothesis. The protein leverage hypothesis is based on a nutritional geometric framework that was developed to explain how humans and other animals meet their nutritional needs (Raubenheimer et al., 2022). The hypothesis emphasizes that protein is the most essential macronutrient as many of the amino acids cannot be synthesized de novo and must be acquired from one's diet. This implies that humans need to consume a target level of protein on a regular basis (i.e., on the order of days) resulting in a strong appetite for protein (Raubenheimer et al., 2015). The protein leverage mechanism is proposed to regulate energy intake towards this protein target through appetite (Simpson & Raubenheimer, 2005) and, as such, the proportion of protein in a given diet is suggested to influence food intake more strongly than total calories consumed. Accordingly, if a food environment is "protein diluted", (i.e., contains a lower proportion of dietary protein than a balanced diet), excess energy will be consumed in an effort to reach the protein target (Simpson & Raubenheimer, 2005). Recent analyses find that the US food supply has indeed become protein diluted over recent decades, and though this dilution is small (~1% of available calories), modeling suggests a significant effect on development of obesity in the adult population (Hall, 2019).

Several studies, including randomized controlled trials (Gosby et al., 2011; Martens et

al., 2013) and observational studies (Honfo et al., 2023; Saner et al., 2023; Saner et

al., 2020; Zhang et al., 2025; Zhang et al., 2023), find support for the energy

regulating effects of the protein leverage mechanism in specific cohorts including

preschool children prone to obesity, children and adolescents living with obesity, and

older adults. To our knowledge, few observational studies have examined protein

leverage effects in adults and only one (Zhang et al., 2025) has been conducted in

European populations.

The dilution of protein in the human food supply (Raubenheimer & Simpson, 2023)

may be linked to the increase in availability and popularity of ultra-processed foods

(UPFs) . The human obesity epidemic has developed concomitantly with the

emergence of these foods, and now UPFs dominate the market in several high-

income countries. For example, in the UK, US, and Canada more than half of the

energy consumed comes from these foods (Monteiro et al., 2019). In the context of

protein leverage, high UPF diets that tend to be low in dietary protein should lead to

an increase in total calories consumed. Indeed, this has been found in national

representative samples of the Australian (Grech et al., 2022a) and US (Martínez

Steele et al., 2018) populations, which show that high UPF diets are related to a

lower proportion of dietary protein intake and a heightened energy intake. In addition,

a carefully planned 14-day RCT (Hall et al., 2019) found overcompensation of energy

intake on a UPF diet (~16% protein) when compared to an non-UPF diet (~19%

protein). This increased energy intake was accompanied by an increase in

bodyweight where participants gained 0.9 kg when on the UPF diet and lost 0.9 kg

when on the non-UPF diet, implying that weight gain observed in high UPF diets

could result from unbalanced protein levels. In fact, replotting of data from Hall et al.

(2019) in Raubenheimer and Simpson (2023) suggests that the protein leverage

mechanism accounts for up to 100% of the observed change in energy intake

between the UPF and non-UPF diets. Similar results were found in an RCT where

participants gained more weight and increased energy intake on a UPF diet than on

a non-UPF diet, while no significant differences were found in protein intake between

the diets (Hamano et al., 2024).

The biological processes that underpin protein leverage are still under study, but one

potential mechanism is variation in levels of the hormone fibroblast growth factor 21

(FGF21). Studies in humans and rodents have found that FGF21 increases in

response to protein restriction (Laeger et al., 2014; Volcko & McCutcheon, 2022). In

mice, FGF21 has also been found to increase protein intake (Solon-Biet et al., 2023)

and protein preference without increasing total food intake (Hill et al., 2019). In

humans, plasma FGF21 levels have been found to be negatively associated with

percent protein in the diet (Gosby et al., 2016; Nicolaisen et al., 2025). FGF21

increases manyfold in response to a low-protein overfeeding diet (Hollstein et al.,

2019), indicating that its release is not just dependent on the caloric content of the

food. Additionally, FGF21 levels after exposure to a low-protein overfeeding diet are

positively related to 24 hour energy expenditure (Vinales et al., 2019). Similarly,

energy requirements increased concomitantly with FGF21 levels in lean men on a

protein restricted diet (Nicolaisen et al., 2025). As such, the relationship between

FGF21, protein intake, and bodyweight is complex.

In our study, we aimed to replicate and extend findings on the protein leverage

mechanism by testing the predicted relationships between dietary protein, energy

intake, and body mass index (BMI) in middle-aged and older adults through two

preregistered hypotheses (osf.io/4g62f). These were that (1) the proportion of dietary

protein will be negatively associated with total energy intake, and (2) the proportion of

dietary protein will be negatively associated with BMI. In addition, we assessed

properties of diets high in UPFs. Finally, we investigated the relationship between

habitual dietary protein intake and FGF21.

2. Methods

2.1 Data source and sample

The current study includes data from the Tromsø Study, a large-scale population-

based study conducted in the municipality of Tromsø, Norway. To date, seven

surveys have been carried out and the eight survey is ongoing (Tromsø1–Tromsø8,

1974–2026). Data collection includes questionnaires and interviews, biological

sampling and clinical measurements. This analysis includes data from Tromsø7

(2015–2016), where all inhabitants of Tromsø municipality aged 40 years and older

(N = 32,591; 50.7 % women) were invited, and 21,083 (65%; 52.5 % women)

attended. Details of the study design, sample, and data collection methods for

Tromsø7 have been described elsewhere (Hopstock et al., 2022).

The current study was approved by the Regional Committee for Medical Research

Ethics Northern Norway (ref. 789456, 14/11/2024), and the Norwegian Agency for

Shared Services in Education and Research (ref. 935030, 9/10/2024). All participants

gave written consent at study attendance. Data from participants that later withdrew

their consent were not used in the analysis. The current study analyzed data from

11,152 participants. The exclusion criteria are visualized in the flow chart depicted in

Figure 1. Data exclusion plan and two hypotheses were preregistered on Open

Science Framework (osf.io/4g62f).

2.2 Food Frequency Questionnaire

All participants were invited to complete a food frequency questionnaire (FFQ)

designed to collect data on habitual dietary patterns by assessing frequency and

quantity of intake of 261 of single foods, beverages, dishes and dietary supplements

(Carlsen et al., 2010). Details of the FFQ used in Tromsø7 have been described

elsewhere (Lundblad et al., 2019). The FFQ was developed and validated (Carlsen et

al., 2010) at the University of Oslo, and is designed to detect dietary habits in

Norwegian adults. Food, energy and nutrient intakes were calculated using the food

database KBS AE14 and KBS software system (KBS, version 7.3.), based on the

Norwegian food composition tables from 2014 to 2015.

2.3 Demographics and Health Related Information

Height and weight were measured by trained personnel using a Jenix DS-102 scale

(DongSahn Jenix, Seoul, Korea). BMI was calculated as kg/m^2 and categorized in the

standard groups normal (<25 kg/m^2), overweight (25–<30 kg/m^2), and obese (≥30

kg/m^2). The group normal includes a low proportion of underweight participants (≤1

%) as reported in Løvsletten et al. (2020).

Demographic and health information was collected by questionnaires. Level of leisure

time physical activity was measured with the Saltin-Grimby Physical Activity Level

Scale (Grimby et al., 2015) including sedentary, light, moderate, and vigorous

physical activity levels. Education level included primary/partly secondary education

(≤10 years), upper secondary (10 + minimum 3 years), tertiary short

(college/university less than 4 years), and tertiary long (college/university 4 years or

more). Current or previous disease included the four main non-communicable

diseases (NCD) from the NCD framework (World Health Organization, 2013):

cardiovascular diseases (myocardial infarction and/or stroke), cancer, chronic

respiratory diseases (asthma and/or chronic bronchitis/emphysema/chronic

obstructive lung disease) and/or diabetes, in addition to psychiatric problems the

participant reported to have sought help for. Presence of chronic pain was defined by

the question "Do you have persistent or constantly recurring pain that has lasted for 3

months or more?" with answer options "yes" or "no". Smoking status was defined by

the question "Do you/did you smoke daily" with answer options "yes, now", "yes,

previously", or "never".

2.4 Fibroblast Growth Factor 21

Non-fasting blood samples were obtained with a light tourniquet released after

venipuncture (Hopstock et al., 2022). In a randomly selected subsample ($n = 1,144$),

FGF21 was analyzed as part of the OLINK Target Inflammation panel which uses

Proximity Extension Assay (PEA) technology to measure relative protein levels.

2.5 NOVA Classification of Food Processing Level

The food items from the FFQ were categorized based on the NOVA classification system (Monteiro, 2009) and the procedure was adapted from Huybrechts et al. (2022) (Huybrechts et al., 2022). The NOVA system classifies foods and dishes into four mutually exclusive groups (Monteiro, 2009). Based on the categorization of the FFQ, a binary variable was made for (0) non-ultra-processed foods (NOVA groups 1-3) and (1) ultra-processed foods (NOVA group 4). For further details, see Supplementary Materials section 1.

2.6 Statistical Analyses

Data pre-processing and statistical analyses were performed using RStudio version 4.2.2. Data visualizations were made in Python version 3.9.

The protein leverage mechanism was tested through a power function (equation 1.1) and indexed by the L-value,

$$TEI = P.\,p^L \quad (1.1)$$

where TEI = total energy intake per day, P = a constant obtained from the analysis, p = the proportion of protein in the diet, and L = the strength of leverage. An L-value of -1 indicates complete protein leverage, while an L-value of 0 indicates that protein has no control over total energy intake (Senior et al., 2022). A positive L-value indicates that protein increases energy intake, i.e., the opposite of the predicted pattern. An L-value between -1 and <0 indicates partial protein leverage. The variables TEI and p are taken from the data set, while P and L are derived from the

model when Equation 1.1 is log-transformed to a linear regression equation (equation

1.2),

$log(TEI) = log(P) + L.log(p)$ (1.2)

To assess the individual contributions of each macronutrient to total energy intake,

energy density, and food dry weight, mixture models were run with the R package,

Mixexp (Lawson & Willden, 2016). The package demands that the predictors (i.e., the

three macronutrients) sum to one, which was accomplished by removing the

contribution of alcohol and fiber to total energy intake. The mixture models were run

based on the framework described by Lawson and Willden (2016), starting with a

linear model and increasing the polynomial complexity (for details see Supplementary

Materials section 2). The output from the mixture model was used to make mixture

triangle surface plots, which plotted predicted outcome variables (total energy intake,

energy density, and food dry weight) per participant based on macronutrient

composition.

The proportion of UPFs in the diet was calculated as intake of UPFs in kJ/day divided

by total intake of food in kJ/day (i.e., UPF proportion (%E) = UPF kJ /day/ total intake

of food kJ/day). Analyses were conducted on UPF intake classified into quintiles, and

for the effect of UPF intake on protein leverage, UPF intake was categorized as low

or high via a median split. For relevant analyses, significant effects were followed up

with contrast analyses using the *emmeans* package in RStudio.

Multiple linear regression models with relevant covariates were run with BMI as the

outcome. This includes the following binary variables, which were entered into the

model in the given order: age (over vs. under 60 years), sex (female vs. male),

physical activity level (sedentary vs. three active levels), health conditions (presence

vs. absence of conditions), education level (lowest vs. three highest levels), and

smoking status (not current vs. current smokers). BMI was positively skewed and

was therefore log-transformed prior to analyses.

For mixture modeling and BMI analyses, the Akaike Information Criterion (AIC)

(Akaike, 1992) was used to choose between models, and if several models were

within two AIC points of each other, the simplest model was favored.

2.7 Code and data availability

The data that support the findings of this study are available upon application to the

Tromsø Study. Restrictions apply to the availability of these data, which were used

under license for this study. All analysis code is available at

https://github.com/jaimemcc/proteinleverage.

3. Results

3.1 Descriptive Statistics

Descriptive statistics for the full sample ($n = 11,152$, 53.3% women) are presented in

Table 1. Information about BMI was missing for 27 participants, and all analyses on

BMI are therefore based on $n = 11,125$. Participants with valid FGF21 data were $n =$

880 (51% women).

3.2 The Protein Leverage Mechanism

Analyses of the strength of leverage protein has over energy intake found support for a partial protein leverage effect with the analysis showing that L = -0.36 (p < .001). The negative L-value indicates that protein was negatively associated with energy intake (Figure 2A), in line with our hypothesis. The power function was fitted for carbohydrate and fat and the analysis revealed L-values of ~0 (p = .966) and 0.33 (p < .001), respectively. This indicates that the proportion of carbohydrates in the diet had no control over energy intake (Figure 2B), while the proportion of fat was positively associated with energy intake (Figure 2C).

3.3 Mixture Models for Protein, Carbohydrate, and Fat

The AIC (Table S1) favored the second model (Table S2) which is visualized in the mixture triangle in Figure 3A which shows that the change in total energy intake mainly followed the protein axis, but fat also had an effect. As the proportion of protein increases from e.g., 10% to 30%, the predicted total energy intake decreases (as seen by the color change from yellow to purple). As the proportion of fat increases from e.g. 10% to 70%, the color changes from purple to orange-yellow. Additional analyses were run for energy density (Table S3) and dry weight of food (Table S4) to further probe macronutrient associations. Energy density (Figure 3B) is greatest in high fat/low protein diets, and lowest in high protein diets, regardless of the level of fat. Dry weight of food (Figure 3C) appears unaffected by fat but to be negatively associated with protein, and is thus highest in low protein diets, supporting a leveraging of protein on appetite.

3.4 Properties of Diets High in Ultra-Processed Foods

Several linear regression models were run to assess how changes in proportion of

UPFs in the diet were related to (1) proportion of dietary protein, (2) dietary protein in

absolute terms, and (3) total energy intake. The output from the regression models

and estimated marginal mean comparisons are reported in the Supplementary

Materials (Tables S5-S10). Proportion of UPFs in the diet was negatively related to

proportion of dietary protein (Table S5) and positively related to total energy intake

(Table S9), while the amount of protein in absolute terms stayed relatively stable

between UPF quintiles (Table S7). Figure 4A summarizes these results.

Moreover, inspection of Figure 4A suggests that the leveraging effect of protein was

greater for those consuming low levels of UPFs vs. those consuming high levels. As

an exploratory analysis to test this, we performed a median split to divide participants

into high and low UPF consumers and calculated the strength of protein leverage in

each subpopulation (Figure 5A). Our analysis revealed that leverage was stronger in

the low UPF group (L = -0.40, p < .001) than in the high UPF group (L = -0.29, p <

.001). We confirmed that this difference was significant by shuffling labels which

revealed the observed difference in L values to be greater than expected by chance

(Figure 5B; p = 0.009).

3.5 Associations between Body Mass Index and Dietary Intake

The prediction that dietary protein is negatively associated with BMI builds on the

assumption that energy intake will be negatively associated with BMI, we therefore

tested this. We ran linear regression models of increasing complexity with covariates

described under Demographics and Health Related Information and used AIC to

choose among models. The AIC favored the model presented in the Supplementary

Materials (Table S11), which revealed a weak and negative relationship between total

energy intake and BMI (Table S12). Potential reasons for this lack of association are

given in the Discussion. The lack of relationship means that these data are not

suitable for testing associations between mechanisms mediating increased energy

intake (i.e., protein leverage) and BMI. As the link between protein leverage and BMI

was one of our pre-registered hypotheses we have nevertheless placed this analysis

in Supplementary Materials (Table S14), however, results should be treated with

caution.

In the analysis of the relationship between proportion of UPFs and BMI, the AIC

(Table S15) favored the model that included UPF quintiles as well as the covariates

age, sex, and physical activity level (see Table S16). The estimated marginal means

were calculated for the quintiles of UPF intake while keeping the covariates in the

model constant as the reference category (i.e., age = under 60 years, sex = female,

physical activity level = sedentary) and while adjusting the p-value using the Dunnett

method. This yielded a trend towards higher UPF quintiles being associated with a

higher BMI as compared to the first UPF quintile (Table S17). The estimated marginal

means per quintile of UPF are visualized in Figure 4B.

3.6 FGF21 and Dietary Protein

To assess the relationship between FGF21 and dietary protein, several multiple linear

regression models were run, and the AIC-favored (Table S18) model is presented in

Table 3 (R^2 = .04, $F(7, 757)$ = 5.9, $p < .001$). The model reveals a negative,

significant, relationship between dietary protein and FGF21, in line with our

expectations. Several other covariates were also significantly associated with FGF21

including age, physical activity level, disease and smoking status.

312 4. Discussion

In this study of 11,152 women and men from a general population in Norway, we

examined predictions of the protein leverage hypothesis. We found strong support for

the protein leverage mechanism (i.e., the energy intake regulating effects of dietary

protein; $L = -.36$) in line with previous studies in the field (Honfo et al., 2023; Saner et

al., 2023; Saner et al., 2020; Zhang et al., 2023). We did not find the predicted

negative relationship between dietary protein and BMI, but given that there was no

positive association between energy intake and BMI in our study, this was not

surprising, as is discussed in more detail below. Our analyses of the relationship

between dietary protein, total energy intake, and habitual UPF intake revealed

relationships that are predicted to arise from the protein leverage hypothesis, namely,

that a higher UPF intake was associated with greater energy intake but lower

proportion of dietary protein. But again, our analysis failed to establish a concordant

effect of UPF intake on BMI. Finally, in an exploratory analysis, we examined the

relationship between FGF21 and dietary protein and this revealed the expected

negative relationship.

328 Positive leveraging effect of fat on energy intake

Our power functions not only revealed a partial protein leverage mechanism, but also

a positive L-value for fat, again in line with previous studies (Honfo et al., 2023; Saner

et al., 2023). The association between fat and energy intake may be related to

several factors. A key consideration is that is that as protein leverages food intake,

increases in fat will have a disproportionate effect on energy intake due to fat being

calorically dense. This effect will be exacerbated in places where habitual diets are

relatively high in fat. In Norway, as well as many other Western countries, fat makes

up >30% of energy intake. In addition, other properties of fat may directly or indirectly

augment this effect. For example, fat has a lower satiating effect than the other two

macronutrients (Blundell & Macdiarmid, 1997), which could lead to larger meals and

thus an excess energy intake when consuming diets high in fat. The high palatability

of fat-rich foods (Mizushige et al., 2007) may also play a role in driving energy intake.

As such, people may eat more when consuming diets high in fat because they taste

better, or being exposed to high fat diets may lead to changes in dietary patterns

such as larger meals or more snacking. The protein leverage hypothesis is based on

a nutritional geometric framework and emphasizes that it is the balance of the

macronutrients that are of importance for energy intake and potential weight gain and

our results for fat and protein attest to this.

Energy intake, protein and their association with BMI

Our second, key pre-registered hypothesis was that protein intake would be

negatively correlated with BMI. This hypothesis rests on an assumption that protein

leverages energy intake (as shown by hypothesis 1) and that this increased energy

intake in turn leads to increased BMI. In our data, however, we failed to find support

for this assumption, namely the relationship between total energy intake and BMI.

The implication is that finding a link between protein and BMI would be highly unlikely

– and indeed we did not find this. What then are the reasons for energy intake not

positively predicting BMI in this sample? Importantly, there are several limitations

when using self-reported food intake (e.g., Dhurandhar et al., 2015; Prentice et al.,

2025), even when using a validated FFQ. Perhaps most significant, are the inherent

difficulties in relating short-term dietary intake (as the FFQ measures) to long-term

consequences such as BMI. This may be especially the case when examining a

middle-aged to older sample, as in the present study. Weight gain resulting in

increased BMI accumulates over a lifetime while the FFQ is designed to measure

habitual food intake in a short, recent period. Moreover, people living with a higher

BMI have been found to underreport energy intake to a larger degree, on a group

level, than people in the normal BMI range (Howes et al., 2024; Poslusna et al.,

2009). As a large proportion of the participants in our study are either living with

overweight or obesity, this could have impacted our data. Indeed, calculations based

on the Goldberg method (Black, 2000 as cited by; Lundblad et al., 2019) indicate that

about 20% of the people in the Tromsø7 sample underreported energy intake when

filling out the FFQ (Lundblad et al., 2019).

Relationship between FGF21, energy expenditure and BMI

Ours and several other studies fail to find that low protein diets are related to a

higher BMI, a key relationship predicted by the protein leverage mechanism (Honfo et

al., 2023; Saner et al., 2023; Saner et al., 2020; Zhang et al., 2023). The role of the

liver-derived hormone FGF21 in energy balance illustrates some important

considerations in interpreting interactions between energy intake, macronutrient

(im)balance, and resulting weight change. Here, we found that dietary protein was

negatively associated with FGF21, an expected finding in line with results in prior

studies (Gosby et al., 2016; Nicolaisen et al., 2025). Moreover, FGF21 is also

suggested to upregulate energy expenditure (Nicolaisen et al., 2025). This has been

shown most robustly in rodent models (Hill et al., 2019; Laeger et al., 2014) although

one study has established this association in humans (Vinales et al., 2019). As such,

the effect of low protein diets on energy balance may seem contradictory at first

(protein leverage increases intake while inducing FGF21 which increases energy

expenditure). However, these two effects can be tightly linked due to the established

effect of surplus energy intake (induced by leverage) on triggering compensatory

increases in expenditure. Thus, a key question - not just for protein leverage but for

all mechanisms of obesity - is to determine the extent to which compensatory

expenditure is able to offset increased intake. Moreover, what our the conditions or

parameters that lead to this compensation being inadequate such that weight gain

and obesity result from excess energy intake.

Potential role of ultra-processed foods

A higher proportion of energy from UPFs was accompanied by a higher energy intake

and a lower proportional protein intake. As detailed in the Results, a change in

percent dietary protein was observed between the highest (16.4%) and lowest

(18.5%) quintiles. This reduction by a mere 2% may seem modest, but this may have

large effects on energy intake via leveraging. A previous RCT found that dilution by

3.5% of protein for a short period (14 days) may have contributed to significant

increases in energy intake and subsequent weight gain in participants (Hall et al.,

2019). Replotting of these RCT data by Raubenheimer and Simpson (2023) suggests

that in this controlled setting, protein leverage could explain almost all increased

energy intake associated with a UPF diet.

Subtle differences in dietary protein at a population-level have been associated with

obesity. For example, by analyzing food balance sheets and available epidemiology,

it was estimated that dilution of dietary protein coupled to perfect protein leverage

could explain as much as two-thirds of observed weight gain in US adult population

since the 1970s (Hall, 2019). In the current study, the *L*-value for protein was -0.36,

which despite being less than perfect leverage, still means that protein leverage is a

significant potential mechanism.

Whether the low protein nature of many UPFs is what leads these diets to cause

excess energy intake is still under debate. Importantly, several other UPF properties

could interact with protein content to increase energy intake. For example, UPF

formulations, including the combination of macronutrients, encourage consummatory

patterns that lead to high energy intake (Fazzino et al., 2019). In particular, high fat

content may increase energy intake, independent of carbohydrate, given the positive

L-value for fat found in our earlier analysis. Another possibility is that the umami and

salt flavors associated with many UPFs may drive energy intake up (Grech et al.,

2022b) due to the protein decoy effect. Foods rich in protein often have these savory

flavors, and people in search of protein may use these cues to guide them towards a

potentially nutritious food source. Dietary fiber (and energy density of food, which is

inversely proportional), is low in UPFs, influences energy intake, and can reduce the

protein leverage effect by limiting intake. Finally, UPFs are designed to increase food

intake due to their industrialized, cosmetic, ingredients (Monteiro et al., 2019). All

these factors may play a role in increasing energy intake in diets high in UPF content,

in addition to the diluted protein content.

Strengths and limitations

A major strength of the current study is the large sample size from a general

population, high attendance, and data collection using standard methods and

validated measurements. Another strength is that several of the findings replicate

other studies, thus further strengthening the implications of the protein leverage

hypothesis. Major limitations, as discussed above, are issues relating to self-reported

food intake including potential underreporting and the difficulties in associating short-

term estimation of food intake with long-term health outcomes, such as obesity.

Conclusion

We found strong support for the protein leverage mechanism through several

analysis methods, i.e., fitted power functions and mixture modeling. An interesting

and related finding was the positive effect of proportion of fat on energy intake. Our

study revealed that a diet consisting of a high proportion of UPFs was related to

heightened energy intake and a lowered proportion of dietary protein, while protein

intake in absolute terms remained relatively stable between UPF diets. Finally, we did

not find a positive association between energy intake and BMI meaning that we were

not able to properly test the predicted consequence of protein leverage increasing

BMI.

5. Acknowledgements

The authors would like to acknowledge the staff involved in data collection and

administration of the Tromsø Study and the participants that volunteered to take part.

We would also like to acknowledge the manuscript's reviewers for insightful and

helpful comments during the review process. Funding was provided by Tromsø

Research Foundation to JEM (19-SG-JMcC). The data that support the findings of

this study are available upon application to the Tromsø Study. Restrictions apply to

the availability of these data, which were used under license for this study.

6. References

Akaike, H. (1992). Information theory and an extension of the maximum likelihood principle. In E. Parzen, K. Tanabe, & G. Kitagawa (Eds.), *Selected Papers of Hirotugu Akaike.* (pp. 610–624). United States: Springer. https://doi.org/10.1007/978-1-4612-0919-5_38

Black, A. E. (2000). Critical evaluation of energy intake using the Goldberg cut-off for energy intake: Basal metabolic rate. A practical guide to its calculation, use and limitations. *International Journal of Obesity*, *24*(9), 1119–1130. https://doi.org/10.1038/sj.ijo.0801376

Blundell, J. E., & Macdiarmid, J. I. (1997). Fat as a risk factor for overconsumption: Satiation, satiety, and patterns of eating. *Journal of the American Dietetic Association*, *97*(7), S63–S69. https://doi.org/10.1016/S0002-8223(97)00733-5

Carlsen, M. H., Lillegaard, I. T. L., Karlsen, A., Blomhoff, R., Drevon, C. A., & Andersen, L. F. (2010). Evaluation of energy and dietary intake estimates from a food frequency questionnaire using independent energy expenditure measurement and weighed food records. *Nutrition Journal*, *9*(1), 37–37. https://doi.org/10.1186/1475-2891-9-37

Dhurandhar, N. V., Schoeller, D., Brown, A. W., Heymsfield, S. B., Thomas, D., Sørensen, T. I. A., Speakman, J. R., Jeansonne, M., & Allison, D. B. (2015). Energy balance measurement: When something is not better than nothing. *International Journal of Obesity*, *39*(7), 1109–1113. https://doi.org/10.1038/ijo.2014.199

Fazzino, T. L., Rohde, K., & Sullivan, D. K. (2019). Hyper-palatable foods: Development of a quantitative definition and application to the US food system database. *Obesity (Silver Spring)*, *27*(11), 1761–1768. https://doi.org/10.1002/oby.22639

Gosby, A. K., Conigrave, A. D., Lau, N. S., Iglesias, M. A., Hall, R. M., Jebb, S. A.,
Brand-Miller, J., Caterson, I. D., Raubenheimer, D., & Simpson, S. J. (2011). Testing
protein leverage in lean humans: A randomised controlled experimental study. *PloS*
One, *6*(10), e25929–e25929. https://doi.org/10.1371/journal.pone.0025929

Gosby, A. K., Lau, N. S., Tam, C. S., Iglesias, M. A., Morrison, C. D., Caterson, I. D.,
Brand-Miller, J., Conigrave, A. D., Raubenheimer, D., & Simpson, S. J. (2016).
Raised FGF-21 and Triglycerides Accompany Increased Energy Intake Driven by
Protein Leverage in Lean, Healthy Individuals: A Randomised Trial. *PloS One*, *11*(8),
e0161003–e0161003. https://doi.org/10.1371/journal.pone.0161003

Grech, A., Rangan, A., Allman-Farinelli, M., Simpson, S. J., Gill, T., & Raubenheimer,
D. (2022a). A comparison of the Australian dietary guidelines to the NOVA
classification system in classifying foods to predict energy intakes and body mass
index. *Nutrients*, *14*(19), 3942. https://doi.org/10.3390/nu14193942

Grech, A., Sui, Z., Rangan, A., Simpson, S. J., Coogan, S. C. P., & Raubenheimer, D.
(2022b). Macronutrient (im)balance drives energy intake in an obesogenic food
environment: An ecological analysis. *Obesity (Silver Spring)*, *30*(11), 2156–2166.
https://doi.org/10.1002/oby.23578

Grimby, G., Börjesson, M., Jonsdottir, I. H., Schnohr, P., Thelle, D. S., & Saltin, B.
(2015). The "Saltin-Grimby Physical Activity Level Scale" and its application to health
research. *Scandinavian Journal of Medicine and Science in Sports*, *25*(S4), 119–125.
https://doi.org/10.1111/sms.12611

Hall, K. D. (2019). The potential role of protein leverage in the US obesity epidemic.
Obesity (Silver Spring), *27*(8), 1222–1224. https://doi.org/10.1002/oby.22520

Hall, K. D., Ayuketah, A., Brychta, R., Cai, H., Cassimatis, T., Chen, K. Y., Chung, S.
T., Costa, E., Courville, A., Darcey, V., Fletcher, L. A., Forde, C. G., Gharib, A. M.,
Guo, J., Howard, R., Joseph, P. V., McGehee, S., Ouwerkerk, R., Raisinger,
K.,…Zhou, M. (2019). Ultra-processed diets cause excess calorie intake and weight
gain: An inpatient randomized controlled trial of ad libitum food intake. *Cell*
Metabolism, *32*(4), 690–690. https://doi.org/10.1016/j.cmet.2020.08.014

Hamano, S., Sawada, M., Aihara, M., Sakurai, Y., Sekine, R., Usami, S., Kubota, N.,
& Yamauchi, T. (2024). Ultra-processed foods cause weight gain and increased
energy intake associated with reduced chewing frequency: A randomized, open-label,
crossover study. *Diabetes, Obesity and Metabolism*, *26*(11), 5431–5443.
https://doi.org/10.1111/dom.15922

Hill, C. M., Laeger, T., Dehner, M., Albarado, D. C., Clarke, B., Wanders, D., Burke, S.
J., Collier, J. J., Qualls-Creekmore, E., & Solon-Biet, S. M. (2019). FGF21 signals
protein status to the brain and adaptively regulates food choice and metabolism. *Cell*
Reports, *27*(10), 2934–2947. e2933. https://doi.org/10.1016/j.celrep.2019.05.022

Hollstein, T., Ando, T., Basolo, A., Krakoff, J., Votruba, S. B., & Piaggi, P. (2019).
Metabolic response to fasting predicts weight gain during low-protein overfeeding in
lean men: further evidence for spendthrift and thrifty metabolic phenotypes. *American*
Journal of Clinical Nutrition, *110*(3), 593–604. https://doi.org/10.1093/ajcn/nqz062

Honfo, S. H., Senior, A. M., Legault, V., Presse, N., Turcot, V., Gaudreau, P.,
Simpson, S. J., Raubenheimer, D., & Cohen, A. A. (2023). Evidence for protein

leverage on total energy intake, but not body mass index, in a large cohort of older
adults. *International Journal of Obesity*. https://doi.org/10.1038/s41366-023-01455-6

Hopstock, L. A., Grimsgaard, S., Johansen, H., Kanstad, K., Wilsgaard, T., & Eggen,
A. E. (2022). The seventh survey of the Tromsø Study (Tromsø7) 2015–2016: Study
design, data collection, attendance, and prevalence of risk factors and disease in a
multipurpose population-based health survey. *Scandinavian Journal of Public Health*,
50(7), 919–929. https://doi.org/10.1177/14034948221092294

Howes, E. M., Laskaridou, E., Davy, K. P., & Hedrick, V. E. (2024). Representation of
adults with class III obesity in studies assessing validity of self‑reported energy intake
using doubly labeled water: A systematic review. *Obesity Reviews*, *25*(5), e13713–
n/a. https://doi.org/10.1111/obr.13713

Huybrechts, I., Rauber, F., Nicolas, G., Casagrande, C., Kliemann, N., Wedekind, R.,
Biessy, C., Scalbert, A., Touvier, M., Aleksandrova, K., Jakszyn, P., Skeie, G.,
Bajracharya, R., Boer, J. M. A., Borné, Y., Chajes, V., Dahm, C. C., Dansero, L.,
Guevara, M.,…Levy, R. B. (2022). Characterization of the degree of food processing
in the European Prospective Investigation into Cancer and Nutrition: Application of
the Nova classification and validation using selected biomarkers of food processing.
Frontiers in Nutrition, *9*, 1035580–1035580.
https://doi.org/10.3389/fnut.2022.1035580

Laeger, T., Henagan, T. M., Albarado, D. C., Redman, L. M., Bray, G. A., Noland, R.
C., Munzberg, H., Hutson, S. M., Gettys, T. W., Schwartz, M. W., & Morrison, C. D.
(2014). FGF21 is an endocrine signal of protein restriction. *The Journal of clinical
investigation*, *124*(9), 3913–3922. https://doi.org/10.1172/JCI74915

Lawson, J., & Willden, C. (2016). Mixture experiments in R using mixexp. *Journal of
statistical software*, *72*(Code Snippet 2), 1–20. https://doi.org/10.18637/jss.v072.c02

Lundblad, M. W., Andersen, L. F., Jacobsen, B. K., Carlsen, M. H., Hjartåker, A.,
Grimsgaard, S., & Hopstock, L. A. (2019). Energy and nutrient intakes in relation to
National Nutrition Recommendations in a Norwegian population-based sample: The
Tromsø Study 2015-16. *Food & Nutrition Research*, *63*, 1–10.
https://doi.org/10.29219/fnr.v63.3616

Løvsletten, O., Jacobsen, B. K., Grimsgaard, S., Njølstad, I., Wilsgaard, T., Løchen,
M.-L., Eggen, A. E., & Hopstock, L. A. (2020). Prevalence of general and abdominal
obesity in 2015–2016 and 8-year longitudinal weight and waist circumference
changes in adults and elderly: The Tromsø Study. *BMJ Open*, *10*(11), e038465–
e038465. https://doi.org/10.1136/bmjopen-2020-038465

Martens, E. A., Lemmens, S. G., & Westerterp-Plantenga, M. S. (2013). Protein
leverage affects energy intake of high-protein diets in humans. *The American Journal
of Clinical Nutrition*, *97*(1), 86–93. https://doi.org/10.3945/ajcn.112.046540

Martínez Steele, E., Raubenheimer, D., Simpson, S. J., Baraldi, L. G., & Monteiro, C.
A. (2018). Ultra-processed foods, protein leverage and energy intake in the USA.
Public Health Nutrition, *21*(1), 114–124. https://doi.org/10.1017/S1368980017001574

Mizushige, T., Inoue, K., & Fushiki, T. (2007). Why is fat so tasty? Chemical reception
of fatty acid on the tongue. *Journal of Nutritional Science and Vitaminology*, *53*(1), 1–
4. https://doi.org/10.3177/jnsv.53.1

Monteiro, C. A. (2009). Nutrition and health. The issue is not food, nor nutrients, so
much as processing. *Public Health Nutrition*, *12*(5), 729–731.
https://doi.org/10.1017/S1368980009005291

Monteiro, C. A., Cannon, G., Levy, R. B., Moubarac, J.-C., Louzada, M. L. C.,
Rauber, F., Khandpur, N., Cediel, G., Neri, D., Martinez-Steele, E., Baraldi, L. G., &
Jaime, P. C. (2019). Ultra-processed foods: What they are and how to identify them.
Public Health Nutrition, *22*(5), 936–941. https://doi.org/10.1017/S1368980018003762

Nicolaisen, T. S., Lyster, A. E., Sjøberg, K. A., Haas, D. T., Voldstedlund, C. T.,
Lundsgaard, A.-M., Jensen, J. K., Madsen, E. M., Nielsen, C. K., & Bloch-Ibenfeldt,
M. (2025). Dietary protein restriction elevates FGF21 levels and energy requirements
to maintain body weight in lean men. *Nature metabolism*, 1–15.
https://doi.org/10.1038/s42255-025-01236-7

Poslusna, K., Ruprich, J., de Vries, J. H. M., Jakubikova, M., & van't Veer, P. (2009).
Misreporting of energy and micronutrient intake estimated by food records and
24 hour recalls, control and adjustment methods in practice. *British Journal of*
Nutrition, *101*(S2), S73–S85. https://doi.org/10.1017/S0007114509990602

Prentice, R. L., Aragaki, A. K., Zheng, C., Manson, J. E., Tinker, L. F., Schoeller, D.
A., Ravelli, M. N., Raftery, D., Gowda, G. A. N., Navarro, S. L., Huang, Y., Mossavar-
Rahmani, Y., Wallace, R. B., Johnson, K. C., Lampe, J. W., & Neuhouser, M. L.
(2025). Energy intake is associated with dietary macronutrient densities: inversely
with protein and monounsaturated fat and positively with polyunsaturated fat and
carbohydrate among postmenopausal females. *American Journal of Clinical Nutrition*,
Advance online publication. https://doi.org/10.1016/j.ajcnut.2025.03.011

Raubenheimer, D., Machovsky-Capuska, G. E., Gosby, A. K., & Simpson, S. (2015).
Nutritional ecology of obesity: From humans to companion animals. *British Journal of*
Nutrition, *113*(S1), S26–S39. https://doi.org/10.1017/S0007114514002323

Raubenheimer, D., Senior, A. M., Mirth, C., Cui, Z., Hou, R., Le Couteur, D. G.,
Solon-Biet, S. M., Léopold, P., & Simpson, S. J. (2022). An integrative approach to
dietary balance across the life course. *iScience*, *25*(5), 104315–104315.
https://doi.org/10.1016/j.isci.2022.104315

Raubenheimer, D., & Simpson, S. J. (2023). Protein appetite as an integrator in the
obesity system: The protein leverage hypothesis. *Philosophical Transactions of the*
Royal Society of London. Series B: Biological Sciences, *378*(1888), 20220212–
20220212. https://doi.org/10.1098/rstb.2022.0212

Saner, C., Senior, A. M., Zhang, H., Eloranta, A.-M., Magnussen, C. G., Sabin, M. A.,
Juonala, M., Janner, M., Burgner, D. P., Schwab, U., Haapala, E. A., Heitmann, B. L.,
Simpson, S. J., Raubenheimer, D., & Lakka, T. A. (2023). Evidence for protein
leverage in a general population sample of children and adolescents. *European*
Journal of Clinical Nutrition, *77*(6), 652–659. https://doi.org/10.1038/s41430-023-
01276-w

Saner, C., Tassoni, D., Harcourt, B. E., Kao, K. T., Alexander, E. J., McCallum, Z.,
Olds, T., Rowlands, A. V., Burgner, D. P., Simpson, S. J., Raubenheimer, D., Senior,
A. M., Juonala, M., & Sabin, M. A. (2020). Evidence for protein leverage in children

and adolescents with obesity. *Obesity (Silver Spring)*, *28*(4), 822–829.
https://doi.org/10.1002/oby.22755

Senior, A. M., Raubenheimer, D., & Simpson, S. J. (2022). Testing the protein-
leverage hypothesis using population surveillance data. *Royal Society Open Science*,
9(9), 220756. https://doi.org/10.1098/rsos.220756

Simpson, S. J., & Raubenheimer, D. (2005). Obesity: The protein leverage
hypothesis. *Obesity Reviews*, *6*(2), 133–142. https://doi.org/10.1111/j.1467-
789X.2005.00178.x

Solon-Biet, S. M., Clark, X., Bell-Anderson, K., Rusu, P. M., Perks, R., Freire, T.,
Pulpitel, T., Senior, A. M., Hoy, A. J., & Aung, O. (2023). Toward reconciling the roles
of FGF21 in protein appetite, sweet preference, and energy expenditure. *Cell
Reports*, *42*(12). https://doi.org/10.1016/j.celrep.2023.113536

Vinales, K. L., Begaye, B., Bogardus, C., Walter, M., Krakoff, J., & Piaggi, P. (2019).
FGF21 is a hormonal mediator of the human "thrifty" metabolic phenotype. *Diabetes*,
68(2), 318–323. https://doi.org/10.2337/db18-0696

Volcko, K. L., & McCutcheon, J. E. (2022). Protein preference and elevated plasma
FGF21 induced by dietary protein restriction is similar in both male and female mice.
Physiology and Behavior, *257*, 113994.
https://doi.org/10.1016/j.physbeh.2022.113994

World Health Organization. (2013). *Global action plan for the prevention and control
of noncommunicable diseases 2013-2020*. World Health Organization.
https://www.who.int/publications/i/item/9789241506236

World Health Organization. (2021). *Obesity and overweight*. Retrieved 11.08.2023
from https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight

Zhang, H., Senior, A. M., Saner, C., Koemel, N. A., Simpson, S. J., Raubenheimer,
D., & Heitmann, B. L. (2025). Maternal protein intake during pregnancy and obesity
risk in mothers and offspring: a prospective cohort study. *The American journal of
clinical nutrition*, *121*(6), 1415–1423. https://doi.org/10.1016/j.ajcnut.2025.04.011

Zhang, H., Senior, A. M., Saner, C., Olsen, N. J., Larsen, S. C., Simpson, S. J.,
Raubenheimer, D., & Heitmann, B. L. (2023). Evidence for the protein leverage
hypothesis in preschool children prone to obesity. *Clinical Nutrition*, *42*(11), 2249–
2257. https://doi.org/10.1016/j.clnu.2023.09.025

Tables

Table 1 | Descriptive statistics for the full sample from the Tromsø Study 2015–2016

Variable	Group	Count
Age (groups of 10 years)	40–49	28.6%
	50–59	28.8%
	60–69	27.7%
	70–79	12.7%
	80+	2.3%
Body mass index (kg/m^2)	Normal (<25)	33.4%
	Overweight (25–<30)	44.0%
	Obese (≥30)	22.7%
Total energy intake(kJ/day)		9,722 (3,048)
Protein intake (%E)		17.6 (2.5)
Carbohydrate intake (%E)		42.2 (6.2)
Fat intake (%E)		34.5 (5.8)
UPF intake (%E)		54.8 (9.7)
Education	Primary/partly secondary education	20.4%
	Upper secondary	27.2%

Variable	Group	Count
	Tertiary short	20.6%
	Tertiary long	31.9%
Physical activity level	Sedentary	12.9%
	Low	58.7%
	Moderate	25.4%
	Vigorous	3.0%
Health status (% with conditions)	Non-communicable diseases	31.2%
	Chronic pain	21.8%
	Psychiatric problems	3.9%
Current smoker		12.5%

Note. n = 11,152. Numbers are means and standard deviations for continuous variables and percentages for categorical variables. The age group 80+ includes participants aged between 80–99. Health status does not sum to 100% because not all participants had health challenges. %E = percent of total energy intake.

Table 2 | Output from power functions with percent protein as predictor

Total energy intake (kJ)		Energy density (kJ/g)		Food dry weight (g)	
L	*p*	*L*	*p*	*L*	*p*
-0.36	< .001	- .48	< .001	-0.31	< .001

Note. Energy density is kJ/ wet weight of food; food dry weight is dry weight of all energy yielding nutrients.

Table 3 | Regression model for fibroblast growth factor 21 by dietary protein

Variable	b	SE	t	p
Intercept	5.30	0.35	15.2	< .001
Protein, %E	-0.05	0.02	-3.2	.001
Age	0.28	0.09	3.0	.002
Sex	-0.12	0.08	-1.4	.168
Physical activity	-0.28	0.13	-2.1	.035
Disease	0.20	0.08	2.4	.016
Education level	-0.02	0.098	-0.2	.848
Smoking status	0.34	0.13	2.7	.008

Note. $n = 880$. Covariates were dichotomous and coded as follows: age = 0 (under 60 years), sex = 0 (female), physical activity level = 0 (sedentary), disease = 0 (no disease), education level 0 (≤10 years).

Figures

Figure 1 | Flowchart of exclusions leading to final sample. Tromsø7 data was collected in 2015-2016. Food frequency questionnaire (FFQ). % indicates percent change since the last step. To increase the validity of the data we excluded (1) participants that completed less than 90% of the FFQ and (2) participants with extreme intakes of energy, i.e., the 1st (below 3,950 kJ/day) and 99th (above 21,249 kJ/day) percentile in total energy intake (TEI). Pregnancy may interfere with habitual eating patterns and body composition and we therefore excluded (3) participants that reported being pregnant at the time of the data collection. Finally, we excluded (4) participants with extreme BMI values, i.e., groups of BMI with less than 10 participants, ensured by binning the BMI data in bins of one BMI unit and excluding bins with fewer than 10 observations.

Figure 2 I Power functions between total energy intake and proportional intake of each macronutrient. (A) shows protein, (B) shows carbohydrate, and (C) shows fat. Dashed lines show regression lines fitted with Equation 1.1. Data on individual participants are presented as a density heat map (darker colors indicate higher density of observations). *L* is the leverage strength of each macronutrient on energy intake where a positive number indicates that total energy intake increases as percentage of the macronutrient increases and a negative value indicates that total energy intake decreases as percentage of the macronutrient increases.

Figure 3 | Equilateral mixture triangle for predicted total energy intake by macronutrient proportions. Predicted total energy intake based on the mixture model shown in Table S2 is shown as colored contours. Negatively sloped lines indicate percent protein, positively sloped lines indicate percent fat, and horizontal lines indicate percent carbohydrate. As percent protein increases, predicted energy intake decreases and as percent fat increases, predicted energy intake increases. Changes to percent carbohydrate have only a minor effect on predicted energy intake.

Figure 4 | Relationship between ultra-processed foods (UPFs), dietary patterns, and BMI. (A) Descriptive bi-coordinate intake plot of protein versus non-protein energy intake (MJ) by quintiles (Qs) of UPF intake. Dots indicate mean intake of protein energy and non-protein energy (i.e., the combination of carbohydrate and fat) per quintile of UPF intake. The quintile dots most closely follow the protein prioritization line. Solid grey lines represent different predictions of intake for Q1 based on energy prioritization (negatively sloped line), protein prioritization (vertical line) or non-protein energy prioritization (horizontal line). Dashed grey lines show shift in total energy intake (negatively sloped) and protein intake (positively sloped) for Q5. *n* = 11,152. (B) Relationship between BMI and UPF intake plotted as quintiles. Data are estimated marginal means from regression model that control for sex, age, and physical activity. Error bars are 95%CI. *n* = 11,125.

Figure 5 | Protein leverage in low and high consumers of ultra-processed foods (UPFs). (A) Plotting of power function showing that the strength of leverage that protein exerts over total energy intake is greater in low vs. high consumers of UPFs after median split. (B) Histogram showing the distribution of the differences in L values between low and high UPF consumers after randomly shuffling labels 10,000 times. Dashed vertical line shows the observed difference from the real data.