Estimating Realized Access to Obstetric Care in Georgia: A Discrete Choice Modeling Analysis

Jingyu Li, MS[1], Margaret Carrel, PhD[2], Stephanie Radke, MD, MPH[3], Lauren N. Steimle, PhD[1]

1. H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, Georgia, USA
2. Department of Geographical & Sustainability Sciences, University of Iowa, Iowa City, Iowa, USA
3. Department of Obstetrics & Gynecology, University of Iowa Hospitals & Clinics, Iowa City, Iowa, USA

Corresponding Author

Lauren N. Steimle, PhD

Telephone: (404) 894-4659

Email: steimle@gatech.edu

Address: 755 Ferst Dr NW, Atlanta, Georgia 30318

Statements and Declarations

Data Availability

We are not able to share Georgia birth records data due to our data use agreement.

Ethics Approval

This study was conducted in accordance with the ethical principles outlined in the Declaration of Helsinki (https://www.wma.net/policies-post/wma-declaration-of-helsinki/). IRB approval was obtained from the Georgia Institute of Technology (Protocol H23091), and the need for consent was waived by the IRB.

Funding Information

This study was supported by the Georgia Clinical and Translational Science Alliance and National Center for Advancing Translational Sciences of the National Institutes of Health under Award number UL1TR002378 and the Harold R. and Mary Anne Nash endowment to the Georgia Institute of Technology. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing Interests

The authors declare no completing interests.

Authors' Contributions

Jingyu Li: conceptualization, data curation, methodology, formal analysis, validation, writing – original draft, visualization.

Margaret Carrel: validation, writing – review and editing.

Stephanie M. Radke: validation, writing – review and editing.

Lauren N. Steimle: data acquisition, conceptualization, methodology, writing – original draft, project administration, funding acquisition, supervision.

Acknowledgements

We want to thank Dr. Stephane Hess, the co-author of the Apollo package, for his help with the initial software setup and troubleshooting. We also want to thank the Georgia Department of Public Health for providing birth records data.

Abstract

The ongoing closure of obstetric units in the United States poses challenges to maternal care access, especially in rural communities. These closures leave patients with limited options for obstetric services and may force them to travel longer distances or rely on low-volume facilities, both associated with adverse maternal outcomes. While policymakers have proposed interventions to support obstetric care in underserved areas, evaluating their potential impact requires understanding how birthing patients select among available facilities and how those selections affect delivery volume and access patterns. This study develops a revealed preference discrete choice modeling framework to estimate realized access to obstetric facilities—the extent to which patients actually use available obstetric services—based on patient characteristics and facility attributes. Using 2016-2019 birth records in the state of Georgia, we developed three discrete choice models: a distance-only model, a multinomial logit model based on facility attributes, and a latent class model that considers heterogeneous weights patients place on facility attributes. We evaluated model performance in predicting low-volume facilities and regional access. The latent class model outperformed others and identified two patient classes: "less distance-sensitive" patients (34.5%) and "more distance-sensitive" patients (65.5%) with differing sensitivity to distance and levels of care. Our findings reveal regional patterns where patients disproportionately seek care far away or at low-volume facilities, and how these patterns may change as obstetric units close. We highlight the practical use of our model for forecasting underutilized facilities and informing policies to better distribute patient volume across the region.

Keywords

Discrete choice modeling, patient behavior, maternal health, rural health, access to care

Highlights

- We analyzed how pregnant people in Georgia, USA sought obstetric care using statewide birth records from 2016-2019

- We developed and compared three discrete choice models (distance-only model that assumes patients select nearest obstetric facilities, multinomial logit model, and latent class model) that considered patient characteristics and facility attributes to predict patients' utilization of birth facilities and validated these models using out-of-sample datapoints.

- Travel distance, facility level of care, rurality of facility location, and health system affiliation significantly influenced patients' selection of delivery facilities. We also identified two classes of patients who varied in their social determinants of health and placed heterogenous weights on facility attributes that affected their selection of obstetric facilities.

- Targeted interventions to improve maternal care access in underserved areas should consider patients' underlying social determinants of health, as geographic proximity alone is not predictive of patients' selection of delivery facility.

- We demonstrate how our model can be applied to forecast low-volume obstetric facilities and access to care and to inform policies that avoid underutilized facilities and better distribute patient volume across the region.

1. Introduction

Closures of obstetric (OB) units in the United States (U.S.) present a critical challenge to maternal care access, particularly in rural communities. Between 2006 and 2020, over 400 OB units shut down nationwide [1]. The disappearance of OB units has been particularly devastating in rural areas, where 267 facilities, nearly a quarter of all rural OB units in the U.S., stopped providing childbirth services between 2011 and 2021 [2]. These closures have left rural populations with increasingly limited options for OB care. As a result, some patients, especially those residing in rural areas, must travel long distances to reach a delivery facility. Others may rely on a more local option for care, which is often a low-volume facility for rural populations. Far travel distance and delivery in low-volume rural facilities are both associated with increased rates of maternal morbidity and mortality [3, 4].

The reasons for OB unit closures have been well-documented. Hospital administrators commonly cite financial viability and workforce shortages as reasons for closing OB units [5]. Operating an OB unit incurs large fixed costs each year, which hospitals seek to make up through reimbursements. However, facilities with low patient volume may struggle to offset the fixed cost associated with the OB service lines [6], and thus to maintain sufficient patient volume for financial viability and patient safety. This challenge is compounded by patient *bypass*, as estimates suggest between 11% and 55% of patients choose to bypass their local birthing facility in favor of one farther away [7, 8]. These decisions may reflect a preference for higher-quality or more risk-appropriate care [9, 10], dissatisfaction with local services [11], or desire for culturally concordant care [12, 13]. When closures occur, they can trigger a negative cycle as departing obstetricians and nurse-midwives further exacerbate the existing workforce shortages.

Thus, as policymakers seek to address gaps in access to OB care and to support rural OB facilities, they must understand how birthing people make decisions about where to deliver, given the birthing facilities available. Traditional measures of access to OB care, such as geographic proximity to birthing facilities, overlook behavioral dimensions of access—how patients respond to available options, as patients may bypass nearby

facilities in response to perceived quality, distance, or facility attributes. Simply

assessing historical volume trends is insufficient for guiding forward-looking initiatives,

as this approach fails to capture how patients will make selections of their birthing

facility as the available OB units change. To understand how patients seek birthing

facilities and how these choices may change in response to closures or consolidations,

we use revealed preference (RP) discrete choice modeling to understand *realized*

access to OB care in the state of Georgia, that is, where patients actually seek care as

opposed to simply how far away they live from the closest birthing facility.

1.1. Literature Review

1.1.1 Access to OB care

Access to care is a multidimensional concept. In this study, we focus specifically on

geographic access, which is often assessed through two distinct perspectives: *potential*

access and *realized access* [14]. Potential access refers to the spatial availability of

care services, while realized access captures the actual utilization of those services by

patients given their location relative to care. In the context of OB care, potential access

is typically measured by indicators such as provider-to-population ratios, travel distance

to the nearest delivery facility, or the presence of a birthing facility in some geographic

unit. For example, the "maternity care desert" framework developed by the March of

Dimes identifies counties with limited or no access to OB services based on the

availability of maternity care providers and facility-based OB services (March of Dimes).

Another measure is the distance to facilities offering critical care OB (CCO) services [15,

16]. While these measures are valuable for identifying structural gaps, particularly in

underserved areas, they fail to capture how individuals use available services. Such

measures assume patients will seek care at the nearest facility and overlook the

variations in preferences, perceived quality, transportation access, and other social

determinants of health.

In contrast, realized access reflects patients' actual use of healthcare services when

seeking care. For instance, some patients bypass their closest OB facility to seek care

at a farther OB unit. Recent studies have documented this behavior empirically in the

U.S. Thorsen et al. reported that 11% of births in Montana occurred at facilities farther from patients' closest labor and delivery (L&D) units [8]. In Iowa, more than half of patients bypassed the nearest OB provider when delivering [7]. Several factors are associated with increased likelihood of bypassing including complex medical diagnoses, higher income, younger age, private insurance coverage, and expanded Medicaid eligibility [7, 9, 17–21]. Conversely, patients with lower income, inadequate prenatal care, or public insurance are more likely to deliver locally [22, 23].

Despite these insights, existing studies often fall short in capturing the full complexity of patient decision-making. Many rely on descriptive or regression-based analyses that do not adequately model the choice process or account for variation across individuals. In addition, many of these studies do not incorporate facility-level and patient-level characteristics simultaneously, or fail to validate findings on out-of-sample data. Their utility for informing policy design and intervention planning needs further investigation.

1.1.2 Background on discrete choice modeling

Understanding what drives patients' selection of birthing facilities is a key component for accurately estimating patients' realized access to OB care and thus identifying regions that may warrant targeted support. In recent years, discrete choice modeling has emerged as an approach to analyze such decisions. Discrete choice models, grounded in behavioral theory, estimate the probability that a decision-maker (DMs) selects a particular option from a defined set of distinctive alternatives and are well-suited for forecasting choices under hypothetical scenarios [24]. They have been applied across a range of contexts, including transportation and travel behavior [25, 26], marketing and retail [27, 28], public policy [29–31], and operations management [32]. A detailed review of this subject is provided by Train 2009 [33].

A central distinction in discrete choice modeling is between revealed preferences (RP) and stated preferences (SP). RP are derived from actual choices DMs have made in real-world settings. In contrast, SP are elicited through hypothetical scenarios such as surveys or experiments where individuals are asked to indicate their choices [34]. RP

data reflect real behavior but may lack variation in key variables as those data are pre-

collected, while SP data can be designed to capture such variation but the consistency

of choices may be influenced by how a question is asked.

The theoretical foundation of discrete choice modeling lies in utility maximization theory,

which posits that individuals act rationally to maximize their utility, subject to constraints.

In the context of discrete choices, utility is composed of an observable component

(based on measurable attributes of the alternatives and DM characteristics) and an

unobservable random component. Multinomial logit (MNL) model is one of the most

widely used discrete choice models and the basis for many other random utility

maximization (RUM) models. The key assumption of MNL model is that the random

component is independently and identically distributed with a Gumbel distribution. This

assumption allows for a closed-form choice probability but imposes the independence of

irrelevant alternatives (IIA) property, which may be restrictive especially when

alternatives share unobserved similarities [35]. To address the limitations of MNL

models, latent class (LC) model is introduced to handle unobserved heterogeneity in

individual taste. This model is built on the idea of a finite set of distinct "patient classes,"

allowing attribute level coefficients to remain consistent within each class but vary

between classes [36].

Prior literature has demonstrated discrete choice modeling as a promising approach to

understand patients' choices in health services research. Studies have consistently

identified travel distance and quality of care as key factors affecting patients' healthcare

choices in both U.S. and international contexts [37–41]. More recent studies have

emphasized that patients are not a homogeneous group, and that preferences vary by

demographic, socioeconomic, and clinical characteristics [37, 42–45]. For example,

several studies have identified subgroups of patients who are more willing to travel

longer distances to access higher-quality care [37, 45].

To our knowledge, the most closely related studies to ours are that of Hwang et al.,

2017 and Jang et al, 2017 who used MNL models to estimate patients' realized

accessibility to OB care in South Korea [46, 47]. Others have formulated a gravity-

based nonlinear optimization model to estimate access at rural OB hospitals in Montana

[48]. However, these studies typically rely only on facility-level attributes and do not

incorporate patient-level heterogeneity, which limits their ability to capture variation in

behavior that may be driven by patient-level characteristics. Another study used a

latent-class choice model to understand the impact of social determinants of health on

the choice of delivery facilities among pregnant women in low-income country, which is

a very different context for OB care and provides limited understanding into facility

closures in the U.S. [44].

Our study presents a novel way to understand patients' delivery facilities choices that

includes both facility and patient-level covariates in the unique context of OB unit

closure. This RP approach allows us to account for heterogeneity in how patients

evaluate facility attributes. We further validate our model using out-of-sample data and

assess performance on outcomes directly relevant to access measure—such as

proximity to the delivery facility and use of low-volume facilities.

1.2 Contributions

In this paper, we present a novel application of discrete choice modeling to estimate and

predict realized access to birthing facilities. To do so, we conducted a revealed-

preference case study in the state of Georgia. We focus on a single-state study

because although general healthcare regulations are imposed at the national level

under the U.S. healthcare governance structure, there is substantial variation across

states, as many specific decisions are made at the state level. Georgia was selected as

a highly relevant case study due to its high rates of maternal morbidity and mortality and

large rural population. Our study makes several contributions:

• First, we demonstrate that realized access differs significantly from access

defined by geographic proximity alone. By incorporating factors such as facility

level of care, health system affiliation, and location into choice modeling, we

provide a more accurate representation of how patients select delivery facilities in

the U.S. context. This result further suggests the limitations of proximity-based

forward-planning approaches and emphasizes the importance of considering

patients' actual choices when evaluating maternal care access.

•   Second, we assess model performance against several critical measures,

including (1) geographic proximity between the patients' residence and delivery

facility, and (2) whether patients deliver at low-volume hospitals, which are

associated with poorer maternal outcomes. These validation steps demonstrate

the practical utility of our model in identifying geographic areas and

subpopulations where targeted intervention to improve OB access are most

needed.

•   Third, we distinguish different types of patients that weigh facility attributes

differently. This finding suggests interventions to improve maternal care access

should be tailored to the needs of different patient subgroups.

•   Lastly, we make innovative use of statewide birth records as an RP dataset.

These records provide a high-resolution, population-level view of patients' actual

facility selection of birthing facilities. Such practice can be easily adapted for use

in other states with different healthcare regulatory environments or patterns of

healthcare access disparities.

2. Methods

2.1 Sample

We analyzed birth records for all births to Georgia residents from January 1st, 2016, to

December 31st, 2019, in the state of Georgia. Figure 1 provides an overview of our

exclusion criteria of birth records. First, we excluded deliveries that had missing records

of residence or birth facilities. Then, we excluded deliveries that involved a maternal

transport as these cases do not reflect patients' original selection of delivery facilities.

Additionally, we excluded deliveries at non-OB facilities that are not included in the 2017

Georgia guidelines for perinatal regionalization, since we expected that these cases

likely reflect situations where delivery was urgent and there was limited to no patient

control over birth facility selection [49]. Lastly, we further excluded deliveries that

occurred more than 120 miles away from the birthing person's residential address, as

selecting a facility at such a distance is highly unlikely under typical circumstances. We
suspected that such cases were likely due to emergencies that occurred while the
patient was traveling or to high-risk, complex deliveries requiring specialized care
available only at distant facilities. Since 99.75% of all deliveries occurred within 120
miles of the birthing person's residence, excluding these observations is expected to
have minimal impact on the analysis. The final sample included 489,512 deliveries. The
Georgia Tech Institutional Review Board approved this study (Protocol H23091). We
fitted our models using 2016-2018 birth records in Georgia and validated our models
using 2019 birth records.

2.2 Discrete choice modeling framework

In this section, we describe the data used to fit our discrete choice models and our
modeling process.

2.2.1. Factors included in discrete choice models

We considered both facility-level attributes and patient characteristics as factors in our
discrete choice models. For patient characteristics, we included all medical risk factors
(for example, number of fetuses, presence of gestational diabetes, and or hypertension)
and sociodemographic variables (for example, age, race, and ethnicity) available in the
birth records. A detailed description of those variables can be found in **Appendix Table
A1**. To guide selection of facility-level attributes, we adapted a conceptual framework
that identified seven domains of provider characteristics that influence patient decision-
making [46, 50]. Specifically, we considered: availability (availability of the provider,
such as open hours and variety of options which might be constrained by patients'
insurance plan), accessibility (whether the provider is easily accessible by patients' own
or public transportation), type and size, staff, organization of care, cost, and provider
demographics (demographics of individual doctors). Due to data availability, we
excluded cost and provider demographics from model fitting.

For availability, we assumed patients were free to choose from the set of all operating
OB facilities within 120 miles of their residence. As proxies for accessibility, we used

road distance between patients' residences and birthing facilities, as well as whether the

facility is in an urban area. We calculated road distance between patients' residence,

represented by the 2020 census block group population-weighted centroids and OB

facilities where deliveries occurred using Open Source Routing Machine (OSRM) API in

Python. As a proxy for type and size, we used *level of care* which represents a facility's

capabilities and resources for handling maternal and fetal risks [51]. To represent staff,

we included whether a facility offered midwifery services, defined as having at least one

delivery attended by a midwife. For the organization of care, we included two additional

factors from public sources, including Baby-Friendly designation [52], which indicates

that a facility adheres to evidence-based practices for breastfeeding and maternal-infant

care, and health system affiliation of each facility, which was obtained for each facility

each year.

To avoid multi-collinearity among factors, we calculated variance inflation factors for all

candidate variables and confirmed that none of them were correlated with each other

(**Appendix Table A2**).

2.2.2 Multinomial Logit Model

The MNL model represents a patient i's true utility of OB facility j (U_{ij}) among a set of

available OB facilities (J) as the sum of two parts: an observable utility (V_{ij}) and a

random error term (ϵ_{ij}). The observable utility can depend on both facility-related

attributes (X_{ij}) and patient-related characteristics (X_i), while the error term follows a

Gumbel distribution [35]. Let $D(i)$ denote patient i's decision, then the probability of

patient i choosing facility j (P_{ij}) can be calculated using a closed-form expression.

334
$$U_{ij} = V_{ij} + \epsilon_{ij}$$

336
$$V_{ij} = \beta_0 + \beta X_{ij} + \gamma_j X_i + \delta_j X_{ij}$$

338
$$P_{ij} = P\big(D(i) = j | X_{ij}, X_i\big) = \frac{e^{V_{ij}}}{\sum_{k \in J} e^{V_{ik}}}$$

Since choice sets available to each patient can vary, we do not estimate any alternative-

specific coefficients ($\beta_0 = \gamma_j = \delta_j = 0 \ \forall \ j \in J$).

To examine whether or not a LC model might be appropriate due to heterogeneity in

patient preference weights, we first fitted MNL models for subgroups of patients.

Specifically, we estimated separate models for groups defined by key patient

characteristics, such as race, ethnicity, education level, payer status, and rural versus

urban residence. For instance, we fitted separate models for rural and urban patients

and determined if there were differences in their coefficients. By comparing results for

rural and urban patients, we could evaluate whether facility attributes such as proximity

to residence and level of care influenced facility choice differently depending on where

patients lived.

2.2.3 Latent Class Model

The LC model takes into account heterogeneity in the weights patients might place on

the factors that influence their selection. The model includes two components: a class

membership model and a class-specific choice model. Let $Q(i)$ denote the index of the

class of the individual i. The class membership model calculates the likelihood of a

patient in each class c based on patient characteristics $P(Q(i) = c | X_i)$. Meanwhile, the

class-specific choice model estimates the likelihood of selecting an OB facility given

each class c. Following the same notation, the probability of patient i choosing facility j

can be calculated as followed.

363
$$P_{ij} = \sum_{c \in C} P(Q(i) = c | X_i) P(D(i) = j | X_{ij}, X_i, Q(i) = c)$$

For this analysis, we began by including all relevant sociodemographic and medical risk

factors in our class membership model. A detailed description of the tested variables is

included in **Appendix Table A1**. To refine our model, we performed variable selection to

identify variables that were statistically significant and distinct across latent classes. We

also tested multiple number of latent classes and iteratively examined each for model

convergence, statistical significance, model fit, and interpretability. Since log likelihood

always improves as the number of latent classes increases, we used Akaike's

Information Criteria (AIC) and Bayesian Information Criterion (BIC), which have been

suggested in prior literature, to compare model fit during model selection process [53,

54].

Due to memory limitations of the working protected health data environment in which

the birth records were stored, we randomly sampled 10,000 datapoints from 2016-2018

records to fit MNL and LC models. To ensure the robustness of our sampling, we also

estimated a simplified model using all available data from 2016–2018. This model

included only three predictors: road distance, facility level of care, and whether the

facility is in an urban area. The coefficients estimated from the sampled dataset, which

have slightly wider 95% confidence intervals (CIs) due to the smaller sample size, still

narrowly encompass the corresponding estimates from the full dataset (**Appendix**

Tables A3 & A4). This suggests that our downsampling approach did not significantly

affect the estimated relationships, because the effects of the predictors observed in the

sampled data are statistically compatible with those in the full dataset.

2.3 Out-of-sample Validation

We compared and validated our fitted discrete choice models using 2019 data against

the distance-only model which assumes that patients choose the closest facility. We

used four evaluation metrics to compare these models. First, we used logarithmic score,

which is a widely adopted strictly proper scoring rule, to assess how well each model

predicted individual selection of OB facilities [55, 56]. This metric measures the choice

prediction performance by comparing the predicted choice probabilities with the actual

observed choices. Logarithmic score is unbounded from below, and a higher score

value indicates better probabilistic prediction. Second, we compared the performance of

predicting low-volume facilities of each model, measured by accuracy, specificity,

sensitivity, and F1 score. We defined a low-volume facility as an OB facility with less

than 500 annual delivery volume [57]. We also conducted a sensitivity analysis by

applying alternative thresholds of 250, 300, and 400 annual delivery volume per year.

In addition to evaluating predictive accuracy, we used the models to forecast key

dimensions of geographic access to OB care. However, access to OB care can be

conceptualized and quantified in multiple ways, and there is currently no standardized

measure or metric exists in the U.S. Distance to facility has been used in previous

literature and is linked to adverse maternal outcomes [3, 58, 59]. Therefore, we

compared the predicted average travel distance to the chosen facility at the census tract

level. Regions with higher predicted distances may indicate areas with reduced access.

Another important dimension of access to care is the facility volume due to volume

outcome relationships [4]. We then compared the predicted percentage of patients

delivering at low-volume facilities (defined as predicted to have less than 500 annual

deliveries) with the observed percentage (in facilities that actually had less than 500

annual deliveries). Finally, we calculated the Spearman's correlation coefficient between

the predicted and observed percentages across census tracts to assess model

performance in capturing spatial variation in access.

To facilitate interpretation across models, all facility attributes were translated into "road

distance equivalent" (RDE) units by dividing each attribute's coefficient by the (negative)

coefficient of the road distance. This allows the impact of non-distance attributes to be

interpreted as equivalent reductions in travel distance. RDE simplifies comparisons

across models and supports assessing the relative importance of each facility attribute

in terms of travel burden [60].

All discrete choice models were fitted in R (version 4.4.1) using package Apollo (version

0.3.3) [61]. All other analyses were performed in Python (version 3.11.9). We used a p-

value of 0.05 as the significance threshold throughout all analyses.

3. Results

3.1 Sample

A total of 489,512 deliveries met the inclusion criteria shown in Figure 1. Of these, 368,519 deliveries occurred between 2016 and 2018 (during modeling period) and 120,993 occurred in 2019 (during validation period). The analysis included 79 OB facilities, two of which closed their OB units during the study period—one in 2017 and the other in 2018. Among 77 operational OB facilities in 2019, 19 (24.68%) had annual volume less than 500, 18 (23.38%) had annual volume less than 400, 12 (15.56%) had annual volume less than 300, and 9 (11.69%) had annual volume less than 250. Descriptive statistics for the included facilities are presented in **Table 1**.

To ensure the modeling sample was representative of the population, we compared the distribution of patient characteristics and facility volumes in the sampled dataset and the full dataset using Chi-square test of independence. We observed no significant difference, suggesting that our sampled dataset had good fit of the full dataset.

3.2 MNL model

Table 2 presents the results of the estimated MNL model. RDE interpretation for each variable is detailed in **Table 4**. All facility-related attributes were statistically significant and were retained in the final model. The negative coefficient for road distance indicates that patients viewed longer travel distances unfavorably. Patients preferred facilities offering higher levels of care compared to level I facilities, with an increase in utility observed from birth centers to level III facilities (p < 0.001). When interpreted in RDE, patients would be equally willing to choose a birth center that is 20.39 miles closer, an RPC that is 9.09 miles farther, a Level II facility that is 6.79 miles farther, a Level III facility that is 13.21 miles farther, than a Level I facility. In addition, patients were willing to travel an additional 5.99 miles for facilities located in urban areas, 1.54 miles for those offering midwifery services, 2.14 miles for Baby-Friendly designated hospitals, and 3.76 miles for facilities affiliated with a multi-facility health system.

Fitted MNL models by subgroups are shown in **Appendix Figure B1-B5**. The results
further confirmed that patients placed heterogeneous importance weights on facility
attributes. For example, patients with commercial insurance and Medicaid would be
equally willing to choose a level III facility located 17.78 and 10.15 miles further than a
level 1 facility, respectively. Therefore, an LC model, which can account for exogeneity
in patients might better address that there is a dimension to the input data that is not
captured.

3.3 LC model

3.3.1 Estimates

We started with 2-class LC models and included all potential patient-level variables as
detailed in **Appendix Table A1**. We noticed that the coefficient for birth center
approached negative infinity since there were only 2 birth centers and very few
deliveries occurred at such facilities. To resolve this convergence issue, we assigned
the MNL coefficient for birth centers and held it as fixed during model estimation. We did
not test LC models with 4 classes or more because the estimation for 3-class models
was unstable and poor (**Appendix Table B1**).

Table 3 shows the results of the final estimated LC model, which had 2 classes. Their
RDE interpretation is detailed in **Table 4**. Coefficients and their 95% CIs for detected
latent classes are visualized in **Figure 2**. We accepted this model because it offered
good interpretability and had the best model fit (AIC = 33686.81; BIC = 33867.07;
adjusted ρ^2 = 0.5231) among all candidate models that identified statistically significant
and distinct patient-level variables across latent classes. We observed notable
differences in class-specific choice models across two identified latent classes. These
differences confirmed that distinguishing heterogeneity in importance weights for each
class was important. Specifically, road distance to OB facilities had more negative
coefficients in class 1 compared to class 2, which shows that patients in class 1 viewed
longer distances to facilities less favorably. To distinguish between these classes, we
refer to class 1 as "more distance-sensitive," and class 2 as "less distance-sensitive".
Both groups favored facilities with higher levels of care compared to level I facilities, as

suggested by positive coefficients for Level II, Level III, and RPCs for both groups.

When interpreted in terms of RDE, Class 1 patients would be equally willing to choose a

birth center that is 12.89 miles closer, a Level II facility 5.27 miles farther, a Level III

facility 6.74 miles farther, and an RPC 3.17 miles farther than a Level I facility. In

contrast, Class 2 patients would accept significantly longer travel distances: 33.74 miles

closer for a birth center, 17.43 miles farther for a Level II facility, 45.43 miles farther for a

Level III facility, and 47.24 miles farther for an RPC. This suggests that "less distance-

sensitive" patients showed much stronger preference to level III facilities and RPCs

compared to "more distance-sensitive" patients. Relying solely on a single-class model

as in MNL might have underestimate these effects, as the computed sample-wide

"average" coefficient could underestimate preference for level III and RPCs for "less

distance-sensitive" patients.

The class membership model revealed patient characteristics associated with each

latent class, as shown in **Table 3**. The model estimated that a randomly selected patient

would fall into the "more distance-sensitive" class with a probability of 65.54% and "less

distance-sensitive" with a probability of 34.46%. Patients with greater number of medical

risk factors (estimate: 0.243, 95% CI: 9.107-0.378, p = 0.001), those with commercial

insurance compared to those Medicaid-insured (estimate: 0.943, 95% CI: 0.754-1.133,

p<0.001) and those who had a college degree or higher compared to those had less

than 9th grade (estimate: 1.045, 95% CI: 0.500-1.589, p<0.001) were more likely to be

"less distance-sensitive".

3.3.2 Class profile

Table 5 details class profiles for the two latent classes (Class 1: "more distance-

sensitive" patients: 65.54%; Class 2: "less distance-sensitive" patients: 34.46%). We

calculated expected values for patient-level characteristics. All variables included in the

final LC class membership model showed distinct distribution across the two latent

classes. Specifically, the expected number of medical risk factors was 0.354 for the

"less-distance sensitive" group compared to 0.285 for the "more distance-sensitive"

group. 54.4% of the "more distance-sensitive" group were Medicaid insured, while only

30.0% of the "less distance-sensitive" group were Medicaid insured. Compared to Class

2, Class 1 had higher rates of commercially insured patients (57.4% vs. 31.0%),

patients with college or higher education (70.4% vs. 48.4%), and had lower rates of

patients with inadequate or intermediate level of prenatal care utilization index (18.4%

vs. 23.9%).

We also compared the expected values of sociodemographic variables for members of

each class and found slight differences among the classes in terms of age, race, and

ethnicity. Older, White, Asian, and non-Hispanic patients were more likely to be "less

distance-sensitive", while younger, Black or African-American, and Hispanic patients

were more likely to be "more distance-sensitive".

3.4 Out-of-sample Validation

3.4.1 Choice prediction

We validated the models' ability to predict individual choice of delivery facility (**Table 6**).

LC model yielded a higher logarithmic score of -1.73, compared to a logarithmic score

of -1.77 by MNL model and a logarithmic score of -35.54 by the distance-only model.

3.4.2 Forecast low-volume facilities

Figure 3 shows true volume vs. volume predicted by the LC model in 2019 by volume

decile. We also calculated and compared the LC model's predictive performance with

the distance-only model and the MNL model in forecasting low-volume facilities as

shown in **Table 7**. The LC model consistently outperformed the distance-only and MNL

models in terms of accuracy, sensitivity, and F1 score across all thresholds. At

thresholds of 250, 300, and 400, the LC model achieved the highest F1 scores of 57.1,

66.7, and 81.3, respectively, compared to lower scores from the other models. At the

threshold of 500, the LC model led in all metrics, with an accuracy of 93.5, sensitivity of

89.5, and F1 score of 87.2.

3.4.3 Measure access to care

Figures 4(a)-(d) show the average travel distance to the chosen facility from their residence by census track, observed during the validation period (a) and predicted by our three models (b)-(d). The distance-only model predicted average travel distances with a Spearman correlation of 0.77 compared to observed values. Average travel distance predicted by both the MNL and the LC models aligned better with the observed pattern, as indicated by higher correlation coefficients of 0.83 and 0.84 between the observed average travel distance per census track and the average travel distance per census track (**Table 8**).

Figure 5(a)-(d) show the percentage of patients who delivered at a low-volume facility with less than 500 annual birth volumes. All models identified similar regions where high percentage of patients received OB care at a facility predicted as a low-volume facility by each model. We observed that predicted pattern by the LC model again aligned the best with the observed pattern among all fitted models, indicated by the highest Spearman correlation coefficient of 0.42 between the observed percentage and the predicted percentage by the LC model (**Table 8**).

4. Discussion

In this article, we demonstrated a discrete choice modeling approach to estimate patients' realized access to delivery facilities. Our findings suggested that distance, facility level of care, location of facilities, and affiliation with a multi-facility health system significantly affected patients' selection of delivery location. Importantly, our study emphasized the importance of accounting for patient heterogeneity when evaluating access to care and designing policies to improve OB service delivery. Through a latent class analysis, we identified a small subgroup of "less distance-sensitive" patients (34.46%) who exhibited a stronger preference for facilities with higher levels of care relative to road distance to facilities compared to the majority of the "more distance-sensitive" group (65.54%). Our final LC model achieved an adjusted ρ^2 of 0.52, which exceeded the conventional threshold of 0.4 for a good model fit [62]. Additionally, we evaluated the predictive performance of our models using out-of-sample data. Fitted

models were applied to forecast patients' facility choices, identify hospitals at risk of low delivery volumes, and evaluate maternal care access under two key measures: (1) average travel distance to a birthing facility and (2) the proportion of deliveries occurring in predicted low-volume hospitals. The LC model that incorporated both facility attributes and patient characteristics outperformed other models across every evaluation metric.

The smaller subgroup of "less distance-sensitive" patients who placed high preference for facilities with higher level and specialized care exhibited characteristics consistent with patterns reported in prior studies. Our findings align with evidence that patients with higher socioeconomic status were more likely to bypass nearby facilities in favor of facilities perceived to provide higher quality care [7, 17, 63]. Interestingly, while this subgroup showed higher utilization of facilities with advanced levels of care, they were less likely to deliver at "Baby-Friendly" designated hospitals. Although Baby-Friendly designation is intended to promote breastfeeding and maternal-infant bonding, hospitals with such designation were relatively rare in Georgia during the study period (only 5 out of 77 facilities from 2016 to 2019) and may not have been co-located with facilities offering higher levels of maternal and neonatal care. For patients with multiple medical risk factors and needing more advanced care, Baby-Friendly facilities may not have met their clinical needs.

Our study adds to the literature by providing insights into hospital choices and OB bypassing. We identified factors associated with choosing a facility farther away and quantified the tradeoffs patients make between facility attributes and travel distance. These findings have important policy implications. First, the strong influence of socioeconomic status on patients' selection of birthing facilities emphasizes the need to design policy interventions that address patients' social determinants of health. Simply expanding high-quality maternal care will not ensure equitable access if systemic barriers, such as transportation limitations, restrictive insurance networks, and health literacy gaps remain unaddressed. Additionally, our results confirm the importance of risk-appropriate care where patients should seek care at a facility that is suited to

handle their risk. Matching patients to facilities with appropriate levels of care is critical to reducing adverse maternal outcomes, as high-risk deliveries at lower-level facilities are associated with poorer outcomes. However, availability of higher quality services cannot impact maternal outcomes if women do not choose to give birth in high-quality facilities. Policymakers should focus on mechanisms that facilitate appropriate matching between patient risk and facility capability, such as regionalized care systems, referral pathways, and patient navigation programs

Our proposed measure of access to care, based on the percentage of patients delivering at low-volume facilities, offers a different perspective on access to maternal care amid OB unit closures. While measures like travel distance to delivery facilities (such as Maternity Care Deserts or distance to CCO services) and others (such as regions with high maternal transport rate) identified similar at-risk regions in central east Georgia, these areas have a low percentage of patients delivering at low-volume facilities [15, 16, 64–66]. This highlights the trade-off between keeping more OB units open to reduce geographic barriers and the potential for lower OB volume due to spreading out service providers. Policymakers must weigh the benefits of proximity against the risks associated with low-volume care, particularly in rural regions. If distance were the main factor in patients' choice of delivery facilities, we would expect the number of low-volume facilities predicted by a distance-only model to match the actual number. However, the predicted number was lower when distance was the only factor considered. This confirms the risk of patients bypassing nearby facilities for more appealing ones, potentially leaving local facilities with low volumes.

Our discrete choice modeling framework provides a practical, data-driven tool for identifying facilities at risk of underutilization and informing policies to better distribute patient volume across the region. For example, a Level I facility in Northeast Georgia, which was only correctly identified as low-volume by both our MNL and LC models by taking the realized access perspective, later confirmed to have closed its OB unit in 2025. In contrast, the distance-only model which reflects potential rather than actual access, failed to identify this facility as underutilized. Our discrete choice models are

also highly adaptable and can be incorporated in other advanced models for future

policy analysis. One such use case is demonstrated in Meredith and Steimle 2025,

where a choice model is incorporated in an optimization framework to identify regions

that were vulnerable to closures [67].

There are opportunities to further improve the modeling approaches described in this

study. First, although LC model demonstrates superior predictive performance

compared to the distance-only and MNL models, our model underestimated failed to

capture a significant proportion of delivery volume at the facility with the highest annual

volume. Additionally, rurality of patients' residence was not chosen in our final LC model,

counter-intuitive to the belief that rural and urban patients would have different

preferences for healthcare facilities. We initially included this variable into LC model, but

this variable turned out to be insignificant or the model that included this variable had

worse fit and predictive performance. We suspect that the difference of facility choices

between urban and rural residents could be explained by other factors, such as distance

to the facility, patients' socioeconomic characteristics, and payers.

Our study has several limitations. First, Georgia birth records do not include deliveries

of out-of-state residents in Georgia facilities. This omission could underestimate delivery

volumes at facilities near state borders, although the proportion of out-of-state residents

delivering in Georgia facilities is expected to be small. Because our modeling population

included only Georgia residents, the absence of out-of-state patients from the dataset

should not bias our estimates of realized access to birthing facilities, as those patients

were not part of the modeling population. Future research can consider linking birth

records with data from the American Hospital Association to obtain accurate annual OB

volume by both in-state and out-of-state residents. Second, we used driving distance as

a measure of proximity from patients' residences to OB facilities. This approach could

underestimate barriers to OB access for those with low socioeconomic status, as they

might lack access to a car or reliable transportation. Additionally, our analysis is limited

to deliveries that occurred at OB facilities within 120 miles from the birthing person's

residence. Future research should examine non-facility births, deliveries occurring at

non-OB facilities, and births taking place far from the patient's residence, as these

cases may reveal additional barriers to accessing facility-based OB care. In particular,

studying patients who travel exceptionally long distances for specialized care could

provide insights into access challenges faced by individuals with complex medical

needs or limited local options for high-risk OB care. Lastly, our analysis relied on RP to

estimate patients' choices. This method does not allow us to differentiate between

active choices, where patients deliberately select a facility, and passive choices, where

external constraints such as availability or referral patterns dictate the decision.

In conclusion, patients placed heterogeneous weights on OB facility attributes. Patients

who have more severe medical conditions, have commercial insurance compared to

Medicaid, and are highly educated are more likely to be "less distance-sensitive" and

select facilities with higher levels of care more often. Incorporating this heterogeneity

into predictive models improves the identification of communities at risk, specifically,

areas where a high proportion of patients must travel long distances not by choice or

deliver at low-volume facilities. Efforts to improve maternal care access should go

beyond addressing geographic barriers alone. Effective policies aimed at enhancing

access should also consider broader social determinants, including differences in

insurance coverage and educational attainment, which affect patients' ability and

willingness to reach higher-quality care. Lastly, our study emphasizes the importance of

a realized access perspective when measuring access to care. Our model can serve as

a practical tool for identifying underutilized facilities and informing future policies that

better distribute patient volume across the region.

References

1. Sonenberg A, Mason DJ (2023) Maternity Care Deserts in the US. JAMA Health Forum 4:e225541. https://doi.org/10.1001/jamahealthforum.2022.5541

2. The Chartis Group (2023) Rural America's OB Deserts Widen in Fallout From Pandemic. The Chartis Group

3. Minion SC, Krans EE, Brooks MM, et al (2022) Association of Driving Distance to Maternity Hospitals and Maternal and Perinatal Outcomes. Obstet Gynecol 140:812–819. https://doi.org/10.1097/AOG.0000000000004960

4. Kozhimannil KB, Leonard SA, Handley SC, et al (2023) Obstetric Volume and Severe Maternal Morbidity Among Low-Risk and Higher-Risk Patients Giving Birth at Rural and Urban US Hospitals. JAMA Health Forum 4:e232110. https://doi.org/10.1001/jamahealthforum.2023.2110

5. Hung P, Kozhimannil KB, Casey MM, Moscovice IS (2016) Why Are Obstetric Units in Rural Hospitals Closing Their Doors? Health Serv Res. https://doi.org/10.1111/1475-6773.12441

6. Kozhimannil KB, Interrante JD, Fritz AH, Carroll C (2024) Information for Rural Stakeholders About Access to Maternity and Obstetric Care: A Community-Relevant Synthesis of Research

7. Carrel M, Keino BC, Novak NL, et al (2023) Bypassing of nearest labor & delivery unit is contingent on rurality, wealth, and race. Birth 50:5–10. https://doi.org/10.1111/birt.12712

8. Thorsen ML, Harris S, Palacios JF, et al (2023) American Indians travel great distances for obstetrical care: Examining rural and racial disparities. Soc Sci Med 325:115897. https://doi.org/10.1016/j.socscimed.2023.115897

9. Kozhimannil KB, Casey MM, Hung P, et al (2016) Location of childbirth for rural women: implications for maternal levels of care. Am J Obstet Gynecol 214:661.e1-661.e10. https://doi.org/10.1016/j.ajog.2015.11.030

10. Lorch SA, Rogowski J, Profit J, Phibbs CS (2021) Access to risk-appropriate hospital care and disparities in neonatal outcomes in racial/ethnic groups and rural–urban populations. Semin Perinatol 45:151409. https://doi.org/10.1016/j.semperi.2021.151409

11. Adams EK, Wright GE (1991) Hospital Choice of Medicare Beneficiaries in a Rural Market: Why Not the Closest? J Rural Health 7:134–152. https://doi.org/10.1111/j.1748-0361.1991.tb00715.x

12. Sanders SR, Erickson LD, Call VRA, et al (2015) Rural Health Care Bypass
Behavior: How Community and Spatial Characteristics Affect Primary Health Care
Selection. J Rural Health 31:146–156. https://doi.org/10.1111/jrh.12093

13. Johnson MB (2020) Prenatal Care for American Indian Women. MCN Am J Matern
Nurs 45:221. https://doi.org/10.1097/NMC.0000000000000633

14. Andersen R, Aday LA (1978) Access to Medical Care in the U.S.: Realized and
Potential. Med Care 16:533–546

15. Brantley MD, Davis NL, Goodman DA, et al (2017) Perinatal regionalization: a
geospatial view of perinatal critical care, United States, 2010-2013. Am J Obstet
Gynecol 216:185.e1-185.e10. https://doi.org/10.1016/j.ajog.2016.10.011

16. Kroelinger CD, Brantley MD, Fuller TR, et al (2021) Geographic access to critical
care obstetrics for women of reproductive age by race and ethnicity. Am J Obstet
Gynecol 224:304.e1-304.e11. https://doi.org/10.1016/j.ajog.2020.08.042

17. Tai W-TC, Porell FW, Adams EK (2004) Hospital Choice of Rural Medicare
Beneficiaries: Patient, Hospital Attributes, and the Patient–Physician Relationship.
Health Serv Res 39:1903–1922. https://doi.org/10.1111/j.1475-6773.2004.00324.x

18. Radcliff TA, Brasure M, Moscovice IS, Stensland JT (2003) Understanding Rural
Hospital Bypass Behavior. J Rural Health 19:252–259.
https://doi.org/10.1111/j.1748-0361.2003.tb00571.x

19. Basu J, Mobley LR (2007) Illness severity and propensity to travel along the urban-
rural continuum. Health Place 13:381–399.
https://doi.org/10.1016/j.healthplace.2006.03.002

20. Bronstein JM, Morrisey MA (1991) Bypassing rural hospitals for obstetrics care. J
Health Polit Policy Law 16:87–118. https://doi.org/10.1215/03616878-16-1-87

21. Bronstein JM, Morrisey MA (1990) Determinants of Rural Travel Distance for
Obstetrics Care. Med Care 28:853

22. Von Reichert C, McBroom WH, Reed FW, Wilson PB (1995) Access to health care
and travel for birthing: Native American-white differentials in Montana. Geoforum
26:297–308. https://doi.org/10.1016/0016-7185(95)00035-6

23. Goldsteen RL, Falcone DJ, Broyles RW, et al (1994) Local Factors Affecting the
Tendency to Bypass Local Hospitals for Inpatient Mental Health Care: An
Exploratory Analysis. J Rural Health 10:89–96. https://doi.org/10.1111/j.1748-
0361.1994.tb00215.x

24. Brathwaite T (2018) The Holy Trinity: Blending Statistics, Machine Learning and
Discrete Choice, with Applications to Strategic Bicycle Planning. UC Berkeley

25. Ben-Akiva M, Lerman SR (2018) Discrete Choice Analysis: Theory and Application
to Travel Demand. MIT Press, Cambridge, MA, USA

26. Lurkin V, Garrow LA, Higgins MJ, et al (2017) Accounting for price endogeneity in
airline itinerary choice models: An application to Continental U.S. markets. Transp
Res Part Policy Pract 100:228–246. https://doi.org/10.1016/j.tra.2017.04.007

27. Berry ST (1994) Estimating Discrete-Choice Models of Product Differentiation.
RAND J Econ 25:242–262. https://doi.org/10.2307/2555829

28. Berbeglia G, Garassino A, Vulcano G (2022) A Comparative Empirical Study of
Discrete Choice Models in Retail Operations. Manag Sci 68:4005–4023.
https://doi.org/10.1287/mnsc.2021.4069

29. Lagarde M, Blaauw D (2009) A review of the application and contribution of discrete
choice experiments to inform human resources policy interventions. Hum Resour
Health 7:62. https://doi.org/10.1186/1478-4491-7-62

30. Mandeville KL, Lagarde M, Hanson K (2014) The use of discrete choice
experiments to inform health workforce policy: a systematic review. BMC Health
Serv Res 14:367. https://doi.org/10.1186/1472-6963-14-367

31. Wang N, Tang L, Pan H (2017) Effectiveness of policy incentives on electric vehicle
acceptance in China: A discrete choice analysis. Transp Res Part Policy Pract
105:210–218. https://doi.org/10.1016/j.tra.2017.08.009

32. Wang R (2021) Discrete Choice Models and Applications in Operations
Management. In: Emerging Optimization Methods and Modeling Techniques with
Applications. INFORMS, pp 199–226

33. Train K (2009) Discrete Choice Methods With Simulation

34. Louviere JJ, Hensher DA, Swait JD (2000) Stated Choice Methods: Analysis and
Applications. Cambridge University Press, Cambridge

35. McFadden D (1974) Conditional logit analysis of qualitative choice behavior.
Academic Press, New York

36. Greene WH, Hensher DA (2003) A latent class model for discrete choice analysis:
contrasts with mixed logit. Transp Res Part B Methodol 37:681–698.
https://doi.org/10.1016/S0191-2615(02)00046-2

37. Gutacker N, Siciliani L, Moscelli G, Gravelle H (2016) Choice of hospital: Which
type of quality matters? J Health Econ 50:230–246.
https://doi.org/10.1016/j.jhealeco.2016.08.001

38. Smith H, Currie C, Chaiwuttisak P, Kyprianou A (2018) Patient choice modelling:
how do patients choose their hospitals? Health Care Manag Sci 21:259–268.
https://doi.org/10.1007/s10729-017-9399-1

39. Lendado TA, Bitew S, Elias F, et al (2022) Effect of hospital attributes on patient
preference among outpatient attendants in Wolaita Zone, Southern Ethiopia:
discrete choice experiment study. BMC Health Serv Res 22:661.
https://doi.org/10.1186/s12913-022-07874-x

40. Ranjbar M, Bazyar M, Pahlevanshamsi F, et al (2024) What do Iranians value most
when choosing a hospital? Evidence from a discrete choice experiment. PloS One
19:e0293241. https://doi.org/10.1371/journal.pone.0293241

41. Ferrari A, Seghieri C, Giannini A, et al (2023) Driving time drives the hospital
choice: choice models for pelvic organ prolapse surgery in Italy. Eur J Health Econ
HEPAC Health Econ Prev Care 24:1575–1586. https://doi.org/10.1007/s10198-
022-01563-6

42. Gao Y, Pan H, Xie Z, Habib KN (2023) Understanding patients heterogeneity in
healthcare travel and hospital choice - A latent class analysis with covariates. J
Transp Geogr 110:103608. https://doi.org/10.1016/j.jtrangeo.2023.103608

43. Nicolet A, Perraudin C, Krucien N, et al Preferences for innovations in healthcare
delivery models in the Swiss elderly population: a latent class, choice modelling
study

44. Yorlets RR, Iverson KR, Leslie HH, et al (2019) Latent class analysis of the social
determinants of health-seeking behaviour for delivery among pregnant women in
Malawi. BMJ Glob Health 4:. https://doi.org/10.1136/bmjgh-2018-000930

45. Pitkänen V, Linnosmaa I (2021) Choice, quality and patients' experience: evidence
from a Finnish physiotherapy service. Int J Health Econ Manag 21:229–245.
https://doi.org/10.1007/s10754-020-09293-z

46. Hwang K, Jang H, Lee T, et al (2017) A Choice Model for Estimating Realized
Accessibility: Case Study for Obstetrics Care in Korea. In: Cappanera P, Li J, Matta
A, et al (eds) Health Care Systems Engineering. Springer International Publishing,
Cham, pp 73–85

47. Jang H (2021) A model for measuring healthcare accessibility using the behavior of
demand: a conditional logit model-based floating catchment area method. BMC
Health Serv Res 21:660. https://doi.org/10.1186/s12913-021-06654-3

48. Harris S, McGarvey RG, Thorsen A, Thorsen M (2024) Inferred attractiveness
gravity-based models for estimating realized access at rural hospitals. J Oper Res
Soc 1–16. https://doi.org/10.1080/01605682.2024.2406236

49. Georgia Department of Public Health (2017) Core Requirements and
Recommended Guidelines for Designated Regional Perinatal Centers. Georgia
Department of Public Health

50. Victoor A, Delnoij DM, Friele RD, Rademakers JJ (2012) Determinants of patient
choice of healthcare providers: a scoping review. BMC Health Serv Res 12:272.
https://doi.org/10.1186/1472-6963-12-272

51. American College of Obstetricians and Gynecologists (2019) Levels of Maternal
Care: Obstetric Care Consensus No, 9. Obstet Gynecol 134:e41.
https://doi.org/10.1097/AOG.0000000000003383

52. Baby-Friendly USA. In: Baby-Friendly USA. https://www.babyfriendlyusa.org/.
Accessed 24 Oct 2025

53. Roeder K, Lynch KG, Nagin DS (1999) Modeling Uncertainty in Latent Class
Membership: A Case Study in Criminology. J Am Stat Assoc 94:766–776.
https://doi.org/10.1080/01621459.1999.10474179

54. Shen J (2009) Latent class model or mixed logit model? A comparison by transport
mode choice data. Appl Econ 41:2915–2924.
https://doi.org/10.1080/00036840801964633

55. Shuford EH, Albert A, Massengill HE (1966) Admissible Probability Measurement
Procedures. Psychometrika 31:125–145. https://doi.org/10.1007/BF02289503

56. Rahnasto I, Hollestelle M (2024) Comparing discrete choice and machine learning
models in predicting destination choice. Eur Transp Res Rev 16:43.
https://doi.org/10.1186/s12544-024-00667-9

57. Handley SC, Passarella M, Herrick HM, et al (2021) Birth Volume and Geographic
Distribution of US Hospitals With Obstetric Services From 2010 to 2018. JAMA
Netw Open 4:e2125373. https://doi.org/10.1001/jamanetworkopen.2021.25373

58. Thorsen ML, Harris S, McGarvey R, et al (2022) Evaluating disparities in access to
obstetric services for American Indian women across Montana. J Rural Health
38:151–160. https://doi.org/10.1111/jrh.12572

59. Rayburn WF, Richards ME, Elwell EC (2012) Drive Times to Hospitals With
Perinatal Care in the United States. Obstet Gynecol 119:611.
https://doi.org/10.1097/AOG.0b013e318242b4cb

60. Hoetker G (2007) The use of logit and probit models in strategic management
research: Critical issues. Strateg Manag J 28:331–343.
https://doi.org/10.1002/smj.582

61.   Hess S, Palma D (2019) *Apollo*: A flexible, powerful and customisable freeware
package for choice model estimation and application. J Choice Model 32:100170.
https://doi.org/10.1016/j.jocm.2019.100170

62.   Ortúzar J de D, Willumsen LG Modelling Transport, 4th Edition | Wiley, 4th ed

63.   Tayyari Dehbarez N, Raun Mørkbak M, Gyrd-Hansen D, et al (2018) Women's
Preferences for Birthing Hospital in Denmark: A Discrete Choice Experiment.
Patient - Patient-Centered Outcomes Res 11:613–624.
https://doi.org/10.1007/s40271-018-0313-9

64.   March of Dimes (2024) Nowhere to go: Maternity Care Deserts across the U.S. In:
March Dimes. https://www.marchofdimes.org/maternity-care-deserts-report.
Accessed 6 Sep 2024

65.   Li J, Radke SM, Steimle LN (2025) Examining perinatal regionalization in practice:
a network analysis of maternal transport in Georgia. BMC Health Serv Res 25:862.
https://doi.org/10.1186/s12913-025-13025-9

66.   Meredith ME, Radke SM, Steimle LN (2024) The implications of using maternity
care deserts to measure progress in access to obstetric care: a mixed-integer
optimization analysis. BMC Health Serv Res

67.   Meredith ME, Steimle LN Identifying Regions Vulnerable to Obstetric Unit Closures
using Facility Location Modeling with Patient Behavior

  Tables and Figures

  Figure 1. Data inclusion criteria.

(a) Class 1: "more distance-sensitive" group   (b) Class 2: "less distance-sensitive" group

893
894   Figure 2. Estimated results of the LC model. 2(a): Class 1: "more distance-sensitive" patient; 2(b): Class 2: "less distance-
895   sensitive" patient.

(a) Calibration plot by decile

(b) Calibration plot by decile – low volume (annual volume < 500)

Figure 3. Calibration plots by decile comparing true annual volume vs. predicted volume from the latent class model on the 2019 data: (a) shows calibration plot by decile for all facilities; (b) shows calibration plot by decile for predicted low-volume facilities (decile mean < 500). Each bar is positioned at its mean value for the corresponding decile. Green triangles indicate decile mean; orange lines indicate decile median; the red line represents y=x line (perfect calibration).

914   Table 1. Descriptive statistics of facilities

	Total (N = 79)
Average Annual Volume	
Mean (SD)	1550 (2020)
Median [Min, Max]	1010 [109, 15800]
Level of Care, N (%)	
Birth Center	2 (2.5)
1	26 (32.9)
2	26 (32.9)
3	18 (22.8)
RPC	7 (8.9)
RUCC Classification, N (%)	
Rural	9 (11.4)
Suburban	19 (24.1)
Urban	51 (64.6)

917 Table 2. Estimated results for the multinomial logit model

Variable	Coefficients	SE	95% CI
Road Distance	-0.111***	0.001	(-0.113, -0.109)
Level of Care (ref: Level I)[a]			
Birth center	-2.254***	0.243	(-2.729, -1.778)
Level II	0.751***	0.052	(0.649, 0.854)
Level III	1.463***	0.053	(1.359, 1.566)
RPC	1.005***	0.058	(0.891, 1.118)
Facility Location (ref: Non-Urban)			
Urban	0.662***	0.071	(0.523, 0.802)
Midwife Service (ref: No)			
Yes	0.170***	0.041	(0.090, 0.250)
Baby-Friendly Designation (ref: No)			
Yes	0.236***	0.040	(0.159, 0.314)
Multi-Facility Health Systems (ref: No)			
Yes	0.415***	0.035	(0.346, 0.485)

Model fit: log-likelihood = -17340.48; AIC =34698.95; BIC =34763.85; adjusted ρ^2 = 0.5088 (equally-likely base)

Abbreviations: SE, standard error; CI, confidence interval; ref, reference level; RPC: regional perinatal center.

a. Coefficients of reference level are held fixed at zero during fitting.

*: significant at p=0.05; **: significant at p=0.01; ***: significant at p=0.001

919   Table 3. Estimated results for the final latent class model

Variable	Class 1 "More Distance-Sensitive" Patient			Class 2 "Less Distance-Sensitive" Patient		
Class Probability	65.54			34.46		
Variable	**Coefficients**	**SE**	**95% CI**	**Coefficients**	**SE**	**95% CI**
Class Membership Model						
ASC	-	-	-	-1.998***	0.313	(-2.611, -1.385)
Number of Medical Risk Factors	-	-	-	0.243***	0.069	(0.107, 0.378)
Payer (ref: Medicaid)						
Tricare	-	-	-	-0.231	0.219	(-0.660, 0.199)
Commercial insurance	-	-	-	0.943***	0.097	(0.754, 1.133)
Self-pay	-	-	-	0.593**	0.188	(0.225, 0.962)
Other insurance, other government assistance	-	-	-	0.865***	0.200	(0.473, 1.256)
Education Level (ref: Less than 9th grade)						
9th through 11th grade	-	-	-	-0.220	0.316	(-0.839, 0.399)
High school diploma or GED	-	-	-	0.649*	0.278	(0.104, 1.194)
College or higher	-	-	-	1.045***	0.278	(0.500, 1.589)
Class-specific Choice Model						
Road Distance	-0.175***	0.006	(-0.187, -0.163)	-0.067***	0.003	(-0.072, -0.062)
Level of Care (ref: Level I)						
Birth center	-2.254	-	-	-2.254	-	-
Level II	0.922***	0.086	(0.753, 1.091)	1.164***	0.157	(0.857, 1.471)
Level III	1.180***	0.087	(1.010, 1.350)	3.035***	0.200	(2.642, 3.427)
RPC	0.555***	0.092	(0.374, 0.736)	3.155***	0.239	(2.687, 3.623)
Facility Location (ref: Non-Urban)						
Urban	0.983***	0.152	(0.685, 1.281)	-0.116	0.174	(-0.458, 0.225)
If Midwife Services Are Available (ref: no)						
Yes	0.227***	0.067	(0.096, 0.359)	0.036	0.118	(-0.195, 0.267)
Baby-Friendly Designation (ref: No)						
Yes	0.892***	0.080	(0.736, 1.048)	-1.395***	0.194	(-1.775, -1.014)
If Belongs to a Multi-Facility Health System (ref: No)						
Yes	0.175**	0.057	(0.064, 0.286)	1.169***	0.125	(0.925, 1.413)

Model fit: log-likelihood = -16818.4; AIC = 33686.81; BIC = 33867.07; adjusted ρ^2 = 0.5231 (equally-likely base)

Abbreviations: SE, standard error; CI, confidence interval; ref, reference level; RPC: regional perinatal center; ASC: alternative specific constant.

a. Coefficients of reference level were held fixed at zero during fitting.

b. Medical risk factors: pre-pregnancy diabetes, gestational diabetes, eclampsia, pre-pregnancy hypertension, gestational/pregnancy-associated hypertension, previous preterm birth, and previous Cesarean delivery.
 *: significant at p=0.05; **: significant at p=0.01; ***: significant at p=0.001

922 Table 4. Road distance equivalent of the fitted multinomial logit and latent class models

Variable	MNL RDE (in miles)	LC class 1 RDE (in miles)	LC class 2 RDE (in miles)
Level of Care (ref: Level I)			
Birth center	-20.39	-12.89	-33.74
Level II	6.79	5.27	17.43
Level III	13.21	6.74	45.43
RPC	9.09	3.17	47.24
Facility Location (ref: Non-Urban			
Urban	5.99	5.62	1.74
Midwife Service (ref: No)			
Yes	1.54	1.30	0.54
Baby-Friendly Designation (ref: No)			
Yes	2.14	5.10	20.88
Multi-facility Health System (ref: No)			
Yes	3.76	1.00	17.50
Abbreviations: RDE, road distance equivalent.			

924 Table 5. Class profiles of two latent classes

Variable	Class 1 ("More Distance-Sensitive": 65.54%)	Class 2 ("Less Distance-Sensitive": 34.46%)
Age in Years (mean, SD)	27.75 (5.907)	29.28 (5.659)
Race[a]		
White	0.554	0.591
Black or African-American	0.375	0.323
Asian	0.043	0.059
American Indian or Alaska Native	0.002	0.001
Native Hawaiian or other Pacific Islander	0.001	0.001
Multiracial	0.023	0.022
Unknown	0.004	0.004
Ethnicity		
Hispanic	0.155	0.117
Non-Hispanic	0.837	0.874
Unknown	0.008	0.009
Education Level*		
Less than 9th grade	0.039	0.016
9th through 11th grade	0.136	0.038
High school diploma or GED	0.333	0.238
College or higher	0.484	0.704
Unknown	0.009	0.004
Rurality of Residence[b]		
Urban	0.825	0.855
Suburban	0.135	0.111
Rural	0.040	0.035
Number of Medical Risk Factors*[c] **(mean, SD)**	0.285 (0.550)	0.354 (0.616)
Plurality[d] **(plurality =1, singleton=0)**	0.034	0.037
Payer*		
Tricare	0.046	0.025
Medicaid	0.544	0.300
Commercial insurance	0.310	0.574
Self-pay	0.062	0.052
Other insurance, other government assistance	0.037	0.049
Unknown	0.001	0.001

APNCU[e]		
Inadequate	0.170	0.124
Intermediate	0.069	0.060
Adequate	0.316	0.345
Adequate plus	0.382	0.417
Unknown	0.063	0.054
BMI Category[f]		
Underweight	0.035	0.030
Healthy	0.383	0.403
Overweight	0.267	0.265
Obese	0.301	0.288
Unknown	0.014	0.015

a. For categorial variables, we reported the percentage of each level in each group.
b. Rurality of residence: we categorized patients' residence by linking 2020 census block group to the Rural-Urban Continuum Codes (RUCCs): urban (RUCCs of 1-3), suburban (RUCCs of 4-6), and rural (RUCCs of 7-9).
c. Medical risk factors: pre-pregnancy diabetes, gestational diabetes, eclampsia, pre-pregnancy hypertension, gestational/pregnancy-associated hypertension, previous preterm birth, and previous Cesarean delivery.
e. The Adequacy of Prenatal Care Utilization Index. This index determines the adequacy of prenatal care utilization based on two parts: the month in which prenatal care is initiated and the number of visits from initiation of care until delivery. It is categorized into four categories: inadequate, intermediate, adequate, and adequate plus.
f. BMI category: Underweight (BMI <18.5), Normal (BMI 18.5-24.9), Overweight (BMI 25-29.9), Obese (BMI >=30)
* Variables included in the final class membership model.

927 Table 6. Choice prediction measured by logarithmic score

	Logarithmic Score
Distance-Only	-35.54
MNL	-1.77
LC	**-1.73**

930   Table 7. Predictive performance of low-volume facilities

	Accuracy	Specificity	Sensitivity	F1 score
Threshold = 250				
Distance-Only	88.31	1	0	0
MNL	90.91	**98.53**	33.33	46.15
LC	**92.21**	98.53	**44.44**	**57.14**
Threshold = 300				
Distance-Only	83.12	**96.92**	8.33	13.33
MNL	89.61	**96.92**	50	60
LC	**90.91**	96.92	**58.33**	**66.67**
Threshold = 400				
Distance-Only	84.42	**98.31**	38.89	53.85
MNL	90.91	96.61	**72.22**	78.79
LC	**92.21**	98.31	**72.22**	**81.25**
Threshold = 500[a]				
Distance-Only	81.82	93.10	47.37	56.25
MNL	92.21	**94.83**	84.21	84.21
LC	**93.51**	94.83	**89.47**	**87.18**

a. Low-volume obstetric facility is defined as an obstetric facility with an annual volume less than 500 (https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2784792). We also tried different thresholds as the cutoff for low-volume facility.

932 Table 8. Spearman correlation coefficients between observed and predicted metrics

Model	Average Travel Distance by Census Track	Low Volume by Census Track
Distance-Only	0.77	0.37
MNL	0.83	0.41
LC	**0.84**	**0.42**

933