# High-dose glucocorticoids, pain, and adverse events. Protocol for an analysis of a natural experiment of patients undergoing total hip or knee arthroplasty patients

Version 1, Tuesday, 11 November 2025. *Any changes made after publication of the protocol will be reported here. Any deviances from the protocol will be reported in the resulting manuscript.*

**Public protocol available at:** *MedRvix doi inserted here*

**Authors:**

Jens Laigaard[1,2,*], MD, https://orcid.org/0000-0001-5297-3769

Marcus Ølgaard Møller[3], MD, https://orcid.org/0009-0006-9637-1106

Søren Overgaard[1,2], MD, PhD, https://orcid.org/0000-0001-6829-4787

Ole Mathiesen[2,4], MD, PhD, https://orcid.org/0000-0003-4544-0619

Anders Peder Højer Karlsen[2,3], MD, PhD, https://orcid.org/0000-0002-0338-3751

**Affiliations:**

[1]Department of Orthopaedic Surgery and Traumatology, Copenhagen University Hospital Bispebjerg, Denmark

[2]Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

[3]Department of Anaesthesia and Intensive Care, Copenhagen University Hospital, Bispebjerg and Frederiksberg, Copenhagen, Denmark

[4]Centre for Anaesthesiological Research, Department of Anaesthesiology, Zealand University Hospital Koge, Koege, Denmark

*Corresponding author's contact information: Email: jens.holm.laigaard@regionh.dk, Phone +45 4261 7377, Address: 2. floor, Nielsine Nielsens vej 6, 2400 Copenhagen NV, Denmark

**Roles**: Jens Laigaard and Anders Peder Højer Karlsen drafted the protocol with inputs from Ole Mathiesen and Søren Overgaard. Jens Laigaard will handle the data, conduct and interpret analyses, supervised by Anders Peder Højer Karlsen. Jens Laigaard will draft the manuscript, which will be critically revised by all authors.

**Funding:** No external funding was received for this specific project.

**Conflict of interest**: No authors have any conflicts of interests to declare in relation to this project.

**Keywords**: Target trial, Causal inference, Quasi-experiment, Real-world data, Hip replacement, Knee replacement, Prosthetic joint infection, Surgical site infection, Dexamethasone, Methylprednisolone

## Abstract

**Introduction:** High-dose glucocorticoids are increasingly used in patients undergoing primary total hip arthroplasty (THA), unicompartmental knee arthroplasty (UKA), and total knee arthroplasty (TKA). However, the evidence supporting the use of high-dose glucocorticoids to manage acute postoperative pain is based mainly on randomised controlled trials, which may have limited external clinical validity due to strict exclusion criteria. In Denmark, the treatment was implemented stepwise, comprising a natural experiment.

**Objective**: To examine the real-world effect of a single dose of high-dose glucocorticoids on postoperative opioid consumption and adverse events in patients undergoing THA, UKA, and TKA.

**Methods:** This protocol describes a natural experiment study that will be reported as a target trial emulation, i.e. by attempting to mimic a randomised clinical trial.

Inclusion and exclusion criteria: All adults (>18 years) undergoing primary, elective THA, UKA, and TKA in the Capital Region and Region Zealand in Denmark are eligible. At each centre, patients who underwent surgery within one year before and one year after the implementation of high-dose glucocorticoids are screened for eligibility.

Intervention: The intervention is the ***intended use*** of a single high dose of glucocorticoids, administered after induction of anaesthesia, as part of the standard treatment.

Comparator: The comparator is no administration of glucocorticoids. Patients in the control arm may have received a lower dose of glucocorticoids for postsurgical nausea and vomiting

Assignment of intervention: For each hospital, patients who underwent surgery **before** the implementation of glucocorticoids serve as controls, while patients operated on **after** the implementation constitute the treatment arm.

Outcomes: The primary outcome is opioid consumption, expressed in morphine equivalent doses (MEQs) in the first 24 hours after surgery. Secondary outcomes are two *adverse effect-outcomes*: incidence of opioid-related adverse events and serious adverse events, and three *efficacy-outcomes:* length of postoperative observation area-stay, length of hospital stay, and worst pain intensity-score measured.

Data collection: This study uses data from the TRIPLE-A project ([www.triplea.dk](www.triplea.dk)), comprising validated electronic health record data.

Statistical analyses: The analyses will be based on a pre-defined model adjusted for important presurgical (and thus pre-intervention) variables using stabilised inverse probability of treatment weighting (SIPTW). A difference of 5 MEQs between treatment arms is considered clinically important.

**Knowledge dissemination**: The results will be shared at conferences and made publicly available.

**Registration**: *MedRvix doi inserted here*

**Introduction**

Total hip arthroplasty (THA), unicompartmental knee arthroplasty (UKA), and total knee arthroplasty (TKA) are common orthopaedic procedures in patients with osteoarthritis. Postoperative multimodal pain management is important for patient recovery and satisfaction. This includes high-dose glucocorticoids, which have been investigated for their ability to reduce inflammation and pain. After two randomised controlled trials (RCTs) showed benefit,[1,2] high-dose glucocorticoids have been implemented as routine treatment for THA, UKA, and TKA in Denmark.[3] However, due to concerns about adverse effects, e.g. increase risk of infection, the treatment was only step-wise implemented, as more evidence emerged.[4–7] Still, the efficacy and adverse event profile is primarily based on data from RCTs, often with strict exclusion criteria and missing data,[8,9] rather than 'real-world' evidence.[10]

In a parallel project,[11] we have mapped the time of implementation of high-dose glucocorticoids at Danish orthopaedic departments to evaluate if the treatment reduced long-term opioid use and affected the number of days alive and out of hospital. For that project, we are unable to determine if the patients did in fact receive the treatment. Moreover, we did not plan examine short-term outcomes because we were unable to include in-patient data from the electronic health record systems. Here, using data from the AI and Automation in Anaesthesia (TRIPLE-A) database, we can also investigate the effect of high-dose glucocorticoids on important short-term outcomes.[12]

We aim to examine the real-world effect of high-dose glucocorticoids on postoperative opioid consumption and adverse events in patients undergoing primary THA, UKA, and TKA.

**Methods**

This protocol describes the methods for a natural experiment study. The study will be reported following the target trial emulation framework.[10,13] The study was listed at the Capital region of Denmark's regional research listing (identifier p-2025-20115), the Ethical Committee of the Capital Region of Denmark waived ethical approval for this study (identifier F-25000312, 2 January 2025), and we were granted access to individual patient files approval from the Capitol Region Team for Medical Records Data (identifier R-25000339, 20 March 2025, with update for the specific subproject on 21 August 2025).

Study setting: The patients considered for this study underwent primary THA, UKA, or TKA surgery at public hospitals in the Capital Region and Region Zealand in Denmark from 2017 to 2025. We will include only hospitals that changed practice during this period, i.e. hospitals that implemented high-dose glucocorticoids for either THA, UKA or TKA. All residents in Denmark have equal access to free-of-charge healthcare financed by general taxes. Each year, around 10.000 patients undergo primary THA, UKA or TKA at public hospitals in the Capital Region and Region Zealand.[14,15] Enhanced Recovery After Surgery (ERAS) guidelines are widely implemented and half of the patients are discharged home the day after surgery. [14–16]

Inclusion and exclusion criteria: All adults (>18 years) patients undergoing primary, elective THA, UKA, and TKA in the Capital Region and Region Zealand in Denmark are eligible. Patients can only enter the study once. Hence, each subsequent THA, UKA or TKA surgery will not be considered for analysis.

At each centre, patients operated within one year before and one year after the implementation of high-dose glucocorticoids will be screened for eligibility. We choose to assess only this period to avoid unnecessary conflation with other changes to practice which could dilute or bias our results. However, we will adjust for calendar time, as well as important co-interventions, to account for possible time bias. Moreover, high-dose glucocorticoids were relatively abruptly implemented and there is no 'training' or washout period required for simple pharmacological interventions.

Intervention: The intervention the ***intended use*** of a single high dose of glucocorticoids, administered after induction of anaesthesia, as part of the standard treatment. A high-dose steroid dose was defined as IV/oral administration of ***at least*** 0.2 mg/kg or 16 mg dexamethasone/dexamethasone phosphate, 80 mg methylprednisolone, or 105 mg prednisolone (https://www.mdcalc.com/calc/2040/steroid-conversion-calculator).

The intervention was given along with other analgesic interventions that were routinely given to THA and TKA patients in Denmark during the study period, including paracetamol, NSAIDs, gabapentin, opioids, and, for TKA patients, local infiltration analgesia.

Comparator: The comparator is no administration of glucocorticoids during anaesthesia. Patients in the control arm may have received a lower dose of glucocorticoids for postsurgical nausea and vomiting. Typical anti-emetic doses are 4-8 mg dexamethasone.[17]

Data sources: For this project, we will use data available from the TRIPLE-A database.[12] The TRIPLE-A database include hundreds of variables on well over one million surgical procedures performed in the Capital Region and Region Zealand in Denmark from 2017 to 2025, extracted from the electronic health record system (EHR) Sundhedsplatformen (Epic Systems Corporation, WI, USA).

Assignment of intervention: For each hospital, patients who underwent surgery ***before*** implementation of glucocorticoids serve as controls, while patients operated ***after*** implementation constitute the treatment arm, regardless of their actual treatment.[18,19] Thus, treatment allocation is non-random and open label. Because certain patients (for example, those with insulin-dependent diabetes) may have been excluded from receiving glucocorticoids according to local protocols, we will report the proportion of patients in the treatment arm who did not receive the treatment (i.e. "protocol violations").

Outcomes: **Primary outcome:** Opioid consumption, expressed in morphine equivalent doses (MEQs) in the first 24 hours after surgery. We included opioid doses administered pre-emptively at the end of surgery, adjusting for time interval between administration and the actual end of surgery using the

drug-specific half-life.[12] We used the opioid conversion table that is implemented in the TRIPLE-A platform for conversions to MEQs.

**Secondary outcomes**: We chose five secondary outcomes:
- Worst postoperative pain intensity during 0-24 hours after surgery, measured with the 0-10 numerical rating scale (NRS).
- Proportion of patients with at least one opioid-related adverse event (ORADE) during 0-24 hours after surgery.
- Length of stay in the post-anaesthesia care unit (PACU), measured in minutes.
- Length of hospital stay, measured in hours from end of surgery to hospital discharge.
- Days alive and out of hospital at 30 days

Data collection: Please see the TRIPLE-A project website (currently www.cepra.nu/triple-a, until www.triplea.dk is launched in January 2026) for how each variable is assessed.[12] In brief, the data quality is dependent on the registrations in the electronic health record system. Administration of medications, surgeries, admissions, readmissions, and length of stay-variables are generally of very high quality, while health-care worker-dependent variables, such as pain scores, can be subject to errors and bias. For this project, we manually validated the treatment variable (administration of high-dose glucocorticoids) and important co-intervention variables (administration of paracetamol, NSAID, opioids) in 50 randomly selected patients without finding any errors.

Sample size: The sample size is based on convenience rather than power and the confidence intervals (CIs) will reflect the uncertainty of the estimate. However, using the CRTsize package in R,[20] we anticipate a power of 95% (5% false-negative rate) for detection of a 5 MEQ difference, assuming a standard deviation of 10 MEQ, 14 paired clusters (7 sites) with 50 patients, an intraclass coefficient of 0.1, and a 5% false-positive rate.

Statistical analysis plan: The data will be stored and analysed at a logged encrypted drive with the statistical software R (R Core Team, Vienna, Austria).[21] The dataset will be deleted after publication of the manuscript. We will check for the proportion of missing outcome data before conducting analysis. The analysis for a given outcome will not be conducted if the missingness is unbalanced. If the missingness is balanced between treatment arms, we will impute missing data using multiple

imputation with the mice package.[22] The number of patients contributing data to each analysis will be reported for each analysis.

The analyses will be based on a pre-defined model adjusted for important presurgical (and thus pre-intervention) variables using stabilised inverse probability of treatment weighting (SIPTW).[23,24]

First, we will apply a logistic mixed-effects model, using the lme4 package[25], to generate propensity scores[26] using available variables that may be linked to the assessed outcomes. However, only variables and interventions that are assessed before the end of anaesthesia are used, as post-anaesthetic variables influence by the intervention. The variables included in the model will include:

- Department (categorical variable), included as a random effect
- Calendar time (continuous variable), included as a fixed effect
- Age (continuous variable), included as a fixed effect
- Sex (binary variable), included as a fixed effect
- Type of surgery (binary variable: THA/TKA), included as a fixed effect
- Type of Anaesthesia (binary variable: General/Spinal), included as a fixed effect
- Paracetamol, NSAID and/or gabapentin as premedication or given intraoperatively (continuous variable with 0-3 medications), included as a fixed effect
- Local infiltration analgesia (binary variable), included as a fixed effect
- Preoperative chronic opioid use (binary variable), included as a fixed effect
- Number of psychiatric diagnoses on admission (continuous variable), included as a fixed effect

Each patient is given weight by the inverse probability of being in their respective arm (treatment/control). That is, patients in the treatment arm are assigned weights as (propensity score)$^{-1}$, while controls are assigned (1 - propensity score)$^{-1}$. Hence, patients in the treatment arm with low propensity scores and controls with high propensity scores are given more weight. Finally, these weights are 'stabilised' by multiplying each unit's weight by the proportion of units in their treatment group.[24]

To enhance interpretability, these weights are used to calculate the average marginal estimates with the *marginaleffects* package in R.[27] Binary outcomes are reported as mean relative and absolute risk differences with confidence intervals, while continuous outcomes are reported as mean differences with confidence intervals. The minimal important difference is considered a 5 MEQ absolute

difference in opioid consumption within the first 24 hours after surgery.[28] For the primary outcome, a P-value below 0.05 is considered statistically significant and the outcome will be reported with 95% confidence interval. For secondary analyses, we will apply Bonferroni correction to manage the risk of false-positive, i.e P-values below 0.05/5 = 0.01 are considered statistically significant. Secondary outcomes are reported with 99% confidence intervals.

We will conduct sensitivity analyses, excluding patient groups that may not have received the intervention immediately after implementation due to specific concerns, e.g. concerns over spikes in blood glucose in patients with diabetes.

Tables and Figures: We will present a table of demographic information, intraoperative, and perioperative details for both groups, including – but not limited to – age, sex, ASA-scores, BMI, type of surgery, type anaesthesia, propofol and remifentanil infusion rates, and the proportion of patients treated with high-dose glucocorticoids, paracetamol, NSAID and LIA.

To visualise and examine the findings and underlying trends, we will use statistical process control in control charts with monthly data relative to the time of implementation. We will chart all assessed outcomes, the proportion of patients who are administered high-dose glucocorticoid, and the proportion of patients who are administered important co-interventions, such as pre- or intraoperative paracetamol, NSAIDs, LIA, and spinal anaesthesia. We will apply the Anhoej rules ('unusually' long runs and 'unusually' few crossings of the median) and Shewhart's 3-sigma rule (values outside +/- 3 SDs) to assess for non-random patterns in the distribution of data points. For continuous measures, we will plot the monthly mean (I-chart). For binary measures, we will plot the monthly proportion of patients (P-chart).

The outcome estimates will be presented in text, tables, forest plots, or similar, whichever we find most appropriate.

Data sharing: We refer to the TRIPLE-A project group regarding data availability.[12] The statistical code can be obtained from the corresponding author upon reasonable request.

Knowledge dissemination: The results will be shared at conferences and made publicly available either through open-access publication or through publication at a preprint server, e.g. MedRvix.org.

References

1. Lunn, T. H. *et al.* Effect of high-dose preoperative methylprednisolone on pain and recovery after total knee arthroplasty: a randomized, placebo-controlled trial. *Br. J. Anaesth.* **106**, 230–238 (2011).

2. Lunn, T. H. *et al.* Effect of high-dose preoperative methylprednisolone on recovery after total hip arthroplasty: a randomized, double-blind, placebo-controlled trial. *Br. J. Anaesth.* **110**, 66–73 (2013).

3. Jorgensen, C. C., Pitter, F. T., Kehlet, H., Lundbeck Foundation Center for Fast-track, H. & Knee Replacement Collaborative, G. Safety aspects of preoperative high-dose glucocorticoid in primary total knee replacement. *Br J Anaesth* **119**, 267–275 (2017).

4. Køppen, K. S. *et al.* Systemic glucocorticoids as an adjunct to treatment of postoperative pain after total hip and knee arthroplasty: A systematic review with meta-analysis and trial sequential analysis. *Eur. J. Anaesthesiol.* **40**, 155–170 (2023).

5. Corcoran, T. B. *et al.* Dexamethasone and Surgical-Site Infection. *N Engl J Med* **384**, 1731–1741 (2021).

6. Steiness, J. *et al.* Non-opioid analgesic combinations following total hip arthroplasty (RECIPE): a randomised, placebo-controlled, blinded, multicentre trial. *Lancet Rheumatol.* **6**, e205–e215 (2024).

7. Gasbjerg, K. S. *et al.* Effect of dexamethasone as an analgesic adjuvant to multimodal pain treatment after total knee arthroplasty: randomised clinical trial. *BMJ* **376**, e067325 (2022).

8. Pedersen, C. *et al.* Differences in patient characteristics and external validity of randomized clinical trials on pain management following total hip and knee arthroplasty: a  systematic review. *Reg. Anesth. Pain Med.* **45**, 709–715 (2020).

9. Rønsbo, T. N., Laigaard, J., Pedersen, C., Mathiesen, O. & Karlsen, A. P. H. Adherence to participant flow diagrams in trials on postoperative pain management after total hip and knee arthroplasty: a methodological review. *Trials* **22**, 280 (2021).

10. Hernan, M. A. & Robins, J. M. *Causal Inference: What If*. (CRC Press, 2023).

11.      Laigaard, J. *et al.* Effect of high-dose glucocorticoids on persistent opioid use 3 to 12 months after primary total hip or knee arthroplasty. Protocol for a target trial emulation using observational data from Danish registries. *medRxiv* 2023.10.31.23297517 (2023) doi:10.1101/2023.10.31.23297517.

12.      Karlsen, A. P. H. *et al.* The AI and Automation in Anaesthesia (TRIPLE-A) Perioperative Database in Eastern Denmark 2017–: A Platform for Epidemiology, AI-Driven Prediction, Quality Control and Automated Data Retrieval. *Acta Anaesthesiol. Scand.* **69**, e70116 (2025).

13.      Cashin, A. G. *et al.* Transparent Reporting of Observational Studies Emulating a Target Trial-The TARGET Statement. *JAMA* **334**, 1084–1093 (2025).

14.      Søren Overgaard *et al.* Danish Hip Arthroplasty Register, National Annual Report for 2021. *RKKP* (2022).

15.      Martin Lindberg-Larsen *et al.* Danish Knee Arthroplasty Register, Annual Report 2022. *RKKP* (2023).

16.      Jensen, C. B. *et al.* 10-year evolution of day-case hip and knee arthroplasty: a Danish nationwide register study of 166,833 procedures from 2010 to 2020. *Acta Orthop.* **94**, 178–184 (2023).

17.      Weibel, S. *et al.* Drugs for preventing postoperative nausea and vomiting in adults after general anaesthesia: a network meta‐analysis. *Cochrane Database Syst. Rev.* https://doi.org/10.1002/14651858.CD012859.pub2 (2020) doi:10.1002/14651858.CD012859.pub2.

18.      Moher, D. *et al.* CONSORT 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *Int J Surg* **10**, 28–55 (2012).

19.      Hemming, K., Haines, T. P., Chilton, P. J., Girling, A. J. & Lilford, R. J. The stepped wedge cluster randomised trial: rationale, design, analysis, and reporting. *BMJ* **350**, h391 (2015).

20.      Rotondi, M. A. CRTSize: Sample Size Estimation Functions for Cluster Randomized Trials. (2023).

21.        R Core Team (2023). R: A language and environment for statistical computing. R Foundation for Statistical Computing.

22.        Buuren, S. van *et al.* mice: Multivariate Imputation by Chained Equations. (2025).

23.        Rosenberger, D. C. & Pogatzki-Zahn, E. M. Chronic post-surgical pain – update on incidence, risk factors and preventive treatment options. *BJA Educ.* **22**, 190–196 (2022).

24.        Austin, P. C. & Stuart, E. A. Moving towards best practice when using inverse probability of treatment weighting (IPTW) using the propensity score to estimate causal treatment effects in observational studies. *Stat. Med.* **34**, 3661–3679 (2015).

25.        Bates, D. *et al.* lme4: Linear Mixed-Effects Models using 'Eigen' and S4. (2023).

26.        ROSENBAUM, P. R. & RUBIN, D. B. The central role of the propensity score in observational studies for causal effects. *Biometrika* **70**, 41–55 (1983).

27.        Arel-Bundock [aut, V. *et al.* marginaleffects: Predictions, Comparisons, Slopes, Marginal Means, and Hypothesis Tests. (2023).

28.        Cook, J. A. *et al.* Specifying the target difference in the primary outcome for a randomised controlled trial: guidance for researchers. *Trials* **16**, 12 (2015).