1 **Primary Findings of Nationwide Home-based Test to Treat Program for COVID-19**
2 **and Influenza**
3

4 Apurv Soni on behalf of Home Test to Treat Investigators
5
6
7 Corresponding Author:
8 Apurv Soni, MD, PhD – Program in Digital Medicine, UMass Chan Medical School, 55
9 Lake Avenue N, S6-757, Worcester, MA 01655 (Email: apurv.soni@umassmed.edu)

10
11

The authors thank the numerous partners of the Home Test-to-Treat program whose
contributions were critical to the successful implementation of the program but did not
rise to the level of co-authorship for the evaluation manuscript. In particular, we
acknowledge Michael Wolfson, PhD (NIH/NIBIB) and Bruce J. Tromberg, PhD
(NIH/NIBIB) for conceptual support and program guidance; Eric Gogstad (CDC) for
expertise in program design and providing influenza antivirals from CDC; Dina Pasricha
Passman, MD, Lisa Tung, MPH, and Chris Crabtree, MPH (ASPR) for conceptual
design and implementation support; Michael Mina, MD, PhD (eMed) and the eMed team
for modifying the digital platform and implementing the care model; VentureWell for
primary program coordination and contracting. and the community engagement teams
contracted by VentureWell for coordinating outreach to over 5,000 community
organizations. We also thank all the telehealth clinicians, pharmacists, and staff who
made the HTTT program possible. Most of all, we are grateful to the HTTT program
enrollees for their trust and participation.
26
Funding: This project has been funded in whole or in part with Federal funds from the
National Institute of Biomedical Imaging and Bioengineering (NIBIB), National Institutes
of Health, Department of Health and Human Services, under Contract No.
75N92022D00010. In addition, the Center for Disease Control (CDC) and the
Administration for Strategic Preparedness and Response (ASPR) provided resources
for antiviral treatments and over-the-counter tests. The content is solely the
responsibility of the authors and does not necessarily represent the official views of the
National Institutes of Health.

Abstract

Importance: Timely initiation of antivirals for SARS CoV 2 and influenza depends on rapid diagnosis and access to prescribing. A national home based test to treat service was implemented to address these barriers.

Objective: To evaluate reach, use, and timeliness of prescribing in the Home Test to Treat program.

Design: Observational cohort analysis of program data from August 15, 2023 through April 17, 2024. Multivariable models assessed factors associated with test kit receipt, telehealth use and timing from symptom onset, antiviral prescribing and timing, and modality of medication fulfillment.

Setting: Nationwide, remote program providing free at home tests, on demand telehealth, and oral antivirals with pharmacy pickup or home delivery.

Participants: 66,169 adults who enrolled in the program and were eligible for analysis. Participation pathways included 1) proactive testing before symptoms, 2) on demand testing after symptom onset without a diagnosis, and 3) treatment only after a positive test.

Exposures: Program pathway and participant characteristics, including insurance status and prior difficulties accessing care.

Main Outcomes and Measures: Receipt of test kits, telehealth use and timing from symptom onset, and prescription of guideline directed oral antivirals within 1 day and within 5 days of symptom onset; choice of home delivery versus local pharmacy pickup.

Results: Enrollees represented all 50 states and 891 of 896 three-digit ZIP Code areas. Proactive testers comprised 52.6%; 15.0% of the analytical sample reported a positive result. Among positives, 80.7% used the treatment only pathway; 83.2 percent chose telehealth and 76.3% of consults occurred outside local business hours. Antivirals were prescribed to 6,821 enrollees; 59.8% received oral antivirals within 1 day and 92.8% within 5 days of symptom onset. Prescribing within 1 day did not differ notably by age, insurance, race or ethnicity, or prior difficulty accessing care. Of those prescribed, 1,092 enrollees, 16.0%, selected home delivery rather than pharmacy pickup. Home delivery was more common among enrollees reporting more reasons for delaying care, difficulty accessing care, lacking insurance, or enrolled in Medicaid ($P < .05$).

Conclusions and Relevance: Home Test to Treat delivered timely prescribing, including for adults with prior access barriers, by combining home based testing, on demand telehealth, and options for medication delivery across the country.

Key Points

Question: Does a national home test-to-treat program deliver timely antiviral treatment for COVID-19 and influenza across diverse adults?

Findings: Among 66,169 enrollees in all 50 states, 83% of positives used telehealth; 82% received antivirals; 60% were treated within 1 day and 93% within 5 days of symptoms; most visits occurred outside business hours; timeliness did not differ meaningfully by age, insurance, or race/ethnicity once enrolled.

Meaning: An integrated home testing and on-demand telehealth pathway achieved rapid, equitable treatment at national scale and provides a practical model to strengthen access and preparedness.

1 **Introduction**

2 The COVID-19 pandemic exposed significant gaps in healthcare access, particularly in

3 timely testing and treatment for at-risk and vulnerable populations.[1] Early intervention

4 with oral antivirals can prevent severe, adverse outcomes, yet differences in access to

5 care were identified.[2–7] Individuals from rural areas, racial and ethnic minorities, and

6 vulnerable communities face reduced access to testing and treatment. Non-Hispanic

7 Black and Hispanic patients received outpatient antivirals at substantially lower rates

8 (approximately 30% less) than non-Hispanic White patients.[1,8] Further, less than a

9 quarter of Medicaid recipients were prescribed oral antivirals during an infection. These

10 findings were persistent despite wide availability in oral antivirals and clear guidelines

11 for indications by the Center for Disease control. The gaps in guideline-directed medical

12 therapy are driven by structural barriers in current public health infrastructure that affect

13 timely recognition of the disease, access to a healthcare provider, assessment of

14 treatment eligibility, and provisioning of treatment. Commonly, the cascade of care

15 requires interaction with different entities: laboratory, clinician, and pharmacists and

16 attrition can emerge at multiple steps.

17 Emerging care models that co-locate and integrate services promise to overcome the

18 challenges in care continuity. The federal government launched test and treat centers

19 for SARS-CoV-2 as one-stop shop but an estimated 59% of rural residents lived over 60

20 minutes from the nearest in-person Test-to-Treat site.[9] The site-based test-to-treat

21 programs improved access to some extent, especially in urban centers, individuals in

22 rural and high–social vulnerability index (SVI) areas experienced delays in care.[2] The

23 emergence of home-testing, telehealth, and e-prescriptions can allow for decentralized

testing and treatment capabilities. However, programs have focused predominantly on

either testing access or antiviral distribution in isolation, without examining the full

continuum from diagnosis to treatment.[10–15] The effectiveness of such a comprehensive

home-based test-to-treat approach remains unknown. Potentially, this care model could

mitigate known disparities involving the elderly, uninsured, and people of color.[16]

In this study, we evaluate the nationwide Home Test to Treat (HTTT) program, which

integrated at-home COVID and influenza testing, telehealth consultations, and

medication delivery for COVID-19 and influenza at no cost to the enrollees. The pilot

program was designed to address gaps in care for underserved populations through a

decentralized model, providing timely diagnosis and treatment via telehealth without in-

person visits. By analyzing enrollment patterns, testing outcomes, telehealth uptake,

and treatment data, we describe the program's reach and effectiveness for providing

timely guideline directed medical treatment.

Methods

Study Design

We conducted a convenience-sampling cohort study of enrollees in the Home Test to

Treat (HTTT) program. This national initiative provided at-home COVID-19 and

influenza testing, telehealth consultations, and direct medication delivery. Initially

conceptualized by a White House task force, HTTT was federally supported by the

National Institutes of Health, the Administration for Strategic Preparedness and

Response, and the Centers for Disease Control and Prevention.[17,18] VentureWell, LLC

served as the prime contractor and issued a call to coalesce a unique partnership

among governmental, non-profit, academic, and private entities. VentureWell contracted

UMass Chan Medical School as the research/evaluation partner and eMed, LLC as the

technology and implementation partner. Full program details, including pilot phases, are

described elsewhere. This report evaluates the main program implementation from

August 15, 2023, to April 17, 2024. The study received full ethical approval from the

Western Copernicus Group Institutional Review Board (Protocol # 1354231).

Participants

Individuals U.S. could voluntarily enroll via the program website (www.test2treat.org) or

by calling the program hotline (Figure 1). Program enrollment was available to adults

aged 18 or older in all 50 U.S. states, the District of Columbia, Puerto Rico, and the U.S.

Virgin Islands. Enrollees were classified in one of three pathways: (1) **Proactive test**

and treat users: those who enrolled in advance without current symptoms or known

infection; (2) **On-demand test and treat users**: individuals who enrolled to request

over-the-counter test kits from the program and potential treatment after onset of

respiratory symptoms; or (3) **On-demand telehealth users**: individuals who were

already COVID-19 or flu positive and enrolled to request telehealth and treatment.

(Figure 1) social media, community networks, word of mouth, and planned

programmatic press releases were used to spread awareness about the program

(Supplementary Figure 1).

Data Collection

<u>Data Sources</u>

All data were collected via the HTTT program's digital platform operated by eMed.

Platform metadata provided timestamps for test kit orders, test results reporting,

telehealth consultations, and medication prescription. The Health Professional Shortage

Area (HPSA) score was obtained from Health Resources and Services Administration

(www.data.hrsa.gov).

<u>Measures</u>

Upon enrollment, individuals completed a questionnaire on demographics (age, gender,

race/ethnicity), recent COVID-19 or flu symptoms or test results, chronic conditions,

insurance status, testing and care options, and how they heard about the program.

Difficulty in accessing routine clinical care outside of the program was assessed by

asking, "If you needed a medical visit tomorrow, how confident are you that you could

get it?" Responses on a 4-point Likert scale were dichotomized to "No" (not at all or a

little confident) vs. "Yes" (somewhat or very confident). Enrollees were also asked if, in

the past year, they had delayed medical care due to limited provider availability,

concerns about cost, lack of transportation, or inability to take time off; each reason was

recorded as a yes/no response. Primary outcomes were related to treatment provision:

(1) the proportion of enrollees who tested positive and received a telehealth

consultation, and (2) the proportion of telehealth users who were prescribed an oral

antiviral. Secondary outcomes focused on timeliness and access: (1) the proportion of

telehealth consultations that began during regular local business hours (9 AM–5 PM

weekdays for local time); (4) time from symptom onset to medication prescription; (5)

the proportion of enrollees receiving a prescription within 1 and 5 days of symptom

onset; and (6) the proportion of enrollees who received medication via home delivery

(versus local pharmacy pickup). In the rare event that an enrollee had distinct episodes

of positive test results, telehealth encounters, or prescriptions, only the first instance

was considered.

Statistical Analysis

Descriptive statistics were calculated to summarize participant demographics and

outcome distributions. Chi-square and ANOVA tests compared differences between the

three enrolled groups (proactive HTTT users, on-demand HTTT users, on-demand

telehealth seekers) for categorical and continuous variables respectively. Multivariable

logistic regression assessed the association between participant characteristics (e.g.,

race, insurance status, age, gender) and key outcomes such as receipt of telehealth

services and timely medication delivery. Adjusted odds ratios (aORs) and 95%

confidence intervals (CIs) were reported for associations. All statistical analyses were

conducted using R 4.3.3. Significance level was 0.05 (two-sided).

Results

Of 104,280 individuals who started enrollment process, 80,783 (77.5%) were eligible

and completed enrollment from 891 out of 896 zip-code 3 regions in the US (Figure 2).

Among those, 14,704 (18.2%) reported a recent positive SARS CoV-2 or influenza test

at entry but demonstrated suspicious behavior i.e. only were interested in a research

substudy instead of seeking telehealth or treatment. Their data were excluded. The

remaining 66,169 enrollees comprised the analytic cohort for this study (Figure 3). More

than half of the enrollees had public insurance (Medicare: 31.4%, Medicaid 13.1%) or were uninsured (13.9%), reflecting the program's emphasis on at-risk and vulnerable population (Table 1). Nearly half of the cohort (44.7%) self-reported lacking confidence in getting needed care within 24 hours outside of the program and more than half had one or more reason to delay healthcare despite a need due to a barrier. Overall, 52.6% enrolled proactively,15.2% enrolled as on-demand test-and-treat users, and the rest (32.2%) were on-demand telehealth seekers. Key differences emerged between the groups: 56.0% of on-demand test-and-treat users, 46.1% of proactive users, and 37.1% of on-demand telehealth seekers indicated difficulty accessing timely care outside of the program. Proactive test and treat users skewed older (38.6% were age ≥65). The on-demand treatment group had a higher proportion of males than the proactive group (39.9% vs. 26.8% respectively). Enrollees who learned of the program via community outreach or friends/family were more likely to enroll as on-demand treatment seekers.

A total of 9,940 enrollees (15.0%) reported a positive COVID-19 or flu test. Across all enrollment pathways, 83.2% of those who tested positive proceeded to receive a telehealth consultation through HTTT. Only 23.7% of these telehealth visits began during regular weekday hours (9 AM–5 PM, local time). Among telehealth users, 82.4% were prescribed an antiviral medication. Overall, 59.8% of enrollees who were given a prescription received it within 1 day of symptom onset, and 92.8% received it within 5 days (Table 2).

In multivariable adjusted analyses, we observed only limited differences based on participant and regional characteristics. Enrollees reporting difficulty accessing care were more likely to utilize telehealth when positive (aOR 1.35, 95% CI 1.20–1.51). In

contrast, older adults (aOR 0.81, 95% CI 0.65–1.00) and Medicare beneficiaries

(aOR 0.71, 95% CI 0.58–0.88) were less likely to opt for a telehealth consultation after

testing positive. Black enrollees and those with access barriers had higher telehealth

uptake, whereas seniors and Medicare-insured enrollees had lower uptake (Figure 3).

Among telehealth users, men had higher odds of being prescribed an antiviral treatment

(aOR 1.29, 95% CI 1.13–1.48). Conversely, both Medicare beneficiaries (aOR 0.69,

95% CI 0.55–0.88) and Hispanic enrollees (aOR 0.75, 95% CI 0.62–0.91) were less

likely to receive an antiviral prescription during their telehealth encounter (Figure 4a).

However, there were no meaningful differences observed among antiviral prescription

rates within one day of symptom of onset (Figure 4b).

Although few enrollees chose home delivery of medication (16.0% overall), this option

was more commonly utilized by those facing access barriers. Enrollees who reported

difficulty accessing care (aOR 1.27, 95% CI 1.09–1.47), who had any recent delays in

seeking care (aOR 1.31, 95% CI 1.21–1.42), or who were uninsured (aOR 1.29, 95% CI

1.08–1.53) or enrolled in Medicaid (aOR 1.63, 95% CI 1.32–2.01) had higher odds of

opting for mail-order medication delivery rather than local pharmacy pickup (Figure 5).

Discussion

The Home Test-to-Treat (HTTT) program achieved high engagement and rapid antiviral

treatment across a large, diverse population of adults. Over 83% of enrollees who

tested positive for COVID-19 or flu proceeded to use the program's telehealth service,

and more than 80% of those were prescribed guideline-recommended antivirals.

Consequently, a majority of treated patients began antiviral therapy promptly after

symptom onset: 59.8% started medication within one day of symptom onset and over

92% within five days. Most telehealth consultations took place during evenings or

weekends (less than a quarter occurred during weekday business hours), underscoring

that HTTT successfully delivered care at times when traditional clinics are often

unavailable. Together, these results demonstrate that a decentralized, at-home test-to-

treat model can achieve prompt, widespread uptake of telehealth and same-day

treatment initiation in practice.

The HTTT program's treatment reach far exceeded what has been observed in

traditional healthcare settings. Only about 24% of high-risk Veterans with COVID-19

infections received antiviral treatment.[4,19] Other studies similarly reported antiviral

uptake well under 35% in eligible patients.[20] In contrast, HTTT prescribed oral antivirals

to over 80% of infected enrollees, illustrating the impact of integrating services. The self-

selected enrollees likely represent a more activated group of individuals. However, a

study among predominantly vaccinated and activated patients, found that less than 20%

of individuals who test positive received oral antivirals. Thus, the program's one-stop

approach: combining at-home testing, on-demand telehealth prescribing, and pharmacy

or home delivery likely minimized the drop-off that occurs at multiple steps in usual care

and increased adherence to guideline directed medical therapy as well as sped up the

care cascade. This is critical because early antiviral therapy dramatically improves

outcomes: for example, starting medication within 24–48 hours of symptoms can

substantially reduce the risk of severe illness.[21–26] Indeed, HTTT often enabled same-

day diagnosis and treatment, an achievement that likely prevented a number of

hospitalizations or complications.

A major finding is that HTTT provided equitable care on many key metrics. We observed

no significant differences by race/ethnicity, insurance status, or prior access to care in

the likelihood of receiving timely treatment once enrolled. By removing cost and

transportation barriers and offering services remotely, HTTT appears to have narrowed

or eliminated many of these disparities. In fact, the program reached groups that often

struggle with access: enrollees who reported difficulty accessing healthcare were more

likely to use telehealth when they became positive (aOR 1.35), suggesting HTTT

successfully engaged those with unmet needs. At the same time, some differences

persisted, highlighting areas for improvement. Older adults and Medicare beneficiaries

were somewhat less likely to initiate a telehealth visit when positive, reflecting known

digital access and literacy challenges among seniors. Additionally, among those who did

use telehealth, antiviral prescribing rates were slightly lower for Hispanic patients and

Medicare-insured patients. These residual gaps could stem from language or

communication barriers, hesitancy to accept treatment, or potential provider biases in

virtual care. Importantly, because HTTT removed many structural obstacles (free tests,

free telehealth, no travel needed), the remaining disparities likely point to nuanced

challenges such as technology comfort and cultural concordance. To fully close the

remaining gaps, future programs could incorporate enhancements: for example,

providing digital literacy support or phone-based options for older adults, and offering

culturally and linguistically tailored outreach during telehealth encounters for minority

patients. Such targeted improvements would build on HTTT's strengths so that no

subgroup is left behind in the test-to-treat process.

HTTT provides a proof of concept for decentralized care when the diagnostic and treatment cascade is clear, time sensitive, and protocolized. COVID-19 and influenza are appropriate use cases because accurate home tests are widely available, eligibility criteria for antivirals are standardized, and benefit depends on rapid initiation. Within this context, HTTT linked at-home testing, on-demand telehealth assessment, and swift medication fulfillment, achieving high telehealth uptake and timely prescribing at national scale, often outside business hours. Across many demographic categories the program delivered comparable timeliness once patients engaged the service, consistent with the view that structural access interventions can reduce inequities that persist in routine care. To translate this operational success to the broader crisis in American healthcare, similar services should be integrated with, rather than isolated from, existing systems through bidirectional exchange with EHRs and health information exchanges, automatic notification to usual clinicians, shared escalation pathways, closed-loop pharmacy fulfillment, and common quality and safety measures.

This study also had several limitations. First, as enrollees self-selected into the HTTT program (through convenience sampling), they likely differ from the general population in ways that could overestimate the feasibility and success of home-based interventions. Enrollees may have better digital access or health motivation than the general population. Second, key timeline data (symptom onset and test dates) were self-reported, introducing potential recall bias. This limits the accuracy of testing and treatment time periods. Third, the program was delivered via a single telehealth platform

and pharmacy system, which may limit generalizability to other settings with different

technologies or workflows. Fourth, we lacked detailed data on individuals who enrolled

but did not utilize telehealth or did not report any positive test. Thus, we could not fully

examine why some enrollees did not progress through the test-to-treat pathway (such

as asymptomatic courses, competing priorities, or distrust).

In summary, HTTT offers an operational blueprint for rapid, equitable treatment at scale.

However, this opportunity must be clear-eyed in the the objective to strengthen primary

care capacity: routing straightforward, time-sensitive episodes through a coordinated

decentralized pathway can free primary care teams to focus on longitudinal

management and diagnostic complexity while extending after-hours coverage. Payment

approaches should recognize triage, coordination, and supervisory time so that health

systems and clinicians have aligned incentives to participate and to lead. If this model is

extended beyond acute viral illness to chronic disease management or more intensive

diagnostic assessment and titration, pairing the decentralized platform with a local,

trusted workforce will be essential. Community health workers, pharmacists, nurses,

and paramedicine teams can support home-based examination, specimen collection,

imaging logistics, medication adjustment, culturally concordant counseling, and reliable

follow-up, with clear handoffs, shared care plans, and safety nets for deterioration, as

well as routine surveillance for adverse events and total cost. Given anticipated growth

in Medicaid enrollment and the uninsured, this approach is actionable at federal, state,

and county levels; public purchasers and health departments can procure integrated

test-to-treat services, align them with FQHC and safety-net networks, and minimize

patient out-of-pocket costs. The Rural Health Transformation Program can provide

scaffolding for demonstration projects that adapt decentralized care to rural settings by

supporting cross-sector partnerships, digital and broadband infrastructure, workforce

development, and evaluation of integration, safety, equity, cost, and interoperability.[27] A

pragmatic agenda to build on this model across additional acute and chronic conditions,

compare performance and timeliness with usual care, standardize minimum data

elements and referral pathways, and identify which functions should remain local versus

centralized is desperately needed.

References

1. Hill L, Artiga S, Rouw A, Kates J. How Equitable is Access to COVID-19 Treatments? KFF. June 23, 2022. Accessed November 8, 2025. https://www.kff.org/covid-19/how-equitable-is-access-to-covid-19-treatments/

2. Gold JAW, Kelleher J, Magid J, et al. Dispensing of Oral Antiviral Drugs for Treatment of COVID-19 by Zip Code-Level Social Vulnerability - United States, December 23, 2021-May 21, 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71(25):825-829.

3. Murphy SJ, Samson W, Sommers BD. COVID-19 Antivirals Utilization: Geographic and Demographic Patterns of Treatment in 2022. Published online 2022. https://aspe.hhs.gov/sites/default/files/documents/a19600d0dccfee0e9c595730d73fd66d/covid-antivirals-report.pdf

4. Monach PA, Anand ST, Fillmore NR, La J, Branch-Elliman W. Underuse of antiviral drugs to prevent progression to severe COVID-19 - Veterans Health Administration, march-September 2022. *MMWR Morb Mortal Wkly Rep*. 2024;73(3):57-61.

5. Wilcock AD, Kissler S, Mehrotra A, et al. Clinical risk and outpatient therapy utilization for COVID-19 in the Medicare population. *JAMA Health Forum*. 2024;5(1):e235044.

6. Shen Y, Robertson MM, Kulkarni SG, et al. Oral COVID-19 antiviral uptake among a highly vaccinated US cohort of adults with SARS-CoV-2 infection between December 2021 and October 2022. *Open Forum Infect Dis*. 2024;11(2):ofad674.

7. Menkir TF, Donnelly CA. The impact of repeated rapid test strategies on the effectiveness of at-home antiviral treatments for SARS-CoV-2. *Nat Commun*. 2022;13(1):5283.

8. Saravolatz LD, Depcinski S, Sharma M. Molnupiravir and nirmatrelvir-ritonavir: oral coronavirus disease 2019 antiviral drugs. *Clinical Infectious Diseases*. 2023;76(1):165-171.

9. Khazanchi R, Strumpf A, Essien UR, Powers SD, McManus KA. Geographic accessibility of COVID-19 test to treat sites by race, ethnicity, age, and rurality. *JAMA Netw Open*. 2022;5(11):e2241144.

10. Narayanasamy S, Veldman TH, Lee MJ, et al. RADx-UP Testing Core: Access to COVID-19 diagnostics in community-engaged research with Underserved Populations. *J Clin Microbiol*. 2023;61(8):e0036723.

11. Soni A, Herbert C, Baek J, et al. Association of Mass Distribution of Rapid Antigen Tests and SARS-CoV-2 Prevalence: Results from NIH-CDC funded Say Yes! Covid Test program in Michigan. *medRxiv*. Published online April 2, 2022. doi:10.1101/2022.03.26.22272727

12. Herbert C, Kheterpal V, Suvarna T, et al. Design and Preliminary Findings of Adherence to the Self-Testing for Our Protection From COVID-19 (STOP COVID-19) Risk-Based Testing Protocol: Prospective Digital Study. *JMIR Form Res.* 2022;6(6):e38113.

13. Ciccone EJ, Conserve DF, Dave G, et al. At-home testing to mitigate community transmission of SARS-CoV-2: protocol for a public health intervention with a nested prospective cohort study. *BMC Public Health.* 2021;21(1):2209.

14. Singler L, Uhlenbrauck G, Corbie-Smith G, et al. Say Yes! COVID test: A health communication campaign to encourage use of rapid, at-home antigen testing in underserved and historically marginalized communities. *Inquiry.* 2023;60:469580221146046.

15. Dasari S, Kurian S, Bastani R, et al. Streamlining test-to-treat: A novel care delivery model for covid-19 oral antiviral access to close the equity divide in Los Angeles county. *NEJM Catal Innov Care Deliv.* 2023;4(3). doi:10.1056/cat.22.0328

16. COVID-19 Oral Antivirals: Overcoming Barriers to Access and Education. https://cveep.org/wp-content/uploads/2023/07/CVEEP_OAV-Literature-Review-EN_0624.pdf

17. The White House. FACT SHEET: Biden Administration Announces Launch of First Federally-Supported Test to Treat Site. The White House. May 26, 2022. Accessed November 8, 2025. https://bidenwhitehouse.archives.gov/briefing-room/statements-releases/2022/05/26/fact-sheet-biden-administration-announces-launch-of-first-federally-supported-test-to-treat-site/

18. NIH launches Home Test to Treat, a pilot COVID-19 telehealth program. National Institute of Biomedical Imaging and Bioengineering. Accessed November 8, 2025. https://www.nibib.nih.gov/news-events/newsroom/nih-launches-home-test-treat-pilot-covid-19-telehealth-program

19. Yan L, Streja E, Li Y, et al. Anti-SARS-CoV-2 pharmacotherapies among nonhospitalized US veterans, January 2022 to January 2023. *JAMA Netw Open.* 2023;6(8):e2331249.

20. *Racial and Ethnic Disparities in Outpatient Treatment of COVID-19-United States.*

21. Lin SN, Rui J, Chen QP, et al. Effectiveness of potential antiviral treatments in COVID-19 transmission control: a modelling study. *Infect Dis Poverty.* 2021;10(1):53.

22. Choi HM, Moon SY, Yang HI, Kim KS. Understanding viral infection mechanisms and patient symptoms for the development of COVID-19 therapeutics. *Int J Mol Sci.* 2021;22(4):1737.

23. Hsu WH, Shiau BW, Huang PY, et al. Clinical effectiveness of oral antiviral treatment for non-hospitalized high-risk patients with COVID-19 during Omicron JN.1 subvariant wave: a US-based propensity-matched cohort study. *Pneumonia (Nathan)*. 2025;17(1):12.

24. Bruno G, Giotta M, Perelli S, De Vita G, Bartolomeo N, Buccoliero GB. Early access to oral antivirals in high-risk outpatients: Good weapons to fight COVID-19. *Viruses*. 2022;14(11):2514.

25. Hammond J, Leister-Tebbe H, Gardner A, et al. Oral nirmatrelvir for high-risk, nonhospitalized adults with Covid-19. *N Engl J Med*. 2022;386(15):1397-1408.

26. Dryden-Peterson S, Kim A, Kim AY, et al. Nirmatrelvir plus ritonavir for early COVID-19 in a large U.s. health system : A population-based cohort study: A population-based cohort study. *Ann Intern Med*. 2023;176(1):77-84.

27. Rural Health Transformation (RHT) Program. Accessed November 8, 2025. https://www.cms.gov/priorities/rural-health-transformation-rht-program/overview

Figure 1: Participant experience in the federal Home-based Test to Treat Program (HTTT)

Figure 2: Geospatial coverage of enrollment density in the HTTT program with Health Professional Shortage Area overlay

Figure 3: CONSORT diagram of participants in the federal Home-based Test to Treat Program (HTTT)

104,280 people completed interest form

23,567 people were not eligible to participate in the program at the time

160 became eligible at a later date

80,873 participants enrolled in the program

14,704 participants demonstrated suspicious participation and were removed from analysis

66,169 participants included in the analysis

34, 830 pro-active HTTT users i.e. asymptomatic and without an infection at the time of enrollment

10,032 on-demand HTTT users i.e. symptomatic but without confirmation of an infection at the time of enrollment

21,307 on-demand telehealth users i.e. had a positive test for COVID-19 or Flu at the time of enrollment

9,940 participants confirmed a positive test result

8,273 participants opted to receive telehealth

6,821 participants were prescribed medications

a: After September 13, 2023 enrollment in the program was restricted to people on non-private insurance or without insurance *unless* they had recently tested positive for COVID-19 or Flu. The ineligible participants were placed in a waitlist and allowed to participate if they reported a positive test result

b: suspicious participation refers to participants who enrolled the program after testing positive for COVID-19 and Flu and do not engage in Home-based Test to Treat activities but participated in the research sub-study

Table 1: Key characteristics of participants by group

Variable	Overall (n=66,169)	Proactive test and treat users (n=34,830)	On-demand test and treat users (n=10,032)	On-demand telehealth users (n=21,307)
Insurance				
Private	24,394 (36.9)	8,997 (25.8)	1,522 (15.2)	13,875 (65.1)
Medicaid	8,663 (13.1)	4,840 (13.9)	2,448 (24.4)	1,375 (6.5)
Medicare	20,763 (31.4)	15,610 (44.8)	2,829 (28.2)	2,324 (10.9)
No insurance	9,192 (13.9)	3,831 (11.0)	2,772 (27.6)	2,589 (12.2)
Other	3,157 (4.8)	1,552 (4.5)	461 (4.6)	1,144 (5.4)
Difficulty accessing care	29,562 (44.7)	16,053 (46.1)	5,613 (56.0)	7,896 (37.1)
Number of reasons for delaying care				
0	28,640 (43.3)	18,458 (53.0)	3,506 (34.9)	6,676 (31.3)
1	27,429 (41.5)	11,101 (31.9)	4,091 (40.8)	12,237 (57.4)
2	7,478 (11.3)	3,985 (11.4)	1,711 (17.1)	1,782 (8.4)
3	2,622 (4.0)	1,286 (3.7)	724 (7.2)	612 (2.9)
Age Group				
18-45, N (%)	32,597 (49.3)	14,198 (40.8)	5,954 (59.4)	12,445 (58.4)
46-65, N (%)	15,961 (24.1)	7,162 (20.6)	2,081 (20.7)	6,718 (31.5)
65+	17,595 (26.6)	13,456 (38.6)	1,997 (19.9)	2,142 (10.1)
Gender				
Female	45,415 (68.6)	25,171 (72.3)	7,510 (74.9)	12,734 (59.8)
Male	20,207 (30.5)	9,322 (26.8)	2,373 (23.7)	8,512 (39.9)
Other	547 (0.8)	337 (1.0)	149 (1.5)	61 (0.3)
Gender Minority	15,613 (23.6)	9,248 (26.6)	3,056 (30.5)	3,309 (15.5)
Race/Ethnicity				
Hispanic	5,284 (8.0)	2,418 (6.9)	1,129 (11.3)	1,737 (8.2)
Non-Hispanic Black	7,312 (11.1)	1,966 (5.6)	778 (7.8)	4,568 (21.4)
Non-Hispanic White	45,059 (68.1)	26,031 (74.7)	6,578 (65.6)	12,450 (58.4)
Non-Hispanic Other	8,514 (12.9)	4,415 (12.7)	1,547 (15.4)	2,552 (12.0)
Chronic Condition	33,542 (50.7)	19,884 (57.1)	6,004 (59.8)	7,654 (35.9)
Source of referral (not mutually exclusive)				
Community	7,818 (11.8)	1,762 (5.1)	681 (6.8)	5,375 (25.2)
Social Media	33,104 (50.0)	18,592 (53.4)	4,719 (47.0)	9,793 (46.0)

Variable	Overall (n=66,169)	Proactive test and treat users (n=34,830)	On-demand test and treat users (n=10,032)	On-demand telehealth users (n=21,307)
Friends/Family	21,464 (32.4)	9,167 (26.3)	2,601 (25.9)	9,696 (45.5)
Others	13,515 (20.4)	6,887 (19.8)	2,551 (25.4)	4,077 (19.1)
HPSA Score (mean (SD))	13.12 (4.8)	12.94 (5.00)	13.15 (4.7)	13.40 (4.7)

Difficulty accessing care defined as:
Reasons for delaying care included: limited provider availability, concerns about cost, lackof transportation, or inability to take time off
HPSA: Health Professional Shortage Area Score (HPSA) obtained from data.hrsa.gov

Table 2: Test and treatment outcomes by group

Variable	Overall (n=66,169)	Proactive test and treat users (n=34,830)	On-demand test and treat users (n=10,032)	On-demand treat users (n=21,307)
Reported Positive Test				
Positive for COVID-19 or Flu	9,940 (15.0)	1,247 (3.6)	666 (6.6)	8,027 (37.7)
Days from Symptom Onset to First Positive Report	1.65 (3.5)	1.99 (6.9)	2.87 (8.6)	1.52 (1.7)
Among those who tested positive (n = 9,940)				
Received Telehealth from HTTT	8,273 (83.2)	1,043 (83.6)	553 (83.0)	6,677 (83.2)
Telehealth Started 9 AM-5PM Local Time on Weekdays	1,959 (23.7)	308 (29.5)	143 (25.9)	1,508 (22.6)
Among those who received telehealth from the program (n = 8,273)				
Were prescribed medications	6,821 (82.4)	867 (83.1)	426 (77.0)	5,528 (82.8)
Days from Symptom Onset to Prescription ≤ 5 days	5,908 (92.8)	700 (91.5)	279 (84.5)	4,929 (93.5)
Days from Symptom Onset to Prescription ≤ 1 day	3,808 (59.8)	488 (63.8)	158 (47.9)	3,162 (60.0)
Home delivery of medication	1,092 (16.0)	191 (22.1)	89 (21.0)	812 (14.7)

Figure 4a: Receive a telehealth consult

Figure 4b: Start of telehealth within business hours

Figure 5a Prescribed medication

Figure 5b Prescribed medication within 1 day of symptom onset

Figure 5c: Opted for home-delivery of medication