Effect of higher dose primaquine for the radical cure of *Plasmodium vivax* malaria in Indonesia: a systematic review and individual patient data meta-analysis

Ihsan Fadilah, MSc[a,b] · James A Watson, DPhil[b,c] · Prof Ayodhia P Pasaribu, PhD[d,e] · Prof Inge Sutanto, PhD[f] · Prof Erni J Nelwan, PhD[g,h] · Kartini Lidia, MSc[i] · Megha Rajasekhar, PhD[j] · Iqbal RF Elyazar, DPhil[a] · Prof Walter RJ Taylor, FRCP[b,k] · Prof Kamala Thriemer PhD[l] · Prof Nicholas PJ Day, FRCP[b,k] · Jeanne Rini Poespoprodjo, MD[m,n] · Prof Julie A Simpson, PhD[b,c,j] · Prof Ric N Price, FRCP[b,c,l,k] · Prof J Kevin Baird, PhD[a,b] · Robert J Commons, FRACP[c,l,o]

a. Oxford University Clinical Research Unit Indonesia, Faculty of Medicine Universitas Indonesia, Jakarta, Indonesia
b. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom
c. Infectious Diseases Data Observatory, Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom
d. Department of Pediatrics, Medical Faculty, Universitas Sumatera Utara, Medan, Indonesia
e. Yayasan Penguatan Kesehatan Masyarakat Tridarma (YPKMT)/Tridarma Healthcare Empowerment Foundation (THEMP), Medan, Indonesia
f. Department of Parasitology, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia
g. Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia
h. Division of Tropical Medicine and Infectious Disease, Department of Internal Medicine, Cipto Mangunkusumo Hospital, Jakarta, Indonesia
i. Department of Pharmacology and Therapy, Faculty of Medicine and Veterinary Medicine, Universitas Nusa Cendana, Kupang, Indonesia
j. Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, VIC, Australia
k. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand
l. Global and Tropical Health Division, Menzies School of Health Research and Charles Darwin University, Darwin, Australia
m. Centre for Child Health, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia
n. Yayasan Pengembangan Kesehatan dan Masyarakat Papua, Timika, Papua, Indonesia
o. General and Subspecialty Medicine, Grampians Health Ballarat, Ballarat, Australia

Correspondence to:
Dr Ihsan Fadilah · Oxford University Clinical Research Unit Indonesia, IMERI Building, Faculty of Medicine Universitas Indonesia, Jalan Salemba Raya No. 6, Jakarta 10430, Indonesia · **ifadilah@oucru.org**

A/Prof Robert J Commons · Global and Tropical Health Division, Menzies School of Health Research and Charles Darwin University, Darwin, NT 0810, Australia · **rob.commons@wwarn.org**

Summary

Background *Plasmodium vivax* malaria has diverse transmission and relapse patterns in Indonesia. The optimal dose of primaquine to prevent relapses across the country is unknown. We evaluated the anti-relapse efficacy, gastrointestinal tolerability, and haematological safety of different primaquine regimens in varied endemic settings in Indonesia.

Methods We systematically searched for studies published between 1 January 2000 and 23 July 2024 prospectively enrolling patients with acute uncomplicated *P. vivax* malaria where some patients were treated with primaquine. Individual patient data (IPD) from eligible studies were pooled and harmonised. We fitted one-stage IPD multivariable regression models to estimate the relationship between the weight-adjusted primaquine dose with three separate primary outcomes: (i) the time to first *P. vivax* recurrence (days 7–180), (ii) any gastrointestinal discomfort (days 5–7), and (iii) ≥25% reduction relative to baseline haemoglobin and a reduction to <7 g/dL (days 1–14).

Findings Of ten eligible studies, seven were available for inclusion. Compared with a total dose of 3·5 mg/kg primaquine, patients treated with a total dose of 7 mg/kg had a lower rate of recurrence over 6 months (adjusted hazard ratio 0·53; 95% confidence interval [CI] 0·45 to 0·63; n = 1797); the relative efficacy was consistent across regions, but the absolute benefit varied. Gastrointestinal discomfort was more frequent with higher doses (adjusted risk ratio 1·32 per 0·25 mg/kg daily dose; 95% CI 1·15 to 1·51; n = 952). For haematological safety (n = 822; 788/822 [96%] patients had G6PD activity ≥70%), only one patient developed clinically relevant haemolysis.

Interpretation Across all transmission settings in Indonesia, a total dose of 7 mg/kg halved the rate of recurrent *P. vivax* malaria over a 6-month period compared with the low dose of 3·5 mg/kg. However, increased daily doses slightly increased risks of gastrointestinal discomfort and haemolysis.

Funding NDM Tropical Network Fund, Bill and Melinda Gates Foundation

Keywords Primaquine, *Plasmodium vivax*, malaria, relapse, efficacy, safety, tolerability, Indonesia

Research in context

Evidence before this study Indonesia is a geographically vast country with heterogeneous patterns of periodicity and varying magnitudes of risk for *Plasmodium vivax* relapse. However, the optimal regimen for the radical cure of *P. vivax* is unknown. The national antimalarial guidelines continue to recommend a low-dose regimen of 3·5 mg/kg without routine glucose-6-phosphate dehydrogenase (G6PD) testing. We systematically searched for trials conducted in Indonesia that randomly assigned patients with *P. vivax* malaria to primaquine dosing regimens. Articles published in any language between 1 January 2000 and 23 July 2024, using the terms "vivax" and "primaquine" in MEDLINE, Embase, Web of Science, Scopus, and the Cochrane Library were identified. No published randomised controlled trial directly comparing high-dose (≥5 mg/kg total) versus low-dose (2 to <5 mg/kg total) primaquine in Indonesia was identified. We identified two trials that randomised patients with *P. vivax* malaria to receive either high-dose (target total dose of 7 mg/kg over seven or 14 days) or low-dose (3·5 mg/kg over 14 days) primaquine, compared with placebo, in combination with dihydroartemisinin-piperaquine. One multi-country trial included two study sites in Sumatra: Hanura and Tanjung Leidong. Compared with placebo, high-dose primaquine reduced relapse by 60–89%. In another trial, patients acquired infections in Papua and were followed in malaria-free East Java. Low-dose primaquine reduced relapse by 74% compared with placebo.

Added value of this study This country-specific systematic review and individual patient data meta-analysis included 1797 patients from seven studies evaluating anti-relapse efficacy, gastrointestinal tolerability, and haematological safety. The analysis represents the most comprehensive analysis to date of different primaquine dosing regimens for *P. vivax* malaria in Indonesia. The results suggest that doubling the total dose from 3·5 mg/kg to 7 mg/kg would halve the rate of *P. vivax* recurrence within 180 days, however, the absolute benefit of this reduction depends on the underlying risk of recurrence in different regions of Indonesia. An increase in the daily dose of primaquine resulted in a moderate increase in gastrointestinal discomfort; and only a rare occurrence of clinically relevant haemolysis, given that G6PD activity among patients was mostly ≥70%.

Implication of all the available evidence For Indonesian patients with *P. vivax* malaria, a high total primaquine dose of 7 mg/kg is more efficacious than the currently implemented 3·5 mg/kg regimen. Higher daily doses of primaquine are associated with increased gastrointestinal discomfort, however, these may be mitigated by co-administration with food. Clinically relevant haemolysis is rare in patients with G6PD activity >30%.

Background

Plasmodium vivax is more geographically widespread than other causes of human malaria. In regions where *P. vivax* and *P. falciparum* are co-endemic, *P. vivax* is increasingly the predominant species because conventional malaria control efforts more readily impact *P. falciparum*.[1] As of 2024, nearly 80% of districts in Indonesia had achieved malaria-free status.[2] However, *P. vivax* malaria remains prevalent with hundreds of thousands of cases; most reported from the eastern Papua region.[3,4]

Dormant liver stages (hypnozoites) of *P. vivax* cause repeated blood-stage infections (relapses) and attacks of acute malaria that constitute a major cause of morbidity and mortality, particularly among young children and pregnant women.[5] Relapsing *P. vivax* avoids conventional control efforts aimed at treating acute malaria and minimising mosquito vector contact. Radical cure of acute malaria, by combining blood schizontocidal drugs with an 8-aminoquinoline (primaquine or tafenoquine) that kills dormant hypnozoites, remains the only effective strategy to avert *P. vivax* relapses.

A meta-analysis of malaria studies across *P. vivax*-endemic regions globally published in 2024 suggested that a high total dose of 7 mg/kg can halve the rate of *P. vivax* recurrences compared with a low total dose of 3·5 mg/kg, with only a slight increase in gastrointestinal discomfort.[6] Patients treated with a daily primaquine dose of up to 0·5 mg/kg with glucose-6-phosphate dehydrogenase (G6PD) activity >30% had comparable risks of haemolysis with patients not exposed to primaquine.[7]

Updated 2024 WHO guidelines reflected these findings, with a high total dose of primaquine recommended over either 14 days or 7 days.[8] Indonesian national guidelines currently recommend a low-dose regimen of 3·5 mg/kg total over 14 days without routine G6PD testing, with a higher dose (7 mg/kg over 14 days) recommended for cases of suspected relapse or treatment failure.[9] However, Indonesia is a geographically large country with varied *P. vivax* epidemiological characteristics. For example, in Papua, Indonesia, there is a high rate of *P. vivax* transmission and high risk of relapse, even when high total dose primaquine is given,[10-12] while other regions of Indonesia have reported good efficacy rates with low dose primaquine.[13,14] Thus, the potential heterogeneity in effects, as well as the overall risks and benefits of different primaquine regimens for Indonesia, remains unclear.

This study aimed to assess the anti-relapse efficacy, gastrointestinal tolerability, and haematological safety of the different primaquine doses in patients with *P. vivax* malaria in Indonesia.

Methods

Search strategy and selection criteria

Building on an existing living systematic review[15] of studies of patients with acute uncomplicated *P. vivax* malaria, we conducted an updated systematic search of MEDLINE, Embase, Web of Science, Scopus, and the Cochrane Library for studies conducted in Indonesia and published in any language between 1 January 2000 and 23 July 2024. Search terms are provided in Supplementary Material (**List S1**). Two reviewers (IF and RJC) conducted the review, with discrepancies resolved through discussion. The protocol was pre-registered with PROSPERO (CRD42024580630). We included randomised or non-randomised therapeutic trials with ≥28 days of active follow up for gastrointestinal tolerability and haematological safety, or ≥42 days for anti-relapse efficacy. Eligible studies included at least one group receiving multi-day primaquine for relapse prevention, initiated within seven days of blood schizontocidal treatment (chloroquine, quinine, or common artemisinin-based combination therapies).

Data pooling

Investigators of eligible studies were invited to contribute individual patient data (IPD), including unpublished data when possible. Shared data were uploaded to the Worldwide Antimalarial Resistance Network repository and curated in accordance with the IDDO SDTM Implementation Guide.[16] Patients were excluded if key variables were missing (age, sex, weight, baseline parasitaemia, treatment regimen), they had severe malaria, pregnancy, mixed infections, or received additional antimalarials after schizonticidal treatment. Additional inclusion criteria are described in Supplementary Material (**List S2**). Included data were pseudo-anonymised and had received prior ethical approval locally. No additional approval was required, as per Oxford Tropical Research Ethics Committee guidance. This study is reported according to PRISMA-IPD guidelines[17] (**Table S1**).

Endpoints

Day zero was defined as initiation of antimalarial treatment after enrolment. The primary efficacy endpoint was the time to the first *P. vivax* recurrence, irrespective of symptoms, between days 7–180. Secondary endpoints included the time to the first symptomatic recurrence and the number of recurrences within the same timeframe. The primary gastrointestinal tolerability endpoint was a composite endpoint including any of the following signs of gastrointestinal discomfort occurring between days 5–7 (after the resolution of acute malaria symptoms and completion of schizonticidal treatment): vomiting, anorexia, diarrhoea. Secondary tolerability endpoints included similar symptoms on day 0 and days 1–2, and acute vomiting within 1 hour of primaquine administration. The primary haematological safety endpoint was a ≥25% haemoglobin reduction from baseline to <7 g/dL between days 1–14. Secondary endpoints included (i) maximum haemoglobin drop between days 1–14, 2–3, and 5–7, (ii) incident anaemia (Hb <11 g/dL) during days 2–3 in patients with baseline Hb ≥11 g/dL and normal G6PD activity (≥70% activity[18]; study subgroup of interest), and (iii) severe haemolytic adverse events (Hb <5 g/dL, >5 g/dL drop, blood transfusion, renal failure requiring dialysis, or death) within 14 days.

Exposures

The efficacy exposure was the weight-adjusted (i.e., mg of primaquine base per kg of body weight [mg/kg]) total primaquine dose. The gastrointestinal tolerability and haematological safety exposure was the weight-adjusted daily dose.[6,7] Doses were calculated from daily tablets or the mg administered and if these were unavailable they were inferred from protocols. Primaquine dose was specified in our models as a continuous variable. The effects of different categories of primaquine dose (**List S2**) were explored.

Data analysis

The main analyses estimated the effect of primaquine dose on recurrence, gastrointestinal tolerability, and haemolysis under a one-stage IPD meta-analysis framework.

Anti-relapse efficacy
Cumulative incidence of first recurrence was modelled with a multivariable Cox proportional hazards model including total primaquine dose, age, sex, natural log-transformed baseline parasite density, and study site. A causal directed acyclic graph (DAG) guided model development (**Figure S1**), and a restricted cubic spline for total primaquine dose allowed for nonlinearity (See **List S2** for further methodological details). We evaluated potential differential effects of total primaquine dose by age group (<5 years or ≥5 years) and origin of infection (within or outside Papua; reflecting a higher potential burden of hypnozoites and/or more primaquine-tolerant strains within Papua). Additional analyses, including sensitivity analyses and assessment of residual confounding using negative controls are described in Supplementary Material (**List S2**).

Gastrointestinal tolerability
We summarised the outcome prevalence descriptively across study sites to highlight heterogeneity in the measurement of this endpoint. In the randomised subset of these data, the risk of gastrointestinal discomfort was modelled using multivariable Poisson regression[19] with daily primaquine dose, age, sex, and log baseline parasite density (based on a DAG; **Figure S3**), with dose categories and splines used to assess nonlinear effects. Study site was not included in the model, as the study population for this outcome originated from a single trial conducted under the same protocol. Similar model specification was used to model the risk of acute vomiting. As the outcome prevalence was high, Poisson regression was used to model the binary outcomes to estimate the risk ratio. Effect estimates from modified Poisson models[19] were obtained as part of the sensitivity analysis.

Haematological safety
The presence of clinically significant haemolysis (≥25% haemoglobin reduction from baseline to <7 g/dL) between days 1–14 was described. Multivariable linear regression was used to model the maximum haemoglobin drop from baseline over 14 days, including as covariates: daily primaquine dose, baseline haemoglobin, $\log_2$ G6PD activity, age, sex, log baseline parasite density, study site, and a dose-sex-$\log_2$ G6PD interaction. A DAG guided model specification (**Figure S4**), and splines were used to assess nonlinear effects. Additional analyses, including a subgroup analysis of incident anaemia and analysis of methaemoglobinaemia, are described in Supplementary Material (**List S2**).

238 Risk of bias was assessed using the ROB2 (randomised trials)[20] and ROBINS-I tools (non-
239 randomised studies),[21] adapted for the current study objectives. Statistical analysis was
240 performed using R (v4.3.0) according to a pre-specified plan.[22]

Role of the funding source

243 The study funder had no role in study design, data collection, analysis, interpretation, and
244 manuscript writing.

Results

Between 1 January 2000 and 23 July 2024, 234 *P. vivax* efficacy studies were published, of which ten included Indonesian study sites and were eligible for the pooled analysis. IPD from eight studies were available for analysis. After patient-level exclusion criteria were applied, 1797 patients enrolled into seven studies were included in the analysis (**Table 1**). The seven studies included one trial that individually randomised patients to different primaquine dose regimens,[14] while the remaining six administered the same total primaquine dose within each study (a target total of either 3.5 or 7 mg/kg).[10-13,23,24] There was a low to moderate risk of within study bias (**Table S2**).

The median age of the patients was 17 years (interquartile range [IQR] 10–29) with 172 (9·6%) younger than 5 years. The median weight was 45 kg (IQR 24–57) and 1056 (59%) were male. Overall, 452 (25%) patients acquired their *P. vivax* infection in Papua (**Figure 2**) and 1432 (80%) were treated with dihydroartemisinin-piperaquine (**Table 1**). A total of 196 (11%) patients were treated without primaquine, 464 (26%) with low total dose, and 1137 (63%) with high total dose (**Figures S5, S6, S7, S8).**

Compared with low total dose, patients administered high-dose primaquine were more likely to be treated over seven days, receive dihydroartemisinin-piperaquine, have a higher baseline parasite density, and have actual dosing data available (**Table S3**, **Figure S9**). Eligible studies that were not included[25-27] predominantly enrolled patients in Papua and had a shorter follow-up duration (**Tables S4, S5, S6**).

Compared with patients treated without primaquine (enrolled in a single study conducted at non-Papuan sites), the rate of *P. vivax* recurrence by day 180 was lower in both patients treated with low-dose (3·5 mg/kg) primaquine (adjusted hazard ratio [AHR] 0·33; 95% CI 0·25 to 0·44) and high-dose (7 mg/kg) primaquine (AHR 0·18; 95% CI 0·12 to 0·26). The corresponding rate of recurrence was lower following high-dose primaquine compared to low-dose primaquine (AHR 0·53; 95% CI 0·45 to 0·63; **Figure 3**). Increasing the total dose above 7 mg/kg remains of uncertain efficacy as the interval estimates of the estimated dose-response curve were wide (**Figure 3**). This dose-response relationship was consistent across alternative dose groupings and sensitivity analyses (**Figures S11**, **S12**). Negative control methods suggest residual confounding was minimal (**List S3**). The effect of increasing the dose of primaquine did not differ with age (<5 or ≥5 years old, $p_{interaction}$ = 0·88) or by origin of infection (within Papua or outside Papua, $p_{interaction}$ = 0·45, **Figure S13**).

In a subset of patients (n = 1608, 89%) followed through multiple P. vivax episodes for at least 180 days, those who acquired infections within Papua had a higher 180-day recurrence rate (76 episodes per 120 person-years; crude incidence rate [IR] 0·63 per person-year; 95% confidence interval [CI] 0·50 to 0·79) than those infected outside Papua (140 episodes per 451 person-years; IR 0·31 per person-year; 95% CI 0·26 to 0·37). **Figure 3** shows recurrence rates for both regions by primaquine dose groupings.

Gastrointestinal tolerability on days 5–7 was assessed in 1254 (70%) patients from five studies.[11,13,14,23,24] Of these, 190 (15%) received no primaquine, 203 (16%) received low daily dose, 482 (38%) intermediate daily dose, and 379 (30%) high daily dose primaquine; with 125

(10%) younger than 5 years. Within each primaquine daily dose group, there was substantial heterogeneity in the percentage of patients reporting gastrointestinal discomfort on days 5–7 across studies (**Figure S14, Table S7**), making data pooling for effect estimation unreliable. To estimate the unbiased effect of increasing daily doses, we used subsets of data from one trial[14] in which primaquine daily doses were randomised. The risk of early gastrointestinal symptoms (day 0 and days 1–2) did not vary with the daily dose of primaquine (**Figure S15, Table S9**). However, by days 5–7, when 894/952 (94%) patients were afebrile, the risk of gastrointestinal symptoms increased with the daily dose of primaquine: adjusted risk ratio [ARR] 1·32 (95% CI 1·15 to 1·51) per 0·25 mg/kg/day increment in primaquine dose (**Figure S15, Table S9**). In 767 patients assessable for acute vomiting, there was inconclusive evidence that a higher daily dose of primaquine resulted in an increased risk of acute vomiting within 60 minutes of administration (ARR 1·19; 95% CI 0·84 to 1·67 per 0·25 mg/kg/day increment; **Figure S16, Table S8**).

One multi-country randomised trial[14] contributed 822 patients to the primary haematological safety analysis, of whom 173 (21%) received no primaquine, 6 (<1%) received a low daily dose, 318 (39%) received an intermediate daily dose, and 325 (40%) a high daily dose. G6PD activity was measured at baseline in all eligible patients and was similar between treatment arms; 788 (96%) had normal G6PD activity (≥70%) and 34 (4%) had intermediate activity (30–69%; **Figure S17**).

Severe haemolysis was recorded in one patient, a female in the 5–14-year age group with 37·4% G6PD activity treated with dihydroartemisinin-piperaquine plus a high daily dose of primaquine (0·99 mg base/kg) administered over 7 days. She had a 41% fall in haemoglobin from 11·6 g/dL on day 0 to 6·9 g/dL on day 3 (4·7 g/dL reduction) and recovered (Hb ≥11 g/dL) at least by day 21 without requiring a blood transfusion. No patients had a haemoglobin fall to <5 g/dL, a haemoglobin fall >5 g/dL from baseline, blood transfusion, renal failure requiring dialysis, or death (**Table S10**).

The effect of primaquine daily dose on haemoglobin varied with sex and G6PD activity ($p_{interaction}$ = 0·01). In contrast to females with normal G6PD activity, a higher daily dose of primaquine was associated with a greater absolute fall in haemoglobin in females with intermediate G6PD activity (**Figures 4A, S18**). In males, lower G6PD activity was not associated with a greater absolute decrease in haemoglobin. In 612 patients with normal G6PD activity and a baseline haemoglobin level of ≥11 g/dL, the risk of anaemia on days 2–3 did not increase with higher primaquine daily doses (**Figure S19**, **Table S9**). In 788 patients with normal G6PD activity, haemoglobin levels reached an expected nadir on days 2–3, likely reflecting haemolysis associated with blood-stage acute malaria. Haemoglobin levels returned to baseline approximately two weeks after initiating treatment (**Figure 4B**). A higher primaquine daily dose resulted in higher day 7 methaemoglobin levels and an increased risk of methaemoglobin levels ≥10% between days 1 to 14 (prevalence = 262/1026 [25·5%], **Figure S20**).

Discussion

The optimal anti-relapse regimen to prevent recurrent *P. vivax* malaria must trade-off the absolute number of recurrent malaria episodes averted against the length of treatment duration, its tolerability, and safety. This balance of practicality, tolerability, and safety against absolute benefit will vary across regions primarily because the absolute risk of recurrence varies considerably. Our meta-analysis of 1797 Indonesian patients demonstrates that increasing the total dose of primaquine from 3·5 to 7 mg/kg would halve the recurrence rate. However, the absolute benefit provided by the higher dose will vary substantially as the crude recurrence incidence rate varied substantially between regions (IR 0·63 [95% CI 0·50 to 0·79] per person-year within Papua; IR 0·31 [95% CI 0·26 to 0·37] per person-year outside Papua). Higher daily doses were associated with a greater risk of gastrointestinal discomfort and there is a risk of severe primaquine-induced haemolysis in females with intermediate G6PD activity.[28] Identifying an optimal balance between benefits and harms is key to determining the best regimen for each region within Indonesia.

Our Indonesia-specific estimates for anti-relapse efficacy comparing high- and low-dose primaquine align with those reported by Commons and colleagues using pooled global data.[6] Both analyses included non-randomised comparisons, so confounding bias cannot be excluded. Much of the within-study dose variation was attributable to differences in body weight rather than to random allocation. Similar to Commons et al.,[6] we used a causal diagram to inform adjustment for key confounders in regression analysis. We further applied negative control methods[29] and found no material residual confounding. Consistent with this study, a meta-analysis of study-level data from randomised trials[30-36] directly comparing high- and low-dose primaquine in any country confirmed that high-dose primaquine reduces the rate of first *P. vivax* recurrence by around 50% (**Figures *5*A, *5*B**).

Infections from Papua may involve a higher hypnozoite load and strains may be more tolerant to primaquine, both necessitating a higher primaquine dose.[37-39] However, we found no evidence that higher doses of primaquine in patients infected in Papua would provide greater relative efficacy compared to patients infected outside Papua. The global analysis reported similar findings, showing high relative efficacy of high-dose primaquine in locations with both low- and high-relapse periodicity.[6] Hence, higher relative anti-relapse efficacy of increased primaquine doses is likely consistent and generalisable across different regions and parasite strains. However, given the diverse case-mix and varying baseline risks across populations, heterogeneous treatment-effects on an absolute scale are expected and will be relevant for policy makers and clinicians. When comparing high versus low total dose primaquine, the number needed to benefit to prevent one recurrence for one year in Papua would be two patients (95% CI 2 to 3), compared with seven patients (95% CI 6 to 9) in non-Papua regions (**Figure *5*C**).

A higher daily dose of primaquine led to increased gastrointestinal discomfort, independent of acute malaria symptoms. Reduced tolerability could lead to non-compliance, preventing the target total dose from being achieved. Reduced adherence or inadequate supervision may lead to lower anti-relapse efficacy.[11,40] Although, previous studies have shown that taking primaquine tablets with a meal significantly improves gastrointestinal tolerability,[41] relevant data to address this in our IPD analysis were not available.[42]

379
A lack of routine G6PD screening limits the safe implementation of high total dose primaquine.
The risk of haemolysis is almost entirely in individuals with G6PD deficiency. While our analysis
did not include this population, our analysis included patients with intermediate G6PD activity.
Chu and colleagues reported that despite being labelled as phenotypically normal by point-of-
care tests (≥30% G6PD activity), G6PD heterozygous females can experience significant
haemolysis when exposed to 1 mg/kg/day of primaquine.[28] We observed that females with
G6PD activity between 30 and 69% experienced greater haemoglobin reductions as the daily
dose increased compared to those with G6PD activity ≥70%, however this was not clinically
significant at doses of 0.5 mg/kg/day or less. Primaquine also induced dose-dependent
increases in methaemoglobin levels in our study population. Prior evidence suggests that this
elevation typically occurs without symptoms, and resolves spontaneously within weeks.[41,43]
Increased primaquine-induced methaemoglobinaemia indicates sufficient production of active
primaquine metabolites and hence, greater anti-relapse efficacy.[44,45]
393
In line with the 2024 WHO antimalarial guidelines,[8] our findings suggest that for most patients
with *P. vivax* in Indonesia, an optimal regimen could involve a high total dose of 7 mg/kg
administered over 1–2 weeks, taken with food, and initiated after screening for G6PD
deficiency. Of note, in western Indonesia, there is a low risk of relapse following a low total
dose primaquine,[13] suggesting that the absolute benefit from high total dose primaquine may
be reduced in these areas and different dosing could be considered.
400
Tafenoquine has been recommended for use in endemic countries in South America, where
chloroquine is typically the blood-stage drug of choice.[8] This single-dose anti-relapse treatment
is currently not recommended for patients receiving artemisinin-based combination therapies
(ACTs).[8] In Indonesia, where ACTs are recommended due to a high prevalence of chloroquine-
resistant *P. vivax*, tafenoquine is not currently available. The INSPECTOR trial, conducted in
soldiers returning from Indonesian Papua, found no clinically meaningful benefit of
tafenoquine plus dihydroartemisinin-piperaquine compared with dihydroartemisinin-
piperaquine alone.[25] The reason for this remains unclear, although a drug–drug interaction or
suboptimal dosing have been proposed.[46]
410
We were unable to include two recently published eligible studies in our pooled dataset,
primarily due to unavailability of IPD. These studies would have added <10% more patients,
making it unlikely that the current results would change significantly. One of these studies[25]
randomised patients to receive low-dose primaquine over 14 days versus a placebo, and its
estimated efficacy (HR 0·26; 95% CI 0·16 to 0·43) aligns with our pooled effect-estimates. In
the current study, ~30% of patients exposed to primaquine were estimated from the study
protocol rather than the actual tablets administered. Given an inability to account for reduced
adherence this may have led to the actual dose of primaquine received being less than
expected, which would be expected to bias the anti-relapse efficacy towards the null, making
our efficacy estimates conservative. While the same applies to tolerability and safety (i.e.,
higher daily doses potentially appearing more tolerable and safer), both in practice can be
effectively mitigated by taking food and G6PD screening. Therefore, the impact of potential
quantitative bias is deemed minimal for clinical use. Lastly, although our focus was on anti-
relapse efficacy, the studies (except for those conducting follow-up in malaria-free regions[10,12])
were unable to distinguish whether recurrences were caused by new infections, treatment

failure, or hypnozoite activation. The use of highly efficacious schizonticides made recrudescence unlikely and approximately 80% of vivax recurrences were estimated to result from relapses.[47] If new infections were indeed substantial, they would have diluted our efficacy estimates towards the null, making our results conservative.

In summary, increasing the total primaquine dose from 3·5 mg/kg to 7 mg/kg is likely to provide a substantial reduction in the risk of recurrence, with no evidence of treatment-effect heterogeneity across Indonesia. However, the variable risk of recurrence across the country means that there are substantial differences in the absolute benefit of 7 mg/kg versus 3·5 mg/kg total dose primaquine (e.g., higher benefit in Papua but lower in Sumatra). Completing a high total dose of primaquine in a shorter timeframe by administering a higher daily dose, is preferable for maximising adherence to a full course of treatment, however this may increase the risk of gastrointestinal discomfort or haemolysis. These risks can be mitigated by co-administering primaquine with food, G6PD screening, and monitoring for potential haemolytic events.

Contributors

IF, JAW, RNP, JKB, and RJC conceived the study, analysed and interpreted the data, and drafted the manuscript. IF and RJC accessed and verified the data. JAW, RNP, and JKB provided technical support. APP, IS, EJN, KL, IRFE, WRJT, KT, NPJD, JRP, JAS, RNP, JKB conceived and undertook the individual studies and enrolled the patients. All authors revised the manuscript and were responsible for the decision to submit for publication.

Declaration of interests

JKB and KT report institutional research funding from Medicines for Malaria Venture. JKB also reports institutional research funding from GSK, Wellcome Trust, and Sanaria; participation on the US national Institutes of Health data safety monitoring board; and membership of the editorial board of *Travel Medicine and Infectious Disease* and the guidelines development group for malaria control and elimination, Global Malaria Programme, WHO. RJC has been a technical advisor for the WHO guidelines development group. RJC, JKB, and RNP report contributions to Up-to-Date. All other authors declare no competing interests.

Data sharing

Pseudo-anonymised participant data used in this analysis are available for access via the Infectious Diseases Data Observatory (IDDO) website (https://www.iddo.org). Requests for data access will be reviewed according to IDDO procedures, to ensure that use of data protects the interests of the participants and researchers according to the terms of ethics approval and principles of equitable data sharing. IDDO is registered with the Registry of Research Data Repositories (https://www.re3data.org/). Code for data analysis and visualisation is available at https://github.com/ihsanfadil/wwarn-pq-ina.

Acknowledgements

IF is supported by the Nuffield Department of Medicine Tropical Network Fund. RJC, RNP, JAS and KT are supported by the Australian National Health and Medical Research Council (NHMRC) Investigator Grants (1194702, 2008501, 1196068 and 2033264 respectively). JAW is a Sir Henry Dale Fellow funded by the Wellcome Trust (223253/Z/21/Z). MR is supported by the Australian NHMRC CRE Grant 2024622 (Australian Centre of Research Excellence for Malaria Elimination) and Australian NHMRC Synergy Grant 2018654. We thank all patient volunteers, healthcare workers, and research staff who contributed to the individual studies at all the sites, and the WWARN team for technical and administrative support. This research was supported by grants from Bill and Melinda Gates Foundation. For open access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript arising from this submission. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the US Centers for Disease Control and Prevention.

Figure 1. Study and patient selection.

Databases systematically searched included MEDLINE, Embase, Web of Science, Scopus and the Cochrane Library. Additional analyses (e.g., sensitivity or restricted analyses) were conducted on smaller subsets of the relevant primary dataset.

524 **Table 1. Patient characteristics by study**

Characteristic	Study							Overall
	Hasugian et al. 2007[23]	Pasaribu et al. 2013[13]	Sutanto et al. 2013[12]	Lidia et al. 2015[24]	Nelwan et al. 2015[10]	Taylor et al. 2019[14]	Poespoprodjo et al. 2022[11]	
Number of patients	139	331	39	51	118	963	156	1797
Median age (Q1, Q3; years)	13 (4, 26)	14 (9, 27)	27 (25, 29)	33 (25, 41)	28 (25, 31)	16 (10, 29)	16 (7, 29)	17 (10, 29)
Age group								
Less than 5 years	44 (32%)	26 (8%)	0 (0%)	0 (0%)	0 (0%)	69 (7%)	33 (21%)	172 (9%)
5 to 14 years	29 (21%)	148 (45%)	0 (0%)	0 (0%)	0 (0%)	386 (40%)	42 (27%)	605 (34%)
At least 15 years	66 (47%)	157 (47%)	39 (100%)	51 (100%)	118 (100%)	508 (53%)	81 (52%)	1020 (57%)
Male sex	73 (53%)	186 (56%)	39 (100%)	24 (47%)	118 (100%)	534 (55%)	82 (53%)	1056 (59%)
Body weight (kg)	43 (14, 53)	38 (22, 52)	64 (59, 72)	42 (36, 45)	69 (63, 74)	44 (24, 55)	45 (19, 56)	45 (24, 57)
Baseline parasite density (per μL)	1575 (375, 5244)	760 (320, 3100)	2848 (784, 4432)	2640 (1500, 6560)	880 (148, 2496)	3911 (1170, 8659)	4920 (1680, 11550)	2793 (630, 7130)
Presence or recent history of fever	139 (100%)	321 (97%)	6 (16%)	51 (100%)	NA	850 (88%)	156 (100%)	1523 (91%)
Number of missing data	0	0	2	0	118	0	0	120
Schizonticidal drug								
Dihydroartemisinin-Piperaquine	68 (49%)	164 (50%)	0 (0%)	25 (49%)	58 (49%)	961 (100%)	156 (100%)	1432 (80%)
Artesunate-Amodiaquine	71 (51%)	167 (50%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	238 (13%)
Artesunate-Pyronaridine	0 (0%)	0 (0%)	0 (0%)	0 (0%)	60 (51%)	0 (0%)	0 (0%)	60 (3·3%)
Chloroquine	0 (0%)	0 (0%)	0 (0%)	26 (51%)	0 (0%)	2 (0·2%)	0 (0%)	28 (1·6%)
Quinine	0 (0%)	0 (0%)	39 (100%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	39 (2·2%)
Primaquine total dose (mg base/kg)	3·8 (3·4, 4·3)	4·2 (3·6, 4·8)	7·1 (6·7, 8·2)	4·5 (4·2, 4·9)	7·1 (6·5, 8·1)	7·0 (6·0, 7·9)	7·3 (5·9, 8·2)	6·4 (4·1, 7·5)
Primaquine total dose group								
No primaquine	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	196 (20%)	0 (0%)	196 (11%)
Low	118 (85%)	265 (80%)	0 (0%)	39 (76%)	1 (0.85%)	21 (2.2%)	20 (13%)	464 (26%)
High	21 (15%)	66 (20%)	39 (100%)	12 (24%)	117 (99%)	746 (77%)	136 (87%)	1137 (63%)
Primaquine daily dose (mg base/kg)	0·27 (0·24, 0·31)	0·30 (0·26, 0·34)	0·51 (0·48, 0·58)	0·32 (0·30, 0·35)	0·53 (0·47, 0·61)	0·59 (0·46, 1·00)	0·43 (0·00, 0·56)	0·47 (0·28, 0·63)
Primaquine duration								
No primaquine	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	196 (20%)	0 (0%)	196 (11%)
7 days	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	385 (40%)	0 (0%)	385 (21%)
14 days	139 (100%)	331 (100%)	39 (100%)	51 (100%)	118 (100%)	382 (40%)	156 (100%)	1216 (68%)
Primaquine dose calculation								
No primaquine	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	196 (20%)	0 (0%)	196 (11%)
Actual dosing	0 (0%)	0 (0%)	38 (97%)	0 (0%)	118 (100%)	767 (80%)	156 (100%)	1079 (60%)
Protocol dosing	139 (100%)	331 (100%)	1 (2.6%)	51 (100%)	0 (0%)	0 (0%)	0 (0%)	522 (29%)

Primaquine supervisio								
No primaquine	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	196 (20%)	0 (0%)	196 (11%)
Unsupervised	139 (100%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	67 (43%)	206 (11%)
Partially supervised	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	89 (57%)	89 (5.0%)
Fully supervised	0 (0%)	331 (100%)	39 (100%)	51 (100%)	118 (100%)	767 (80%)	0 (0%)	1306 (73%)
Baseline haemoglobin level (g/dL)	11·0 (8·8, 12·4)	11·9 (10·8, 12·8)	14·3 (13·1, 14·7)	10·3 (9·5, 11·2)	13·8 (13·0, 14·5)	12·6 (11·4, 13·9)	11·6 (10·4, 13·1)	12·3 (11·0, 13·7)
Number of missing data	2	0	0	0	0	0	0	2
Day 7 methaemoglobin level (%)	NA	3·8 (2·7, 5·7)	5·6 (4·1, 7·1)	NA	5·8 (3·7, 8·2)	8·3 (5·4, 12)	NA	6·0 (3·5, 9·4)
Number of missing data	139	28	1	51	0	536	156	911
Transmission intensity								
Low	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Moderate	0 (0%)	331 (100%)	0 (0%)	0 (0%)	0 (0%)	963 (100%)	156 (100%)	1450 (81%)
High	139 (100%)	0 (0%)	39 (100%)	51 (100%)	118 (100%)	0 (0%)	0 (0%)	347 (19%)
Origin of infections								
Within Papua	139 (100%)	0 (0%)	39 (100%)	0 (0%)	118 (100%)	0 (0%)	156 (100%)	452 (25%)
Outside Papua	0 (0%)	331 (100%)	0 (0%)	51 (100%)	0 (0%)	963 (100%)	0 (0%)	1345 (75%)
Study duration								
At least 180 days	0 (0%)	331 (100%)	39 (100%)	0 (0%)	118 (100%)	963 (100%)	156 (100%)	1607 (89%)
Less than 180 days	139 (100%)	0 (0%)	0 (0%)	51 (100%)	0 (0%)	0 (0%)	0 (0%)	190 (11%)

525 Numbers are in median (first quartile [Q1], third quartile [Q3]) or frequency (percentage). NA, not available; kg, kilogram; μL, microlitre; g, gram; dL, decilitre; mg, milligram. Transmission intensity was categorised as low (<1 case per
526 1000 person-years), moderate (1 case to <10 cases per 1000 person-years), and high (≥10 cases per 1000 person-years) according to subnational malaria incidence estimates for the median year of study enrolment.[48] Patients
527 contributing to the tolerability and safety datasets, as well as any sensitivity or restricted analyses, were subsets of the dataset shown in this table.
528

531 **Figure 2. Study sites that contributed to the pooled individual patient data across Indonesia**

532 Each circle represents a study site, and the number of participants included in the pooled data. Primaquine total dose: no primaquine (0 mg/kg), low (≥2 and <5 mg/kg), high
533 (≥5 mg/kg). Basemap shapefile data were obtained from the publicly available Natural Earth project (https://www.naturalearthdata.com) and accessed via the open-source R
534 packages maps[49] and ggplot2.[50] * Malaria-naive soldier patients were deployed and acquired infections in Papua then returned to malaria-free Java for follow-up (also referred
535 to as the soldier relapse model[10,12]).

537
538

Figure 3. Effect of increasing primaquine total dose on the rate of _P. vivax_ recurrence by day 180

(A) The reference value (HR = 1) was set at the target low dose of 3·5 mg/kg. The shaded region shows 95% CI. Estimates were derived from a multivariable Cox proportional hazards model, fitted to the primary efficacy dataset (n = 1797). The histogram along the top margin shows the distribution of primaquine total doses, with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale. (B) Crude incidence rates in the subset of patients (n = 1607) followed for up to 180 days across multiple _P. vivax_ episodes by primaquine total dose and origin of infection. Patients treated without primaquine were only available from outside Papua. Point estimates (solid squares) were calculated as the number of _P. vivax_ episodes divided by person-years at risk, with 95% CI for a Poisson rate obtained using an exact method. The horizontal axis is shown on a logarithmic scale.

551
552
Figure 4. Effect of primaquine daily dose on haematological profile

(A) Maximum absolute change from baseline in haemoglobin levels between days 1 and 14 across primaquine
daily doses, sex, and G6PD activity levels. Estimates were derived from a multivariable linear model fitted to the
primary safety dataset (n = 822, 34 patients with intermediate G6PD activity between 30% and 69%). The plot
shows model-implied predictions at four selected cut-offs (30%, 50%, 70%, and 100%) of G6PD activity to aid
interpretation. The histogram along the top margin shows the distribution of primaquine daily doses in the model
data, with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). (B)
Temporal dynamics of haematological recovery in patients with G6PD activity of at least 70%. Estimates were
derived from a multilevel, multivariable linear model (n = 790). Model-implied predictions at primaquine daily
doses of 0·25 mg/kg and 0·5 mg/kg are shown. Solid curves represent the expected haemoglobin levels over time
following treatment, with thick and thin shaded regions denoting the 95% confidence and prediction intervals,
respectively. AMM, adjusted male median.

Figure 5. Meta-analysis of randomised controlled trials and its comparison to global and Indonesian IPD meta-analyses on the effect of high-dose versus low-dose primaquine

(A) Systematic search for trials that randomised high- and low-dose primaquine was conducted without country restrictions from 1960 to 2025. Estimates from the first four studies[33-36] were derived from the number of events and the period at risk reported in each study to calculate the incidence rate ratio (patients censored before the end of the study were assumed to contribute half of their respective follow-up period). Estimates from the last three studies[30-32] were obtained directly from the hazard ratio reported in each study. In the current context, these effect measures (i.e., the incidence rate ratio and hazard ratio) were assumed to be approximately interchangeable and summarised here as the risk ratio. Vertical dashed line indicates an equal effect (RR = 1). Estimates to the left of the dashed line indicate greater efficacy of high dose primaquine. Estimates to the right indicate greater efficacy of low dose primaquine. (B) Pooled randomised estimates were derived from a study-

level global meta-analysis presented in the top panel. Estimates from Commons 2023[6] were derived from an IPD
meta-analysis of randomised and non-randomised *P. vivax* efficacy studies conducted globally. Estimates from
the current study were derived by comparing the hazard ratio estimate at 7 mg/kg with that at 3·5 mg/kg in the
Indonesian pooled data. The solid square represents a point estimate with a 95% confidence interval. The
horizontal axis is shown on a logarithmic scale. CI, confidence interval. IPD, individual patient data. (C) Expected
number needed to benefit with total high-dose primaquine (compared with total low-dose primaquine, assuming
a risk ratio of 0.53) across varying absolute risks of recurrence over a one-year period.

Supantary Material

List S1. Systematic search terms for the databases

Vivax AND (artefenomel OR arterolane OR amodiaquine OR atovaquone OR artemisinin OR arteether OR artesunate OR artemether OR artemotil OR azithromycin OR artekin OR chloroquine OR chlorproguanil OR cycloguanil OR clindamycin OR coartem OR dapsone OR dihydroartemisinin OR duo-cotecxin OR doxycycline OR halofantrine OR lumefantrine OR lariam OR malarone OR mefloquine OR naphthoquine OR naphthoquinone OR piperaquine OR primaquine OR proguanil OR pyrimethamine OR pyronaridine OR proguanil OR quinidine OR quinine OR riamet OR sulphadoxine OR tetracycline OR tafenoquine)

 List S2. Supplementary methods

<u>**Additional inclusion criteria**</u>

For the haematological safety analysis, only patients with G6PD activity ≥30% of the adjusted
male median (AMM) or a negative qualitative test were included. Reporting of parasite
presence or absence during follow-up, presence of gastrointestinal discomfort (vomiting,
anorexia, diarrhoea) during follow-up, and haemoglobin measured on day 0 and at least once
during follow up were required for anti-relapse efficacy, gastrointestinal tolerability, and
haematological safety analyses, respectively. For Poespoprodjo 2022,[11] we included only
patients for whom actual dose data were available. Protocol-based dose calculation was
considered unreliable in the context of unsupervised primaquine.
655
656 <u>**Categories of total and daily primaquine doses**</u>

Category	Weight-adjusted primaquine exposure (mg/kg)	
	Daily	Total
No primaquine	0	0
Very low	…	>0 and <2
Low	>0 and <0·375	≥2 and <5
Intermediate	≥0·375 and <0·75	…
High	≥0·75	≥5

657
658 <u>**Data analysis: anti-relapse efficacy**</u>

659 Variables violating proportional hazards assumptions were adjusted accordingly. Patients were
660 censored at recurrence, new antimalarial treatment, loss to follow-up, study end, outcome
661 assessment, or a smear gap >67 days.
662
To detect residual confounding, we used negative control methods,[29] using *P. falciparum*
recurrence and patient body weight (see **Figure S2** for the rationale). Sensitivity
analyses restricted the event of interest to symptomatic *P. vivax* recurrent parasitaemia and
excluded unsupervised regimens, as well as data from the sole cluster-randomised trial. In
studies with ≥180 days follow-up that tracked patients through multiple episodes of *P. vivax*
parasitaemia, the incidence of recurrence was also analysed using Poisson regression. Rates
were calculated as episodes per person-year, with censoring consistent with the Cox model.
670
<u>**Data analysis: haematological safety**</u>

The analysis of absolute fall in haemoglobin level was repeated for endpoints on days 2–3 and
5–7. Among patients with normal G6PD activity and baseline haemoglobin ≥11 g/dL, risk of
anaemia by days 2–3 was modelled with Poisson regression using the same covariates,
excluding G6PD level. Haematological recovery in this G6PD group was modelled with a
multilevel linear model using time, daily primaquine dose, baseline haemoglobin, age, sex, log-
transformed baseline parasite density, study site, and patient-specific intercepts. Splines and
interaction terms were used for time and dose. The effect of daily primaquine dose on the day
7 methaemoglobin concentration and the risk of clinical methaemoglobinaemia (i.e., 10% or
above) between days 1 and 14 was modelled using multivariable linear and Poisson regression,
respectively; including age (years), sex (male or female), baseline parasite density (parasites

per μL of blood; natural log transformed), and study site as covariates. We fitted this model by
incorporating a nonlinear effect using a restricted cubic spline.
684
685

   Table S1. PRISMA-IPD checklist

PRISMA-IPD Section/topic	Item No	Checklist item	Reported on page
Title			
Title	1	Identify the report as a systematic review and meta-analysis of individual participant data.	1
Abstract			
Structured summary	2	Provide a structured summary including as applicable:	2
		Background: state research question and main objectives, with information on participants, interventions, comparators and outcomes.	
		Methods: report eligibility criteria; data sources including dates of last bibliographic search or elicitation, noting that IPD were sought; methods of assessing risk of bias.	
		Results: provide number and type of studies and participants identified and number (%) obtained; summary effect estimates for main outcomes (benefits and harms) with confidence intervals and measures of statistical heterogeneity. Describe the direction and size of summary effects in terms meaningful to those who would put findings into practice.	
		Discussion: state main strengths and limitations of the evidence, general interpretation of the results and any important implications.	
		Other: report primary funding source, registration number and registry name for the systematic review and IPD meta-analysis.	
Introduction			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of the questions being addressed with reference, as applicable, to participants, interventions, comparisons, outcomes and study design (PICOS). Include any hypotheses that relate to particular types of participant-level subgroups.	4
Methods			
Protocol and registration	5	Indicate if a protocol exists and where it can be accessed. If available, provide registration information including registration number and registry name. Provide publication details, if applicable.	5
Eligibility criteria	6	Specify inclusion and exclusion criteria including those relating to participants, interventions, comparisons, outcomes, study design and characteristics (e.g. years when conducted, required minimum follow-up). Note whether these were applied at the study or individual level i.e. whether eligible participants were included	5

		(and ineligible participants excluded) from a study that included a wider population than specified by the review inclusion criteria. The rationale for criteria should be stated.	
Identifying studies - information sources	7	Describe all methods of identifying published and unpublished studies including, as applicable: which bibliographic databases were searched with dates of coverage; details of any hand searching including of conference proceedings; use of study registers and agency or company databases; contact with the original research team and experts in the field; open adverts and surveys. Give the date of last search or elicitation.	5
Identifying studies - search	8	Present the full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	5
Study selection processes	9	State the process for determining which studies were eligible for inclusion.	5
Data collection processes	10	Describe how IPD were requested, collected and managed, including any processes for querying and confirming data with investigators. If IPD were not sought from any eligible study, the reason for this should be stated (for each such study).	5
		If applicable, describe how any studies for which IPD were not available were dealt with. This should include whether, how and what aggregate data were sought or extracted from study reports and publications (such as extracting data independently in duplicate) and any processes for obtaining and confirming these data with investigators.	
Data items	11	Describe how the information and variables to be collected were chosen. List and define all study level and participant level data that were sought, including baseline and follow-up information. If applicable, describe methods of standardising or translating variables within the IPD datasets to ensure common scales or measurements across studies.	5
IPD integrity	A1	Describe what aspects of IPD were subject to data checking (such as sequence generation, data consistency and completeness, baseline imbalance) and how this was done.	5
Risk of bias assessment in individual studies.	12	Describe methods used to assess risk of bias in the individual studies and whether this was applied separately for each outcome. If applicable, describe how findings of IPD checking were used to inform the assessment. Report if and how risk of bias assessment was used in any data synthesis.	7
Specification of outcomes and effect measures	13	State all treatment comparisons of interests. State all outcomes addressed and define them in detail. State whether they were pre-specified for the review and, if applicable, whether they were primary/main or secondary/additional outcomes. Give the principal measures of effect (such as risk ratio, hazard ratio, difference in means) used for each outcome.	5–6

Synthesis methods	14	Describe the meta-analysis methods used to synthesise IPD. Specify any statistical methods and models used. Issues should include (but are not restricted to): • Use of a one-stage or two-stage approach. • How effect estimates were generated separately within each study and combined across studies (where applicable). • Specification of one-stage models (where applicable) including how clustering of patients within studies was accounted for. • Use of fixed or random effects models and any other model assumptions, such as proportional hazards. • How (summary) survival curves were generated (where applicable). • Methods for quantifying statistical heterogeneity (such as I^2 and t^2). • How studies providing IPD and not providing IPD were analysed together (where applicable). • How missing data within the IPD were dealt with (where applicable).	5–6
Exploration of variation in effects	A2	If applicable, describe any methods used to explore variation in effects by study or participant level characteristics (such as estimation of interactions between effect and covariates). State all participant-level characteristics that were analysed as potential effect modifiers, and whether these were pre-specified.	5–6
Risk of bias across studies	15	Specify any assessment of risk of bias relating to the accumulated body of evidence, including any pertaining to not obtaining IPD for particular studies, outcomes or other variables.	8
Additional analyses	16	Describe methods of any additional analyses, including sensitivity analyses. State which of these were pre-specified.	5–6
Results			
Study selection and IPD obtained	17	Give numbers of studies screened, assessed for eligibility, and included in the systematic review with reasons for exclusions at each stage. Indicate the number of studies and participants for which IPD were sought and for which IPD were obtained. For those studies where IPD were not available, give the numbers of studies and participants for which aggregate data were available. Report reasons for non-availability of IPD. Include a flow diagram.	8
Study characteristics	18	For each study, present information on key study and participant characteristics (such as description of interventions, numbers of participants, demographic data, unavailability of outcomes, funding source, and if applicable duration of follow-up). Provide (main) citations for each study. Where applicable, also report similar study characteristics for any studies not providing IPD.	8
IPD integrity	A3	Report any important issues identified in checking IPD or state that there were none.	8

Risk of bias within studies	19	Present data on risk of bias assessments. If applicable, describe whether data checking led to the up-weighting or down-weighting of these assessments. Consider how any potential bias impacts on the robustness of meta-analysis conclusions.	8
Results of individual studies	20	For each comparison and for each main outcome (benefit or harm), for each individual study report the number of eligible participants for which data were obtained and show simple summary data for each intervention group (including, where applicable, the number of events), effect estimates and confidence intervals. These may be tabulated or included on a forest plot.	8
Results of syntheses	21	Present summary effects for each meta-analysis undertaken, including confidence intervals and measures of statistical heterogeneity. State whether the analysis was pre-specified, and report the numbers of studies and participants and, where applicable, the number of events on which it is based.	8–9
		When exploring variation in effects due to patient or study characteristics, present summary interaction estimates for each characteristic examined, including confidence intervals and measures of statistical heterogeneity. State whether the analysis was pre-specified. State whether any interaction is consistent across trials.	
		Provide a description of the direction and size of effect in terms meaningful to those who would put findings into practice.	
Risk of bias across studies	22	Present results of any assessment of risk of bias relating to the accumulated body of evidence, including any pertaining to the availability and representativeness of available studies, outcomes or other variables.	8
Additional analyses	23	Give results of any additional analyses (e.g. sensitivity analyses). If applicable, this should also include any analyses that incorporate aggregate data for studies that do not have IPD. If applicable, summarise the main meta-analysis results following the inclusion or exclusion of studies for which IPD were not available.	8–9
Discussion			
Summary of evidence	24	Summarise the main findings, including the strength of evidence for each main outcome.	10
Strengths and limitations	25	Discuss any important strengths and limitations of the evidence including the benefits of access to IPD and any limitations arising from IPD that were not available.	10–12
Conclusions	26	Provide a general interpretation of the findings in the context of other evidence.	10–12
Implications	A4	Consider relevance to key groups (such as policy makers, service providers and service users). Consider implications for future research.	10–12

Funding			
Funding	27	Describe sources of funding and other support (such as supply of IPD), and the role in the systematic review of those providing such support.	7

Table S2. Risk of bias assessment

Study	Analysis	Domain based on ROB2 for randomised trials					Overall
		Randomisation	Intervention	Outcome		Reported results	
				Missingness	Measurement		
Taylor 2019	Efficacy	Low	Low	Low	Low	..	Low
	Tolerability	Low	Low	Low	Low	..	Low
	Safety	Low	Low	Low	Low	..	Low

Study	Analysis	Domain based on ROBINS-I for non-randomised studies of interventions							Overall
		Confounding	Intervention classification	Selection	Intervention deviation	Missingness	Outcome measurement	Reported results	
Hasugian 2007	Efficacy	Low	Moderate	Low	Low	Low	Low	..	Moderate
	Tolerability	Low	Moderate	Low	Low	Moderate	Low	..	Moderate
	Safety	..	..	..	..	..	..	..	..
Pasaribu 2013	Efficacy	Low	Moderate	Low	Low	Low	Low	..	Moderate
	Tolerability	Low	Moderate	Low	Low	Moderate	Low	..	Moderate
	Safety	..	..	..	..	..	..	..	..
Sutanto 2013	Efficacy	Low	Low	Low	Low	Low	Low	..	Low
	Tolerability	..	..	..	..	..	..	..	..
	Safety	..	..	..	..	..	..	..	..
Lidia 2015	Efficacy	Low	Moderate	Low	Low	Low	Low	..	Moderate
	Tolerability	..	..	..	..	..	..	..	..
	Safety	..	..	..	..	..	..	..	..
Nelwan 2015	Efficacy	Low	Low	Low	Low	Low	Low	..	Low
	Tolerability	..	..	..	..	..	..	..	..
	Safety	..	..	..	..	..	..	..	..
Poespoprodjo 2022	Efficacy	Low	Moderate	Low	Low	Low	Low	..	Moderate
	Tolerability	Low	Moderate	Low	Low	Moderate	Low	..	Moderate
	Safety	..	..	..	..	..	..	..	..

690
691 .. (not applicable in this IPD or not eligible in the specified analysis)

693
694

Figure S1. Causal directed acyclic graph showing the relationship between the two factors of interest (red nodes): primaquine mg/kg total dose and the risk of *P. vivax* relapse

Under this causal model, confounding by age, sex, and study site is present. Age, sex, or study site itself is a proxy for other host (body weight, immunity to malaria, patient behaviours, CYP2D6 polymorphisms) and geographical or aetiological (strain, blood-stage drug resistance) factors.

703
704

705
706

Figure S2. Causal directed acyclic graphs illustrating the use of *P. falciparum* infection and body weight as negative controls to detect residual confounding

A. <u>Outcome negative control</u>: The presence of *P. falciparum* re-infection in patients treated for *P. vivax* mono-infection during follow-up in areas with local transmission should be independent of the total primaquine dose received, provided that measured confounding is adequately adjusted for. However, if substantial residual confounding is not sufficiently controlled (either due to inadequate adjustment of measured confounding or the presence of unmeasured confounding), a spurious causal effect of primaquine dose on the risk of *P. falciparum* infection may become apparent.

B. <u>Exposure negative control</u>: Among patients with *P. vivax* who received no primaquine (thereby excluding the essential anti-hypnozoite component of treatment) the risk of *P. vivax* relapse should be independent of the total primaquine dose received, provided that measured confounding is adequately adjusted for. However, if substantial residual confounding is not sufficiently controlled (either due to inadequate adjustment of measured confounding or the presence of unmeasured confounding) a spurious causal effect of body weight on the risk of *P. vivax* relapse may become apparent.

A node within a square represents an adjustment, either through model stratification or restriction in our analysis. A dotted line indicates that two nodes may be related due to some common causes.

724
725

Figure S3. Causal directed acyclic graph showing the relationship between the two factors of interest (red nodes): primaquine mg/kg daily dose and the risk of gastrointestinal (GI) discomfort

Under this causal model, confounding by age, sex, and study site is present. Age, sex, or study site itself is a proxy for other host (body weight, patient behaviours, CYP2D6 polymorphisms) and geographical factors.

731
732

Figure S4. Causal directed acyclic graph showing the relationship between the two factors of interest (red nodes): primaquine mg/kg daily dose and haemolysis

Under this causal model, confounding by age, sex, and study site is present. Age, sex, or study site itself is a proxy for other host (body weight, baseline haemoglobin, patient behaviours, CYP2D6 polymorphisms) and geographical factors.

740
741
742 **Figure S5. Distribution of primaquine total dose by primaquine duration**

743 In the 14-day primaquine regimen, the observed two peaks reflect the targeted total primaquine dose of 3·5 and
744 7 mg base per kg body weight.

747 **Figure S6. Distribution of primaquine daily dose by primaquine regimen**

Table S3. Patient characteristics by primaquine total dose group

Characteristic	Primaquine total dose			Overall
	No primaquine 0 mg/kg	Low dose 2 to <5 mg/kg	High dose ≥5 mg/kg	
Number of patients	196	464	1137	1797
Median age (years)	16 (9, 30)	18 (11, 31)	16 (9, 29)	17 (10, 29)
Age group				
Less than 5 years	13 (6·6%)	50 (11%)	109 (9·6%)	172 (9·6%)
5 to 14 years	76 (39%)	136 (29%)	393 (35%)	605 (34%)
At least 15 years	107 (55%)	278 (60%)	635 (56%)	1020 (57%)
Male sex	111 (57%)	258 (56%)	687 (60%)	1056 (59%)
Body weight (kg)	45 (23, 56)	45 (26, 54)	46 (23, 59)	45 (24, 57)
Baseline parasite density (per µL)	3831 (1200, 9543)	1160 (320, 4130)	3344 (830, 7574)	2793 (630, 7130)
Presence or recent history of fever	171 (87%)	448 (97%)	904 (89%)	1523 (91%)
Number of missing data	0	1	119	120
Schizontocidal drug				
Dihydroartemisinin-Piperaquine	196 (100%)	248 (53%)	988 (87%)	1432 (80%)
Artesunate-Amodiaquine	0 (0%)	196 (42%)	42 (3·7%)	238 (13%)
Artesunate-Pyronaridine	0 (0%)	1 (0·2%)	59 (5·2%)	60 (3·3%)
Chloroquine	0 (0%)	19 (4·1%)	9 (0·8%)	28 (1·6%)
Quinine	0 (0%)	0 (0%)	39 (3·4%)	39 (2·2%)
Primaquine total dose (mg base/kg)	0·00 (0·00, 0·00)	4·04 (3·50, 4·38)	7·21 (6·52, 8·12)	6·38 (4·12, 7·53)
Primaquine daily dose (mg base/kg)	0·00 (0·00, 0·00)	0·29 (0·25, 0·31)	0·58 (0·48, 0·96)	0·47 (0·28, 0·63)
Primaquine duration				
No primaquine	196 (100%)	0 (0%)	0 (0%)	196 (11%)
7 days	0 (0%)	5 (1·1%)	380 (33%)	385 (21%)
14 days	0 (0%)	459 (99%)	757 (67%)	1216 (68%)
Primaquine dose calculation				
No primaquine	196 (100%)	0 (0%)	0 (0%)	196 (11%)
Actual dosing	0 (0%)	42 (9·1%)	1037 (91%)	1079 (60%)
Protocol dosing	0 (0%)	422 (91%)	100 (8·8%)	522 (29%)
Primaquine supervision				
No primaquine	196 (100%)	0 (0%)	0 (0%)	196 (11%)
Unsupervised	0 (0%)	129 (28%)	77 (6·8%)	206 (11%)
Partially supervised	0 (0%)	9 (1·9%)	80 (7·0%)	89 (5·0%)
Fully supervised	0 (0%)	326 (70%)	980 (86%)	1306 (73%)
Baseline haemoglobin level (g/dL)	12·75 (11·60, 14·10)	11·50 (10·30, 12·70)	12·60 (11·30, 14·00)	12·30 (11·00, 13·70)
Number of missing data	0	2	0	2
Day 7 methaemoglobin level (%)	NA	3·9 (2·7, 5·9)	6·9 (4·2, 10·3)	6·0 (3·5, 9·4)
Number of missing data	196	213	502	911
Transmission intensity				
Low	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Moderate	196 (100%)	306 (66%)	948 (83%)	1450 (81%)
High	0 (0%)	158 (34%)	189 (17%)	347 (19%)
Origin of infections				
Within Papua	0 (0%)	139 (30%)	313 (28%)	452 (25%)
Outside Papua	196 (100%)	325 (70%)	824 (72%)	1345 (75%)
Study duration				
At least 180 days	196 (100%)	307 (66%)	1104 (97%)	1607 (89%)
Less than 180 days	0 (0%)	157 (34%)	33 (2·9%)	190 (11%)

754 **Table S4. Studies included in analysis**

Paper	Study site	Latitude	Longitude	Year start	Year end	MAP incidence rate (per 1000 person-years)	Transmission intensity#	Relapse periodicity§
Hasugian 2007[23]	Timika	−4·61	136·85	2005	2005	22.61	High	High
Pasaribu 2013[13]	Tanjung Leidong	2·77	99·98	2011	2011	2·75	Moderate	High
Sutanto 2013[12]	Lumajang	−8·13	113·22	2010	2011	36·88*	High	High
Lidia 2015[24]	Kupang	−10·18	123·61	2013	2013	15·22	High	High
Nelwan 2015[10]	Sragen	−7·42	111·02	2013	2013	42·44*	High	High
Taylor 2019[14]	Hanura	−5·53	105·24	2015	2017	1·01	Moderate	High
Taylor 2019[14]	Tanjung Leidong	2·77	99·98	2015	2017	1·03	Moderate	High
Poespoprodjo 2022[11]	Timika	−4·45	136·98	2015	2016	56·43	High	High

755
756 MAP malaria atlas project. § Relapse periodicity was categorised as high (median relapse periodicity of 47 days or less) and low (median relapse periodicity of more
757 than 47);[51] # Transmission intensity was categorised as low (<1 case per 1000 person-years), moderate (1 case to <10 cases per 1000 person-years), and high (≥10 cases
758 per 1000 person-years) according to subnational malaria incidence estimates for the median year of study enrolment.[48] * Based on the location where patients were
759 infected by *P. vivax* in Papua.
760

761 **Table S5. Studies that were eligible for analysis but not included for data pooling**

Characteristic	Study		
	Maguire et al.[26]*	Arcelia et al.[27]	Sutanto et al.[25]
Year published	2006	2023	2023
Number of treatment arms	2	1	3
Region origin of infection	Papua	Outside Papua	Papua
Number of sites	1	1	2
Follow-up (days)	28	28	180
Randomised	Yes	No	Yes
Recruitment period	1996–1999	2019–2020	2018–2019
Treatment arms	(1) Chloroquine + 14-day, low dose primaquine (2) Mefloquine + 14-day, low dose primaquine	(1) Dihydroartemisinin-Piperaquine + 14-day, low dose primaquine	(1) Dihydroartemisinin-Piperaquine (2) Dihydroartemisinin-Piperaquine + 14-day, low dose primaquine (3) Dihydroartemisinin-Piperaquine + Tafenoquine
P. vivax patients enrolled	125*	60	150
Treated with primaquine	125*	60	50
Primaquine supervision	NA	NA	Yes
Sex (primaquine receiving arm)	68% Male*	NA	100% Male
Age (in years, primaquine receiving arm)	Mean = 22·3*	30 Children (2 to 18 years) 30 Adults	Mean = 28·6
Reason for exclusion	Missing minimum data	Data not available by 28 February 2025	Data not available by 28 February 2025

762 * Based on contributed IPD that may include both randomised trials and prospective cohorts not part of the trial
763 itself. NA, not available; IPD, individual patient data.
764

765 **Table S6. Comparison of patient characteristics who received primaquine between**
766 **included studies and eligible but not available studies**

Characteristic	Included studies (n = 7)	Eligible but not included studies (n = 3)
Region origin of infection, studies (percentage)		
Papua	2 (28·6%)	2 (66·7%)
Outside Papua	5 (71·4%)	1 (33·3%)
Year of enrolment, studies (percentage)		
Pre-2015	5 (71·4%)	1 (33·3%)
2015-2019	2 (28·6%)	2 (66·7%)
Follow up duration in days, studies (percentage)		
42	2 (28·6%)	2 (66·7%)
>42 to <120	0 (0%)	0 (0%)
120	0 (0%)	0 (0%)
>120	5 (71·4%)	1 (33·3%)
Age (weighted-average years)[#,$]	16·9[*]	25·7[§]
Male (weighted percentage)[#]	59·6%[*]	85%[§]

767
768 # Weights approximated by the numbers of *P. vivax* patients treated with primaquine. * Based on included
769 participants in this pooled analysis (n = 1797). $ Study-specific average is the mean or median in vivax patients
770 treated with primaquine. § Based on two studies from Table S5 for which the required summary statistics were
771 available.

Figure S7. Primaquine total dose received across different patient age

A. While the median average across the three age groups was similar, patients under five years old were more likely to receive a higher total dose (relative to a target dose) than older patients, as also depicted in panel B. In B, each dot represents an individual patient. The red curve denotes a fitted regression line between age and primaquine total dose. Dotted line represents the target dose of 3·5 or 7 mg base per kg body weight. This may reflect the relatively greater difficulty of administering primaquine in very young patients. For panel B, the horizontal axis is shown on a logarithmic scale.

Figure S8. Primaquine total dose by schizontocidal drug (A) and study (B)

Figure S9. Primaquine total dose by body weight in patients receiving primaquine

787 Dotted line represents the total dose targets of 3·5 or 7 mg/kg. For Hasugian 2007 and Lidia 2015, the follow up duration was less than 180 days. Event status indicates whether
788 the patient developed a *P. vivax* recurrence. Horizontal and vertical axes are shown on a logarithmic scale.

790
791 **Figure S10. Distributions of the days to first *P. vivax* recurrence by study**

792 For Hasugian 2007 and Lidia 2015, the dotted line represents the end of active follow up. The other studies
793 monitored patients for at least 180 days.

Figure S11. *P. vivax* recurrence at day 180 by different groups

By (A) clinical total dose group in the primary efficacy dataset, n = 1797; (B) Clinical total dose group excluding the cluster randomised trial,[11] n = 1641; (C) Decile with 10 equal sized groups in the primary efficacy dataset, n = 1797; and (D) Excluding two studies[13,24] due to the limited number of events, n = 1415. The reference value, at

which the hazard ratio equals one, was set at 0 mg/kg. The whisker shows 95% confidence intervals. Estimates
were derived from a multivariable Cox proportional hazards model. The vertical axis for A–D is shown on a
logarithmic scale. Based on the model underlying panel (A), panels (E) and (F) display the model-implied relapse-
free probabilities in different total dose groups, separately as examples in Papua and Non-Papua settings,
respectively. To estimate these survival probabilities, continuous covariates were set at their median values, and
sex was set to male.

809
810

812
813
814

815
816

Figure S12. Sensitivity analyses of the hazard ratio across primaquine total dose

(A) Using symptomatic *P. vivax* recurrences as the endpoint event, n = 1794; (B) Excluding the cluster randomised trial,[11] n = 1641; (C) Excluding unsupervised primaquine treatment, n = 1591; (D) Estimating the incidence rate ratio in patients followed for multiple *P. vivax* episodes, n = 1607. The reference value, at which the hazard ratio equals one, was set at the target low dose of 3·5 mg/kg. The shaded region shows 95% confidence intervals. Estimates were derived from a multivariable Cox proportional hazards model (A, B, C) or a multivariable Poisson model (D). A restricted cubic spline with three knots was specified on primaquine total dose to allow for a non-linear trend. The histogram along the top margin shows the distribution of primaquine daily doses in the model data, with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale.

827
828

List S3. Negative control results

Negative control methods for exposure (AHR of *P. vivax* recurrence associated with a 1 kg increase in body weight in patients treated without primaquine = 1.1; 95% CI 0.98 to 1.03; p = 0.60) and outcome (AHR of *P. falciparum* infection associated with a 1 mg/kg increase in primaquine total dose = 1.03; 95% CI 0.94 to 1.14; p = 0.48) suggest that residual confounding in our anti-relapse effect estimates was minimal.

836
837

838
839

Figure S13. Estimated hazard ratios of *P. vivax* recurrence at day 180, assuming effect
modification by (A) origin of infection and (B) age category

The reference value, at which the hazard ratio equals one, was set at the target low dose of 3·5 mg/kg. The shaded
region shows 95% confidence intervals. Estimates were derived from a multivariable Cox proportional hazards
model. A restricted cubic spline with three knots was specified on primaquine total dose to allow for a non-linear
trend. The histogram along the top margin shows the distribution of primaquine daily doses in the model data (n
= 1797), with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). The
vertical axis is shown on a logarithmic scale.

Gastrointestinal discomfort on days 5−7 (%)

Figure S14. Heterogeneity in percentages of gastrointestinal discomfort on days 5–7 by study and primaquine daily dose

This variability may stem from the relatively subjective nature of assessing this endpoint. The solid square and horizontal line represent the point and 95% confidence interval estimates.

Table S7. Pooled counts and percentages of patients who experienced gastrointestinal discomfort

Primaquine daily dose	Days of observation		
	Baseline	1–2	5–7
No primaquine	0/0 (···)	0/0 (···)	6/190 (3·2%)
Low	9/335 (2·7%)	46/335 (13·7%)	38/203 (18·7%)
Intermediate	309/584 (52·9%)	192/583 (32·9%)	45/482 (9·3%)
High	298/376 (79·3%)	189/376 (50·3%)	53/379 (14·0%)
Total	616/1295 (47·6%)	427/1294 (33·0%)	142/1254 (11·3%)

865
866

867
868

Figure S15. Risk ratio of gastrointestinal discomfort (A) between days 5 to 7, (B) between
days 1 to 2, and (C) at day 0 [baseline]

(A) The reference value (RR = 1) was set at the primaquine daily dose of 0 mg/kg. Estimates were derived from a
multivariable Poisson model (n = 952). Two different specifications for primaquine daily dose were shown to
evaluate potential non-linearity across doses: linear trend (solid grey line with shaded region for 95% CI) and
restricted cubic spline with three knots (dashed curve with two dotted curves representing the 95% CI limits).
Inclusion of study site (p = 0.7) in a sensitivity analysis did not alter the effect-estimate for dose. The secondary y-
axis displays the estimated absolute risk of gastrointestinal discomfort as a percentage. These values are derived
by scaling the model-predicted risk ratios by the baseline risk. The histogram along the top margin shows the
distribution of primaquine daily doses in the model data, with the leftmost bar representing patients who were
treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale.
880
(B, C) The reference value, where the risk ratio equals one, was set at the primaquine daily dose of 0·5 mg/kg.
Estimates were derived from a multivariable Poisson model (n in panel A = 767, n in panel B = 767). Two different
specifications for primaquine daily dose were shown to evaluate potential non-linearity across doses: trend (solid
grey line with shaded region for 95% CI) and restricted cubic spline with three knots (dashed curve with two
dotted curves representing the 95% CI limits). The histogram along the top margin shows the distribution of
primaquine daily doses in the model data, with the leftmost bar representing patients who were treated without
primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale. The y-axis is displayed up to a risk
ratio of 16 to enable comparison with the primary gastrointestinal tolerability endpoint.

Table S8. Pooled counts and percentages of patients who experienced acute vomiting within 1 hour of taking primaquine between days 0 and 14

Primaquine daily dose	Days of observation
	0–14
Low	0/206 (0%)
Intermediate	15/642 (2·3%)
High	14/380 (3·7%)
Total	29/1228 (2·4%)

Figure S16. Risk ratio of acute vomiting within 1 hour of taking primaquine between days 0 and 14

The reference value, where the risk ratio equals one, was set at the primaquine daily dose of 0·5 mg/kg. Estimates were derived from a multivariable Poisson model (n = 767). Two different specifications for primaquine daily dose were shown to evaluate potential non-linearity across doses: trend (solid grey line with shaded region for 95% CI) and restricted cubic spline with three knots (dashed curve with two dotted curves representing the 95% CI limits). The histogram along the top margin shows the distribution of primaquine daily doses in the model data. The vertical axis is shown on a logarithmic scale.

Table S9. Poisson model estimates with robust standard errors

Model	RR (95% robust CI)
Tolerability	
Effect of increasing 0·25 mg/kg daily dose on GI discomfort (days 5–7)	1·32 (1·17 to 1·48)
Effect of increasing 0·25 mg/kg daily dose on GI discomfort (day 0)	0·99 (0·96 to 1·02)
Effect of increasing 0·25 mg/kg daily dose on GI discomfort (days 1–2)	1·03 (0·97 to 1·10)
Effect of increasing 0·25 mg/kg daily dose on acute vomiting	1·19 (0·84 to 1·67)
Safety	
Effect of increasing 0·25 mg/kg daily dose on clinical anaemia	0·90 (0·80 to 1·01)

907 Figure S17. (A) Haemoglobin concentrations and (B) G6PD activity levels at patient
908 enrolment in the primary safety dataset (n = 822)

910 Table S10. Haematological adverse events on days 1–14, by dose group

Haematological adverse event	Primaquine daily dose			
	No primaquine	Low	Intermediate	High
Haemoglobin fall to <5 g/dL	0/173	0/6	0/318	0/325
Haemoglobin fall >5 g/dL from baseline	0/173	0/6	0/318	0/325
Blood transfusion	0/173	0/6	0/318	0/325
Renal failure requiring dialysis	0/173	0/6	0/318	0/325
Death	0/173	0/6	0/318	0/325

Figure S18. Maximum absolute change from baseline in haemoglobin levels (A) between days 2 and 3 and (B) between days 5 and 7 across primaquine daily doses, sex, and G6PD activity levels

Estimates were derived from a multivariable linear model fitted to the subsets of the primary safety dataset (n in panel A = 794, n in panel B = 781). An interaction term was included in the model to account for differential treatment effects of primaquine daily dose as sex and G6PD activity vary. A restricted cubic spline was specified for G6PD activity to model a non-linear trend. Primaquine daily dose and G6PD activity were both modelled as continuous covariates. The plot shows model-implied predictions only at four selected cut-offs (30%, 50%, 70%, and 100%) of G6PD activity to aid interpretation. The histogram along the top margin shows the distribution of primaquine daily doses in the model data, with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale. AMM, adjusted male median.

926
927
928

929
930
Figure S19. Risk ratio of developing anaemia (haemoglobin <11 g/dL) between days 2 to 3
in patients with G6PD activity of at least 70% and had baseline haemoglobin level of at
least 11 g/dL

The reference value, where the risk ratio equals one, was set at the primaquine daily dose of 0 mg/kg. Estimates
were derived from a multivariable Poisson model, fitted to a subset of the primary tolerability dataset. Two
different specifications for primaquine daily dose were shown to evaluate potential non-linearity across doses:
linear trend (solid grey line with shaded region for 95% CI) and restricted cubic spline with three knots (dashed
curve with two dotted curves representing the 95% CI limits). The histogram along the top margin shows the
distribution of primaquine daily doses in the model data (n = 612) , with the leftmost bar representing patients
who were treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale.
941
942

Figure S20. Effect of primaquine daily dose on methaemoglobin levels

(A) Effect of primaquine daily dose on day 7 methaemoglobin levels (n = 886), modelled using multivariable linear regression, adjusting for age, sex, baseline parasite density, and study site. A restricted cubic spline with three knots was applied to primaquine daily dose. The solid curve represents the expected day 7 methaemoglobin levels across the range of primaquine daily doses. The thick and thin shaded regions represent the 95% confidence intervals and prediction intervals, respectively. Each dot represents an individual patient's day 7 methaemoglobin level. The dotted horizontal lines have y-intercepts of 10% and 20%, thresholds at which symptoms related to tissue hypoxia (e.g., light-headedness, tachycardia) may become more apparent. (B) Risk ratios of clinical methaemoglobinaemia (i.e., 10% or more) from day 1 to 14 across primaquine daily doses. Estimates were modelled using a multivariable Poisson model with a restricted cubic spline with three knots applied to primaquine daily dose. The solid curve and shaded region represent the point estimates and 95% confidence intervals, respectively. The histogram along the top margin shows the distribution of primaquine daily doses in the model data (n = 1026) , with the leftmost bar representing patients who were treated without primaquine (i.e., 0 mg/kg). The vertical axis is shown on a logarithmic scale.

References

1. Price RN, Commons RJ, Battle KE, Thriemer K, Mendis K. *Plasmodium vivax* in the era of the shrinking *P. falciparum* map. Trends Parasitol 2020; 36(6): 560-70.
2. Kementerian Kesehatan Republik Indonesia. Laporan situasi terkini perkembangan program pengendalian malaria di Indonesia tahun 2024, 2025.
3. Djaafara BA, Sherrard-Smith E, Churcher TS, et al. Spatiotemporal heterogeneity in malaria transmission across Indonesia: analysis of routine surveillance data 2010–2019. BMC Med 2025; 23(1): 136.
4. Fadilah I, Djaafara BA, Lestari KD, et al. Quantifying spatial heterogeneity of malaria in the endemic Papua region of Indonesia: Analysis of epidemiological surveillance data. Lancet Reg Health Southeast Asia 2022; 5.
5. Phyo AP, Dahal P, Mayxay M, Ashley EA. Clinical impact of vivax malaria: a collection review. PLoS Med 2022; 19(1): e1003890.
6. Commons RJ, Rajasekhar M, Edler P, et al. Effect of primaquine dose on the risk of recurrence in patients with uncomplicated *Plasmodium vivax*: a systematic review and individual patient data meta-analysis. Lancet Infect Dis 2023.
7. Rajasekhar M, Simpson JA, Ley B, et al. Primaquine dose and the risk of haemolysis in patients with uncomplicated *Plasmodium vivax* malaria: a systematic review and individual patient data meta-analysis. Lancet Infect Dis 2023.
8. World Health Organization. WHO guidelines for malaria, 30 November 2024, 2024.
9. Kementerian Kesehatan Republik Indonesia. Buku saku tatalaksana kasus malaria; 2023.
10. Nelwan EJ, Ekawati LL, Tjahjono B, et al. Randomized trial of primaquine hypnozoitocidal efficacy when administered with artemisinin-combined blood schizontocides for radical cure of Plasmodium vivax in Indonesia. BMC Med 2015; 13: 294.
11. Poespoprodjo JR, Burdam FH, Candrawati F, et al. Supervised versus unsupervised primaquine radical cure for the treatment of falciparum and vivax malaria in Papua, Indonesia: a cluster-randomised, controlled, open-label superiority trial. Lancet Infect Dis 2022; 22(3): 367-76.
12. Sutanto I, Tjahjono B, Basri H, et al. Randomized, Open-Label Trial of Primaquine against Vivax Malaria Relapse in Indonesia. Antimicrob Agents Chemother 2013; 57(3): 1128-35.
13. Pasaribu AP, Chokejindachai W, Sirivichayakul C, et al. A randomized comparison of dihydroartemisinin-piperaquine and artesunate-amodiaquine combined with primaquine for radical treatment of vivax malaria in Sumatera, Indonesia. J Infect Dis 2013; 208(11): 1906-13.
14. Taylor WRJ, Thriemer K, Von Seidlein L, et al. Short-course primaquine for the radical cure of *Plasmodium vivax* malaria: a multicentre, randomised, placebo-controlled non-inferiority trial. Lancet 2019; 394(10202): 929-38.
15. Commons RJ, Thriemer K, Humphreys G, et al. The Vivax Surveyor: Online mapping database for *Plasmodium vivax* clinical trials. Int J Parasitol Drugs Drug Resist 2017; 7(2): 181-90.
16. Infectious Diseases Data Observatory (IDDO). IDDO SDTM implementation manual. 2023. https://www.iddo.org/tools-and-resources/data-tools (accessed accessed: 01/04/2024.

17. Stewart LA, Clarke M, Rovers M, et al. Preferred Reporting Items for Systematic Review and Meta-Analyses of individual participant data: the PRISMA-IPD Statement. JAMA 2015; 313(16): 1657-65.

18. World Health Organization. Technical consultation to review the classification of glucose-6-phosphate dehydrogenase (G6PD), 2022.

19. Zou G. A modified poisson regression approach to prospective studies with binary data. American journal of epidemiology 2004; 159(7): 702-6.

20. Sterne JA, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ 2019; 366.

21. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. BMJ 2016; 355.

22. WorldWide Antimalarial Resistance Network (WWARN). Primaquine Indonesia study group. 2023. https://www.iddo.org/primaquine-indonesia-study-groupaccessed: 27/02/2025).

23. Hasugian A, Purba H, Kenangalem E, et al. Dihydroartemisinin-piperaquine versus artesunate-amodiaquine: superior efficacy and posttreatment prophylaxis against multidrug-resistant *Plasmodium falciparum* and *Plasmodium vivax* malaria. Clin Infect Dis 2007; 44(8): 1067-74.

24. Lidia K, Dwiprahasto I, Kristin E. Therapeutic Effects of Dyhidroartemisinin Piperaquine Versus Chloroquine for Uncomplicated Vivax Malaria in Kupang, East Nusa Tenggara, Indonesia. Age 2015; 13: 50.

25. Sutanto I, Soebandrio A, Ekawati LL, et al. Tafenoquine co-administered with dihydroartemisinin-piperaquine for the radical cure of *Plasmodium vivax* malaria (INSPECTOR): a randomised, placebo-controlled, efficacy and safety study. Lancet Infect Dis 2023.

26. Maguire JD, Krisin, Marwoto H, Richie TL, Fryauff DJ, Baird JK. Mefloquine is highly efficacious against chloroquine-resistant *Plasmodium vivax* malaria and *Plasmodium falciparum* malaria in Papua, Indonesia. Clin Infect Dis 2006; 42(8): 1067-72.

27. Arcelia F, Pasaribu AP, Yanni GN. Effectiveness of dihydroartemisinin-piperaquine after 10 years as treatment for vivax malaria in Indonesia. J Infect Dev Ctries 2023; 17(05): 700-6.

28. Chu CS, Bancone G, Moore KA, et al. Haemolysis in G6PD heterozygous females treated with primaquine for *Plasmodium vivax* malaria: a nested cohort in a trial of radical curative regimens. PLoS Med 2017; 14(2): e1002224.

29. Lipsitch M, Tchetgen ET, Cohen T. Negative controls: a tool for detecting confounding and bias in observational studies. Epidemiol 2010; 21(3): 383-8.

30. Chamma-Siqueira NN, Negreiros SC, Ballard S-B, et al. Higher-dose primaquine to prevent relapse of *Plasmodium vivax* malaria. N Engl J Med 2022; 386(13): 1244-53.

31. EFFORT Trial Investigators. EFFORT Trial: Unpublished results. American Society of Tropical Medicine Annual Meeting 2024; 2024.

32. Eng V, Lek D, Sin S, et al. 14 days of high-dose versus low-dose primaquine treatment in patients with *Plasmodium vivax* infection in Cambodia: a randomised, single-centre, open-label efficacy study. Lancet Infect Dis 2025.

33. Krudsood S, Tangpukdee N, Wilairatana P, et al. High-dose primaquine regimens against relapse of *Plasmodium vivax* malaria. Am J Trop Med Hyg 2008; 78(5): 736.

34. Pukrittayakamee S, Imwong M, Chotivanich K, Singhasivanon P, Day NP, White NJ. A
comparison of two short-course primaquine regimens for the treatment and radical cure
of *Plasmodium vivax* malaria in Thailand. Am J Trop Med Hyg 2010; 82(4): 542.
35. Rajgor D, Gogtay N, Kadam V, et al. Antirelapse efficacy of various primaquine regimens
for *Plasmodium vivax*. Malar Res Treat 2014; 2014(1): 347018.
36. Saravu K, Tellapragada C, Kulavalli S, et al. A pilot randomized controlled trial to compare
the effectiveness of two 14-day primaquine regimens for the radical cure of vivax malaria
in South India. Malar J 2018; 17: 1-11.
37. Collins WE, Jeffery GM. Primaquine resistance in *Plasmodium vivax*. Am J Trop Med Hyg
1996; 55(3): 243-9.
38. Garrison PL, Hankey DD, Coker WG, et al. 2. CURE OF KOREAN VIVAX MALARIA WITH
PAMAQUINE AND PRIMAQUINE. J Am Med Assoc 1952; 149(17): 1562-3.
39. White NJ. Determinants of relapse periodicity in *Plasmodium vivax* malaria. Malar J 2011;
10(1): 297.
40. Mehdipour P, Rajasekhar M, Dini S, et al. Effect of adherence to primaquine on the risk
of *Plasmodium vivax* recurrence: a WorldWide Antimalarial Resistance Network
systematic review and individual patient data meta-analysis. Clin Infect Dis 2023; 22(1):
306.
41. Fryauff D, Baird K, Basri H, et al. Randomised placebo-controlled trial of primaquine for
prophylaxis of falciparum and vivax malaria. Lancet 1995; 346(8984): 1190-3.
42. Clayman CB, Arnold J, Hockwald RS, Yount EH, Edgcomb JH, Alving AS. 3. Toxicity of
primaquine in caucasians. J Am Med Assoc 1952; 149(17): 1563-8.
43. Baird JK, Hoffman SL. Primaquine therapy for malaria. Clin Infect Dis 2004; 39(9): 1336-
45.
44. Fadilah I, Commons RJ, Chau NH, et al. Methaemoglobin as a surrogate marker of
primaquine antihypnozoite activity in Plasmodium vivax malaria: A systematic review and
individual patient data meta-analysis. PLoS Med 2024; 21(9): e1004411.
45. White NJ, Watson JA, Baird JK. Methaemoglobinaemia and the radical curative efficacy
of 8-aminoquinoline antimalarials. Br J Clin Pharmacol 2022; 88(6): 2657-64.
46. Watson JA, Commons RJ, Tarning J, et al. The clinical pharmacology of tafenoquine in the
radical cure of *Plasmodium vivax* malaria: An individual patient data meta-analysis. Elife
2022; 11.
47. Commons RJ, Simpson JA, Watson J, White NJ, Price RN. Estimating the Proportion of
Plasmodium vivax Recurrences Caused by Relapse: A Systematic Review and Meta-
Analysis. Am J Trop Med Hyg 2020; 103(3): 1094-9.
48. Battle KE, Lucas TCD, Nguyen M, et al. Mapping the global endemicity and clinical burden
of *Plasmodium vivax*, 2000–17: a spatial and temporal modelling study. Lancet 2019;
394(10195): 332-43.
49. Becker RAW, A. R.; Brownrigg, R.; Minka, T. P.; Deckmyn, A. maps: Draw Geographical
Maps. 3.4.1 ed: R package; 2023.
50. Wickham HC, W.; Henry, L.; Pedersen, T. L.; Takahashi, K.; Wilke, C.; Woo, K.; Yutani, H.
ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics. 3.5.1 ed: R
package; 2023.
51. Battle KE, Karhunen MS, Bhatt S, et al. Geographical variation in *Plasmodium vivax*
relapse. Malar J 2014; 13(1): 144.
1094