Cochlear Place Specificity of the Auditory Brainstem Response to Narrowband Chirp versus 2-1-2 stimuli: High Pass Noise/Derived Responses.

Ronald Nkansah Adjekum[1]

David R. Stapells[1]

[1]School of Audiology & Speech Sciences, The University of British Columbia,

2177 Wesbrook Mall, Vancouver, BC CANADA V6T 1Z3

Financial disclosures/conflicts of interest

This study was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant to Susan Small (RGPIN-2018-04655). Financial support for R.N.A. was provided by: (i) a Four-Year Doctoral Fellowship from the University of British Columbia (UBC), and (ii) the UBC School of Audiology and Speech Sciences. There are no conflicts of interest, financial, or otherwise.

All correspondence should be addressed to:

David R. Stapells, School of Audiology & Speech Sciences, The University of British Columbia, 2177 Wesbrook Mall, Vancouver, BC CANADA V6T 1Z3
E-mail: stapells@audiospeech.ubc.ca

Author contribution statement

R.N.A. and D.R.S. designed the experimental procedures and analyses. R.N.A. recruited

participants and ran the experiments, analyzed the ABR waveforms and managed results'

databases. R.N.A. and D.R.S. analyzed and discussed the research findings and wrote the article

content.

Ethical approval

Ethics approval was obtained from the University of British Columbia Clinical Research Ethics

Board.

Acknowledgements

This study was carried-out in Dr. Susan Small's laboratory. A strong supporter of this research,

due to her illness and subsequent death, Dr. Small was unable to participate in the planning or

completion of this study. This study was supported by a Natural Sciences and Engineering

Research Council of Canada Discovery Grant to Susan Small (RGPIN-2018-04655). Financial

support for R.N.A. was provided by: (i) a Four-Year Doctoral Fellowship from the University of

British Columbia (UBC), and (ii) the UBC School of Audiology and Speech Sciences.

1   **ABSTRACT**

Objective: In recent years, many researchers have recommended using narrowband chirp

(NBchirp) stimuli for Auditory Brainstem Response (ABR) audiometry instead of more-standard

2-1-2 cycles linear-gated tones, primarily because NBchirps often result in larger ABR wave V

amplitudes. However, the acoustic frequency spectra of currently recommended NBchirps are

wider than those for 2-1-2 tones, and it is currently not known whether ABRs to these NBchirps

have similar (or poorer) cochlear place specificity compared to 2-1-2 tones. The current study

used the high-pass noise/derived response technique to assess the cochlear regions contributing

to ABRs evoked by NBchirp versus 2-1-2 stimuli.

Design: A total of 24 adults with normal hearing participated (N=12 for each stimulus

frequency). Stimuli were 60-dB peSPL 500- and 2000-Hz NBchirps and 2-1-2 tones mixed with

high-pass (HP) filtered masking noise. The level of broadband (pink) noise required to mask the

ABR was determined individually, then the broadband noise at this level was HP filtered at ½-

octave intervals. Three ABR replications were obtained for each condition, with recordings

stopped when the residual noise level of each replication was reduced to 40 nanovolts. Derived

responses (DRs) representing 1-octave-wide or ½-octave-wide cochlear regions were calculated

by subtracting ABRs recorded in HP noise.

Results: Non-masked ABR amplitudes in response to NBchirps were significantly larger than

those to 2-1-2 stimuli, averaging 55% larger for 500 Hz and 81% larger for 2000 Hz. For both

500- and 2000-Hz stimuli, HP noise masking produced significant amplitude decreases,

occurring 1 to ½ octave higher for ABRs to NBchirps versus 2-1-2 tones. One-octave-wide and

½-octave-wide DR amplitude profiles for the ABRs to 2-1-2 tones showed good cochlear place

specificity, as described in previous studies. DR results for the NBchirps were similar but

showed important differences. The profiles for the 2000-Hz NBchirps showed significantly

larger amplitudes in the 4- and 1-kHz DRs compared to the 2-1-2 stimuli. Many more responses

were seen 1-octave away for the 2000-Hz NBchirp compared to 2-1-2 tone. DR results for 500-

Hz tones showed similar patterns but differences did not quite reach statistical significance,

except amplitudes to NBchirps were larger at DR354, DR500 and DR707. A measure of the

width of the 1-octave-wide and ½-octave-wide DR amplitude profiles ($BW_{0.075}$, in Hz) showed

the 500- and 2000-Hz NBchirp profiles were significantly wider (32% to 77%) compared to

those for 2-1-2 stimuli. As the cochlear area able to respond decreased, wave V amplitudes to

NBchirp stimuli decreased more than those for 2-1-2 stimuli, with no difference between stimuli

for ½-octave-wide responses.

Conclusion: ABRs to narrowband chirps reflect wider cochlear contributions than those to 2-1-2

tones. Responses to NBchirps arise from cochlear regions as far as one octave away from the

stimulus frequency. In contrast, responses to 2-1-2 tones arise from cochlear regions primarily

within approximately ±0.5 octaves of the stimulus frequency. Further research in individuals

with hearing loss is required to determine whether the wider bandwidths for NBchirps result in

threshold mis-estimations, and whether NBchirp amplitude advantages over more-standard

stimuli remain with hearing loss.

1   **INTRODUCTION**

2   For several decades, researchers studying auditory brainstem response (ABR) audiometry

have investigated the best choice of stimuli to estimate frequency-specific behavioral thresholds

rapidly and accurately in patients who cannot be tested with conventional pure-tone audiometry.

Many ABR studies have demonstrated that 2-1-2 (cycles) tones and other brief tones with similar

duration are good choices for predicting the behavioral pure-tone audiogram using the ABR (for

review: Small & Stapells 2017). However, many have noted concerns about 2-1-2 and other

similar brief tones, suggesting that ABR thresholds to these stimuli are too high and/or they lack

frequency specificity (e.g., Scherg & Volk 1983; Sohmer & Kinarti 1984; Rodrigues et al. 2013;

Laukli 2014; Ferm & Lightfoot 2015; Leusin Mattiazzi et al. 2024). In recent years, some have

recommended using a newer stimulus, narrowband chirps (NBchirp), for threshold ABRs instead

of 2-1-2 cycles linear-gated tones (Elberling & Don 2010), primarily because NBchirps often,

though not always, result in larger ABR wave V amplitudes (Cobb & Stuart 2016; Ferm &

Lightfoot 2015; Dzulkarnain et al. 2018; Pinto et al. 2022). NBchirps are brief tonal stimuli

uprising in frequency which attempt to compensate for temporal dispersion along the cochlear

partition by presenting the low-frequency components within the stimulus before the high-

frequency components, according to cochlear travelling-wave delay models. Theoretically, if

delay models are correct, aligning the arrival time of each frequency component in the stimulus

to its place of maximum excitation along the basilar membrane should produce larger amplitude

responses (Elberling et al. 2007; Gøtsche-Rasmussen et al. 2012).

21   NBchirps at four frequencies (500, 1000, 2000, and 4000 Hz) derived from the broadband

CE-Chirp® (level independent) developed by Elberling and colleagues are currently the most

researched (and clinically utilized) NBchirp stimuli [1]; they were constructed based on a delay

model originally determined from derived-band ABR latencies (Elberling et al. 2007; Elberling

& Don 2008). Based on the research cited above, these NBchirp stimuli are expected to elicit

significantly larger ABR wave V amplitudes (compared to standard 5-cycle tonal stimuli).

Nevertheless, it has also been argued that a "1-octave" wide region of the basilar membrane may

be too small an area for the NBchirp to result in a significant increase in the ABR amplitude

(Wegner & Dau 2002).

Most of the studies advocating the use of NBchirps have focussed on amplitude

advantages over 2-1-2 stimuli, which would lead to improved signal-to-noise ratio, making ABR

recordings faster and responses easier to detect (e.g., Bell et al. 2002; Ferm et al. 2013; Ferm &

Lightfoot 2015; Cobb & Stuart 2016; Dzulkarnain et al. 2018; Talaat et al. 2019; Pinto et al.

2022). Nonetheless, the amplitude advantage of NBchirps is not consistently present for all

stimulus frequencies and/or intensities (e.g., Bal 2022; Cobb & Stuart 2016; Dzulkarnain et al.

2018; Rodrigues et al. 2013; Megha et al. 2019; Ceylan et al. 2025). Some of the inconsistencies

may be related to delay model issues. Most of these studies have been carried-out in adults with

normal hearing; few have assessed NBchirp/standard tone ABR amplitude differences in either

infants or individuals with hearing loss. More study is required.

---

[1] The current NB CE-Chirp® LS narrowband chirps are based on the level-independent broadband CE-Chirp® (as opposed to the level-specific broadband CE-Chirp® LS). NB CE-Chirp® LS narrowband chirps are level-specific only in their onset timings are adjusted such that wave V latencies should be about the same independent of intensity. NB CE-Chirp® LS narrowband chirp durations and spectra are not affected by intensity (Adjekum et al. 2024).

Previous studies have noted that the acoustic spectra of NBchirps appear wider than those

of the 2-1-2 standard brief-tone stimuli (Bell et al. 2002; Cobb & Stuart 2016; Adjekum et al.

2024). Although current NBchirps are "octave band" stimuli (i.e., their spectrum is 1-octave-

wide when measured at -3 dB), Adjekum and colleagues recently argued that bandwidth

measured at -20 dB is a more realistic measure of the effective bandwidth of 2-1-2 or NBchirp

spectra. NBchirp spectra bandwidths, measured at -20 dB, are 1.5 octaves, compared to 0.9

octaves for 2-1-2 stimuli (Adjekum et al. 2024). However, it is currently not known whether

ABRs to these NBchirps have worse (or similar) cochlear place specificity compared to 2-1-2

tones. Because NBchirps have wider frequency spectra, it is likely the frequency and place

specificity of ABRs to these stimuli will differ from 2-1-2 tones and may not be acceptable for

clinical use. In other words, the ABRs to NBchirps may include contributions from unwanted

frequency places (Bell et al. 2002; Adjekum et al. 2024). The wider spectral content of NBchirps

compared to 2-1-2 stimuli may also explain the larger ABR amplitudes to these stimuli.

The high-pass noise/derived response (HPDR) technique has proven to be a useful

method for quantifying cochlear contributions to an ABR to different stimuli. Teas, Eldridge, and

Davis (1962) were the first to employ the HPDR subtraction technique to record auditory evoked

potentials in animals. Later, Elberling (1974) and Eggermont (1976) employed this technique for

electrocochleographic recordings in humans; Don and Eggermont (1978), as well as Parker and

Thornton (1978) used this technique to assess the ABR in humans. The HPDR technique is

applied by first recording ABRs in quiet and in the presence of broadband masking noise at an

effective masking level, and then steeply high-pass (HP) filtering the noise at this intensity. The

subtraction of responses recorded in HP noise with a lower cutoff frequency from responses

recorded in HP noise with a higher cutoff frequency yield "derived responses" (DRs) that reflect

the responses to stimulation of the basilar membrane approximately between the two cutoff

frequencies (Don & Eggermont 1978; Parker & Thornton 1978; Picton et al. 1981; Evans &

Elberling 1982; Stapells & Fok 2023). In a recent "masking within the derived band" study,

Stapells and Fok (2023) confirmed the validity of the ABR HPDR method, with their results

indicating 1-octave-wide separations in the HP noise settings resulted in responses reflecting 1-

octave-wide bands with center frequencies closest to the lower HP cutoff in the HPDR

calculation. The HPDR technique has proven to be an extremely valuable tool for investigation

of mechanisms underlying the ABR, and, to a lesser extent, for clinical investigations. Most

relevant to the current study, by assessing amplitudes of DRs representing narrow frequency

bands in response to a stimulus, the HPDR technique provides a direct assessment of the

cochlear contributions (i.e., "cochlear place specificity") to the auditory evoked potential by that

stimulus (Kramer 1992; Nousak & Stapells 1992; Oates & Stapells 1997b; Herdman et al. 2002;

Stoll & Maddox 2024)

Several studies using the HPDR technique have shown good place specificity for the

ABR to 2-1-2 tones and other 5-cycle tones (such as exact-Blackman windowed) (Kramer 1992;

Nousak & Stapells 1992; Oates & Stapells 1997b). These studies have demonstrated ABR

amplitude profiles showing (i) derived response (DR) response maxima at or within ±0.5 octave

of the nominal stimulus frequency, indicating a place-specific response from the cochlea, (ii)

sharp drop-offs above and below the DR frequency with maximum response amplitude,

indicating little contribution from frequencies one or more octaves away from stimulus

frequency, and (iii) center frequency and bandwidth measures which indicate place-specific

contributions from the frequencies of interest. A similar assessment of cochlear place specificity

has not been investigated for the ABR to NBchirps.

The current study used the HPDR method to investigate whether the acoustic spectral

differences seen between NBchirp and 2-1-2 stimuli result in ABRs to the two stimuli showing

differing cochlear contributions; specifically, whether the wider NBchirp spectra result in wider

cochlear contributions to the ABR.

8   **MATERIALS AND METHODS**

Subjects

Twenty-four adults (N=12 per 500- and 2000-Hz stimulus frequency) with normal

hearing were recruited for the study. The participants had no significant otologic or neurologic

histories and were between the ages of 18 to 40 years. Subjects were assigned to groups

randomly, without consideration of age or sex. Normal hearing required a pure-tone behavioral

threshold of 15 dB HL (ANSI, 2023) or better at the octave frequencies of 250, 500, 1000, 2000,

4000, and 8000 Hz in the test ear. Tympanometry (226-Hz probe tone) was used to assess

middle-ear function. A normal tympanogram was defined as that for which peak compensated

static admittance, tympanogram width, and tympanometric peak pressure suggested normal

middle-ear pressure and tympanic membrane mobility (i.e., Type A tympanogram). For the

ABR study, only one ear was tested (randomly selected between left and right ears).

Stimuli and Calibration

The 2-1-2 and NBchirp stimuli were initially recorded from the research version of an

Interacoustics Eclipse ABR system (EPx5 4.4 software). The Eclipse ABR system was set to

deliver NB CE-Chirp® LS and 2-1-2 stimuli at 100 dB pe SPL. The stimulus electrical output

was fed directly to Sigview software (version 3.2) using a digitization rate of 44,100 Hz and a

95-ms analysis window. An average of 100 time-domain samples was obtained for each stimulus

and then edited to include only the stimulus portion. These Sigview-recorded stimuli were then

converted by an Intelligent Hearing Systems (IHS) SmartEP system to the SmartEP STM

stimulus format (16-bit resolution at a rate of 40000 Hz). Figure 1 shows the acoustic spectra of

these NBchirp and 2-1-2 stimuli. As previously demonstrated (Cobb & Stuart 2016; Adjekum et

al. 2024), the NBchirp spectra are 70-80% wider (at -20 dB) than the 2-1-2 spectra. Comparisons

of spectra of the stimuli presented by the IHS SmartEP and the Interacoustics Eclipse showed no

difference between the stimuli (for both NB CE-Chirp® LS and 2-1-2 stimuli).

11
12   < INSERT FIGURE 1 ABOUT HERE >
13

During ABR recordings, 500- and 2000-Hz stimuli (NBchirp and 2-1-2) were presented

monaurally at 60 dB pe SPL via Etymotic Research ER-3A insert earphones. This level is

equivalent to 40 (500 Hz) and 37 (2000 Hz) dB nHL for NBchirps (Gøtsche-Rasmussen et al.

2012)[2] and to 38-41 (500 Hz) and 38-40 (2000 Hz) dB nHL for 2-1-2 stimuli (Fedtke & Richter

2007; Bagatto, 2020; Hatton et al. 2022). Stimuli were presented at a rate of 38.9/s with

alternating stimulus polarity. Acoustic calibrations of the 60 dB pe SPL NBchirp and 2-1-2

stimuli were checked before each test using a GRAS RA0113 2-cc coupler and a 1-inch

---

[2] After correction for the use of different couplers: IEC 60126 2-cc vs IEC 60711 (Haughton 2006)

microphone (Larson Davis, Model 2575) and a PRM902 preamplifier with a Larsen Davis

System 824 sound level meter.

Broadband and High-Pass Noise Masking

Broadband pink noise (General Radio Company 1382 random-noise generator) was HP

filtered using a cascade of two Stanford Research Systems (model SR650, each 115dB/octave

slope) and then attenuated. HP noise cutoffs were selected to provide ½-octave separations. For

500-Hz stimuli, cutoffs were 8000, 4000, 2830, 2000, 1410, 1000, 707, 500, 354, and 250 Hz.

For 2000-Hz stimuli, cutoffs were 16000, 8000, 5660, 4000, 2830, 2000, 1410, 1000, 707, and

500 Hz.[3]

Effective masker levels using broadband pink noise (HP filters both set at 0.001 Hz) were

determined for each participant (and each stimulus). Initially, the levels of masking noise

required to completely mask the stimuli behaviorally were obtained. Subsequently, levels

required to mask the ABR were obtained starting at 5 dB above the behavioral masking level and

then increasing the masker intensity using 1-dB steps until the ABR was determined to be

masked, then a "down-2-dB-up-1-dB" bracketing method was used to establish the effective

masking level/threshold. Once determined for a participant and stimulus, the effective masking

level obtained for broadband pink noise was maintained for all ABR conditions with HP noise.

---

[3] The ER-3A earphone response output is limited to frequencies below about 10,000 Hz (Etymotic Research, 2025; Elberling et al. 2012). ER-2 earphones have wider bandwidth and were initially considered for this study. However, ER-3A earphones were finally chosen as they are the earphones most used clinically. The "HP16000" HP noise condition was kept for the study with the possibility it might provide additional insight concerning HP noise filter characteristics.

Mean (±SD) behavioral masking levels for the 500-Hz stimuli were 64±2.1 dB SPL for

NBchirps and 62±1.6 dB SPL for 2-1-2 stimuli. Behavioral masking levels for the 2000-Hz

stimuli were 63 ±3.2 dB SPL for NBchirps and 62±1.3 dB SPL for 2-1-2 stimuli. Mean (±SD)

ABR effective masking levels for 500-Hz stimuli were 77±2.5 dB SPL for NBchirps and 75±2.8

dB SPL for 2-1-2 stimuli. ABR effective masking levels for 2000-Hz stimuli were 76±2.7 dB

SPL for NBchirps and 73±3.7 dB SPL for 2-1-2 stimuli.

All ABR recordings were obtained using the IHS SmartEP evoked potential system.

Gold-plated electrodes were placed on the vertex (Cz) and mastoid (M1 or M2, ipsilateral to

stimulated ear), with a low forehead (Fpz) electrode acting as ground. Interelectrode impedances

were 3000 Ohms or less. The EEG recordings were amplified (100,000x), filtered (50-1500 Hz,

12 dB/octave), and digitized over a time window of 24.7 ms using a sampling rate of 20,000 Hz.

An online residual noise (RN) calculation was calculated over a 10-ms window that included the

wave V-V' response but excluded stimulus artifact: 10.5-20.5 ms for 500 Hz and 6.5-16.5 ms for

2000 Hz. Individual recordings were continued until the RN was equivalent to ≤40 nanovolts

(denoted as 0.080 µV by the SmartEP; Hatton et al. 2022).

Three replications of ABR waveforms, each with a RN ≤40 nV, when averaged together

give an overall ABR waveform with a RN of approximately 23 nV. Because the derived-

response subtraction procedure increases the noise in the ABR waveform by the square root of 2,

the RN of the DRs (average of 3 replications) will be approximately 33 nV, which allows us to

assess wave V-V' responses with peak-to-peak amplitudes as low as 0.07 µV - 0.10 µV (i.e., 2-

3x the RN) (e.g., Picton et al. 1983; Picton & Hink 1974; Don & Elberling 1996).

Procedure

2   For most participants (22 of 24), this study involved one recording session of

approximately 3 hours; 2 participants required two sessions to complete the study. Prior to

conducting the experiments, ethics approval was obtained from our university's Clinical

Research Ethics Board. During ABR testing, participants were seated in a recliner chair and

encouraged to relax and sleep. All testing was conducted in a double-walled sound-treated booth.

Each participant was tested with the NBchirp and 2-1-2 stimuli in quiet, in the presence of

broadband pink noise, and nine HP masking noise conditions. Each participant was tested for

only one stimulus frequency using both stimuli. The order of NBchirp versus 2-1-2 stimuli and

HP noise cutoff frequencies was randomized for each participant.

Derived Responses

Derived responses were obtained by subtracting, in succession, ABRs to the stimulus in

HP noise at one cutoff frequency from the responses to the same stimulus in HP noise with a

cutoff frequency either 1-octave or ½-octave higher. Thus, DRs for 1-octave- and ½-octave -

wide analyses were obtained at each test frequency. The subtractions for the 1-octave and ½-

octave DRs each resulted in 8 separate DRs, respectively. Calculations of the center frequencies

of derived noise bands, formed by subtracting HP noise spectra using 1-octave and ½-octave HP

noise separations, indicate that the center frequencies of these noise bands are closest to the

frequency of the lower HP noise used in the DR subtractions.[4] Recently, Stapells and Fok

(2023) presented masked-ABR results which support the use of the lower HP noise cutoff to

designate the center frequency of 1-octave-wide DRs. We therefore designated the lower

frequency of the two HP noise cutoff frequencies used in the subtraction procedure as the DR

center frequency, similar to previous studies by Stapells and colleagues (Picton et al. 1981;

Nousak & Stapells 1992; Oates & Stapells 1997b; Herdman et al. 2002) and others (e.g., Don et

al. 1979; Eggermont & Don 1980; Kramer 1992).

Response Identification and Measurement

Three judges experienced in tone-ABR recordings determined the presence or absence of

ABR Wave V, with agreement required by at least two judges. ABR waveforms for each

condition/subject were organized into waveform "sets," with each set including (i) superimposed

replications (usually three, but sometimes up to five replications when not all replications had

RN $\leq$ 40 nV), (ii) a "grand average" of the replications, and (iii) the "split buffers" showing two

superimposed waveforms, each comprised of half the total trials. All three judges reviewed all

results for stimulus frequency (500 or 2000 Hz), type (NBchirp vs 2-1-2 stimuli), the HP noise

cutoff frequency, and center frequencies of the DRs. Results for the actual study showed

reasonable agreement between the judges (Fleiss' Kappa: 0.74), with response presence/absence

agreement between all three judges of 81% across all conditions.

---

[4] The calculations of center frequencies of the derived (i.e., subtracted) noise bands, as well as the nominal and measured HP noise cutoffs, are presented in Supplemental Data Content 1.

ABR wave V amplitudes were measured for each subject for all conditions with present

responses. If no response was present (as determined by the judges), then the waveform's

overall RN (for the average of all replications) was entered as the amplitude. Wave V amplitude

was measured from wave V to the following wave V' negativity. Wave V was defined as the

maximum vertex-positive peak occurring between 6 and 20 ms after stimulus onset; in the case

of multiple peaks of equal amplitude occurring between 6 and 20 ms, the peak that occurs before

the largest negativity shift was selected. Wave V' was defined as the largest negative point

occurring within 8 ms following wave V (Stapells & Picton 1981; Oates & Stapells 1997a; Oates

& Stapells 1997b).

From the DR amplitude profiles, we also determined the bandwidth (in Hz) for a wave V

amplitude at 0.075 µV ($BW_{0.075}$), a level that represents a near-threshold response for both

stimuli. Bandwidths were calculated by determining the frequency for a wave V-V' amplitude of

0.075 µV on either side of the DR amplitude profile (using interpolation between frequencies),

with the difference (in Hz) between the low and high frequencies calculated as bandwidth, and

the geometric mean of these two frequencies calculated as the profile's centre frequency.

Supplemental Digital Content 2 provides a schematic example of one subject's $BW_{0.075}$

calculation.

Statistical Analyses

The percentage of subjects showing present responses for NBchirp and 2-1-2 stimuli was

calculated for each test condition. The percent of present ABRs for NBchirp versus 2-1-2 stimuli

were compared using chi-square tests for DR results at ±1 octave away from the stimulus

frequency.[5] ABR wave V amplitudes to NBchirp and 2-1-2 stimuli in each condition (no-mask,

HP Noise, 1-octave DR, and ½-octave DR) were summarized using means and standard

deviations (SD). Wave V latencies were not assessed in the present study.

Dependent samples t-tests were calculated to analyze wave V amplitudes to NBchirp

versus 2-1-2 tones in the no-mask condition. Repeated-measures ANOVAs were carried out on

the HP noise and DR results. Two-way repeated-measures ANOVAs were calculated separately

for the 500- and 2000-Hz stimuli to examine the effect of HP noise cutoff frequency on the

amplitudes of ABR wave V-V' to NBchirp versus 2-1-2 stimuli. Similarly, 2-way repeated-

measures ANOVAs were computed separately for the 500- and 2000-Hz stimuli to analyze the

effects of stimulus type and DR frequency on wave V amplitudes to NBchirp and 2-1-2 stimuli.

One-octave-wide and ½-octave-wide DR amplitude results were analyzed separately. DR

bandwidths were assessed using 2-way mixed-model ANOVAs (with stimulus frequency as a

between factor and stimulus type as a within factor) were also calculated for the 500-and 2000-

Hz 1-octave-wide bandwidths (in octaves), and for the ½-octave-wide DR bandwidths (in

octaves) to NBchirp versus 2-1-2 stimuli. Finally, the effect of limiting cochlear responding

bandwidth (CBW) on wave V amplitudes for the no-masking condition and the 1-octave-wide

and ½-octave-wide DR conditions corresponding to the stimulus frequencies (i.e., DR500 and

DR2000) were assessed using mixed-model ANOVA (STIMFRQ as a between-subject effect;

---

[5] Observing responses within ±0.5 octave of the stimulus frequency would likely not be a concern for audiometry. However, obtaining responses as far as 1-octave away from the stimulus frequency would be very problematic clinically. Beyond 1-octave away, there were too few responses for either stimulus.

STIM type and CBW as within-subjects effects), carried out for those participants with results

for all three CBW conditions.

Where appropriate, Huynh-Feldt epsilon corrections were applied to the ANOVA degrees

of freedom to compensate for any violations of sphericity (Huynh & Feldt 1976; Glass &

Hopkins 1996). Neuman-Keuls *post hoc* analyses were applied where appropriate to analyze

significant main effects and interactions in the ANOVA results (Glass & Hopkins 1996). With

the exception of chi-square tests, results of all statistical testing were considered significant if p <

.05. As there was a total of four chi-square tests carried-out on the percent-present results each

for 1-octave and ½-octave wide DRs, after Bonferroni correction, significant chi-square results

required p<.0125.

13   **RESULTS**

No-Mask Condition

The top of Figure 2 presents the grand-mean waveforms for the no-mask condition. Clear

waves V are seen in the waveforms in the no-mask conditions in response to both stimuli for

both frequencies. ABR waveforms for all participants for the No-Mask condition (as well as for

the 1-octave-wide and ½-octave-wide DRs representing the stimulus frequency) are presented in

Supplemental Digital Content 3. Mean (SD) wave V amplitudes to non-masked 500-Hz stimuli

were 0.19 (0.06) µV for NBchirps and 0.13 (0.04) µV for 2-1-2 stimuli. Mean amplitudes to non-

masked 2000-Hz stimuli were 0.27 (0.07) µV for NBchirps and 0.15 (0.04) µV for 2-1-2 stimuli.

The larger amplitudes in the no-mask condition for NBchirp compared to 2-1-2 stimuli were

statistically significant [500 Hz: $t(11) = 3.8$, p < .001; 2000 Hz: $t(11) = 9.53$, p < .001]. Almost

all participants showed larger amplitudes to NBchirps (500 Hz: 11/12; 2000 Hz: 12/12) with

mean NBchirp amplitude advantages being 55% for 500 Hz and 81% for 2000 Hz.

3
4   < INSERT FIGURE 2 ABOUT HERE >
5

High-Pass Noise

Figure 2 shows grand-mean waveforms across all subjects to the 500- and 2000-Hz

NBchirp and 2-1-2 stimuli for all 10 HP noise conditions (plus the no-mask condition). In

general, the grand-mean waveforms indicated (i) larger responses to NBchirps, at least until the

HP cutoff is within approximately ½ octave above the stimulus frequency, (ii) little change in

responses until HP noise cutoff was within approximately 1 octave of the stimulus frequency,

and (iii) large decrease in wave V amplitude once the HP noise cutoff was ½-octave above the

stimulus frequency and below, with grand-mean waveforms showing little-or-no response once

the HP noise cutoff is ½-octave below the stimulus frequency.

15
Figure 3A presents the percent of subjects showing present responses for each stimulus

and HP noise condition. For responses to 500-Hz stimuli, response presence remained high (92-

100%) for HP noise cutoffs down to 707 Hz, substantially decreasing (to 33-58%) for 500-Hz

HP noise, and to 8% (i.e., N=1 subject with response) or less below. Similar results were seen for

2000-Hz stimuli, except that response presence did not substantially decrease until the HP noise

cutoff was ½ octave *below* the stimulus frequency. No clear differences were seen in response

presence between the two stimuli at either stimulus frequency.

Figure 3B presents mean (and SD) amplitude results for the effects of HP cutoff on

responses to NBchirp and 2-1-2 stimuli for the two stimulus frequencies. In general, the

amplitude profiles suggested (i) larger amplitudes to NBchirp vs 2-1-2 stimuli, (ii) larger

responses to 2000-Hz vs 500-Hz stimuli (especially for NBchirps), and (iii) either no change or

only small decreases in amplitude until the HP noise cutoff reaches approximately ½ to 1 octave

above the stimulus frequency, with amplitudes then sharply decreasing below this. Two-way

repeated-measures ANOVAs were carried-out separately for the 500-Hz and 2000-Hz stimulus

frequencies. Results of these ANOVAs are shown at the top of Table 1.

< INSERT FIGURE 3 AND TABLE 1 ABOUT HERE >
8

The ANOVA results for each stimulus frequency indicates the significant main effect for

HP noise cutoff frequency (HPFRQ) seen in the grand-mean waves and amplitude graphs.

Similarly, the significant main effect for Stimulus Type (STIM) reflected the significantly larger

amplitudes in response to NBchirps. However, the significant interaction (HPFRQ x STIM)

indicated the above main effects may not hold across all conditions.

Neuman-Keuls *post hoc* analysis of the HPFRQ x STIM interaction for 500-Hz stimuli

indicated: (i) amplitudes for NBchirp stimuli were significantly larger than 2-1-2 stimuli only for

HP noise cutoffs of 1000 Hz and higher, (ii) amplitudes for NBchirp stimuli significantly

decreased between 1000- and 707-Hz HP noise, and between 707- and 500-Hz HP noise cutoffs,

whereas those for 2-1-2 stimuli only significantly decreased between 707- and 500-Hz cutoffs,

and (iii) amplitudes for NBchirps significantly decreased from no-mask to HP noise 8000 Hz, a

decrease which was not seen for 2-1-2 stimuli (with no significant decreases seen between 8000-

through 1000-Hz HP noise cutoffs).

For the 2000-Hz stimuli, Neuman-Keuls *post hoc* analysis of the HPFRQ x STIM

interaction indicated: (i) amplitudes for NBchirp stimuli were significantly larger than 2-1-2

stimuli only for HP noise cutoffs of 2000 Hz and higher (although statistically significant, the

amplitude differences between stimuli were very small at HP2000 and HP2830), (ii) amplitudes

for NBchirp stimuli significantly decreased between 5660- and 4000-Hz HP noise cutoffs,

between 4000- and 2830-Hz HP noise cutoffs, between 2830- and 2000-Hz HP noise cutoffs, and

between 2000- and 1410-Hz HP noise cutoffs, whereas those for the 2-1-2 stimuli only

significantly decreased between 2830- and 2000-Hz HP noise cutoffs, and between 2000- and

1410-Hz HP noise cutoffs, and (iii) amplitudes for NBchirps significantly decreased from the no-

mask to HP16000 conditions, a decrease which was not seen for 2-1-2 stimuli (with no

significant decreases seen between 16000- through 5660-Hz HP noise cutoffs). Overall, the

amplitude results with HP noise indicated responses to NBchirp stimuli were significantly

reduced with HP noise cutoffs 0.5 to 1 octave higher than those seen for 2-1-2 stimuli.

1-Octave-Wide Derived Responses

Figure 4 shows grand-mean waveforms (across all subjects) in response to the 500- and

2000-Hz NBchirp and 2-1-2 stimuli for all eight 1-octave-wide derived-response conditions. As

noted above, ABR waveforms for all participants for the 1-octave-wide DR500 and DR2000

conditions are presented in Supplemental Digital Content 3. These grand-mean waveforms

showed: (i) larger response amplitudes to NBchirps than to 2-1-2 stimuli, (ii) largest ABR

amplitudes to both NBchirp and 2-1-2 stimuli in the DR bands within a ½ octave of the stimulus

frequency, with amplitudes decreasing as DR frequency is further away from the stimulus

frequency, (iii) for responses to the 2000-Hz tones, NBchirp results showed a clear wave V in the

DR 1-octave above the stimulus frequency (i.e. DR4000), whereas there was no clear response

for the 2-1-2 stimuli in the DR4000 waveform (both stimuli showed responses for DR1000), and

(iv) for responses to the 500-Hz tones, differences between NBchirp and 2-1-2 stimuli were

smaller and less-obvious, at least in the grand-mean waveforms. The grand-mean waveforms for

500-Hz stimuli showed small responses to both stimuli at 1-octave above (i.e., DR1000) but not

1-octave below (DR250) the stimulus frequency.

< INSERT FIGURE 4 ABOUT HERE >

Figure 5A presents the percent of subjects showing present responses to the NBchirp and

2-1-2 stimuli in each 1-octave-wide derived-response condition. For responses to the 500-Hz

stimuli, response presence was 100% for the DR frequency representing stimulus frequency and

remained high (92-100%) for DR conditions ½-octave away from the center frequency, then

substantially decreased 1-octave away from the center frequency (DR250: 42% for NBchirp,

50% for 2-1-2; DR1000: 58% for NBchirp: 33% for 2-1-2). For responses to the 2000-Hz

stimuli, similar to results for 500-Hz stimuli, response presence was 100% for the DR frequency

representing stimulus frequency and remained high at DR conditions ½-octave away from the

center frequency. In contrast to results for 500-Hz stimuli, however, at 1-octave away from the

stimulus frequency (DR4000 and DR1000), DR results for 2000-Hz NBchirps continued to show

100% response presence, whereas DR results for 2000-Hz 2-1-2 stimuli showed substantially

reduced response presence (DR1000: 100% for NBchirp, 58% for 2-1-2; DR4000: 100% for

NBchirp: 25% for 2-1-2). The small differences in response presence one octave away for the

500-Hz stimuli did not reach statistical significance [DR250: χ^2 (1, N=12) = 0.17, p=0.682;

DR1000: χ^2 (1, N=12) = 1.52, p=0.219]; however, differences for 2000-Hz stimuli were

significant [DR1000: χ^2 (1, N=12) = 6.32, p=0.0120; DR4000: χ^2 (1, N=12) = 14.40, p<0.001],

with NBchirps showing significantly more responses present.

Figure 5B presents mean (and SD) amplitude results for the effects of 1-octave-wide DR

frequency on wave V amplitudes to NBchirp and 2-1-2 stimuli for the two stimulus frequencies.

In general, the amplitude profiles showed: (i) both NBchirp and 2-1-2 amplitude profiles had

largest mean amplitudes in the DR frequency representing the stimulus frequency, with mean

amplitudes steeply decreasing in DRs above and below the stimulus frequency, especially when

greater than ½-octave away, and (ii) NBchirp wave V amplitudes were greater than 2-1-2

amplitudes, especially within ½ octave of the stimulus frequency. Differences between stimuli

decreased for DR frequencies ≥1-octave away from the stimulus frequency. Two-way repeated

measures ANOVAs were carried-out separately for the 500- and 2000-Hz stimuli, with results

shown in the middle of Table 1.

< INSERT FIGURE 5 ABOUT HERE >
12

For 500-Hz stimuli, the significant main effect for DR frequency (DRFRQ) was

consistent with the results seen in the grand-mean waves (Figure 4) and amplitude profiles

(Figure 5B): response amplitudes for both stimuli were significantly larger for the DR closest to

the stimulus frequency and significantly decreased as DR frequency goes further away from 500

Hz. Although the main effect for stimulus type (STIM) was not significant, the significant

DRFRQ x STIM interaction indicated significant differences existed between the two two stimuli

in some conditions. Neuman-Keuls *post hoc* analysis of this interaction for 500-Hz stimuli

indicated significantly larger amplitudes in response to NBchirp compared to 2-1-2 stimuli for

DR707, DR500, and DR354 conditions, with no differences between stimuli beyond these

frequencies. Neuman-Keuls *post hoc* analysis also indicated response amplitudes for both stimuli

significantly decreased for DR frequencies above 707 Hz and below 354 Hz.

For 2000-Hz stimuli, the significant main effect for DR frequency was also consistent

with the results seen in the grand-mean waves (Figure 4) and amplitude profiles (Figure 5B):

response amplitudes were significantly larger for the DR closest to the stimulus frequency. As

with the 500-Hz results, amplitude decreased significantly as the DR frequency went further

away from 2000 Hz. The significant STIM (stimulus type) main effect indicated amplitudes of

DRs to NBchirps were larger than those to 2-1-2 stimuli; however, the significant

DRFRQxSTIM interaction indicated this was not be the case for all DR conditions. Neuman-

Keuls *post-hoc* analysis of the significant DRFRQ x STIM interaction indicated the amplitudes

in response to NBchirps were significantly larger than those to the 2-1-2 stimuli for DR4000,

DR2830, DR2000, DR1410, and DR1000 Hz DR conditions. Similar to the results for 500-Hz

stimuli, the Neuman-Keuls analyses for 2000-Hz stimuli also revealed that NBchirp amplitudes

showed significant decreases starting ½-octave above and below the stimulus frequency, whereas

amplitudes for 2-1-2 stimuli showed a significant decrease from DR2000 to DR1410 but only a

non-significant trend ($p < .1$) to a smaller amplitude from DR2000 to DR2830.

The NBchirp amplitude profiles in Figure 5B appeared to be wider than those for the

2-1-2 amplitude profiles. Bandwidths (and center frequencies) of each subject's 1-octave-wide

amplitude profiles were determined for a wave V-V' amplitude of 0.075 µV on either side of the

profile ($BW_{0.075}$). This was possible for all 12 subjects at 2000 Hz but not for one subject at 500

Hz, whose amplitudes for 2-1-2 stimuli did not allow us to determine a lower-side frequency for

0.075 µV. Bandwidth results for the 1-octave-wide DRs are shown at the top of Table 2. A

mixed-model ANOVA (stimulus frequency as a between effect; stimulus type as a within effect)

was calculated for the 500-and 2000-Hz 1-octave-wide DR bandwidths (converted to octaves)

for NBchirp versus 2-1-2 stimuli. The ANOVA results indicated bandwidths of the NBchirp

profiles were significantly wider than those for the 2-1-2 stimuli [STIM main effect: p=.328, F(1,

21) = 51.75, p < .001, η^2_p = 0.71]. Expressed in octaves, there was no significant difference

between the bandwidths of the profiles for 500- versus 2000-Hz stimuli [STIMFRQ main effect:

p=.328 F(1, 21) = 1.00, p = .328, η^2_p = 0.05]. There was a nonsignificant trend for NBchirp

bandwidths, which were larger than 2-1-2 stimuli for both frequencies, to be even larger for

2000-Hz versus 500-Hz stimuli [STIMFRQ x STIM interaction: F(1, 21) = 3.71, p = .068, η^2_p =

0.15]. Also shown in Table 2, the center frequencies of the 500- and 2000-Hz 1-octave-wide DR

amplitude profiles were very close to the stimulus frequency, with no obvious differences

between NBchirp and 2-1-2 stimuli.

< INSERT TABLE 2 ABOUT HERE >
11

½-Octave-Wide Derived Responses

To obtain a more frequency-specific evaluation of the contributions to the ABR by the

500- and 2000-Hz NBchirp and 2-1-2 stimuli, with better certainty of the center frequencies of

derived bands, DRs representing ½-octave-wide bands were calculated. One subject in each of

the 500- and 2000-Hz groups provided ½-octave DR results which were too difficult to interpret,

likely due to the poorer signal-to-noise ratio of these responses. Thus, the sample size for the ½-

octave results was 11 subjects per stimulus frequency. Figure 6 presents grand-mean waveforms

to the 500- and 2000-Hz NBchirp and 2-1-2 stimuli for all eight ½-octave-wide DR conditions.

[ABR waveforms for all participants for the ½-octave-wide DR500 and DR2000 conditions are

presented in in Supplemental Digital Content 3.] The ½-octave waveforms appeared noisier

because wave V amplitudes in the ½-octave-wide DRs were smaller compared to the 1-octave-

wide DRs (note the scale difference between Figures 4 and 6), which resulted in a smaller signal-

to-noise ratio. The decrease in amplitude was due, at least in part, to the ½-octave-wide DRs

originating from contributions of smaller portions of the cochlea. The residual electrical noise

(RN) in the waveforms, however, was approximately the same between the ½- and 1-octave-

wide DRs (on average, 31 vs 30 nV). In general, the grand-mean waveforms in Figure 6 were

consistent with the observations of the grand-mean waveforms for the 1-octave-wide DR

conditions (Figure 4). Specifically, the grand-mean ½-octave DR waveforms in Figure 6

indicated: (i) somewhat larger amplitudes in response to NBchirps than 2-1-2 stimuli, at least for

2000-Hz stimuli, (ii) the largest response amplitudes to both stimuli were within a half octave of

the stimulus frequency, (iii) for the 2000-Hz stimuli, the grand-mean waveform to NBchirps

showed a clear wave V response in the DR4000 grand mean (1-octave higher than the stimulus

frequency) whereas there was no clear response to the 2-1-2 stimuli at this DR frequency, and

(iv) for the 500-Hz stimuli, any differences in the pattern of response amplitudes between

NBchirp versus 2-1-2 stimuli were relatively small, except the largest grand-mean NBchirp

response was for DR707 versus DR500 for 2-1-2 stimuli.

< INSERT FIGURE 6 ABOUT HERE >
16

Figure 7A presents the percent of subjects showing present responses to the NBchirp and

2-1-2 stimuli at each ½-octave derived-response condition. For responses to 500-Hz stimuli,

response presence was high (91-100%) at DR500 and DR707, with response presence decreasing

1-octave away and greater from the stimulus frequency. NBchirps response presence for DR1000

was higher than that for 2-1-2 stimuli (91% vs 64%); however, differences between response

presence for NBchirp and 2-1-2 stimuli 1-octave away from 500 Hz did not reach statistical

significance [DR250: χ^2 (1, N=11) = 0.21, p=0.647; DR1000: χ^2 (1, N=11) = 2.33, p=0.127]. For

2000-Hz stimuli, response presence for both stimuli was 100% at DR2000 and DR2830. At 1-

octave above the stimulus frequency (DR4000), results for 2000-Hz NBchirps continued to show

high (91%) response presence, whereas results for 2-1-2 stimuli showed substantially reduced

response presence (27%). At 1-octave below the 2000-Hz stimulus frequency (DR1000), both

stimuli showed low percent response presence (NBchirp: 46%; 2-1-2: 18%). The differences

between NBchirp and 2-1-2 percent-present results did not reach statistical significance for

DR1000 [χ^2 (1, N=11) = 1.89, p=0.170], but 2000-Hz NBchirps showed significantly more

responses for DR4000 [χ^2 (1, N=11) = 9.21, p<0.001].

Figure 7B presents mean (and SD) amplitude results for the effects of DR frequency on

½-octave-wide DRs to NBchirp and 2-1-2 stimuli for the two frequencies. In general, the

amplitude profiles showed: (i) generally, larger amplitudes to 2000-Hz tones for NBchirps

compared to 2-1-2 stimuli, but only small differences between the stimuli in response to 500-Hz

tones, (ii) largest mean amplitudes in the DRs within ½-octave of the stimulus frequency, with

amplitudes substantially decreasing in DRs at least 1-octave away from the stimulus frequency,

(iii) in response to 2000-Hz tones, relatively large mean amplitude to NBchirps in DR4000,

whereas results for 2-1-2 stimuli in DR4000 showed very small mean amplitude (reflecting

mostly "no-response" results for 2-1-2 stimuli at DR4000), and (iv) the amplitudes in response to

NBchirps reflected the wider range of DR frequencies with responses than the 2-1-2 stimuli,

consistent with the grand-mean waveforms (Figure 6) and the response presence results (Figure

7A). Two-way repeated measures ANOVAs were conducted separately for amplitude results for

the 500-Hz and 2000-Hz stimulus frequencies, with results shown in the bottom of Table 1.

< INSERT FIGURE 7 ABOUT HERE >

For the ½-octave-wide DRs to 500-Hz stimuli, the significant main effect for DR

frequency (DRFRQ) was consistent with the results seen in the grand-mean waves (Figure 6) and

amplitude profiles (Figure 7B). Both stimuli showed maximum responses within ½ octave of the

stimulus frequency, with amplitudes significantly decreasing above and below. Neither the main

effect for STIM nor the DRFRQxSTIM interaction reached statistical significance. The non-

significant trends STIM and DRFRQxSTIM were likely due to the NBchirp mean amplitude

being a little larger than that for 2-1-2 tones in the DR707 ½-octave -wide DR; otherwise, mean

amplitudes were essentially the same in all other ½-octave -wide DRs, including DR500.

For the ½-octave -wide DRs to 2000-Hz stimuli, the two-way ANOVA showed the

significant main effects of stimulus type (STIM), DR frequency (DRFRQ), as well as a

significant DRFRQxSTIM interaction, which was consistent with the results seen in the grand-

mean waves (Figure 6), the percent-present histograms (Figure 7A), and the amplitude graphs

(Figure 7B) for the ½-octave -wide DRs. Amplitudes decreased significantly as the DR

frequency was further away from 2000 Hz. The STIM significant main effect of indicated

significantly larger responses to NBchirps compared to 2-1-2 stimuli; however, this did not hold

across all DR conditions. Neuman-Keuls *post hoc* analysis of the DRFRQxSTIM interaction

indicated that the amplitudes in response to NBchirps were significantly larger than those to

2-1-2 stimuli in the DR4000, DR2000, and DR1410 conditions (but not in the DR2830

condition). Further, these analyses indicated that DR amplitudes to NBchirps significantly

decreased between DR2000 and DR2830 but not between DR2830 and DR4000. In contrast,

amplitudes in response to 2-1-2 stimuli did not significantly decrease between DR2000 and

DR2830 but did significantly decrease between DR2830 and DR4000. Amplitudes to both

stimuli significantly decreased when DR frequency was increased above DR4000 or below

DR2000.

Similar to the 1-octave-wide DR results, the ½-octave -wide NBchirp amplitude profiles

in Figure 7B appeared to be wider than those of the 2-1-2 amplitude profiles. Bandwidth (and

center frequencies) of each subject's ½-octave-wide DR amplitude profiles were determined for a

wave V-V amplitude of 0.075 µV on either side of the profile. Summary results are shown in the

bottom of Table 2. A mixed-model ANOVA (stimulus frequency as a between effect; stimulus

type as a within effect) was carried-out for the 500-and 2000 Hz ½-octave-wide $BW_{0.075}$ (in

octaves) to NBchirp versus 2-1-2 stimuli. As with the 1-octave-wide results, the ANOVA

indicated no significant differences in the width of the ½-octave-wide DR amplitude profiles

between 500- and 2000-Hz stimuli when expressed in octaves [STIMFRQ main effect: $F(1, 20)$

$= 0.00$, $p = .987$, $\eta^2_p = 0.00$]. However, the bandwidths of the ½-octave-wide DR NBchirp

amplitude profiles were again significantly wider than those for the 2-1-2 stimuli [STIM main

effect: $F(1, 20) = 11.97$, $p = .003$, $\eta^2_p = 0.37$], and this held for both stimulus frequencies, as

indicated by the nonsignificant interaction [STIMFRQ x STIM interaction: $F(1, 20) = 1.17$, $p =$

$.293$, $\eta^2_p = 0.06$]. As with the 1-octave-wide DR results, the ½-octave-wide DR amplitude

profiles' center frequencies were similar between the NBchirp and 2-1-2 stimuli.

Effect of Cochlear Bandwidth: No-Mask, 1-Octave-Wide DR and ½-Octave-Wide DR

Insight into the effects of limiting the extent of the cochlear regions available to respond

(cochlear bandwidth, CBW) to the NBchirp and 2-1-2 stimuli is provided by comparing wave V

amplitudes in the no-mask condition with the 1-octave-wide and ½-octave-wide DR conditions

with the same center frequency as the stimulus frequency (i.e., DR500 and DR2000), as shown

in Table 3. A mixed-model ANOVA (STIMFRQ as a between effect; STIM type and CBW as

within effects) was carried out for those participants with results for all three CBW conditions.

The key finding was a significant CBW x STIM interaction [F(2, 40) = 38.87, ε = 1.00, p < .001,

η^2_p = 0.66]. Wave V amplitudes in response to NBchirps compared to 2-1-2 stimuli, evaluated

using Neuman-Keuls *post hoc* comparisons, showed greater decreases as cochlear bandwidth

was narrowed from no-mask to 1-octave-wide to ½-octave wide DRs. Responses to 2-1-2 stimuli

showed no significant change from no-mask to 1-octave-wide DRs, with a small albeit

significant decrease in amplitude from 1-octave-wide to ½-octave-wide DRs. As already noted,

wave V amplitudes in the no-mask condition were significantly larger in response to NBchirps

compared to 2-1-2 stimuli; this amplitude advantage decreased in the 1-octave-wide condition,

and was absent when responses are restricted to ½ octave around the stimulus frequency (i.e.,

there was no significant different between amplitudes to NBchirp versus 2-1-2 stimuli in the ½-

octave-wide DR conditions corresponding to the stimulus frequencies). NBchirps showed larger

responses to 2000-Hz stimuli, which was not seen with the 2-1-2 stimuli, as reflected by a

significant STIM x STIMFRQ interaction [F(1, 20) = 5.01, ε = 1.00, p = .037, η^2_p = 0.20]. The

CBW x STIM x STIMFRQ interaction was not significant [F(2, 40) = 1.13, ε = 1.00, p = .367,

η^2_p = 0.05]. The difference in ½-octave-wide DR2000 amplitudes was not significant in this

more-limited analysis compared to the significant Neuman-Keuls result reported above in the

analysis which included all ½-octave-wide DR frequencies but only results for 2000-Hz stimuli.

Nevertheless, the difference between mean amplitudes was very small and of little practical

significance (NBchirp: 0.13 µV; 2-1-2: 0.11µV).

The results also demonstrate another effect of restricting the cochlear bandwidth:

although larger in amplitude measured in microvolts (at least, in the no-mask and some 1-octave

DR conditions), relative to their non-masked amplitude, DR amplitudes (in percent-non-masked)

for NBchirps decreased more compared to amplitudes for responses to 2-1-2 stimuli. A mixed-

model ANOVA (STIMFRQ as a between effect; STIM type and CBW as within effects) was

carried out on the percent-of-non-masked-amplitude results for those participants with results for

both the 1-octave and ½-octave DR conditions. As shown in Table 3, expressed as a percent of

non-masked amplitude, NBchirp DRs were significantly smaller than those for 2-1-2 stimuli

[STIM main effect: $F(1, 20) = 22.58$, $\varepsilon = 1.00$, $p < .001$, $\eta^2_p = 0.53$]. Percent-amplitudes were

also significantly smaller in the ½-octave DR condition compared to the 1-octave condition

[CBW main effect: $F(1, 20) = 55.52$, $\varepsilon = 1.00$, $p < .001$, $\eta^2_p = 0.74$]. No other main effects or

interactions were statistically significant [STIMFRQ: $F(1, 20) = 1.07$, $\varepsilon = 1.00$, $p = .31$, $\eta^2_p =$

0.05; CBW x STIMFRQ: $F(1, 20) = 0.09$, $\varepsilon = 1.00$, $p = .76$, $\eta^2_p = 0.09$; STIM x STIMFRQ: $F(1,$

$20) = 0.11$, $\varepsilon = 1.00$, $p = .74$, $\eta^2_p = 0.01$; CBW x STIM: $F(1, 20) = 2.85$, $\varepsilon = 1.00$, $p = .11$, $\eta^2_p =$

0.12; CBW x STIM x STIMFRQ: $F(1, 20) = 0.65$, $\varepsilon = 1.00$, $p = .43$, $\eta^2_p = 0.03$] .

< INSERT TABLE 3 ABOUT HERE >

17   **DISCUSSION**

Many studies have indicated larger ABR wave V (and Auditory Steady State Response,

ASSR) amplitudes to NBchirp stimuli compared to more-standard stimuli. Though not

consistently present for all frequencies and/or intensities, the larger amplitudes in response to

NBchirp stimuli have been primarily attributed to the timing of frequencies within these stimuli

based on cochlear travelling wave delays. However, the larger amplitudes seen with NBchirps

might simply be due to these stimuli having wider spectra (Figure 1) and, thus, wider cochlear

contributions. The current study is the first study to directly assess cochlear contributions to

ABR wave V to current NBchirps using the high-pass noise/derived response (HPDR) technique.

Results showed significantly wider cochlear contributions to ABR wave V to NBchirp compared

to 2-1-2 stimuli.

ABR to NBchirp and 2-1-2 Stimuli in High-Pass Noise

ABR wave V-V' amplitudes in response to non-masked 60 dB pe SPL 500- and 2000-Hz

NBchirps were significantly larger than those to 2-1-2 stimuli and almost all participants (23/24)

showed larger responses to non-masked NBchirps. This finding is consistent with some previous

studies showing significantly larger wave V amplitudes to 500- and 2000-Hz NBchirps

compared to standard 5-cycle stimuli. However, the literature is quite inconsistent: some studies

have shown larger wave V amplitudes to NBchirps at low ($\leq$ 40 dB nHL) stimulus levels (e.g.,

Ferm et al. 2013; Rodrigues et al. 2013; Ferm & Lightfoot 2015). Other studies, however, have

reported nonsignificant differences in the amplitude of responses to NBchirp and 2-1-2 stimuli at

low stimulus levels (e.g., Dzulkarnain et al. 2018; Pinto et al. 2022; Ceylan et al. 2025).

Additionally, some reported no differences to higher stimulus levels (e.g., Megha et al. 2019; Bal

2022) and some have reported no differences (or responses to standard tones larger) specifically

to 500-Hz stimuli (e.g., Rodrigues et al. 2013; Cobb & Stuart 2016). There is a large diversity

across these studies in study subject populations (adults, older children, infants) and, especially,

ABR recording parameters, which may explain some of the inconsistency. Despite this, it seems

most studies conclude wave V amplitudes are larger in response to NBchirps. This study's HP

noise and DR results provided some insight concerning these response amplitudes.

In the current study, averaged across all HP noise conditions, NBchirp stimuli produced

larger responses than 2-1-2 stimuli. However, the amplitude difference between responses

substantially decreased once the HP noise cutoff was within ½-octave above the stimulus

frequency. Thus, the large gain in amplitude by NBchirps over 2-1-2 stimuli required

contributions from cochlear frequency regions more than ½-octave higher than the stimulus

frequency.

The effects of HP noise cutoff frequency on wave V amplitudes showed differences

between NBchirp and 2-1-2 stimuli in the frequency at which HP noise cutoffs produced

significant ABR amplitude changes. ABR amplitudes to NBchirp stimuli were significantly

reduced with HP noise cutoffs one-half octave (500-Hz stimuli) to 1-octave (2000-Hz stimuli)

higher than those noise cutoffs which resulted in significant decreases in ABR amplitude for

2-1-2 stimuli. This suggests that the ABR to NBchirps receive contributions from wider cochlear

frequency places than the responses to 2-1-2 stimuli, at least at 60 dB pe SPL.

In the current study, the HP noise had a very steep slope but only 40-45 dB stopband

attenuation. The effects of HP masking suggested some leakage of noise occurred into

frequencies below the HP cutoffs. Noise leakage effects were primarily decreases in response

amplitude in the 16000-Hz and/or 8000-Hz HP noise conditions. As 16,000 Hz is well above the

output limits of the ER-3A earphones, any response changes are likely due to noise leakage

below the 16,000-Hz cutoff (noise at 73-77 dB SPL with a 50-dB stopband attenuation could still

see 23-27 dB SPL noise leaking into the stopband). Given it is unlikely that 60 dB peSPL 2000-

Hz stimuli (NBchirp or 2-1-2) would produce responses from such high-frequency cochlear

regions (nor, indeed, from 8000-Hz regions for 500-Hz stimuli), the more likely explanation is

noise leakage into the stopband. However, significant amplitude decreases in the 16000- or

8000-Hz HP noise conditions were only observed for NBchirps and not for 2-1-2 stimuli. One

would expect leakage to affect responses to both stimuli equally. ABRs to NBchirps originating

from wider frequency regions might explain this finding.

Derived Responses to NBchirp versus 2-1-2 Stimuli

5   Cochlear place specificity of ABR wave V to NBchirps and 2-1-2 stimuli is demonstrated

by the amplitude profiles (graph of amplitude vs DR frequency) of the 1-octave-wide and ½-

octave-wide DRs. In general, these amplitude profiles showed that responses to both NBchirp

and 2-1-2 stimuli originated primarily from cochlear contributions representing frequencies

within one octave of the stimulus frequency. Thus, both stimuli demonstrate cochlear place

specificity. More-detailed assessment of the results, however, suggested significantly better

cochlear place specificity for the ABR to 2-1-2 stimuli compared to NBchirps.

12   The DR grand-mean waveforms, percent of subjects showing present responses at each

DR condition, and DR amplitude profiles, center frequencies, and bandwidth measures all

indicated good place specificity of the ABR to the 500- and 2000-Hz 2-1-2 stimuli. Specifically,

these measures showed: (i) response presence limited primarily to DRs within ± 0.5 octave of the

stimulus frequency, (ii) response amplitude maxima at the nominal stimulus frequency or within

a half octave of the nominal stimulus frequency, (iii) a sharp drop-off in the amplitude of

responses approximately one-half to 1-octave above and below the DR frequency with the

maximum response amplitude (usually at the stimulus frequency), indicating that the cochlear

contributions to the ABR wave V to the 2-1-2 stimuli are primarily within approximately within

±0.5 octaves of the stimulus frequency, and (iv) the mean center frequencies calculated for the 1-

octave-and ½-octave-wide DR amplitude profiles to the 2-1-2 stimuli were close to the nominal

stimulus frequencies of the 2-1-2 stimuli. NBchirp amplitude profiles were similar in shape to

those for 2-1-2 stimuli, with center frequencies close to the stimulus frequency. However,

NBchirps showed more responses present at frequencies one octave higher (and sometimes one

octave lower) as well as significantly larger amplitudes one octave above and below the stimulus

frequency.

A measure of the width (in Hz) of the amplitude profiles provided an estimate of the

bandwidth of the responses to each stimulus and, thus, an easy comparison between stimuli. In

the current study, we determined bandwidth from the amplitude profiles for a wave V amplitude

of 0.075 µV, a level that represents a near-threshold response for both stimuli. Mean ½-octave-

wide DR bandwidths obtained for the ABRs to 2-1-2 stimuli in the current study were 1.04

octaves for 500 Hz and 0.87 octaves for 2000 Hz, thus approximating a 1-octave-wide basilar

membrane place. In comparison, at 1.38 octaves for 500 Hz and 1.54 octaves for 2000 Hz, mean

½-octave-wide DR bandwidths for the NBchirp amplitude profiles were significantly wider than

those for 2-1-2 stimuli, and well over 1-octave-wide. Bandwidths measured from the 1-octave-

wide DR amplitude profiles were also significantly wider for NBchirps compared to 2-1-2

stimuli. Thus, NBchirps produce larger contributions from a wider cochlear region compared to

2-1-2 stimuli.

In previous studies using a variety of masking methods, the bandwidths for ABRs to the

more-standard 500- and 2000-Hz tones have demonstrated largest cochlear contributions

originate from frequency places within approximately ±0.5 octave of the stimulus nominal

frequency (e.g., Klein 1983; Horiuchi 1989; Wu & Stapells 1994; Abdala & Folsom 1995; Oates

& Stapells 1997b; Oates & Purdy 2001). Using the HPDR technique, Oates showed wave V DR

(½-octave-wide) amplitude profile bandwidths of 622 and 2058 Hz for 500- and 2000-Hz 2-1-2

stimuli (at 80 dB peSPL), respectively (Oates 1996). In the current study, ½-octave-wide DR

bandwidths for ABRs to 2-1-2 stimuli (428 and 1367 Hz) were much smaller than those reported

by Oates, likely due to the lower stimulus intensity employed in the current study (60 pe SPL).

Although larger in amplitude measured in microvolts, DR amplitudes for NBchirps

decreased more relative to their non-masked amplitude than DR amplitudes for 2-1-2 stimuli as

cochlear bandwidth was decreased. The significant differences in the DR percent amplitudes

between NBchirp and 2-1-2 stimuli likely cannot be attributed to differences in noise masking

levels, as approximately the same levels (within 2-3 dB) of masking noise were used for masking

the ABR to the NBchirp and 2-1-2 stimuli. A more likely explanation for the NBchirp DR

percent amplitudes being significantly lower than those for 2-1-2 stimuli is that the responses to

non-masked NBchirps originate from a wider cochlear frequency range (as demonstrated by this

study's HP noise and DR results), thus response contributions to the non-masked ABR are

spread out over more DRs, and individual DRs are smaller expressed as a percentage of the no-

mask response amplitude.

NBchirp Amplitude Advantage versus Cochlear Bandwidth

The ABR wave V-V' amplitudes in response to non-masked 60 dB pe SPL 500- and

2000-Hz NBchirps were significantly larger than those to 2-1-2 stimuli. This amplitude

advantage for NBchirps has been shown by many studies, and is the primary reason for

recommending NBchirps over more-standard tones, although it is not consistently present in all

intensity/frequency conditions (see above). However, as the results of the current study clearly

demonstrate, the NBchirp amplitude advantage requires a response from a wider cochlear range.

When cochlear bandwidth was reduced by the HPDR technique to 1 octave, the advantage

remained but was significantly decreased. This counters the suggestion by Wegner and Dau

(2002) that a 1-octave wide bandwidth is not sufficient to produce a chirp amplitude advantage.

When the cochlear bandwidth was limited to a ½-octave-wide region, the NBchirp amplitude

advantage was essentially absent.

3   The results of the current study cannot unequivocally determine whether the larger

amplitude of responses to non-masked NBchirps is due to their wider acoustic spectrum (Figure

1) or to the optimized timing of frequencies within the NBchirp to overlap responses from

different frequency regions such that they constructively superimpose and result in a larger

response. Reducing the cochlear bandwidth of the response using the HPDR technique would

effectively limit and perhaps eliminate the effects of the optimized timing, especially for the ½-

octave-wide DR condition. Even if the effects of optimized timing are absent, ½-octave-wide DR

results indicate NBchirps produce responses from a wider cochlear region than 2-1-2 stimuli.

Similar results were seen when HP noise cutoffs were within 0.5-1 octave above the stimulus

frequency. Thus, as cochlear bandwidth decreased (from wideband to 1-octave-wide to ½-

octave-wide), there was less cochlear "space" for NBchirps to utilize any amplitude advantage of

dispersing frequencies according to travelling wave delays. From these results, one could

postulate that the ½-octave-wide DRs for NBchirps reflected primarily spectral contributions

rather than spectral *plus* optimization of frequency timing, and thus, little-or-no difference

existed between response amplitudes to the two stimuli.

Clinical Implications

20   This the first study to directly assess the cochlear place specificity of the ABR to

NBchirps. As already noted, many previous studies have assessed the cochlear place specificity

of the ABR to more-standard tonal stimuli, such as 2-1-2 linear-gated tones and 5-cycle

Blackman-windowed tones. Previous research has employed pure-tone masking, HP noise

masking, and notched-noise masking techniques (e.g., Klein 1983; Horiuchi 1989; Wu &

Stapells 1994; Abdala & Folsom 1995; Oates & Stapells 1997a; Oates & Stapells 1997b; Oates

& Purdy 2001). Others have studied 2-1-2 and similar stimuli in individuals with steeply sloping

cochlear hearing loss (Davis et al. 1984; Purdy & Abbas 2002). In general, results from these

studies suggest the ABR to standard tones such as 2-1-2 stimuli show adequate cochlear place

specificity, at least up to about 80 dB pe SPL. However, when audiogram configurations show

particularly steep hearing losses (≥50-60 dB/octave slope), standard tones may misestimate

hearing thresholds for the frequencies with elevated thresholds (Davis et al. 1984; Horiuchi

1989; Stapells et al. 1990; Purdy & Abbas 2002).

To our knowledge, no previous NBchirp studies have assessed the place specificity of the

ABR to NBchirps by systematically varying masker frequencies. A number of studies, however,

have assessed NBchirps in individuals with sensorineural hearing loss using the ABR and/or

ASSR (e.g., Seidel et al. 2015; Lee et al. 2016; Sininger et al. 2018; Talaat et al. 2019; Eder et al.

2020; Ehrmann-Müller et al. 2021; Leusin Mattiazzi et al. 2024; Khan & Vanaja 2025; Galhoum

et al. 2023; Michel & Jørgensen 2017), with most showing reasonably good threshold estimation

results for NBchirps. NBchirp threshold results for several individual cases with mostly gently or

moderately sloping hearing loss (15-35 dB/octave slope) plus 1 case with steeply sloping loss

have also been published (Seidel et al. 2015; Mühler et al. 2012; Khan & Vanaja 2025).

However, none of these studies have assessed NBchirp results in groups of individuals with

steeply/precipitously sloping audiogram configurations (≥40 dB/octave slope). Testing ABR

thresholds in only individuals with flat or mildly sloping audiograms does not adequately assess

cochlear place specificity (Davis et al. 1984; Purdy & Abbas 2002). Indeed, only two studies

have directly compared ABR to NBchirps vs standard tones in individuals with hearing loss

(Talaat et al. 2019; Khan & Vanaja 2025), none in infants. Results of these two studies indicate

good threshold estimates using either NBchirp or standard tone stimuli, with results from Talaat

and colleagues also suggesting faster test times and apparently larger amplitudes for NBchirps,

though results details were not provided (Talaat et al. 2019). Importantly, based on the mean

audiograms presented by these 2 studies, most individuals with hearing loss had flat or gently

sloping audiogram configurations and thus would not provide a test of any frequency or place

specificity differences between stimuli (Talaat et al. 2019; Khan & Vanaja 2025). Due to

NBchirp's wider cochlear activation, as indicated by the results of the present study, ABR

thresholds for NBchirp stimuli may be more likely to misestimate the severity of hearing

thresholds, compared to 2-1-2 stimuli. Misestimation of thresholds would most likely be an issue

primarily when there are large differences in thresholds between adjacent octave (and especially

half-octave) audiometric frequencies; that is, steeply falling (or falling) audiograms. Further

research is needed comparing NBchirp and 2-1-2 stimuli ABR thresholds in individuals,

especially infants and young children, with steeply sloping hearing loss.

As noted above, the ABR amplitude advantage for NBchirps over standard tones is the

primary motivation for using NBchirps for ABR audiometry. This advantage, however, has not

been shown to be consistently present for all stimulus frequency/intensity conditions. Moreover,

as the current study shows, as the available response bandwidth narrows (to 1 octave or to ½

octave), the amplitude advantage decreases and even disappears. It is currently an open question

whether sensorineural hearing losses remove/reduce this amplitude advantage (perhaps due to

the audiogram configuration and/or to having cochlear traveling wave properties different from

those used in the model to produce the NBchirps). Clearly, more research is needed comparing

ABR results (amplitudes and thresholds) to NBchirp and 2-1-2 stimuli in individuals with

hearing loss.

How wide of a spectrum is acceptable for ABR stimuli is a longstanding matter of debate

within Audiology (Davis 1976; Stapells & Picton 1981; Davis et al. 1984; Sohmer & Kinarti

1984; Laukli & Mair 1986; Gorga & Thornton 1989; Laukli 2014; McCreery et al. 2015). In

cases with normal hearing or hearing loss with flat or mildly sloping/rising configurations, ABR

thresholds to 2-1-2 or NBchirp stimuli, for the most part, represent the pure-tone behavioral

thresholds at the stimulus frequencies. When hearing losses show a steeper configuration, this is

not always the case (Davis et al. 1984; Horiuchi 1989; Stapells et al. 1990; Purdy & Abbas

2002). The ±½-octave cochlear place bandwidth of the ABR to 2-1-2 tones seems to be accepted

within the field, given most early hearing programs use either standard brief tones (such as 2-1-2

linear-gated or 5-cycle Blackman-windowed tones) or NBchirps.[6] Based on the current study,

the ABR to 2-1-2 stimuli may reflect contributions from cochlear places representing frequencies

within one-half octave of the stimulus frequency (e.g., from 1414 to 2828 Hz in response to

2000-Hz stimuli). On the other hand, the much wider bandwidth of ABRs to NBchirps – as wide

as 2 octaves (Table 2) -- would seem to be too wide. As this study demonstrated, the 2000-Hz

NBchirp produced responses from the 4000-Hz cochlear region in most subjects (compared to

few for 2-1-2 stimuli). Concern about NBchirp frequency-specificity has led some researchers to

---

[6] Current NBchirp stimuli, such as the NB CE-Chirp® LS, have been described in many publications as being "octave-band" stimuli, which is essentially true when measured only at -3 dB (Elberling & Don 2010; Gøtsche-Rasmussen et al. 2012; Adjekum et al. 2024).

suggest embedding NBchirp stimuli in HP noise or notched noise (Corona-Strauss et al. 2012;

Baljić et al. 2017b). However, as the current study shows, restricting the responding cochlear

region using masking noise reduces (or eliminates) the wave V amplitude advantage of NBchirps

over 2-1-2 stimuli. Should future research show reduced (or absent) amplitude advantages for

NBchirps in individuals with hearing loss, there would seem little reason to using NBchirp

stimuli with such a wide bandwidth. Further research in individuals with hearing loss is needed.

Study Limitations

The current study investigated cochlear place specificity for a relatively low stimulus

intensity (60 dB pe SPL, approximately 40 dB nHL). Although the frequency and place

specificity of the ABR to standard tonal stimuli at higher stimulus intensities has previously been

studied (e.g., Klein 1983; Horiuchi 1989; Oates & Stapells 1997b; Oates & Purdy 2001), studies

are clearly needed on the place specificity of the ABR to higher intensity NBchirps (e.g., 80-90

dB pe SPL). Additionally, this study only investigated the place specificity of the ABR to 500-

and 2000-Hz NBchirps. There is a need to conduct further studies on the place specificity of the

ABR to 1000- and 4000-Hz NBchirps. For example, it may be that the place specificity of the

ABR to 4000-Hz NBchirps will be different from that of the ABR to 2000-Hz NBchirps because

of the reduction in the frequency content of the 4000-Hz NBchirp by the ER-3A earphone. The

current study showed smaller differences between NBchirp vs 2-1-2 results for 500-Hz stimuli; it

is not known whether results for 1000-Hz NBchirps would be similar to the place specificity

results seen in the present study for 2000-Hz vs 500-Hz stimuli.

This study only assessed results from 12 normal-hearing participants for each stimulus

frequency. This relatively small sample size may have reduced this study's ability to detect small

differences between stimuli and conditions (for example, response presence differences 1 octave

away from stimulus frequency).

3   The current study assessed results only from adult participants with normal hearing.

Although these results are likely to be applicable (at least partially) to individuals with cochlear

hearing loss, research in individuals (especially infants) with hearing loss is required. The

NBchirp delay models are derived from travelling wave data of the normal adult cochlea, and

these delay models may not represent delays found in infant and/or pathological cochleae. This

could result in suboptimal amplitude enhancement or even response cancellation (i.e., response

amplitudes to NBchirps no different from, or even smaller than, those to 2-1-2 stimuli).

Importantly, further studies are needed to investigate whether individuals with steep (rising or

falling) cochlear hearing loss configurations show differences in amplitudes or thresholds for

ABR thresholds to NBchirp and 2-1-2 stimuli.

Conclusions

Although both stimuli demonstrate cochlear place specificity, ABRs to NBchirps show

significantly poorer place specificity compared to ABRs to 2-1-2 stimuli. As the relative

amplitude advantage for NBchirps over 2-1-2 stimuli decreases as the region of the cochlea

contributing to the response decreases, it is not yet clear whether NBchirps amplitude advantages

over more-standard stimuli exist when there is cochlear hearing loss. Further research, especially

in individuals with steep cochlear hearing loss, is required.

REFERENCES

Abdala, C., Folsom, R. (1995). The development of frequency resolution in humans as revealed
by the auditory brain-stem response recorded with notched-noise masking. *J. Acoust. Soc.*
Am., 98, 921–930.

Adjekum, R., Small, S.A., Chan, S., Stapells, D.R. (2024). Frequency specificity of narrowband
chirp and 2-1-2 stimuli: Spectral analyses. *medRxiv*, 2024.11.09.24317036.

ANSI S3.6-2018 (R2023) (2018). *Specification for audiometers.*, New York: Acoustical Society
of America.

Bagatto, M. (2020). Ontario Infant Hearing Program Protocol for Auditory Brainstem Response-
based Audiological Assessment (ABRA). Version 2018 02. Available at:
https://ir.lib.uwo.ca/cgi/viewcontent.cgi?article=1000&context=nca [Accessed November
17, 2021].

Bal, N.G.T., Özge; Hüsam, Hilal; Özdemir, Sümeyye; Lien, Bartlomiej Dao (2022). Optimizing
auditory brainstem response test with narrow band level-specific CE-Chirps. *Ann. Otol.*
Neurotol., 5, 050–054.

Bell, S.L., Allen, R., Lutman, M.E. (2002). An investigation of the use of band-limited chirp
stimuli to obtain the auditory brainstem response: *Int. J. Audiol.*, 41, 271–278.

Ceylan, S., İşlek, A., Baba, P., Özkul, Y. (2025). The superiority of narrow band CE-Chirp
auditory brainstem response for predicting the behavioral hearing thresholds in healthy
adults. *Egypt. J. Otolaryngol.*, 41, 32.

Cobb, K.M., Stuart, A. (2016). Neonate auditory brainstem responses to CE-Chirp and CE-Chirp
octave band stimuli I: Versus click and tone burst stimuli. *Ear Hear.*, 37, 710–723.

Davis, H. (1976). Principles of electric response audiometry. *Ann. Otol. Rhinol. Laryngol.*, 85,
1–96.

Davis, H., Hirsh, S.K., Popelka, G.R., Formby, C. (1984). Frequency selectivity and thresholds
of brief stimuli suitable for electric response audiometry. *Audiology*, 23, 59–74.

Don, M., Eggermont, J.J. (1978). Analysis of the click-evoked brainstem potentials in man using

high-pass noise masking. *J. Acoust. Soc. Am.*, 63, 1084–1092.

Don, M., Eggermont, J.J., Brackmann, D.E. (1979). Reconstruction of the audiogram using

brainstem responses and high-pass noise masking. *Ann. Otol. Rhinol. Laryngol.*, Suppl,

1–20.

Don, M., Elberling, C. (1996). Use of quantitative measures of auditory brain-stem response

peak amplitude and residual background noise in the decision to stop averaging. *J.*

Acoust. Soc. Am., 99, 491–499.

Dzulkarnain, A.A.A., Abdullah, S.A., Ruzai, M.A.M., Ibrahim, S.H.M.N., Anuar, N.F.A.,

Rahim, 'Afaf Ezzaty Abdul (2018). Effects of different electrode configurations on the

narrow band Level-Specific CE-Chirp and tone-burst auditory brainstem response at

multiple intensity levels and frequencies in subjects with normal hearing. *Am. J. Audiol.*,

27, 294–305.

Eder, K., Schuster, M.E., Polterauer, D., Neuling, M., Hoster, E., Hempel, J.M., Semmelbauer,

S. (2020). Comparison of ABR and ASSR using NB-chirp-stimuli in children with severe

and profound hearing loss. *Int. J. Pediatr. Otorhinolaryngol.*, 131, 109864.

Eggermont, J.J. (1976). Electrocochleography. In W. D. Keidel & W. D. Neff, eds. *Handbook of*

Sensory Physiology, Volume V, Part 3. (*pp. 625–704*). Berlin: Springer-Verlag.

Eggermont, J.J., Don, M. (1980). Analysis of the click-evoked brainstem potentials in humans

using high-pass noise masking. II. Effect of click intensity. *J. Acoust. Soc. Am.*, 68,

1671–1675.

Ehrmann-Müller, D., Shehata-Dieler, W., Alzoubi, A., Hagen, R., Cebulla, M. (2021). Using

ASSR with narrow-band chirps to evaluate hearing in children and adults. *Eur. Arch.*

Otorhinolaryngol., 278, 49–56.

Elberling, C. (1974). Action potentials along the cochlear partition recorded from the ear canal in

man. *Scand. Audiol.*, 3, 13–19.

Elberling, C., Don, M. (2010). A direct approach for the design of chirp stimuli used for the
2   recording of auditory brainstem responses. *J. Acoust. Soc. Am.*, 128, 2955–2964.

3   Elberling, C., Don, M. (2008). Auditory brainstem responses to a chirp stimulus designed from
4   derived-band latencies in normal-hearing subjects. *J. Acoust. Soc. Am.*, 124, 3022–3037.

5   Elberling, C., Don, M., Cebulla, M., Stürzebecher, E. (2007). Auditory steady-state responses to
6   chirp stimuli based on cochlear traveling wave delays. *J. Acoust. Soc. Am.*, 122, 2772–
7   2785.

8   Elberling, C., Kristensen, S.G.B., Don, M. (2012). Auditory brainstem responses to chirps
9   delivered by different insert earphones. *J. Acoust. Soc. Am.*, 131, 2091–2100.

Etymotic Research ER-3A ABR Insert Earphone Data Sheet - Etymotic Research. *yumpu.com*.
Available at: https://www.yumpu.com/en/document/read/19542993/er-3a-abr-insert-
earphone-data-sheet-etymotic-research [Accessed November 11, 2025].

Evans, E.F., Elberling, C. (1982). Location-specific components of the gross cochlear action
potential: an assessment of the validity of the high-pass masking technique by cochlear
nerve fibre recording in the cat. *Audiology*, 21, 204–227.

Fedtke, T., Richter, U. (2007). Reference zero for the calibration of air-conduction audiometric
equipment using "tone bursts" as test signals. *Int. J. Audiol.*, 46, 1–10.

Ferm, I., Lightfoot, G. (2015). Further comparisons of ABR response amplitudes, test time, and
estimation of hearing threshold using frequency-specific chirp and tone pip stimuli in
newborns: Findings at 0.5 and 2 kHz. *Int. J. Audiol.*, 54, 745–750.

Ferm, I., Lightfoot, G., Stevens, J. (2013). Comparison of ABR response amplitude, test time,
and estimation of hearing threshold using frequency specific chirp and tone pip stimuli in
newborns. *Int. J. Audiol.*, 52, 419–423.

Galhoum, D.H., Ibraheem, O.A.A., Elsaeed, M.A., Elnabtity, N. (2023). Relationship between
behavioral and electrophysiologic hearing thresholds using narrow-band chirp-evoked
auditory brainstem response. *Zagazig Univ. Med. J.*, 29, 796–804.

Glass, G.V., Hopkins, K.D. (1996). *Statistical methods in education and psychology. 3rd*
Edition, Needham Heights, MA: Allyn & Bacon.

Gorga, M.P., Thornton, A.R. (1989). The choice of stimuli for ABR measurements. *Ear Hear.*,
10, 217–230.

Gøtsche-Rasmussen, K., Poulsen, T., Elberling, C. (2012). Reference hearing threshold levels for
chirp signals delivered by an ER-3A insert earphone. *Int. J. Audiol.*, 51, 794–799.

Hatton, J.L., Van Maanen, A., Stapells, D.R. (2022). British Columbia Early Hearing Program:
Auditory Brainstem Response (ABR) Protocol. Available at: http://www.phsa.ca/bc-
early-hearing/Documents/ABR_Protocol.pdf.

Haughton, P. (2006). Insert earphones--a comparison of short-duration signals measured with an
occluded ear simulator and a 2cc coupler. *Int. J. Audiol.*, 45, 60–65.

Herdman, A.T., Picton, T.W., Stapells, D.R. (2002). Place specificity of multiple auditory
steady-state responses. *J. Acoust. Soc. Am.*, 112, 1569–1582.

Horiuchi, Y. (1989). Frequency specificity of ABR evaluated by tuning curves. *Nihon Jibiinkoka*
Gakkai Kaiho, 92, 1895–1910.

Huynh, H., Feldt, L.S. (1976). Estimation of the Box correction for degrees of freedom from
sample data in randomized block and split-plot designs. *J. Educ. Stat.*, 1, 69–82.

Khan, Q., Vanaja, C.S. (2025). Auditory brainstem response and auditory steady state response
elicited using narrowband chirps: A comparison with tone burst auditory brainstem
response. *J. India Inst. Speech Hear.*, 44, 24–36.

Klein, A.J. (1983). Properties of the brain-stem response slow-wave component. II. Frequency
specificity. *Arch. Otolaryngol.*, 109, 74–78.

Kramer, S.J. (1992). Frequency specific auditory brainstem responses to bone-conducted stimuli.
Audiology, 31, 61–71.

Laukli, E. (2014). Frequency specificity and accuracy of ABR and ASSR. *Int. J. Audiol.*, 53,
697–698.

Laukli, E., Mair, I.W.S. (1986). Frequency specificity of the auditory brainstem responses. A
derived band study. *Scand. Audiol.*, 15, 141–146.

Lee, M.Y., Ahn, S.Y., Lee, H.J., Jung, J.Y., Rhee, C.-K., Suh, M.-W. (2016). Narrow band CE-
Chirp auditory steady-state response is more reliable than the conventional ASSR in
predicting the behavioral hearing threshold. *Auris. Nasus. Larynx*, 43, 259–268.

Leusin Mattiazzi, Â., Cóser, P.L., Endruweit Battisti, I.D., Dalcin Pinto, J., Pinto Vieira Biaggio,
E. (2024). Auditory electrophysiological thresholds with different chirps and their
correlation with behavioral thresholds in hearing-impaired children. *Ear Hear.*, 45, 1228–
1240.

McCreery, R.W., Kaminski, J., Beauchaine, K., Lenzen, N., Simms, K., Gorga, M.P. (2015). The
impact of degree of hearing loss on auditory brainstem response predictions of behavioral
thresholds. *Ear Hear.*, 36, 309–319.

Megha, K.N., Divyashree, K.N., Lakshmi, A., Adithya, S., Keerthana, K.P., Pushpalatha, Z.V.,
Konadath, S. (2019). Narrow-band chirp and tone burst auditory brainstem response as an
early indicator of synaptopathy in industrial workers exposed to occupational noise.
Intractable Rare Dis. Res., 8, 179–186.

Michel, F., Jørgensen, K.F. (2017). Comparison of threshold estimation in infants with hearing
loss or normal hearing using auditory steady-state response evoked by narrow band CE-
chirps and auditory brainstem response evoked by tone pips. *Int. J. Audiol.*, 56, 99–105.

Mühler, R., Mentzel, K., Verhey, J. (2012). Fast hearing-threshold estimation using multiple
auditory steady-state responses with narrow-band chirps and adaptive stimulus patterns.
ScientificWorldJournal, 2012, 192178.

Nousak, J.K., Stapells, D.R. (1992). Frequency specificity of the auditory brain stem response to
bone-conducted tones in infants and adults. *Ear Hear.*, 13, 87–95.

Oates, M.A. (1996). *Frequency specificity of the auditory brainstem and middle latency*

responses. Doctoral dissertation. New York, NY: City University of New York.

Available at: (Order No. 9618088). Available from ProQuest Dissertations & Theses

Global. (304291613). Retrieved from https://www.proquest.com/dissertations-

theses/frequency-specificity-auditory-brainstem-middle/docview/304291613/se-2.

Oates, P., Stapells, D.R. (1997a). Frequency specificity of the human auditory brainstem and

middle latency responses to brief tones. I. High pass noise masking. *J. Acoust. Soc. Am.*,

102, 3597–3608.

Oates, P., Stapells, D.R. (1997b). Frequency specificity of the human auditory brainstem and

middle latency responses to brief tones. II. Derived response analyses. *J. Acoust. Soc.*

Am., 102, 3609–3619.

Oates, P.A., Purdy, S.C. (2001). Frequency specificity of the human auditory brainstem and

middle latency responses using notched noise masking. *J. Acoust. Soc. Am.*, 110, 995–

1009.

Parker, D.J., Thornton, A.R.D. (1978). The validity of the derived cochlear nerve and brainstem

evoked responses of the human auditory system. *Scand. Audiol.*, 7, 45–52.

Picton, T.W., Hink, R.F. (1974). Evoked potentials: How? What? and Why? *Am. J. EEG*

Technol., 14, 9–44.

Picton, T.W., Linden, R.D., Hamel, G., Maru, J.T. (1983). Aspects of averaging. *Semin. Hear.*, 4,

327–341.

Picton, T.W., Stapells, D.R., Campbell, K.B. (1981). Auditory evoked potentials from the human

cochlea and brainstem. *J. Otolaryngol. Suppl*, 10, 1–41.

Pinto, J.D., Forneck, L.L.M., Ferreira, L., Cargnelutti, M., Cóser, P.L., Biaggio, E.P.V. (2022).

Auditory brainstem response with the iChirp stimuli in the infant's audiological

diagnosis. *Int. J. Pediatr. Otorhinolaryngol.*, 154, 111042.

Purdy, S.C., Abbas, P.J. (2002). ABR thresholds to tonebursts gated with Blackman and linear
windows in adults with high-frequency sensorineural hearing loss. *Ear Hear.*, 23, 358.

Rodrigues, G.R.I., Ramos, N., Lewis, D.R. (2013). Comparing auditory brainstem responses
(ABRs) to toneburst and narrow band CE-chirp in young infants. *Int. J. Pediatr.*
Otorhinolaryngol., 77, 1555–1560.

Scherg, M., Volk, S.A. (1983). Frequency specificity of simultaneously recorded early and
middle latency auditory evoked potentials. *Electroencephalogr. Clin. Neurophysiol.*, 56,
443–452.

Seidel, D.U., Flemming, T.A., Park, J.J.-H., Remmert, S. (2015). Hearing threshold estimation
by auditory steady-state responses with narrow-band chirps and adaptive stimulus
patterns: implementation in clinical routine. *Eur. Arch. Otorhinolaryngol.*, 272, 51–59.

Sininger, Y.S., Hunter, L.L., Hayes, D., Roush, P.A., Uhler, K.M. (2018). Evaluation of speed
and accuracy of next-generation auditory steady state response and auditory brainstem
response audiometry in children with normal hearing and hearing loss. *Ear Hear.*, 39,
1207.

Small, S.A., Stapells, D.R. (2017). Frequency-specific threshold assessment in infants using
frequency-specific Auditory Brainstem Response and the Auditory Steady-State
Response. In A. M. Tharpe & R. C. Seewald, eds. *Comprehensive Handbook of Pediatric*
Audiology. (*pp. 505–549*). San Diego: Plural Publishing.

Sohmer, H., Kinarti, R. (1984). Survey of attempts to use auditory evoked potentials to obtain an
audiogram. *Br. J. Audiol.*, 18, 237–244.

Stapells, D.R., Fok, M.R. (2023). Derived-band auditory brainstem responses: cochlear
contributions determined by narrowband maskers. *Int. J. Audiol.*, 1–9.

Stapells, D.R., Picton, T.W. (1981). Technical aspects of brainstem evoked potential audiometry
using tones. *Ear Hear.*, 2, 20–29.

Stapells, D.R., Picton, T.W., Durieux-Smith, A., Edwards, C.G., Moran, L.M. (1990).

Thresholds for short-latency auditory evoked potentials to tones in notched noise in

normal-hearing and hearing-impaired subjects. *Audiology*, 29, 262–274.

Stoll, T.J., Maddox, R.K. (2024). Enhanced place specificity of the parallel auditory brainstem

response: an electrophysiological study. *J. Assoc. Res. Otolaryngol.*, 25, 477–489.

Talaat, H.S., Hammad, A., El Abedein, A.M.Z. (2019). Hearing threshold evaluation in children

using narrow band chirp auditory brainstem response and tone burst auditory brainstem

response. *Int. J. Otolaryngol. Head Neck Surg.*, 9, 30–37.

Teas, D.C., Eldridge, D.H., Davis, H. (1962). Cochlear responses to acoustic transients: An

interpretation of whole-nerve action potentials. *J. Acoust. Soc. Am.*, 34, 1438–1459.

Wegner, O., Dau, T. (2002). Frequency specificity of chirp-evoked auditory brainstem responses.

J. Acoust. Soc. Am., 111, 1318–1329.

Wu, C.-Y., Stapells, D.R. (1994). Pure-tone masking profiles for human auditory brainstem and

middle latency responses to 500-Hz tones. *Hear. Res.*, 78, 169–174.

Figure Legends

Figure 1. Acoustic spectra (output from ER-3A earphones) for NBchirp and 2-1-2 stimuli for

500- and 2000-Hz stimulus frequencies.

Figure 2. Grand-mean waveforms across all subjects (N=12 per stimulus frequency) recorded to

60 dB pe SPL 500-Hz (left) and 2000-Hz (right) NBchirp and 2-1-2 stimuli for all 10 HP noise

conditions, as well as the no-mask condition.

Figure 3. Effects of HP noise on response presence and amplitude. A, Percent of subjects

showing "response present" ABR to NBchirp versus 2-1-2 stimuli in HP noise conditions for

500-Hz (left) and 2000-Hz (right) stimuli. N=12 subjects per stimulus frequency. B, Mean and

standard deviation (SD) amplitude results for the effects of HP noise cutoff on ABR wave V-V'

to NBchirp and 2-1-2 stimuli for 500- and 2000-Hz stimulus frequencies. The mean (and SD)

residual noise (RN) across all conditions and subjects is also plotted for each stimulus type and

frequency. NM: no-mask condition; NA: not applicable (not tested).

Figure 4. Grand-mean derived-response waveforms to 500-Hz (left) and 2000-Hz (right)

NBchirp and 2-1-2 stimuli for all eight 1-octave-wide DR conditions. N=12 per stimulus

frequency.

Figure 5. 1-octave-wide DR response presence and amplitude. A, Percent of subjects showing

"response present" ABR to NBchirp versus 2-1-2 stimuli in 1-octave-wide DR conditions for

500-Hz (left) and 2000-Hz (right) stimuli. N=12 subjects per stimulus frequency. B, Mean and

standard deviation (SD) amplitude results for the effects of DR center frequency noise cutoff on

ABR wave V-V' to NBchirp and 2-1-2 stimuli for 500- and 2000-Hz stimulus frequencies. The

mean (and SD) residual noise (RN) across all DR conditions and subjects is also plotted for each

stimulus type and frequency. NA: not applicable (not tested).

Figure 6. Grand-mean derived-response waveforms (across 11 subjects per stimulus frequency)

to 500-Hz (left) and 2000-Hz (right) NBchirp and 2-1-2 stimuli for all eight ½-octave -wide DR

conditions.

Figure 7. 0.5-octave-wide DR response presence and amplitude. A, Percent of subjects showing

"response present" ABR to NBchirp versus 2-1-2 stimuli in 0.5-octave-wide DR conditions for

500-Hz (left) and 2000-Hz (right) stimuli. N=11 subjects per stimulus frequency. B, Mean and

standard deviation (SD) amplitude results for the effects of DR center frequency noise cutoff on

ABR wave V-V' to NBchirp and 2-1-2 stimuli for 500- and 2000-Hz stimulus frequencies. The

mean (and SD) residual noise (RN) across all DR conditions and subjects is also plotted for each

stimulus type and frequency. NA: not applicable (not tested).

Table 1: Results of two-way repeated measures ANOVAs comparing wave V amplitudes to NBchirp versus 2-1-2 stimuli in (i) HP noise conditions, (ii) 1-octave DR conditions and (iii) ½-octave DR conditions

Effect	500-Hz stimuli					2000-Hz stimuli				
	df	F	ε†	p‡	η^2_p	df	F	ε†	p‡	η^2_p
HP NOISE										
HPFRQ	10, 110	56.53	0.32	<.001*	0.84	10, 110	101.97	0.17	<.001*	0.90
STIM	1, 11	69.03	1.00	<.001*	0.86	1, 11	93.55	1.00	<.001*	0.90
HPN x STIM	10, 110	7.56	0.71	<.001*	0.41	10, 110	51.29	0.52	<.001*	0.82
1-OCT DR										
DRFRQ	7, 77	56.34	0.65	<.001*	0.84	7, 77	58.43	0.41	<.001*	0.84
STIM	1, 11	1.46	1.00	.253	0.12	1, 11	82.34	1.00	<.001*	0.88
DRFRQ x STIM	7, 77	3.86	0.68	.005*	0.26	7, 77	5.92	0.68	<.001*	0.35
½-OCT DR										
DRFRQ	7, 70	32.78	0.82	<.001*	0.77	7, 70	26.79	0.77	<.001*	0.73
STIM	1, 10	3.70	1.00	.083	0.27	1, 10	12.17	1.00	.006*	0.55
DRFRQ x STIM	7, 70	2.35	0.53	.076	0.19	7, 70	4.91	1.00	<.001*	0.33

* Significant (p < .05)

† Huynh–Feldt epsilon (ε) correction factor for degrees of freedom

‡ Probability reflects corrected degrees of freedom

DR, Derived Response; HPN, High Pass Noise

Table 2: Mean (and standard deviation) center frequency and bandwidth (BW$_{0.075}$) results for (i) 1-octave-wide and (ii) ½-octave-wide DR amplitude profiles to 500- and 2000-Hz NBchirp and 2-1-2 stimuli.

	500 Hz		2000 Hz	
	NBchirp	2-1-2	NBchirp	2-1-2
1-OCTAVE-WIDE DRs				
N	11	11	12	12
Center Frequency (Hz)	498±65	498±89	1992±364	1866±190
Bandwidth (Hz)	667±138	484±125	3375±1278	1937±918
Bandwidth (octave)	1.81±0.35	1.37±0.40	2.16±0.55	1.39±0.55
½-OCTAVE-WIDE DRs				
N	11	11	11	11
Center Frequency (Hz)	605±81	579±85	2233±372	2260±355
Bandwidth (Hz)	595±153	429±243	2666±1525	1367±536
Bandwidth (octave)	1.38±0.40	1.04±0.57	1.54±0.58	0.87±0.34

DR, Derived Response

Table 3: Effect of cochlear bandwidth on wave V amplitude (in µV and % of no-mask condition). Mean (±SD) results for 1-octave-wide and ½-octave-wide DR conditions with same CF as stimulus frequency (DR500 and DR2000). Results for those participants with complete data (N=11 per stimulus frequency).

Stimulus Frequency		Wide (No-Mask)		1 Octave DR		Half Octave DR	
		NBchirp	2-1-2	NBchirp	2-1-2	NBchirp	2-1-2
	Amp in µV	0.20	0.14	0.16	0.13	0.09	0.10
	(±SD)	±0.06	±0.04	±0.05	±0.05	±0.04	±0.02
500 Hz	Amp %No-Mask (±SD)			82.4% ±16.0	99.7% ±32.4	48.0% ±25.0	77.8% ±30.5
	Amp in µV	0.28	0.16	0.21	0.15	0.13	0.11
	(±SD)	±0.06	±0.04	±0.06	±0.03	±0.03	±0.03
2000 Hz	Amp %No-Mask (±SD)			74.0% ±11.9	92.2% ±13.4	45.8% ±8.7	68.4% ±22.4

DR, Derived Response; SD, Standard Deviation

FIGURE 1

FIGURE 2

| 500 Hz | 2000 Hz |

NO MASK

HPN(Hz)

HPN(Hz)

16000

+0.2 μV

········· NBchirp ——— 2-1-2

25.6 ms

FIGURE 3

FIGURE 4

1-OCTAVE-WIDE DERIVED RESPONSES

FIGURE 5

A

1-OCTAVE-WIDE DERIVED RESPONSES

B

1-OCTAVE-WIDE DERIVED RESPONSES

FIGURE 6

0.5-OCTAVE-WIDE DERIVED RESPONSES

500 Hz	2000 Hz

DR2830 ... DR5660
DR2000 ... DR4000
DR1410 ... DR2830
DR1000 ... DR2000
DR707 ... DR1410
DR500 ... DR1000
DR354 ... DR707
DR250 ... DR500

········ NBchirp
——— 2-1-2

+0.05 μV

25.6 ms

FIGURE 7