Visualizing the Genetic Heterogeneity of Major Depressive Disorder using GDVIS

Anaïs B. Thijssen[1,2,3], Yuri Milaneschi[2,3,4], Meike Bartels[2,5], Brenda W.J.H. Penninx[2,4], Joëlle Pasman[1,3,6], Karin J.H. Verweij[1,2,3] and Wouter J. Peyrot[2,4]

[1]Amsterdam UMC location University of Amsterdam, Department of Psychiatry, Meibergdreef 9, Amsterdam, The Netherlands
[2] Amsterdam Public Health, Amsterdam, The Netherlands
[3]Amsterdam Neuroscience, Amsterdam, the Netherlands
[4]Amsterdam UMC location Vrije Universiteit Amsterdam, Department of Psychiatry, Boelelaan 1117, Amsterdam, The Netherlands
[5]Department of Biological Psychology, Faculty of Behavioral and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands
[6]Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden

Corresponding authors:
Anaïs B. Thijssen: a.b.thijssen@amsterdamumc.nl
Wouter J. Peyrot: w.peyrot@amsterdamumc.nl

ABSTRACT

Investigating the genetic heterogeneity of psychiatric disorders, such as Major Depressive Disorder (MDD) is a key area of research, aimed at increasing etiological insights and ultimately improving clinical outcomes via personalized treatment. Heterogeneity is typically investigated by analyzing disorder subtypes (e.g. depression with/without comorbid anxiety), where heritabilities of the genome-wide association study (GWAS) of each subtype versus controls are estimated, followed by estimation of genetic correlations between these GWASs and of these GWASs with external traits. However, it is hard to synthesize the interconnectedness of these heritability and genetic correlation estimates into a simple overarching understanding of disorder heterogeneity, especially when considering multiple subtype definitions and external traits. Here, we introduce a novel method, Genetic Distance Visualization (GDVIS), which aids interpretation of genetic differences between disorder subtypes, as well as their relation to external traits. GDVIS transforms heritabilities to genetic distances and genetic correlations to angles of a triangular geometric representation of the subtypes. GDVIS requires as input only the two genome-wide association study (GWAS) results of the two subtypes versus controls. All relevant heritabilities (resp. genetic distances) and genetic correlations (resp. angles) are derived based on geometric properties, including the subtype vs. subtype comparison (that was not part of the input). We applied GDVIS to seven subtypes of MDD using data from the UK Biobank. We show that the GDVIS-derived heritabilities and genetic correlations accurately correspond to LDscore-regression estimates. GDVIS illustrates how genetic distances between subtypes can arise from either qualitatively different genetic effects (e.g. for the subtypes of MDD with and without childhood trauma, which only share a limited portion of genetic liability), or from quantitatively different genetic effects (e.g. for the subtypes of MDD with and without recurrence, which completely share genetic risk, but with stronger effects in the subtype with recurrence). Second, we applied GDVIS to relate the MDD subtype-definitions to fifteen external traits. GDVIS visualizations show that subtype vs subtype comparisons can reveal whether the subtypes are differentially associated with the external trait, information that cannot be distilled from the usual subtype vs control comparisons. In conclusion, GDVIS is an effective tool to intuitively and comprehensively visualize the genetic heterogeneity of disorder subtypes.

INTRODUCTION

The last decades have seen a tremendous increase in sample sizes for discovery of genetic variants associated with Major Depressive Disorder, resulted in new knowledge about the underlying genetic architecture and biology[1, 2]. Increases in sample size are most efficiently achieved by adopting a simple design of contrasting individuals with a diagnosis of MDD to healthy controls. However, a diagnosis of Major Depression can be made on the basis of many different constellations of symptoms[3], making depression a far from uniform construct. This phenotypic heterogeneity complicates both fundamental research on the biological substrates of depression and likely contributes to the high variability of treatment success. To address this issue, research on the genetics of MDD has focussed on investigating subgroups of individuals that share key symptoms or clinical features. Analysing more homogenous samples can help with finding new, and more specific genetic risk loci, even at the cost of reducing sample sizes[4]. Moreover, it has the potential to help understanding the aetiology of different types of depression and aid in personalized medicine[5]. Significant progress has been made in distinguishing subtypes of MDD that have different underlying genetic influences[6-8].

Genetic heterogeneity is typically investigated by analyzing disorder subtypes (e.g. depression with/without comorbid anxiety), based on GWASs of each subtype versus controls. Typically, these two GWASs are then compared in terms of their heritability and genetic correlation with each other, and with other traits. Despite the great progress in this research[7, 8], these types of analyses face two problems. First, direct genetic comparisons of subtypes are generally overlooked, while these contain important information about the genetic distinctiveness of these subtypes. Second, when considering several subtype-definitions, the interpretation of the vast amount of heritability and genetic correlation estimates is complex, and this complexity is further increased when comparing subtypes to other traits.

Here, we introduce the Genetic Distance Visualization (GDVIS) tool to investigate direct subtype comparisons, and to intuitively visualize the heritability and genetic correlation estimates in geometrical representations. We apply GDVIS to 7 previously defined subtypes of MDD[7], contrast these subtypes to 15 external traits, and contrast these subtypes to each other.

RESULTS

Overview of methods

GDVIS visualizes the genetic relationships between two subtypes of a disorder (subtype1-cases and subtype2-cases) and controls, aiding interpretation of their genetic differences and, optionally, differences in the relationship to an external trait. GDVIS computes these relationships efficiently; to infer seven parameters including the heritabilities of subtype1-cases vs. controls, subtype2-cases vs. controls, subtype1-cases vs. subtype2-cases, and all-cases vs. controls, as well as the three genetic correlations between each comparison, GDVIS requires only the two GWAS results of subtype1-cases vs. controls and of subtype2-cases vs. controls. GDVIS-visualizations aid interpretation of the genetic correlations and heritabilities for a single subtype definition (e.g., MDD with/without comorbid anxiety), and also facilitates comparing across multiple subtype definitions (e.g., MDD with/without comorbid anxiety and MDD with/without recurrent episodes).

GDVIS transforms SNP-heritabilities and genetic correlations to Euclidian distances

and -angles, respectively (Figure 1, Figure S1). Specifically, we show that the square root of the heritability on the observed scale with 50/50 case-control ascertainment ($h^2_{obs|50/50}$) naturally corresponds to distance between the (sub)groups compared (see Methods; Figure 1a). The scale with 50/50 ascertainment provides a natural unified measure to evaluate the different subtype vs. control and subtype vs. subtype comparisons, because it does not rely on different population prevalences (as with the liability scale) or unbalanced ascertainment (as with the regular observed scale). For a comparison of the liability scale to the observed scale with 50/50 ascertainment based on[9], see Figure S2. Second, we show that the genetic correlation between the GWASs of two comparisons corresponds to the inverse cosine of the angle formed by the lines of these comparisons[10]. For example, a genetic correlation of 0 corresponds to an angle of 90 degrees (see Methods; Figure 1b).

GDVIS uses nine steps to compute the geometric representation of genetic relations (Figure 2). First, LDSC[11, 12] is applied to compute $h^2_{sub1.cases\_controls}$ based on the GWAS results of subtype1-cases vs controls ($GWAS_{sub1.cases\_controls}$), $h^2_{sub2.cases\_controls}$ based on the GWAS subtype2-cases vs controls ($GWAS_{sub2.cases\_controls}$), and the genetic correlation $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ between these two comparisons. All heritabilities presented in this paper are on the observed scale with 50/50 ascertainment, which is attained by applying $N_{effective}$ in LDSC. Second, the genetic distance of each subtype to the controls follows from the square roots of their heritabilities and defines two line segments (orange and green line in Figure 2). Third, the angle formed between these line segments is determined by the inverse cosine of the genetic correlation (black dot in Figure 2). Fourth, the genetic distance between subtype1-cases and subtype2-cases (purple line in Figure 2) is computed with the law of cosines; note that squaring this genetic distance gives the heritability of this comparison. Fifth, the two remaining angles of the triangle (at subtype1-cases and subtype2-cases; white and gray dots in Figure 2) are computed by again applying the law of cosines; note that the cosine of these angles will give the genetic correlation (i.e., $r_{g|sub1.cases\_controls,sub1.cases\_sub2.cases}$ and $r_{g|sub2.cases\_controls,sub1.cases\_sub2.cases}$). Sixth, the group all-cases (i.e., the combined set of subtype1-cases and subtype2-cases; black square in Figure 2) lies on the line segment between subtype1-cases and subtype2-cases, where the distance between all-cases and subtype2-cases is inversely proportional to the proportion of subtype2-cases among all-cases (i.e., the larger the proportion of subtype2-cases, the closer all-cases is to subtype2-cases; Figure S1). Seventh, the genetic distance from all-cases to controls can be inferred using the law of cosines (gray line in Figure 2). Eighth, with all the distances known, the remaining angles (i.e., $r_{g|sub1.cases\_controls,all.cases\_controls}$ and $r_{g|sub2.cases\_controls,all.cases\_controls}$ that are created with the line segment from all-cases to controls can be inferred from the law of cosines as well. Ninth, the population mean lies on the line segment between all-cases and controls, where the distance between population mean and all-cases is inversely proportional to the prevalence of all-cases (i.e., the higher the prevalence, the closer all-cases is to is to the population mean; asterisk in Figure 2). GDVIS employs filters on the input to only visualize subtypes that are genetically different based on $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ being significantly different from 1 and/or significant different values of the heritabilities $h^2_{sub1.cases\_controls}$ and $h^2_{sub2.cases\_controls}$. All GDVIS-derived values of the heritabilities and genetic correlations are validated with empirical analyses (see below).

The geometric GDVIS-representation of the subtypes can be extended to include an

external trait (ext), resulting in a 3D-visualization. For this, GDVIS additionally requires the GWAS of the external trait, which can be either a case-control trait (e.g. schizophrenia) or a continuous trait (e.g. BMI). First, h^2_{ext}, $r_{g|ext,sub1.cases\_controls}$ and $r_{g|ext,sub2.cases\_controls}$ are estimated with LDSC and converted to distances and angles respectively (see above, Figure 1, and Methods). While the angles have a clear interpretation for continuous traits, we note that the length of the line segment has no direct interpretation as genetic distance for continuous traits. The line segment of the external trait will intersect the subtype triangle at the population mean, resulting in a different placement of the controls of the external trait relative to the controls of the subtype triangle, reflecting that these groups of controls are not equal to each other. For both initial trait and external trait, the population mean lies on the respective case-control line segments, with its distance from the cases inverse proportional to the population prevalence. When the external trait is continuous, the population mean lies at the midpoint of the external trait line segment. GDVIS will derive estimates for $r_{g|ext,all.cases\_controls}$ and $r_{g|ext,sub1.cases\_sub2.cases}$, which are validated in empirical analyses (see below).

GDVIS can be used to compare two subtype-definitions to each other in a 3D-visualization (e.g. subtype A: MDD with/without sleep or weight increase vs. subtype B: MDD with/without comorbid anxiety). The required input of this GDVIS extension are the GWASs of subtype1-cases vs. controls and subtype2-cases vs. controls, for subtype-definition A and subtype-definition B (i.e. 4 GWASs in total). First, GDVIS computes the representations for both single subtype definitions as above. Next, the genetic correlations between the subtypes are used to infer the relation between them, and are translated into corresponding angles to form the geometrical representation. GDVIS will derive estimates for $r_{g|subA1.cases\_subA2.cases,subB1.cases\_subB2.cases}$, which are validated in empirical analyses (see below).

We applied GDVIS to seven previously identified genetic subtype definitions of depression from the UK-Biobank, **with** (subtype1-cases)/ **without** (subtype2-cases) the following features: childhood trauma, comorbid anxiety, early onset (first episode < age 30 years), recurrent episodes, high severity, sleep or weight increase, and suicidality (adapted from[7, 13]). Depression cases and controls were defined using the MHQ questionnaire[14], yielding 30,431 MDD cases and 83,202 controls (Table S1). The subtype with the largest case-control unbalance was MDD with/without suicidality ($N_{sub1.cases}$ = 19,998; $N_{sub2.cases}$ = 10,433), while the most balanced was MDD with/without childhood trauma ($N_{sub1.cases}$ = 14,146; $N_{sub2.cases}$ = 16,285). For each subtype we ran two GWASs (with-feature vs. controls and without-feature vs. controls). To empirically validate the GDVIS-derived values, we also ran the GWASs with-feature vs. without-feature, and the GWAS all-cases vs. controls. We considered GWASs of the following 15 external traits (Table S2), including binary traits: anxiety disorder (ANX)[13], bipolar disorder (BIP)[15], major depressive disorder (MDD)[2], schizophrenia (SCZ)[16], childhood trauma (CT)[17]; and continuous traits: body mass index (BMI)[18], educational attainment (EA)[19], Wellbeing[20], Neuroticism[21]; and five Genomic-SEM cross-disorder factors: substance use dependence factor[22], compulsive factor[22], internalizing factor[22], neurodevelopmental factor[22], thought disorders factor[22], and hierarchical factor[22].

GDVIS-derived heritabilities and genetic correlations of MDD-subtypes correspond to LDSC-estimates

The input for GDVIS are the GWAS results of subtype1-cases vs. controls and subtype2-cases vs. controls, and the proportion of subtype1-cases to subtype2-cases. LDSC is used to estimate corresponding heritabilities and the genetic correlation. Based on these values, GDVIS derives the heritabilities $h^2_{sub1.cases\_sub2.cases}$, $h^2_{all.cases\_controls}$, and the genetic correlations $r_{g|sub1.cases\_controls,sub1.cases\_sub2.cases}$, $r_{g|sub2.cases\_controls,sub1.cases\_sub2.cases}$, $r_{g|all.cases\_controls,sub1.cases\_controls}$, $r_{g|all.cases\_controls,sub2.cases\_controls}$. Applying GDVIS to the depression subtypes, we show that GDVIS-derived values match closely to LDSC-estimated values, with an average absolute difference of 0.002 (range 0 – 0.004) for the heritabilities and 0.013 (range 0 – 0.077) for the genetic correlations (Table 1). The differences were slightly larger for the genetic correlations, with the larger differences corresponding to larger standard errors in the LDSC-estimates. Specifically, the largest difference was observed for the genetic correlation between the subtype sleep or weight increase, where the GDVIS-derived genetic correlation was 0.413, while the LDSC-estimated value was 0.336. We note that this difference of 0.077 is well within the LDSC standard error of 0.227 (all absolute delta's were within 0.59 standard error units of the LDSC estimate). We conclude that GDVIS adequately derives the two heritabilities and three genetic correlations, based on input of only the GWAS results of subtype1-cases vs. controls and subtype2-cases vs. controls.

GDVIS yields insights into several depression subtypes

The heritability of all-cases versus controls ($h^2_{all.cases\_controls}$) was 0.092 (se 0.008). After dividing the cases into subtypes, heritability for the subtype with the feature ($h^2_{sub1.cases\_controls}$ ranging from 0.110 (se 0.010) to 0.165 (se 0.017)) was consistently higher than that for the subtype without the feature ($h^2_{sub2.cases\_controls}$ ranging from 0.047 (se 0.013) to 0.084 (0.010)), indicating that MDD subtypes characterized by the presence of a given subtype-feature exhibit stronger genetic influences than cases in which that same feature is absent. The genetic correlations $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ ranged from 0.601 (se 0.061) for childhood trauma to 1 (se 0.129) for recurrent episodes. GDVIS revealed substantial differences in heritability of the direct comparison between the subtypes ($h^2_{sub1.cases\_sub2.cases}$, ranging from 0.018 for recurrent episodes to 0.105 for childhood trauma), demonstrating that some subtypes provide a more profound separation between MDD cases than others (Figure 3).

The distance between subtype1-cases and subtype2-cases can reflect (1) either qualitative distinct genetic influences, indicating different genetic effects are associated with the different subtypes. This can be as seen with an $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ of less than 1 (e.g., mostly for the subtype-definition of suicidality); (2) a quantitative difference in magnitude of effects reflected in a difference between $h^2_{subtype1.cases\_controls}$ and $h^2_{subtype2.cases\_controls}$ (e.g. subtype definition of high severity); or (3) a combination of both (e.g. subtype definition of early onset). The subtype definition based on childhood trauma showed the greatest genetic distance between subtype1-cases and subtype2-cases ($h^2 = 0.105$), which was even slightly larger than the distance all-cases vs. controls ($h^2 = 0.092$). The GDVIS-visualization is thus able to intuitively condense a large amount of estimates (including the 7x4=28 heritabilities and 7x5=35 genetic correlations) into seven subtype

plots, which aids interpretation of this wealth of information at a glance (see Table 1, Figure 3).

Our main analyses were based on the UK Biobank MHQ definition[14] using without excluding related individuals to optimize sample size and power (see Methods). In secondary analyses we explored the effect of using unrelated individuals and of using a broader depression definition. First, GDVIS visualizations based on unrelated individuals were similar to those of related individuals (Figure S3). Second, we investigated the effect of using the broader depression definition as applied in the study by Nguyen et al.[7], which furthermore uses mostly unscreened controls (see Methods). Here the GDVIS visualizations were overall comparable, with slightly more pronounced differences for the subtypes of MDD with/without early onset and MDD with/without sleep or weight increase (Figure S4).

GDVIS compares subtypes to external traits
GDVIS allows for the comparison of disorder subtypes to an external trait. We applied GDVIS to investigate the genetic relation of the depression subtypes to 15 external traits (Table S2), including four dichotomous traits: anxiety disorder (ANX)[13], bipolar disorder (BIP)[15], major depressive disorder (MDD)[2], schizophrenia (SCZ)[16]; five continuous traits: childhood trauma (CT)[17], body mass index (BMI)[18], educational attainment (EA)[19], Wellbeing[20], Neuroticism[21]; and six Genomic-SEM cross-disorder factors: substance use dependence factor[22], compulsive factor[22], internalizing factor[22], neurodevelopmental factor[22], thought disorders factor[22], and hierarchical factor[22].

The GDVIS input consists of the GWAS of subtype1-cases, subtype2-cases and the additional GWAS for the external trait. GDVIS only generates plots after verifying that the genetic correlation estimates derived from these GWAS are compatible with each other (see Methods). The subtype recurrence with external traits did not pass this requirement, likely due to error terms in LDSC estimates (e.g., $r_{g|sub1.cases\_controls,sub2.cases\_controls} = 1$ (se 0.129), but $r_{g|ext,sub1.cases\_controls} = 0.744$ (se 0.192) and $r_{g|ext,sub2.cases\_controls} = 0.559$ (se 0.223) for anxiety disorder as external trait). The external trait childhood trauma did not pass the checks for the subtype of childhood trauma and high severity. GDVIS builds on the geometric subtype representation described above. To extend to the external trait, the genetic correlations $r_{g|ext,sub1.cases\_controls}$ and $r_{g|ext,sub2.cases\_controls}$ were computed using LDSC. Subsequently, GDVIS derived estimates for $r_{g|ext,sub1.cases\_sub2.cases}$ using the rule of cosines and other geometrical computations (see Methods). The GDVIS-derived estimates corresponded well to LDSC estimates, with an average absolute difference of 0.034 (range 0.0002 – 0.156; Figure 4, Table S3).

Several subtype-external trait pairs showed large differences between the genetic correlations $r_{g|ext,sub1.cases\_controls}$ and $r_{g|ext,sub2.cases\_controls}$ (Figure 5). For example, the correlation of the subtype with weight increase vs. controls with BMI was 0.371 (se 0.032), while the subtype without weight increase vs. controls correlated 0.012 (se 0.036) with BMI. Most genetic correlations $r_{g|ext,sub1.cases\_sub2.cases}$ were non-zero, ranging from -0.878 for the comparison MDD with high severity vs. MDD without high severity to the external trait wellbeing, to 0.854 for the comparison MDD with high severity vs. MDD without high severity to the internalizing factor.

GDVIS-results show that the direct subtype1-cases vs. subtype2-cases comparison provides important additional insight about the distinguishing potential of external traits on subtypes, that cannot be obtained by considering only the subtype1-cases vs. controls and

subtype1-cases vs. controls comparison. For instance, for the subtype of high severity with the internalizing factor, the $r_{g|ext,sub1.cases\_controls}$ and $r_{g|ext,sub2.cases\_controls}$, do not reveal that the internalizing factor is a separating trait (i.e., a trait that is differentially associated with the subtypes), with values of 0.822 and 0.905, respectively. Still, $r_{g|ext,sub1.cases\_sub2.cases}$ is 0.854 (Figure 6A), indicating that the internalizing factor is differentially related to the subtypes. The 3D visualizations intuitively show that these seemingly contradictory findings are compatible, because they take into account how the subtypes interconnect. Specifically, the comparisons subtype1-cases vs. controls (orange line in Figure 6A), subtype2-cases vs. controls (green line), subtype1-cases vs. subtype2-cases (purple line) all form their own angle with the external trait (blue line), with the angle between subtype1-cases vs. subtype2-cases being driven by both the heritabilities of and the genetic correlation between subtypes1-cases vs. controls with subtype2-cases vs. controls. Figure 6B demonstrates the relation of the external trait BMI with the MDD subtype of with/without sleep or weight increase. BMI has a higher genetic correlation with the MDD with sleep or weight increase vs. controls (0.371) than with MDD without sleep or weight increase vs. controls (0.012), which is also reflected in the genetic correlation of the MDD with sleep or weight increase vs. MDD without sleep or weight increase vs. control with BMI (r_g = 0.683). This indicated that the subtype sleep or weight increase is closely genetically related to BMI, but is still distinct from BMI.

Visualizations of 88 subtypes vs. external traits are presented interactively at the GDVIS shiny app: https://gdvis.shinyapps.io/gdvis/ (88 visualizations resulting from 7 subtypes x 15 external traits minus all the comparisons with the subtype recurrence and the two comparisons with the external trait childhood trauma that were not geometrically valid; see Methods).

GDVIS compares two subtype definitions to each other
We further applied GDVIS to investigate how genetically distinct the subtype definitions were from each other. We excluded each subtype definition pair that was not compatible in geometric space resulting in 12 subtype pair comparisons (omitting all of the combinations with recurrence and most of the combinations with high severity; the two subtypes where $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ was not significantly different from 1; see Methods). GDVIS-derived values of $r_{g|subA1.cases\_subA2.cases,subB1.cases\_subB2.cases}$ were compared to LDSC-estimated values (Figure 7, Table S4). The mean absolute difference was 0.05 (range 0.02-0.16), which was larger than the mean absolute difference for the comparisons with external traits; we note that this reflects the larger LDSC standard errors around these estimates due to smaller GWAS sample sizes, and that GDVIS-derived values were all well within these standard error ranges. Results show that most subtypes were genetically divergent, as $r_{g|subA1.cases\_subA2.cases,subB1.cases\_subB2.cases}$ values were not close to 1 (absolute range 0.072-0.732). For example, for the subtype definitions of comorbid anxiety and childhood trauma, $r_{g|subA1.cases\_subA2.cases,subB1.cases\_subB2.cases}$ was 0.463, resulting in a cross-shaped geometric representation (Figure 8). GDVIS visualizations of the 11 other subtype-subtype comparisons can be found in Figure S5 and interactively in the GDVIS shiny app. In secondary analyses, we compared the genetic correlations to phenotypic correlations; differences were found, but all phenotypical correlation estimates were within the 95% confidence interval of the genetic correlation estimates (Figure S6).

DISCUSSION

We developed a novel tool, GDVIS, to visualize the genetic relationship between subtypes of a disorder, their relationship to external traits, and their interrelationship. GDVIS transforms heritabilities to genetic distances, and genetic correlations to angles, resulting in triangular geometric representations of the genetic architecture of the subtypes. In this way, the output of GDVIS synthetizes the multiple parameters describing complex relationships between subtypes and traits into a simple overarching informative overview. When comparing multiple subtypes, GDVIS visualizations provide an intuitive way to identify which subtypes may offer the most potential for future clinical utility based on their genetic distinctiveness. We furthermore showed that all GDVIS-derived estimates, which are based on the geometric representation of genetic relationships, were comparable to LDSC-based estimates. We applied GDVIS to MDD subtypes, and make three main observations. First, by integrating the subtype1-cases vs. subtype2-cases relation with both subtype vs. control comparisons (Figure 3), GDVIS shows that subtype1-cases vs. subtype2-cases genetic differences either arise from qualitative different genetic effects (i.e., $r_{g|sub1.cases\_controls,sub2.cases\_controls} < 1$) or from quantitative different genetic effects (i.e., $h^2_{sub1.cases\_controls} \neq h^2_{sub2.cases\_controls}$). For example, the genetic difference between the subtypes with vs. without recurrent episodes reflects purely quantitative genetic effects, meaning that while the same risk alleles are involved in both subtypes, the magnitudes in the subtype with recurrent episodes are consistently larger across all SNPs. This finding is in line with theoretical predictions about the etiology of recurrent MDD episodes[23]. On the other hand, the most differentiating subtype is that of MDD cases with vs. without childhood trauma. Here, the genetic difference between the subtypes is mostly qualitative, with different genetic influences distinguishing the subtypes ($r_{g|sub1.cases\_controls,sub2.cases\_controls} = 0.60$). While childhood trauma is an informative subtype due to its established clinical relevance[24-26], we note that it is not a feature present only within MDD (as e.g., high severity of MDD), but also in controls, and caution that not all features present in both cases and controls (e.g. height) bear clinical relevance. Second, the subtype1-cases vs subtype2-cases comparison, which contains information directly distinguishing subtypes, provides important added value over the subtype vs. controls comparisons alone (Figure 5 and 6). Specifically, the correlation of this direct comparison to external traits provides insight in how the subtypes are distinguishable—this cannot be uncovered from the respective subtype vs control comparisons, that largely capture the overall MDD signal. Third, while GDVIS-visualizations for single subtype-definitions might seem comparable (e.g. the clinically most severe subtype has the highest heritability across the subtypes), we show that the subtype definitions are genetically distinct (Figure 8). This finding parallels the low phenotypic correlations between the subtypes.

Multiple genetic methods exist to compare subtypes and disorders. First, a method often employed to evaluate genetic differences between subtypes involves estimating the heritabilities of the GWAS of subtype1-cases vs controls and subtype2-cases vs controls, as well as the genetic correlations between these two comparisons with external traits[7, 8]. GDVIS extends this approach by offering the geometrical visualization as well as by deriving and integrating the subtype1-cases vs subtype2-cases comparison, which adds crucial information on the genetic distinctiveness of the subtype. Second, a method applied to investigate genetic heterogeneity of a disorder or to compare different disorders is Genomic Structural Equation Modelling[22, 27, 28]. The goal of GSEM is different as it moves beyond the observed variables to create latent factors that summarize the coherence between symptoms

or disorders. GSEM does not provide the direct geometric representation of heritabilities and genetic correlations like GDVIS. Third, GDVIS builds on findings from $F_{ST,causal}$ (included in CC-GWAS[10]), a previous method based on genetic geometric representations, but is more versatile as it allows for the comparisons of disorder subtypes in addition to the comparison of different disorders, and can extend into 3D geometric space to include the relation to external traits. In contrast to $F_{ST,causal}$, GDVIS infers the genetic distance based on all SNPs and not just independent causal SNPs. Furthermore, GDVIS-derived estimates are empirically validated, while CC-GWAS is a purely analytical construct. Fourth, for subtype-features also present in controls, G x E interactions are a potential method to compare subtypes. For instance, research has addressed the question of whether the genetic risk for MDD interacts with childhood trauma[29]. GDVIS focusses specifically on the difference between subtypes (in comparison to controls), whereas interaction analysis focuses on differences in direction and magnitude across environmental strata.

Findings from GDVIS align with and expand on previous studies investigating genetic subtypes of MDD. First, a study investigating eight different depression subtype definitions relied on 16 subtype vs control GWASs[7]. They showed that three subtype definitions had distinct heritabilities and found differential genetic overlap of all subtypes with external traits. The study concluded that genetic heterogeneity exists within MDD and that genetic evidence shows that subtypes have partially distinct etiology. GDVIS analyses extend this approach by using the subtype vs. subtype comparison and show that MDD subtypes are even more genetically differentiated than previously recognized. Furthermore, GDVIS visualization were shown to be especially useful for intuitive comparison of multiple subgroups. Second, the subtype of typical versus atypical MDD has been of increasing interest in the literature[6, 30]. Research into the genetic underpinnings of this subtype have approximated the phenotype by focusing on hypersomnia and hyperphagia features. No significant heritability differences of the atypical vs control comparison and the typical vs control comparison have been found[8, 31]; however, as GDVIS illuminates, the subtypes are genetically different when simultaneously taking the genetic correlation between these comparisons into account. This is supported by the polygenic score (PGS) analyses conducted in these previous studies, which showed distinct patterns of association with external traits[8, 31]. Third, age at onset of depression was recently investigated using a large Nordic dataset, contrasting early age at onset (< 25 years) with late onset (>50 years)[32]. As expected when comparing more extreme ends of the distribution, the subtypes were more genetically differentiated, with a genetic correlation of 0.58 compared to the genetic correlation of 0.79 that we obtained when splitting the age at onset as below or above 30 years. Consistent with our findings, the heritability of the early onset subtype vs. controls was higher than the late onset subtype vs. controls. The subtypes were compared directly in a subtype vs. subtype GWAS; however, the interconnectedness of the three GWASs was not fully resolved and difficult to intuitively comprehend as the heritabilities were expressed on different liability scales (rather than the observed 50/50 scale) and no genetic correlations were estimated of the subtype vs. subtype GWAS with the subtypes vs. controls GWASs or with external traits.

GDVIS has several limitations. First, GDVIS necessitates the use of heritabilities on the observed scale with 50/50 ascertainment and this scale is less widely used in the literature, limiting intuitive heritability interpretation. However, the benefit of this scale is that it provides a uniform distance measure for disorders with different population

prevalences and for subtype vs. subtype comparisons. We provide Figure S2 to aid intuition of the relation between the 50/50 scale and the liability scale heritabilities. Second, GDVIS does not compute standard errors, limiting the interpretation of the significance of results. However, the purpose of GVIS is not in significance testing, and we note that computing standard errors adds a disproportional layer of complexity due to the intrinsic challenges to compute standard errors of genetic correlations requiring block jackknife within LD-score regression[11]. Third, as has been exemplified by our application of GDVIS to MDD, not all subtypes, subtype combinations, or relations between subtypes and external traits can be visualized. However, this is either because there are no genetic differences between the subtypes, or because the LDSC-estimates are incompatible with each other. Furthermore, we note that empirical and derived values were in correspondence for the subtypes that could be visualized. Fourth, although not a limitation of the method per se, we note that GDVIS relies on which case and subtype definitions are used, which can remain slightly arbitrary and does have some impact on the visualizations (Figure S4). Fifth, although genetic space can be thought of as a >1,000,000-dimensional space (i.e. the number of independent SNPs), visual geometric representations are restricted to 3D space; hence, plotting more than one subtype and one external trait or more than two subtype-definitions is not possible. Sixth, the GDVIS software is currently only able to compare two subgroups of a disorder that together form the complete set of cases. Therefore, GDVIS is not suitable for subgroups that consist of three groups (e.g. low impairment, medium impairment, high impairment). A future research direction would be to extend GDVIS to allow for comparison of three subgroups. Finally, this study made use of only individuals of European ancestry, thereby limiting the generalizability of findings to individuals of other ancestries.

In conclusion, GDVIS provides a novel way of understanding the genetic architecture of disorder heterogeneity. GDVIS is a freely available tool that can be used with any dichotomous subtype of a disorder. GDVIS requires simple and minimal input data, which it uses to efficiently compute the visualizations, while automatically imposing checks and filters to ensure that only valid visualizations are produced. This makes the GDVIS tool an invaluable tool in subtype research.

METHODS

The geometrical GDVIS-representation of heritability and genetic correlation
GDVIS transforms heritabilities to genetic distances, and genetic correlations to angles (Figure 1). As demonstrated by the derivation of $F_{ST,causal}$,[10] the variance explained by a single causal SNP on the 50/50-ascertainment scale of a dichotomous phenotype ($\sigma^2_{50/50}$) corresponds to the squared difference in standardized allele frequency, thus representing a Euclidian distance metric. Specifically, for genotype (G) and dichotomous phenotype (D) with 50% ascertainment,

$$\sigma^2_{50/50} = \beta^2_{D \sim G} = cov(s(G), s(D))^2 \approx \left(\frac{E[G*s(D)]}{\sqrt{2p(1-p)}} \right)^2 \approx \left(\frac{p_1 - p_0}{\sqrt{2p(1-p)}} \right)^2 \quad \text{Eq 1}$$

where p is the population allele frequency, p_1 the frequency in cases, p_0 the frequency in controls, and $s(G)$ and $s(D)$ are the standardization to a mean of 0 and variance of 1 for D and G respectively. Previous work derived this correspondence for independent causal SNPs[10]; here we show via extensive empirical analyses that this extends to a genome-wide

correspondence of the square root of heritability as distance metric (Table 1, Figure 4, Figure 7). Thus, the GWAS of a dichotomous phenotype D can be represented as a line segment with length $\sqrt{h^2_{50/50}}$ in Euclidian space.

When considering two GWASs of two dichotomous phenotypes (D_A and D_B), they form two line-segments in Euclidian space. These two line-segments form an angle α_{D_A,D_B}:

$$\alpha_{D_A,D_B} = \arccos\left(r_{g\,|\,D_A,D_B}\right) \qquad\qquad \text{Eq 2}$$

This equation has previously been derived for independent causal SNPs[10]; here we show via extensive empirical analyses that this extends to a genome-wide correspondence of genetic correlation estimated with LDSC[11] (Table 1, Figure 4, Figure 7).

GDVIS follows the same rationale as the genetic distance measure $F_{ST,causal}$: the analytically derived F_{ST} at causal independent SNPs for cross-disorder case/control comparisons[10]. However, GDVIS differs in the following key aspects. First, GDVIS infers the genetic distance based on all SNPs and not just independent causal SNPs. Second, GDVIS can be applied on disorder subtypes and not only for cross-disorder comparisons. Third, GDVIS-estimates of genetic distances and angles are validated with empirical results of heritabilities and genetic correlation, while $F_{ST,causal}$ is purely an analytical construct used to increase power of cross-disorder case vs. case GWAS.

We note that GDVIS refers to the genetic distance as square root of the heritability, but there is an important conceptual difference between genetic distance and heritability: genetic distances have an interpretation between any two (potentially overlapping) groups, while heritability describes the variance explained by genetic effects in a phenotype in the population. We choose to use the term square root of heritability and genetic distance interchangeably, because of the correspondence in empirical estimates and because of the widespread intuitive understanding of estimates of the heritability.

Thus transforming heritabilities to genetic distances and genetic correlations to angles (Figure 1) allows for geometrical computations based on heritabilities and genetic correlations (see below).

GDVIS method for one subtype-definition
GDVIS uses nine steps to compute the geometric representation of the genetic architecture of a subtype.

Step 1: Estimate heritabilities and genetic correlation
GWAS results of subtype1-cases vs controls ($GWAS_{sub1.cases\_controls}$) and subtype2-cases vs controls ($GWAS_{sub2.cases\_controls}$) are used to compute $h^2_{sub1.cases\_controls}$, $h^2_{sub2.cases\_controls}$ and $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ with LDSC (Bulik-Sullivan 2015a, 2015b). The heritabilities are expressed on the observed scale with 50/50 ascertainment, which is computed by using $N_{effective}$ in LDSC. For unrelated individuals, $N_{effective}$ is computed by $4/(\frac{1}{N_{cases}} + \frac{1}{N_{controls}})$ (refs.[11, 12, 33, 34]). For related individuals, $N_{effective}$ for each variant is computed by $\frac{1}{0.5SE^2 f(1-f)}$, where f is the allele frequency of the genetic variant and SE is the standard error of the effect of the variant in the linear mixed

model from fastGWA[35, 36]. When the LDSC-estimated genetic correlation value exceeds 1, it will be capped at 1 for the geometrical computations.

Step 2: Compute subtype1.cases_controls and subtype2.cases_controls line segments
Taking the square root of the heritabilities obtained from step 1 gives the genetic distance from subtype1-cases to controls (orange line) and subtype2-cases to controls (green line), i.e. the length of the line segments in the GDVIS-representation.

Step 3: Compute the angle between subtype1.cases_controls and subtype2.cases_controls
The inverse cosine of $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ gives the angle formed by the respective line segments at the controls in radians (represented by the black dot). We note that we report angles in degrees.

Step 4: Compute sub1.cases_sub2.cases line segment
Using the length of the line segments from $subtype1.cases\_controls$ and $subtype2.cases\_controls$ and their angle, the length of the line segment $subtype1.cases\_subtype2.cases$ (purple line) is computed with the law of cosines: $\sqrt{A^2 + B^2 - 2 * A * B * \cos(\gamma)}$ where A is the length of $subtype1.cases\_controls$, B is the length of $subtype2.cases\_controls$ and γ is the angle between these lines. Squaring the resulting line length of $subtype1.cases\_subtype2.cases$ gives its GDVIS-derived heritability (which is empirically compared to LDSC estimates; see below and Table 1).

Step 5: Compute angles with sub1.cases_sub2.cases line segment
The angle formed between the line segment of $subtype1.cases\_controls$ (orange line) and $subtype1.cases\_subtype2.cases$ (purple line) can be computed using the law of cosines: $\text{acos}((C^2 + A^2 - B^2)/(2AC))$, where C is the length of the line segment for $subtype1.cases\_subtype2.cases$, A is the length of the line segment for $subtype1.cases\_controls$ and B is the length of the line segment for $subtype2.cases\_controls$. The angle between the line segments of $subtype2.cases\_controls$ (green line) and $subtype1.cases\_subtype2.cases$ (purple line) can be computed similarly. Taking the cosine of the angle will give the genetic correlation.

Step 6: Compute all.cases_sub1.cases and all.cases_sub2.cases line segments
The all-cases group is the union of subtype1-cases and subtype2-cases. Note that for the $all.cases\_controls$ line segment to be equal to the overall disorder-controls line segment, the subtype1-cases and subtype2-cases together need to cover the complete set of cases. The point corresponding to the all-cases group all-cases lies on the line segment of subtype1-cases and subtype2-cases. The proportion of subtype1-cases to subtype2-cases determines where along the line the all-cases point falls. Specifically, the proportion of subtype1-cases to all-cases times the length of $subtype1.cases\_subtype2.cases$ determines the distance from all-cases to subtype2-cases: $length_{sub2.cases\_all.cases} = \frac{N_{sub1.cases}}{N_{all.cases}} *$ $length_{sub1.cases\_sub2.cases}$. For example, a larger proportion of subtype1-cases will result in all-cases being closer to the subtype1-cases point. The length of subtype1.cases to all-cases is the length of $subtype1.cases\_subtype2.cases$ minus the length of $subtype2.cases\_all.cases$.

Step 7: Compute all.cases_controls line segment

To get the length of the line segment from all-cases to controls (gray line), the law of cosines can be used: $\sqrt{B^2 + E^2 - 2BE\cos(\alpha)}$, where B is the length of the line segment from subtype2-cases to controls (obtained in step 2), E is the length of the line segment from all-cases to subtype2-cases (obtained in step 6) and α is the angle between these lines in radians (obtained in step 5). Squaring this distance will give the heritability.

Step 8: Compute angles with all.cases_controls line segment
The angle formed between the line segment $all.cases\_controls$ and $subtype1.cases\_controls$ can be computed with the law of cosines: $\text{acos}((F^2 + A^2 - D^2)/(2*A*B))$, where F is the length of $all.cases\_controls$, A is the length of $subtype1.cases\_controls$ and D is the length of $all.cases\_subtype1.cases$. The angle between the line segment $all.cases\_controls$ and $subtype2.cases\_controls$ can be computed similarly. The resulting angles are converted to genetic correlations by taking the cosine.

Step 9: Compute the population mean
The population mean lies on the line segment from all-cases to controls, where the distance from controls to the population mean is related to the population prevalence of the disorder (analogue to step 6). This distance is calculated by multiplying the length from $all.cases$ to controls with the population prevalence.

Supplementary validation analyses
The $GWAS_{sub1.cases\_sub2.cases}$ will often not be available, and is not necessary as it can be computed from GDVIS. However, when $GWAS_{sub1.cases\_sub2.cases}$ is available, supplementary validation analyses can be performed. The $GWAS_{all.cases\_controls}$ can also be used for validation, which is specifically relevant when it is uncertain if the union of sub1.cases and sub2.cases form to full set of cases (see step 6 above). The GDVIS-software has the option to compare its derived values based on these GWASs to the LDSC-estimated values. Specifically, for $h^2_{sub1.cases\_sub2.cases}$, $h^2_{all.cases\_controls}$,

$r_{g|sub1.cases\_controls,sub1.cases\_sub2.cases}$, $r_{g|sub2.cases\_controls,sub1.cases\_sub2.cases}$,

$r_{g|all.cases\_controls,sub1.cases\_controls}$, and $r_{g|all.cases\_controls,sub2.cases\_controls}$ GDVIS determines what the difference between the GDVIS-derived and the LDSC-estimated values are, and how the magnitude of this difference relates to the magnitude of the standard error of the. We present these validation steps for our analyses in Table 1.

Input filtering: check whether the subtypes are genetically distinct
Before creating the visualization of a subtype, GDVIS checks the input data to make sure that only heritable, genetically different subtypes are plotted. First, GDVIS checks whether $GWAS_{sub1.cases\_controls}$ and $GWAS_{sub1.cases\_controls}$ are significantly different from 0. Next, GDVIS checks whether there are indications that the subtypes are genetically distinct by determining (1) whether the 95% confidence interval of $r_{g|sub1.cases\_controls,sub2.cases\_controls}$ contains 1, and (2) whether the heritabilities of $GWAS_{sub1.cases\_controls}$ and $GWAS_{sub1.cases\_controls}$ are within each other's 95% confidence intervals. If the genetic correlation is not significantly different from 1 and the heritabilities do not differ

significantly, this suggests that the subtypes are not significantly different and no GDVIS visualizations will be plotted.

GDVIS method for one subtype-definition with an external trait
The geometric GDVIS-representation of a subtype can be extended to include an external trait (*ext*). For this, GDVIS additionally requires the GWAS of the external trait. To create the 3D-visualization, GDVIS computes the subtype representation as before, and then follows these steps:

Step 1: Estimate heritability and genetic correlations
The heritability and genetic correlations h^2_{ext}, $r_{g|ext,sub1.cases\_controls}$ and $r_{g|ext,sub2.cases\_controls}$ are estimated using LDSC. If the external trait is a dichotomous trait, h^2_{ext} needs to be on the observed scale with 50/50 ascertainment, which can be computed as described in step 1 above. The external trait can also be a continuous trait, for which the heritability is expressed on its natural scale. For continuous traits, only the GDVIS angles are of interest; the transformation of heritability to genetic distance is less informative because it does not connect defined groups like cases and controls. Next, the genetic correlation of the external trait with $GWAS_{sub1.cases\_controls}$ and $GWAS_{sub2.cases\_controls}$ is calculated using LDSC.

Step 2: Compute the ext.cases_ext.controls or external trait line segment
The line length from the external cases to external controls is obtained by taking the square root of the heritability from step 1. Note that the external controls are not the same as the controls used in the subtype triangle, which is reflected by the two points not being in the same geometrical space. When the external trait is continuous, the line segment will not connect cases to controls, and the ends are not well-defined, which is visually represented by a fading line.

Step 3: Compute the angles between the line segment of ext.cases_ext.controls with the line segments of subtype1.cases_controls and subtype2.cases_controls
The angle formed between the line segment $ext.cases\_ext.controls$ with $subtype1.cases\_controls$ and $subtype1.cases\_subtype2.cases$ can be computed by taking the inverse cosine of the genetic correlations calculated in step 1.

Step 4: Compute the population mean point for the external trait
The line segment of the external trait will intersect the subtype triangle at the population mean. To calculate where on the external trait line segment the population mean lies, the population prevalence is used as described above (in step 9 of the section '*GDVIS method for one subtype-definition*'). For continuous traits, the population mean is represented by the middle of the line segment.

Supplementary validation analyses
If the $GWAS_{sub1.cases\_sub2.cases}$ and/or $GWAS_{all.cases\_controls}$ are available, GDVIS has the option to compare its derived values based on the GWASs to the LDSC-estimated values, also for the external trait. Specifically, for $r_{g|all.cases\_controls,ext}$, and $r_{g|sub1.cases\_sub2.cases,ext}$ it is determined what the difference between the GDVIS-derived and the LDSC-estimated values are, and how the magnitude of this difference relates to the

magnitude of the standard error of the LDSC estimate. We present these validation steps for our analyses in Figure 4 and Table S3.

Input filtering: check compatibility for geometrical representation
Due to inherent variance of LDSC estimates, it may occur that the subtype and external trait are not compatible for geometric representation. Specifically, in some instances $r_{g|sub1.cases\_con,sub2.cases\_con}$ can be very close to 1 (e.g., 0.99) while the correlation with the external trait ($r_{g|sub1.cases\_con,ext}$ and $r_{g|sub2.cases\_con,ext}$) differ considerably (e.g., 0.5 vs 0.3). GDVIS filters for such inconsistencies. First, GDVIS computes a helper triangle and determines whether this is a valid triangle. The helper triangle is the base of a triangular pyramid which has the controls as apex with three sides going down from the controls to the base which are all 1. The three face angles at the controls correspond to $r_{g|sub1.cases\_con,ext}$, $r_{g|sub2.cases\_con,ext}$, and $r_{g|sub1.cases\_con,sub2.cases\_con}$. For the base triangle to be a valid triangle, the sum of its smallest two lines has to be longer than the length of the longest line. If this is not the case, the base is not a valid triangle, indicating that the three angles $r_{g|sub1.cases\_con,ext}$, $r_{g|sub2.cases\_con,ext}$, and $r_{g|sub1.cases\_con,sub2.cases\_con}$ are not compatible with each other, in which case GDVIS will not plot the subtype with the external trait.

GDVIS method to compare two subtype definitions
GDVIS can compare two subtype definitions (definition A and definition B) to each other by simultaneously visualizing two single subtype triangles in the same geometrical space. The required input is the same as the input for the single subtype triangles. GDVIS only visualizes this geometric representation if the two single subtypes are calculated using the same cases and the same controls, which is checked by comparing $h^2_{all.A.cases\_controls}$ to $h^2_{all.B.cases\_controls}$. GDVIS uses several steps to compute this geometric representation of two subtype definitions:

Step 1: *Estimate genetic correlations and compute angles between the subtypes*
LDSC is used to compute $r_{g|subA1.cases\_con,subB1.cases\_con}$, $r_{g|subA2.cases\_con,subB1.cases\_con}$, $r_{g|subA1.cases\_con,subB2.cases\_con}$, and $r_{g|subA2.cases\_con,subB2.cases\_con}$, which are then converted to angles by taking the inverse cosine of these estimates.

Step 2: Rotate the triangle of the second subtype-definition
GDVIS starts by overlaying both individual subtype-definition triangles on top of each other, such that the $all.cases\_controls$ line segments completely align. In principle, each of the four angles based on $r_{g|subA1.cases\_con,subB1.cases\_con}$, $r_{g|subA2.cases\_con,subB1.cases\_con}$, $r_{g|subA1.cases\_con,subB2.cases\_con}$, and $r_{g|subA2.cases\_con,subB2.cases\_con}$ can be used to rotate the second triangle; GDVIS takes the angle corresponding to the genetic correlation estimate with the smallest standard error to perform the rotation. It will rotate the second triangle around the $all.cases\_controls$ line segment to match this angle (e.g., if $r_{g|subA1.cases\_controls,subB1.cases\_controls}$ is used, the second triangle will be rotated in such a way that the angle between the line segments $subtypeA1.cases\_controls$ and $subtypeB1.cases\_controls$ corresponds to the angle from $r_{g|subA1.cases\_controls,subB1.cases\_controls}$)

Supplementary validation analyses
If the $GWAS_{subA1.cases\_subA2.cases}$ and $GWAS_{subB1.cases\_subB2.cases}$ are available, GDVIS will

compare its derived value of $r_{g|subA1.cases\_subA2.cases,subB1.cases\_subB2.cases}$ to the value estimated by LDSC. Specifically, it is determined what the difference between the GDVIS-derived and the LDSC-estimated values are, and how the magnitude of this difference relates to the magnitude of the standard error of the LDSC-estimate value. We present these validation steps for our analyses in Figure 7 and Table S4.

Input filtering: check compatibility for geometrical representation
To check whether the computed values result in a geometrically realizable representation, GDVIS first computes four helper triangles and determines whether they are geometrically valid (also see above). The helper triangles are formed by the points of the four different subtypes (subtypeA1-cases, subtypeA2-cases,subtypeB1-cases,subtypeB2-cases). Each helper triangle connects three subgroup points. The length of the lines of the helper triangles are computed in the same way as the lines of the helper triangle for the subtype with an external trait. When the sum of the two smallest lines are not longer than the longest line, the helper triangle is not valid. If one of the helper triangles is not valid, this indicates that the genetic correlations computed at step 1 are not compatible with each other and the two subtypes cannot be visualized together.

Phenotype and genotype data
Data came from the UK Biobank (UKB)[37], a population-based biobank containing genetic and phenotypic information on approximately 500,000 individuals. Mental health data were collected in multiple ways, including a mental health questionnaire that was available for a subset of individuals, linked hospital records and nurse interviews. The age of the individuals varies from 37 to 73 years and 54% of participants are female. Data were accessed under application 40310. Our main analyses are restricted to individuals of European ancestry for whom genetic and mental health questionnaire data[14] were available (*N*=113,633).

Phenotype definitions
MDD cases and controls
Our main analyses rely on previously defined cases and controls of lifetime MDD[14]. Diagnosis of MDD was based on the mental health questionnaire (MHQ), which included a modified version of the Composite International Diagnostic Interview Short Form (CIDI-SF). Individuals meeting criteria for a lifetime history of depression on the CIDI-SF were classified as MDD cases. Controls were those individuals that did not meet the lifetime history of depression based on the CIDI-SF, did not show signs of depression based on the Patient Health Questionnaire-9 (PHQ9), and did not indicate that they had ever been diagnosed with depression.

In secondary analyses (*N*=374,536), we used a broader definition of depression as used in the study by Nguyen et al[7], which was based on either 1) meeting criteria for a lifetime or current diagnosis of MDD based on the CIDI-SF, a diagnostic instrument of MDD based on International Classification of Diseases version-10 from linked hospital records, or 2) having a probable depression (based on[38]), 3) self-report of a diagnosis of depression, or 4) exhibiting cardinal MD symptoms (prolonged loss of interest in normal activities or prolonged feelings of sadness or depression). Controls were semi-screened, i.e., individuals that had sought help for anxiety, nerves, tension, or depression or had used antidepressants, antipsychotics or lithium were excluded. Results of the secondary analyses using this broader MDD definition are reported in Figure S4. Genetic correlations between the GWASs based

on the MHQ MDD definition and the broader definition of depression from[7] can be found in Table S5.

Subtypes of MDD
We based our subtypes on previously proposed subtypes from[7], with some slight adjustments to make the subtypes comparable to each other (i.e., in the original study, the different subtype-definitions were based on different definitions of cases and controls). We excluded cases that did not have data on all subtypes, to ensure that each subtype-definition was based on the same individuals (N subtype range 10,433-19,998; N controls: 83,202, Table S1). Genetic correlations of our subtypes based on the broader case-control definition of depression with the original GWASs from the study by Nguyen et al[7] (also based on the broader case-control definition) can be found in Table S6.

- *MDD with or without childhood trauma:* In contrast to Nguyen et al[7], we added the subtype of MDD with or without childhood trauma. MDD cases that had experienced at least one of the following five types of trauma were considered to have MDD with childhood trauma: physical abuse, emotional abuse, sexual abuse, emotional neglect, and/or physical neglect. Different cut-offs were used to determine the presence of each type of childhood trauma. Physical abuse was determined to be present if an individual indicated that this happened at least sometimes (UKB field ID: 20488). Emotional abuse was determined to be present if it happened at least sometimes (UKB field ID: 20487). Sexual abuse was determined to be present if it happened at least rarely (UKB field ID: 20490). Emotional neglect was determined to be present if an individual indicated to never, rarely or sometimes have felt loved as a child (UKB field ID: 20489). Physical neglect was determined to be present if an individual indicated they never, rarely or sometimes had someone to take them to a doctor if needed (UKB field ID: 20491).
- *MDD with or without comorbid anxiety*: Cases were considered to either have or not have comorbid anxiety based on self-reported anxiety (UKB field ID: 20544, 20002) or the presence of an ICD-diagnosis of anxiety (UKB field ID: 41202). In the original definition by Nguyen et al[7], controls were excluded if they had an anxiety disorder. In this paper, the controls were not screened for anxiety, but remained the same across all subtypes (i.e., the MDD controls defined above).
- *MDD with or without early onset:* Individuals that reported having their first depressive episode before the age of 30 were considered to have MDD with an early onset, whereas individuals that have reported having their first depressive episode after the age of 30 were considered to have MDD without early onset (UKB field ID: 20433). In the original definition by Nguyen et al[7] individuals with an onset before the age of 30 are considered to have MDD with an early onset and individuals with an onset after the age of 44 are considered individuals with a late onset of MDD. This definition was modified to a binary distinction, because the GDVIS-visualization requires the combination of the subtypes to form the full set of all cases.
- *MDD with or without recurrent episodes:* Recurrence was based either on ICD codes (UKB field ID: 41202) specifying whether the depression was single episode or recurrent, on probable recurrent or single episode depression (UKB field ID: 20126) as defined by[38], or on the self-reported lifetime number of depressed episodes (UKB field ID: 20442). If any of these three variables indicated that the person had more

than one episode, they were considered to have the recurrent subtype. For the rest of the individuals, if any of the criteria indicated a single episode, they were considered to be of the single episode subtype.

- *MDD with or without high severity:* The original definition by Nguyen et al[7] included the question on whether a case had mild or moderate recurrent depression (UKB field ID: 20126). However, because this question is also used in the single versus recurrent subtype, we changed the definition of this subtype. We based the definition on the number of CIDI symptoms that an individual with MDD endorsed (UKB field IDs: 20446-0.0, 20441-0.0, 20536-0.0, 20532-0.0, 20449-0.0, 20450-0.0, 20435-0.0, 20437-0.0). Individuals with 5 or 6 symptoms were defined as having MDD without high severity, while individuals with 7 or 8 symptoms were defined as having MDD with high severity. Note that only 8 questions are asked in the UKB and that the maximum number of symptoms is therefore 8 instead of all 9 CIDI symptoms.

- *MDD with or without sleep or weight increase:* Individuals that experienced weight increase or increased sleep during their worst depressive episode (UKB field IDs: 20536, 20534) were considered to have MDD with sleep or weight increase. This definition differs from the definition by Nguyen et al[7] where the subtype is called MDD with atypical-like features, as they required individuals to have both increased weight and increased sleep. We made this definition less strict to increase the number of individuals with the increased sleep or weight subtype. Individuals that are considered to not have sleep or weight increase could have either experienced decreased sleep and/or weight or remained stable. Further subdividing this subtype would be of interest, but we lacked the sample size to do so.

- *MDD with or without suicidality:* Individuals with suicidality were those that indicated having thoughts of death during their worst depression (UKB field ID: 20437), or having recent thoughts of suicide or self-harm (either several days, more than half of the days or nearly every day; UKB field ID: 20513).

In contrast to Nguyen et al[7] we did not add the subtype of MDD with or without impairment, as a requirement for having MDD using the MHQ definition is that individuals need to experience impairment. We furthermore did not include the subtype of post-partum depression, as the full set of individuals with MDD cannot be as clearly separated into two subtypes (i.e., men and women that have not given birth yet cannot be placed into one of the subgroups) and the controls would need to be limited to women that have given birth as well, making the comparisons between subtypes problematic.

External traits
We investigated the genetic relation of the depression subtypes to 15 external traits (Table S2), including four dichotomous traits: anxiety disorder (ANX)[13], bipolar disorder (BIP)[15], major depressive disorder (MDD)[2], schizophrenia (SCZ)[16]; four continuous traits: childhood trauma (CT)[17], body mass index (BMI)[18], educational attainment (EA)[19], Wellbeing[20], Neuroticism[21]; and six Genomic-SEM cross-disorder factors: substance use dependence factor[22], compulsive factor[22], internalizing factor[22], neurodevelopmental factor[22], thought disorders factor[22], and a hierarchical factor representing overall liability to a psychiatric disorder[22]. GWAS sample sizes and heritabilities are reported in Table S2.

Genome-wide association study
For each subtype-definition, we ran a GWAS of MDD cases with the feature vs. controls,

MDD cases without the feature vs. controls and MDD cases with the feature vs. MDD cases without the feature.

We ran GWASs on related individuals using fastGWA, a mixed linear model for binary traits as implemented in GCTA v1.94.1[36]. We calculated the $N_{effective}$ as follows:

$\frac{1}{0.5SE^2 f(1-f)}$, where f is the allele frequency of the genetic variant and SE is the standard error of the effect of the variant[35].

We extracted HapMap3 SNPs from genotypes imputed by the UKB (UKB field ID: 22828) and performed the following quality control: minor allele frequency (MAF) > .01, Hardy-Weinberg Equilibrium (HWE) p-value > 10^{-10} and missingness > 0.5. The genetic relationship between individuals was accounted for using a sparse genetic relationship matrix constructed with GCTA (see ref.[39] for details). Covariates were age, sex, genotyping array and the first 10 genetic principal components (as determined in[39]).

As a sensitivity analysis to assess the impact of relatedness, we ran GWAS on unrelated individuals using PLINK 2[39]. We calculated the $N_{effective}$ as follows: $4 / (\frac{1}{N_{cases}} + \frac{1}{N_{controls}})$ (refs.[33, 34]). We again used the HapMap3 imputed SNPs. Covariates were age, sex, genotyping array and the first 10 principal components. The genetic correlation results between the GWASs of related individuals with GWASs of unrelated individuals ranged from 0.977 to 1.362 (Table S7).

Figure 1. **Transforming heritabilities to distances and genetic correlations to angles. Panel A**. The square root of the heritability on the observed scale with 50/50 ascertainment (   ) provides a Euclidian distance measure between the two compared groups; the line segment follows the direction from the GWAS of the class coded as 0 to the class coded as 1 (the direction is relevant for the angle). **Panel B**. The genetic correlation between two GWASs corresponds to the inverse cosine of the angle formed by the line segments of these comparisons. Note that the inverse cosine gives the angle in radians, but we report the angle in degrees. GWAS1/GWAS2: GWAS comparing two classes (e.g., cases to controls, subtype vs. subtype or subtype vs. controls).

GDVIS input

GWAS	LDSC h^2 (SE)
Cases with feature vs controls	0.161 (0.01)
Cases without feature vs controls	0.084 (0.01)

Genetic correlation	LDSC r_g (SE)
{Cases with feature vs controls} vs {cases without feature vs controls}	0.601 (0.06)

Other	
Population prevalence of all cases	0.16
Proportion of sub1 (cases with feature) in all cases	0.46

GDVIS-plot

Heritability
— 0.161
— 0.084
— 0.105
— 0.093

r_g (degrees)
● 0.601 (53°)
○ 0.700 (46°)
◐ -0.161 (99°)

■ all cases
✱ population mean

With feature
Without feature
controls

GDVIS-derived values

Heritabilities	GDVIS-derived
Cases with feature vs cases without feature	0.105
All cases vs controls	0.093

Genetic correlations	
{All cases vs controls} vs {cases with feature vs controls}	0.914
{All cases vs controls} vs {cases without feature vs controls}	0.873
{cases with feature vs controls} vs {cases with feature vs cases without feature}	0.700
{cases without feature vs controls} vs {cases with feature vs MDD without feature}	0.151

Figure 2. Overview of the GDVIS workflow to visualize a subtype. We report the GDVIS-input, the GDVIS visualization of one subtype definition, and the GDVIS-inferred values based on geometrical computations. Heritabilities are on the 50:50 scale, the length of the line segments correspond to the square root of the heritability. With feature = subtype1-cases, without feature = subtype2-cases. The estimates in the example are for MDD with/without childhood trauma in Figure 3.

A. Input to the GDVIS-tool

Subtype: with (sub1) - without (sub2)	h^2 with vs. con	h^2 without vs. con	r_g {with vs. con} vs. {without vs. con}	pop.prev	prop with
Childhood Trauma	0.161 (0.014)	0.084 (0.010)	0.601 (0.061)	0.16	0.46
Comorbid anxiety	0.139 (0.015)	0.080 (0.010)	0.845 (0.072)	0.16	0.39
Early onset	0.149 (0.014)	0.077 (0.010)	0.790 (0.071)	0.16	0.42
Recurrent episodes	0.124 (0.011)	0.047 (0.012)	1.000 (0.129)	0.16	0.64
High severity	0.132 (0.013)	0.059 (0.011)	0.953 (0.095)	0.16	0.55
Sleep or weight Increase	0.165 (0.017)	0.071 (0.009)	0.875 (0.077)	0.16	0.35
Suicidality	0.110 (0.010)	0.084 (0.016)	0.814 (0.088)	0.16	0.66

B. Color reference

C. GDVIS-derived values versus LDSC estimated values

Subtype: with (sub1) - without (sub2)	LDSC (SE) estimated	GDVIS-inferred	Delta	LDSC (SE) estimated	GDVIS-inferred	Delta	LDSC (SE) estimated	GDVIS-inferred	Delta
	h^2 with vs. without			h^2 all MDD cases vs. controls			r_g {with vs. con} vs. {with vs. without}		
Childhood Trauma	0.104 (0.016)	0.105	0.001	0.091 (0.008)	0.093	0.002	0.700 (0.039)	0.700	0.000
Comorbid anxiety	0.045 (0.017)	0.041	-0.004	0.091 (0.008)	0.093	0.002	0.650 (0.075)	0.665	0.015
Early onset	0.058 (0.016)	0.057	-0.001	0.091 (0.008)	0.094	0.002	0.683 (0.064)	0.700	0.017
Recurrent episodes	0.019 (0.015)	0.018	-0.001	0.091 (0.008)	0.092	0.000	0.998 (0.318)	1.000	0.002
High severity	0.020 (0.017)	0.023	0.003	0.091 (0.008)	0.094	0.002	0.904 (0.317)	0.873	-0.031
Sleep or weight increase	0.050 (0.019)	0.047	-0.003	0.091 (0.008)	0.094	0.002	0.759 (0.073)	0.802	0.043
Suicidality	0.036 (0.018)	0.038	0.002	0.091 (0.008)	0.093	0.001	0.510 (0.122)	0.496	-0.015
	r_g {without vs. con} vs. {with vs. without}			r_g {all MDD cases vs. con} vs. {with vs.con}			r_g {all MDD cases vs. con} vs. {without vs.con}		
Childhood Trauma	-0.154 (0.098)	-0.151	0.004	0.911 (0.017)	0.914	0.003	0.876 (0.020)	0.873	-0.003
Comorbid anxiety	0.130 (0.187)	0.163	0.033	0.950 (0.024)	0.954	0.003	0.969 (0.015)	0.967	-0.002
Early onset	0.088 (0.162)	0.115	0.027	0.949 (0.021)	0.947	-0.002	0.944 (0.018)	0.945	0.002
Recurrent episodes	1.000 (0.863)	1.000	0.000	1.000 (0.009)	1.000	0.000	1.000 (0.070)	1.000	0.000
High severity	0.724 (0.692)	0.683	-0.042	0.998 (0.013)	0.994	-0.004	0.975 (0.039)	0.980	0.006
Sleep or weight increase	0.336 (0.227)	0.413	0.077	0.965 (0.025)	0.962	-0.003	0.971 (0.016)	0.974	0.003
Suicidality	-0.085 (0.238)	-0.102	-0.017	0.982 (0.010)	0.982	0.000	0.909 (0.042)	0.909	-0.000

Table 1. Input and output values of GDVIS for visualization of the seven major depressive disorder subtypes. Panel A. Input for the visualization for GDVIS. The heritabilities are on the observed scale with 50/50 ascertainment. Heritabilities and genetic correlations are estimated with LDSC. The population prevalence of major depressive disorder (MDD) is the same for each subtype. 'Prop with' refers to the proportion of MDD cases that have the 'with' subtype (e.g., MDD with childhood trauma). **Panel B.** Color reference for the comparisons presented in Panels A and C. **Panel C.** Comparison of GDVIS-derived values vs. the LDSC-estimated values for the heritabilities of   and   , and the genetic correlations of these comparisons with the input GWASs.

Figure 3. GDVIS visualizations of seven major depression disorder subtypes. We present the GDVIS visualization of the following seven subtypes of Major Depressive Disorder (MDD): with (subtype1-cases)/ without (subtype2-cases) childhood trauma, comorbid anxiety, early onset (< 30 yo), recurrent episodes, high severity, sleep or weight increase, suicidality. Numerical values are reported of input and GDVIS-derived values are listed in Table 1.

Figure 4. Comparison of GDVIS-derived values and LDSC-estimated values of the genetic correlations between external traits and subtype1-subtype2 comparisons. We report LDSC-estimated (x-axis) and GDVIS-derived (y-axis) genetic correlation values for the external trait with the subtype1-cases and subtype2-cases comparisons ($r_{g|ext,sub1.cases\_sub2.cases}$). Different colors represent the six different subtypes that were related to 15 external traits (anxiety disorder[13], bipolar disorder[15], major depressive disorder[2], schizophrenia[16], childhood trauma[17], body mass index[18], educational attainment[19], Wellbeing[20], Neuroticism[21], substance use dependence factor[22], compulsive factor[22], internalizing factor[22], neurodevelopmental factor[22], thought disorders factor[22], and hierarchical factor[22]). Error bars represent ± the standard error of the LDSC estimate. The subtype of recurrence was not compatible for any of the 15 external traits and the external trait childhood trauma was not compatible for the subtype-definitions based on childhood trauma and high severity (due to incompatibility of LDSC input values, see main text). Numerical values for these results are reported in Table S4.

Figure 5. Genetic correlations of the subtypes with the external traits. We report the genetic correlation of the GWAS of 15 external trait with the following comparisons of six MDD subtypes (i) subtype1-cases versus controls, (ii) subtype2-cases versus controls, (iii) subtype1-cases versus subtype2-cases, and (iv) all-cases versus controls. The first two are based on LDSC and input to GDVIS, the latter two are GDVIS-derived. The subtype of recurrence was not compatible in geometrical space with external traits, the external trait childhood trauma was not compatible with the subtypes of childhood trauma and high severity (due to incompatibility of LDSC input values, see main text).

Figure 6. GDVIS visualizations of subtypes with external traits. Panel A. GDVIS visualization of the major depressive disorder (MDD) subtype of high severity with the internalizing factor[22] as external trait. **Panel B.** GDVIS visualization of the MDD subtype of sleep or weight increase with the external trait body mass index[18]. Heritabilities are on the 50:50 scale, the length of the line segments correspond to the square root of the heritability. For continuous traits, the length of the line segment has no direct interpretation as there are no clear endpoints. Visualizations exist in 3D geometrical space, as exemplified by the object in the cube. The top (i) and side (ii) views are taken to illustrate the relation of the subtype with the external trait. Interactive 3D visualizations can be accessed at: https://gdvis.shinyapps.io/gdvis/. All numerical values are reported in Table S3.

Figure 7. Comparison of GDVIS-derived values and LDSC-estimated values of the genetic correlation between the subtype1-subtype2 comparisons of two subtype-definitions. We report LDSC-estimated (x-axis) and GDVIS-derived (y-axis) genetic correlation values for the comparisons of $GWAS_{sub1.cases\_sub2.cases}$ for subtype-definition A and subtype-definition B (e.g., $r_{g|MDD.with.CT\_MDD.without.CT,MDD.with.early.onset\_MDD.without.early.onset}$). Error bars represent $\pm$ the standard error of the LDSC estimate. Different colors represent the different combinations of subtype-definitions. We exclude nine pairs of subtypes due to incompatibility for geometrical representation (see Methods). Numerical values are reported in Table S4.

Figure 8. GDVIS visualizations of the comparison of two subtype-definitions. We present the GDVIS visualization of the subtype-definition major depressive disorder (MDD) with/without high severity, and the subtype-definition MDD with/without high severity. The visualizations are in 3D geometrical space, as exemplified by the object in the cube. The top (i) and side (ii) views are taken to illustrate the relation between the subtype-definitions. Heritabilities are on the 50:50 scale, the length of the line segments correspond to the square root of the heritability. There is only one line segment representing the GWAS from MDD controls to all MDD cases, as these groups are the same in each subtype-definition. Interactive 3D visualizations can be accessed at: https://gdvis.shinyapps.io/gdvis/. All numerical values are reported in Table S4.

ACKNOWLEDGEMENTS

A.B.T. is supported by a PhD scholarship from Amsterdam University Medical Center. A.B.T., W.J.P., M.B. and B.W.J.H.P. are supported by Stress in Action. The research project 'Stress in Action': www.stress-in-action.nl is financially supported by the Dutch Research Council and the Dutch Ministry of Education, Culture and Science (NWO gravitation grant number 024.005.010). The work of W.J.P. was partly financially supported by the Amsterdam Cohort Hub, which is part of the Sector Plan 'Accelerating Health' of the Dutch Ministry of Education, Culture and Science. UK Biobank: This study has been conducted using UK Biobank resource under application number 40310. Y.M. is supported by Amsterdam UMC (Starter Grant Ronde 2), Amsterdam Neuroscience (PoC funding 2024-2026), and the ImmunoMIND consortium, funded by UK Research and Innovation as part of the UK national Mental Health Platform. KJHV is supported by the Foundation Volksbond Rotterdam. M.B. is supported by an NWO VICI grant (VI.C.211.054).

CONFLICT OF INTEREST STATEMENT

The authors report no conflicts of interest.

URLs

UK Biobank: https://www.ukbiobank.ac.uk/
GDVIS shiny app: https://gdvis.shinyapps.io/gdvis/
Package availability: https://github.com/ABThijssen/GDVIS/
Code availability: https://github.com/ABThijssen/Depression_heterogeneity/

REFERENCES

1.   Howard DM, Adams MJ, Clarke TK, et al. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci* 2019; 22: 343–352. 20190204. DOI: 10.1038/s41593-018-0326-7.

2.   Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium. Electronic address ameau and Major Depressive Disorder Working Group of the Psychiatric Genomics C. Trans-ancestry genome-wide study of depression identifies 697 associations implicating cell types and pharmacotherapies. *Cell* 2025; 188: 640–652 e649. 20250114. DOI: 10.1016/j.cell.2024.12.002.

3.   Spiller TR, Duek O, Helmer M, et al. Unveiling the Structure in Mental Disorder Presentations. *JAMA Psychiatry* 2024; 81: 1101–1107. DOI: 10.1001/jamapsychiatry.2024.2047.

4.   Traylor M, Markus H and Lewis CM. Homogeneous case subgroups increase power in genetic association studies. *Eur J Hum Genet* 2015; 23: 863–869. 20141001. DOI: 10.1038/ejhg.2014.194.

5.   Hodgson K, McGuffin P and Lewis CM. Advancing psychiatric genetics through dissecting heterogeneity. *Hum Mol Genet* 2017; 26: R160–R165. DOI: 10.1093/hmg/ddx241.

6.   Lojko D and Rybakowski JK. Atypical depression: current perspectives. *Neuropsychiatr Dis Treat* 2017; 13: 2447–2456. 20170920. DOI: 10.2147/NDT.S147317.

7.   Nguyen TD, Harder A, Xiong Y, et al. Genetic heterogeneity and subtypes of major depression. *Mol Psychiatry* 2022; 27: 1667–1675. 20220108. DOI: 10.1038/s41380-021-01413-6.

8.   Milaneschi Y, Lamers F, Peyrot WJ, et al. Genetic Association of Major Depression With Atypical Features and Obesity-Related Immunometabolic Dysregulations. *JAMA Psychiatry* 2017; 74: 1214–1225. DOI: 10.1001/jamapsychiatry.2017.3016.

9.   Lee SH, Wray NR, Goddard ME, et al. Estimating missing heritability for disease from genome-wide association studies. *Am J Hum Genet* 2011; 88: 294–305. 20110303. DOI: 10.1016/j.ajhg.2011.02.002.

10.   Peyrot WJ and Price AL. Identifying loci with different allele frequencies among cases of eight psychiatric disorders using CC-GWAS. *Nat Genet* 2021; 53: 445–454. 20210308. DOI: 10.1038/s41588-021-00787-1.

11.   Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across human diseases and traits. *Nat Genet* 2015; 47: 1236–1241. 20150928. DOI: 10.1038/ng.3406.

12.   Bulik-Sullivan BK, Loh PR, Finucane HK, et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015; 47: 291–295. 20150202. DOI: 10.1038/ng.3211.

13.   Purves KL, Coleman JRI, Meier SM, et al. A major role for common genetic variation in anxiety disorders. *Mol Psychiatry* 2020; 25: 3292–3303. 20191120. DOI: 10.1038/s41380-019-0559-1.

14.   Davis KAS, Coleman JRI, Adams M, et al. Mental health in UK Biobank - development, implementation and results from an online questionnaire completed by 157 366 participants: a reanalysis. *BJPsych Open* 2020; 6: e18. 20200206. DOI: 10.1192/bjo.2019.100.

15.   O'Connell KS, Koromina M, van der Veen T, et al. Genomics yields biological and phenotypic insights into bipolar disorder. *Nature* 2025; 639: 968–975. 20250122. DOI: 10.1038/s41586-024-08468-9.

16.   Trubetskoy V, Pardinas AF, Qi T, et al. Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* 2022; 604: 502–508. 20220408. DOI: 10.1038/s41586-022-04434-5.

17.   Dalvie S, Maihofer AX, Coleman JRI, et al. Genomic influences on self-reported childhood maltreatment. *Transl Psychiatry* 2020; 10: 38. 20200127. DOI: 10.1038/s41398-020-0706-0.

18.   Yengo L, Sidorenko J, Kemper KE, et al. Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet* 2018; 27: 3641–3649. DOI: 10.1093/hmg/ddy271.

19.   Okbay A, Wu Y, Wang N, et al. Polygenic prediction of educational attainment within and between families from genome-wide association analyses in 3 million individuals. *Nat Genet* 2022; 54: 437–449. 20220331. DOI: 10.1038/s41588-022-01016-z.

20.   Baselmans BML, Jansen R, Ip HF, et al. Multivariate genome-wide analyses of the well-being spectrum. *Nat Genet* 2019; 51: 445–451. 20190114. DOI: 10.1038/s41588-018-0320-8.

21.   Nagel M, Jansen PR, Stringer S, et al. Meta-analysis of genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic loci and pathways. *Nat Genet* 2018; 50: 920–927. 20180625. DOI: 10.1038/s41588-018-0151-7.

22.   Grotzinger AD, Singh K, Miller-Fleming TW, et al. Transcriptome-Wide Structural Equation Modeling of 13 Major Psychiatric Disorders for Cross-Disorder Risk and Drug Repurposing. *JAMA Psychiatry* 2023; 80: 811–821. DOI: 10.1001/jamapsychiatry.2023.1808.

23.   Monroe SM and Harkness KL. Major Depression and Its Recurrences: Life Course Matters. *Annu Rev Clin Psychol* 2022; 18: 329–357. 20220225. DOI: 10.1146/annurev-clinpsy-072220-021440.

24.   Childhood Trauma Meta-Analysis Study G. Treatment efficacy and effectiveness in adults with major depressive disorder and childhood trauma history: a systematic review and meta-analysis. *Lancet Psychiatry* 2022; 9: 860–873. 20220922. DOI: 10.1016/S2215-0366(22)00227-9.

25.   Kuzminskaite E, Vinkers CH, Milaneschi Y, et al. Childhood trauma and its impact on depressive and anxiety symptomatology in adulthood: A 6-year longitudinal study. *J Affect Disord* 2022; 312: 322–330. 20220624. DOI: 10.1016/j.jad.2022.06.057.

26.   Watters ER, Aloe AM and Wojciak AS. Examining the Associations Between Childhood Trauma, Resilience, and Depression: A Multivariate Meta-Analysis. *Trauma Violence Abuse* 2023; 24: 231–244. 20210727. DOI: 10.1177/15248380211029397.

27.   Adams MJ, Thorp JG, Jermy BS, et al. Genome-wide meta-analysis of ascertainment and symptom structures of major depression in case-enriched and community cohorts. *Psychol Med* 2024; 54: 3459–3468. 20240926. DOI: 10.1017/S0033291724001880.

28.   Grotzinger AD, Rhemtulla M, de Vlaming R, et al. Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nat Hum Behav* 2019; 3: 513–525. 20190408. DOI: 10.1038/s41562-019-0566-x.

29.   Peyrot WJ, Van der Auwera S, Milaneschi Y, et al. Does Childhood Trauma Moderate Polygenic Risk for Depression? A Meta-analysis of 5765 Subjects From the Psychiatric Genomics Consortium. *Biol Psychiatry* 2018; 84: 138–147. 20170921. DOI: 10.1016/j.biopsych.2017.09.009.

30.   Milaneschi Y, Lamers F, Berk M, et al. Depression Heterogeneity and Its Biological Underpinnings: Toward Immunometabolic Depression. *Biol Psychiatry* 2020; 88: 369–380. 20200128. DOI: 10.1016/j.biopsych.2020.01.014.

31.   Badini I, Coleman JRI, Hagenaars SP, et al. Depression with atypical neurovegetative symptoms shares genetic predisposition with immuno-metabolic traits and alcohol consumption. *Psychol Med* 2022; 52: 726–736. 20200706. DOI: 10.1017/S0033291720002342.

32.   Shorter JR, Pasman JA, Kurvits S, et al. Genome-wide association analyses identify distinct genetic architectures for early-onset and late-onset depression. *Nat Genet* 2025 20251113. DOI: 10.1038/s41588-025-02396-8.

33.   Grotzinger AD, Fuente J, Prive F, et al. Pervasive Downward Bias in Estimates of Liability-Scale Heritability in Genome-wide Association Study Meta-analysis: A Simple Solution. *Biol Psychiatry* 2023; 93: 29–36. 20220608. DOI: 10.1016/j.biopsych.2022.05.029.

34.	Uffelmann E, Major Depressive Disorder Working Group of the Psychiatric Genomics C, Schizophrenia Working Group of the Psychiatric Genomics C, et al. Estimating Disorder Probability Based on Polygenic Prediction Using the BPC Approach. *medRxiv* 2025 20250321. DOI: 10.1101/2024.01.12.24301157.

35.	Zhu Z, Zheng Z, Zhang F, et al. Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun* 2018; 9: 224. 20180115. DOI: 10.1038/s41467-017-02317-2.

36.	Jiang L, Zheng Z, Fang H, et al. A generalized linear mixed model association tool for biobank-scale data. *Nat Genet* 2021; 53: 1616–1621. 20211104. DOI: 10.1038/s41588-021-00954-4.

37.	Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature* 2018; 562: 203–209. 20181010. DOI: 10.1038/s41586-018-0579-z.

38.	Smith DJ, Nicholl BI, Cullen B, et al. Prevalence and characteristics of probable major depression and bipolar disorder within UK biobank: cross-sectional study of 172,751 participants. *PLoS One* 2013; 8: e75362. 20131125. DOI: 10.1371/journal.pone.0075362.

39.	Abdellaoui A, Dolan CV, Verweij KJH, et al. Gene-environment correlations across geographic regions affect genome-wide association studies. *Nat Genet* 2022; 54: 1345–1354. 20220822. DOI: 10.1038/s41588-022-01158-0.