From default mode to action mode: biphasic network reconfiguration with meditation in schizophrenia

Qing Wang[a, b]*, Lizhao Du[a, c], Jialing Sheng[d], Qiao Wang[e], Yuan Shi[a], Ting Xue[a], Zhong Sun[d], Yingying Tang[a, b], Donghong Cui[a,f]*

[a] Shanghai Key Laboratory of Psychotic Disorders, Brain Health Institute, National Center for Mental Disorders, Shanghai Mental Health Center, School of Medicine, Shanghai Jiao Tong University, Shanghai, 200030, China.

[b] NeuroImaging Core, Shanghai Mental Health Center, School of Medicine, Shanghai Jiao Tong University, 600 S. Wanping Rd., Shanghai, 200030, China.

[c] Shanghai Med-X Engineering Research Center, School of Biomedical Engineering, Shanghai Jiao Tong University, Shanghai, China

[d] The First Minzheng Mental Health Center, No. 9999 Zhongchun Rd., Minhang, Shanghai, 201105, China.

[e] Department of Public Health, Brody School of Medicine, East Carolina University, 610 Moye Blvd, Mailstop 660, Greenville, NC 27834, U.S.A.

[f] Brain Science and Technology Research Center, Shanghai Jiao Tong University, Shanghai, China.

Short Title: Meditation brain networks in Schizophrenia

Corresponding Author: Qing Wang, and Donghong Cui

First and last name: Qing and Wang, Donghong and Cui

E-mail address: vincent.w.qing@gmail.com, manyucc@126.com

Author notes:

Qing Wang: https://orcid.org/0000-0002-6843-5516

Lizhao Du: https://orcid.org/0000-0003-4664-0938

Qiao Wang: https://orcid.org/0000-0002-3570-9496

Ting Xue: https://orcid.org/0000-0002-3650-6308

Yingying Tang: https://orcid.org/0000-0002-4705-3682

Keywords: Meditation, schizophrenia, RCT, trajectory, fMRI

1 Abstract

2 Introduction

3 Meditation is widely used to support mental well-being, and recent randomized trials suggest
4 benefits for persistent psychotic symptoms in schizophrenia. However, the magnitude and timing of
5 causal treatment effects, response heterogeneity, and underlying neurobiological mechanisms over
6 clinically meaningful timescales remain unclear.

7 Methods

We analyzed data from an eight-month, parallel-group randomized clinical trial (ChiCTR1800014913)
of 64 male inpatients with chronic schizophrenia, randomized to daily clinician-guided meditation
plus rehabilitation or rehabilitation alone. Prespecified outcomes were PANSS percentage decrease
rate and RBANS increase rate. Linear mixed-effects models estimated time-specific causal average
treatment effects. Cross-lagged panel models examined temporal relations between symptom and
cognitive benefits; latent-class mixed models characterized treatment-response heterogeneity.
Resting-state fMRI at baseline, 3, and 8 months yielded functional components, their complexity
indices, and functional-connectivity predictors of clinical benefit.

16 Results

Meditation produced progressive symptom improvement (average treatment effects on PANSS
decrease rate: 11.8% after 3 months; 20.8% after 8 months) and an early cognitive gain (7.6% after 3
months) that plateaued. Early cognitive improvement predicted, but did not mediate, later symptom
relief. Response trajectories were heterogeneous; marital status and lower antipsychotic burden
characterized high responders. Neuroimaging revealed a biphasic pattern: higher baseline default-
mode–cerebellar complexity predicted short-term benefit, whereas greater 3-month action-mode-
sensorimotor–executive complexity predicted longer-term gains; functional-connectivity models
converged on these findings.

25 Conclusions

26 Clinician-guided meditation, added to rehabilitation, yields robust causal treatment effects on
27 symptoms in schizophrenia. A biphasic shift from default-mode–cerebellar involvement to action-
28 mode engagement provides phase-specific, information-based indicators to guide personalized
29 meditation in severe mental illness.

Introduction

Meditation, encompassing practices such as mindfulness, focused attention, open monitoring, and compassion, yields small to moderate improvements in stress, mood and wellbeing with good acceptability across diverse settings[1,2]. However, its role in severe psychiatric disorders, particularly treatment-resistant schizophrenia, is much less clear. The field is moving beyond the question of "does it work?" toward "for whom, when, and how does it work?", with increasing emphasis on dose, specific components, and delivery format[3].

Across serious medical and psychiatric conditions, meditation and mindfulness-based interventions have shown condition-dependent benefits. Randomized trials report advantages for mood and quality of life in Parkinson's disease[4], reductions in pain-related disability in chronic pain[5] and migraine[6], prevention of depressive relapse[7], and symptom reduction in post-traumatic stress disorder[8]. In schizophrenia, meta-analyses suggest small-to-moderate improvements in total and negative symptoms, functioning, and quality of life, especially with structured, clinician-guided programmes[9,10]. Our group recently reported an eight-month randomized trial of intensive clinician-guided meditation for male inpatients with chronic schizophrenia, showing meaningful improvements in refractory hallucinations, delusions, and quality of life compared with rehabilitation alone[11]. Yet that initial report focused only on primary clinical outcomes, without examining causal treatment effects, symptom–cognition interplay, treatment heterogeneity, and the potential underlying brain mechanisms.

Neuroimaging work in meditation indicates that practice reshapes large-scale brain networks[3,12]. Training tends to reduce activity and dominance of self-referential systems such as the default-mode network (DMN)[13,14], while strengthening executive and frontoparietal control networks at rest[15,16]. Beyond static functional connectivity, meditation appears to increase the temporal complexity of BOLD signals, reflected in scale-free dynamics and entropy-based measures, which index neural flexibility and can distinguish clinical populations, including schizophrenia[17,18]. Recent experimental work in pain demonstrates that meditation engages neural signatures distinct from placebo, reinforcing the idea of mechanism-specific pathways rather than non-specific expectancy effects[19].

Despite this progress, schizophrenia remains under-studied in mechanistic meditation research. Most trials are short term, focus on single endpoints, and rarely model how benefits unfold over time or differ between individuals. Longitudinal neuroimaging within randomized designs is particularly scarce, and prior work seldom addresses: (i) how symptom and cognitive gains interplay over clinically meaningful timescales; (ii) treatment heterogeneity which can be captured by repose

trajectories; and (iii) how large-scale brain networks reorganize from early to later phases of

meditation-based treatment.

In this study, we embedded longitudinal resting-state fMRI within the previously reported eight-

month, parallel-group randomized clinical trial of clinician-guided meditation versus rehabilitation

alone in male inpatients with chronic schizophrenia (ChiCTR1800014913)[11]. Building on the original

clinical report, we re-estimated treatment effects using causal models, quantified the time-specific

improvements in symptoms and cognition, examined whether early cognitive change predicts later

symptom relief, and identified heterogeneous treatment-response trajectories and their pragmatic

predictors.

To generate mechanistic insight, we derived large-scale functional components from resting-state

fMRI and characterized their temporal complexity and functional connectivity across baseline, 3-

month, and 8-month sessions. Guided by contemporary accounts of the brain's default-mode

network[13] and the action-mode network[20], we hypothesized a stage-specific reconfiguration in

which default-mode–cerebellar systems are prominent in early change, followed by later

engagement of sensorimotor–executive, action-oriented networks as clinical gains consolidate. By

linking these network-level markers to individual clinical trajectories, we aim to offer an integrated,

clinically interpretable account of how meditation contributes to recovery in schizophrenia and how

it might be personalized within psychotherapeutic and rehabilitative care.

Methods

Study Design and Participants

We conducted an eight-month, single-center, parallel-group randomized clinical trial in male Han

Chinese inpatients with chronic schizophrenia and persistent positive symptoms at the First

Minzheng Mental Health Center, China (trial registration: ChiCTR1800014913). This study adhered to

the Declaration of Helsinki and was approved by the institutional review board (YJZXLL2017005). Full

details of this trial can be found in our previous report of the primary outcome [11] and the

supplementary CONSORT checklist.

Eligible participants met DSM-IV criteria for schizophrenia, had illness duration ≥ 5 years, and

showed persistent positive symptoms (hallucinations: PANSS P3 ≥ 5 and/or ≥ 2 items among P1, P5,

P6, G1, G3, G9 ≥ 5) despite ≥ 2 adequate antipsychotic trials. Key exclusion criteria included: any

other major Axis I disorder; Mini-Mental State Examination (MMSE) < 24; language barriers

precluding consent or assessment; known neurological disease; major medical illness; and current

substance abuse or dependence. All participants provided written informed consent. Antipsychotic

regimens were maintained under physician supervision and converted to chlorpromazine equivalents

(CPZ). Of 64 randomized participants, all completed 4 months and 63 completed 8 months (one

death in the control arm, adjudicated unrelated to study procedures).

Randomization and Masking

Participants were randomized 1:1 to daily clinician-guided meditation plus rehabilitation or

rehabilitation alone, using individual-level allocation. The randomization sequence was generated by

an independent statistician and implemented by study staff not involved in outcome assessment. The

outcome assessors were masked to group assignment; participants and instructors were unmasked

due to the nature of the interventions.

Procedures

Rehabilitation Group (control arm)

The rehabilitation-only arm received a 90-min general rehabilitation programme (GRP) Monday–

Saturday, comprising manual work, gardening, painting, and reading, plus a 90-min Sunday group

intervention session (GIS) focused on daily-living skills and psychoeducation about schizophrenia,

including self-acceptance.

Meditation Plus Rehabilitation Group (intervention arm)

The meditation arm received 30 min of clinician-guided meditation followed by 60 min of GRP,

Monday–Saturday. The meditation protocol included: (1) Seven-point posture of Buddha Vairocana;

(2) Abdominal breathing; (3) Sustained awareness of and concentration on the breath; (4) Non-

reactive attention to thoughts and mental events. The meditation-arm Sunday GIS comprised

instruction in meditation methods and cultural context, psychoeducation about schizophrenia and

self-acceptance, a 30-min group practice, and discussion of experiences and challenges. GISs were

weekly during the first 8 weeks, then monthly thereafter.

Clinical assessments and MRI were obtained at baseline, 3 months, and 8 months. Adverse events

were monitored throughout; no adverse events were attributed to study procedures.

Outcomes and Assessments

Baseline demographics (age, education, family income, age at onset, illness duration, medications)

were collected via structured questionnaire. The primary clinical endpoint was the Positive and

Negative Syndrome Scale (PANSS) total decrease rate (PDR) with established rescaling[21]:

$$PDR_t = \frac{PANSS_0 - PANSS_t}{PANSS_0 - 30} \times 100\%.$$

Secondary endpoints were rescaled PANSS subscale decrease rates: PPDR, PNDR, and PGDR

(rescaling factors 7, 7, and 16, respectively), and cognition measured as the Repeatable Battery for

the Assessment of Neuropsychological Status (RBANS) increase rate (RIR; computed analogously

without rescaling). Mindfulness (Five Facet Mindfulness Questionnaire, FFMQ) and CPZ were

recorded as contextual measures. Imaging endpoints (temporal complexity and functional

connectivity) were exploratory, designed to generate mechanistic hypotheses and predictive markers

aligned with clinical timescales.

Imaging Acquisition and Preprocessing

Full brain MRI data were acquired on a Siemens 3 T Verio with a 32 channel head coil. T1 weighted

MPRAGE: 1 × 1 × 1 mm³ isotropic voxel, TR = 2530 ms, TE = 3.65 ms. Resting state fMRI

(rs fMRI): gradient echo EPI, 3 × 3 × 3 mm³ isotropic voxel, TR = 1500 ms, TE = 30 ms, flip

angle 77°, duration 6 min (240 volumes); first 10 volumes discarded for nonstationary. Data were

organized in BIDS[22], and preprocessed using fMRIPrep v22.1.1[23]. Post fMRIPrep rs fMRI were

detrended, smoothed with 6 mm FWHM Gaussian kernel, and nuisance regressed (six motion

parameters and the global signal)[24]. Imaging quality control yielded 186 usable rs fMRI datasets

across sessions.

Functional Components and Temporal Complexities

Large scale activity patterns were estimated with group canonical ICA[25] (canICA; k = 20

components) on all quality controlled rs fMRI datasets. Component maps were binarized; mean

time series were extracted per participant and session. For each component we computed sample

entropy (SampEn) and the Hurst exponent (HE) as complementary temporal complexity measures

[17]. Components whose complexity predicted PDR were screened for further analysis using

elastic net with 5 fold cross validation and 10,000 permutations. To summarise component

complexity with consistent directionality, we defined a composite complexity index (CCI) using

weighted *z*-scores:

$$CCI_{itc} = \frac{Z(SampEn_{itc}) + (1 - 2 \cdot \mathbf{1}\{HE_{itc} > 0.5\} \cdot Z(HE_{itc}))}{2},$$

where i indexes participant, t session, and c component; $\mathbf{1}\{HE_{itc} > 0.5\}$ is an indicator equal to 1 if

$HE_{itc} > 0.5$ and 0 otherwise. Higher *CCI* denotes greater overall temporal complexity. To further test

the phase-specific roles of these screened components, we utilized a 2-wave structural equation

model (SEM) with PDR_3 and PDR_8 as outcomes.

Functional connectivity (FC) was computed as the Pearson correlation between AAL2 atlas

regions[26,27]. We evaluated three prediction tasks per arm: (1) 3-month PDR from baseline FC; (2)

8-month PDR from baseline FC; (3) 8-month PDR from FC changes in the first 3 months (ΔFC_{3-0}).

Models used z-scored FC values, elastic net with 5-fold cross-validated hyperparameters, and 10,000

permutations to derive signed importance scores (magnitude=R² contributions; sign=direction of

association with PDR)[28]. Edges with importance > 0.05 are listed as candidate predictors; full
edgewise results are provided in the supplementary materials.

Statistical analysis

All analyses were performed in the intention-to-treat population with available data at each
timepoint. Average treatment effects (ATEs) for PDR and RIR were estimated using linear mixed-
effects models (LMM) with fixed effects for group (meditation vs rehabilitation), time (baseline, 3
months, 8 months), and group × time interaction, and subject-level random effects to account for
within-participant correlation. ATEs at 3 and 8 months are reported as model-based marginal
differences with 95% confidence intervals. The moderator effects from baseline chlorpromazine
equivalents (CPZ) dose, PANSS, RBANS, and FFMQ were also tested by adding the interaction term
group × moderator into the original test.

Temporal relations between symptom and cognitive changes were examined with a cross-lagged
panel model (CLPM) including autoregressive paths, concurrent treatment effects, and cross-lagged
paths between 3-month cognition (RIR_3) and 8-month symptoms (PDR_8) and vice versa. Bootstrap-
based mediation analysis (10,000 resamples) tested the indirect effect of treatment on PDR_8
through RIR_3.

Treatment-response heterogeneity was assessed using latent-class linear mixed models (LCLMM)[29]
applied to PDR trajectories (2–5 classes; Bayesian information criterion for model selection). Baseline
predictors of high-responder membership were identified using elastic-net logistic regression with 5-
fold cross-validation and 10,000 permutations.

Phase-specific imaging contributions were tested using a two-wave structural equation model (SEM)
with PDR at 3 and 8 months as outcomes. The model included: (i) autoregression for CCI; (ii) paths
from baseline CCI to 3-month PDR, *i.e.*, $PDR_{i3} \sim group_i + CCI_{i0(DMN-CIC)} + CCI_{i0(AMN-SEC)} +$
$age_i + PANSS_{i0} + RBANS_{i0} + \varepsilon_{i3}$; and (iii) paths from 3-month CCI to 8-month PDR, *i.e.*,
$PDR_{i8} \sim group_i + CCI_{i3(DMN-CIC)} + CCI_{i3(AMN-SEC)} + age_i + PANSS_{i0} + PDR_{i3} + \varepsilon_{i8}$. Full
specifications, fit indices, and Wald contrasts are provided in the Supplementary Materials.

Missing Data

188   All models used all available observations under a missing-at-random assumption. LMMs naturally
189   accommodate intermittently missing outcome data. SEMs used full-information maximum likelihood
190   with robust (MLR) standard errors and Yuan–Bentler scaled χ^2. Imaging sessions failing quality control
191   were excluded from imaging analyses.

Estimation and Software

193   All analyses were conducted in a Python 3.9.12 and R 4.2.2 environment. Key packages included:
194   statsmodels 0.14.4 for LMM, semopy 2.3.11 for CLPM, lcmm 2.1.0 for LCLMM, HeuDiConv 0.9.0 for
195   BIDS conversion, fMRIPrep 22.1.1 for rs-fMRI preprocessing, nilearn 0.10.4 for imaging analysis, and
196   lavaan 0.6-18 for 2-wave SEM.

Patient and Public Involvement

198 Patients or members of the public were not involved in the design, conduct, reporting, or
199 dissemination plans of this study.

Results

Meditation produces progressive symptom improvement and plateauing cognitive gains

As shown in Fig. 1, linear mixed-effects models indicated a progressive clinical benefit of meditation
on symptoms and a front-loaded cognitive gain. The average treatment effect (ATE) on PANSS
decrease rate (PDR) was 11.8% (95% CI 9.3–14.3; p<0.001) after 3 months of meditation and
increased to 20.8% (95% CI 16.0–25.6; p<0.001) after 8 months (Fig. 1b). Cognition (RBANS increase
rate, RIR) improved 7.6% after 3 months (95% CI 2.4–12.8; p=0.004) but was not significant after 8
months (5.5%, 95% CI −2.1 to 13.1; p=0.153) (Fig. 1c). Baseline CPZ dose, PANSS, RBANS, and
mindfulness did not show robust moderation effects (Tables S3–S12).

Early cognitive gains forecast, but do not mediate later symptom relief

Cross-lagged panel modeling supported temporal precedence of early cognition over later symptoms.
Meditation exerted a direct effect on PDR at 3 months (β=0.118; p<0.001) and a smaller direct effect
at 8 months (β=0.070; p=0.036), with additional indirect influence via earlier symptom change.
Meditation produced 3-month RIR (β=0.076; p=0.011), but no direct treatment effect was observed
for 8-month RIR. Critically, 3-month RIR predicted 8-month PDR (β=0.197; p=0.017), whereas 3-
month PDR did not predict 8-month RIR (β=0.020; p=0.946). Bootstrap mediation provided no
evidence that early cognition mediated the 8-month symptom benefit (indirect effect=0.020; 95% CI
−0.0004 to 0.0479) (Table S14; Fig. S2). Together, these results suggest partly dissociable therapeutic
pathways in which early cognitive improvement forecasts later symptom relief without statistically
explaining it.

Meditation elicits heterogeneous clinical trajectories

Latent-class mixed modeling identified two response trajectories within the meditation arm: a high-
responder subgroup (n=10) and a normal-responder subgroup (n=22) (Fig. 2a). Trajectories were
similar through the first 3 months, after which high responders continued to accrue symptom gains
while normal responders decelerated (both outperforming rehabilitation alone; n=31). Baseline
profiles suggested higher responders were more often married, had lower antipsychotic burden, with
elder age and age of onset etc. (Fig. 2b). In permutation-tested elastic-net classification, marital
status (β=0.681; p=0.003) and lower CPZ dose (β=−0.566; p=0.017) remained robust correlates of
high response (Fig. 2c), nominating pragmatic baseline markers for prospective testing.

Biphasic network complexities align with symptom improvement

Group canICA identified two components (Fig. 3a–b): an action-mode-sensorimotor–executive

component (AMN–SEC) and a default-mode–cerebellar-integrative component (DMN–CIC).

Permutation-validated screening linked symptom benefit to component temporal complexity, with

convergent validation by SHAP analyses (Fig. S3). AMN–SEC comprised bilateral precentral gyri,

superior, middle, and inferior frontal regions, Rolandic opercula, insulae, mid-cingulate, sensorimotor

cortices, supramarginal gyri, precuneus, superior temporal gyri, and inferior motor cerebellar lobules

(VIIb–VIII). It is involved in cognitive control, sensorimotor integration, and goal-directed

behaviors[20]. DMN–CIC included posterior cingulate, calcarine and lingual regions, superior and

middle occipital areas, superior parietal cortex, precunei, middle temporal gyrus, extensive

posterolateral cerebellar hemispheres (Crus I/II, lobules VI–VIII), and vermal lobules IV–IX. DMN–CIC

is involved in high-level integrative processes: self-referential ideation, sensory-cognitive binding,

and internal modeling[13].

Cross-sectional associations between component complexity and concurrent clinical benefit were

small after 3 months but strengthened after 8 months, particularly for AMN–SEC. To test time-

specific contributions, a two-wave SEM focused on PDR after 3 and 8 months showed good fit

$(\chi^2(37) = 39.321, p = 0.366)$ and high explained variance (71.5% and 71.6% for 3-month and 8-

month respectively). Baseline DMN–CIC complexity predicted greater 3-month improvement

(β=0.017; 95% CI 0.005–0.029; p=0.004), whereas 3-month AMN–SEC complexity predicted 8-month

improvement after accounting for carry-over effect of earlier symptom improvement (β=0.032; 95%

CI 0.007–0.057; p=0.013). Within-phase contrasts supported an early DMN–CIC contribution and a

later AMN–SEC contribution (Wald tests: p=0.009 and p=0.047, respectively). The above patterns

were not evident in the rehabilitation alone arm, supporting a biphasic model in which early benefit

aligns with default-mode–cerebellar characteristics, while later gains align with action-mode-

sensorimotor–executive recalibration.

Functional connectivity predicts short- and long-term response under meditation

Elastic-net models with permutation importance identified functional connectivity predictors of

symptom improvement across three tasks (Fig. 4). In the meditation arm, baseline connectivity

contributed to short-term prediction, with stronger baseline DMN–CIC couplings associated with less

improvement (e.g., *Cerebellum_4–5_R–Vermis_1–2 etc.*). For long-term response, prediction was

strongest for action-mode-sensorimotor–executive couplings and early connectivity changes, with

Precentral_L–SMA_L emerging as the dominant marker (baseline importance 0.154; early-change

importance 0.371). In the rehabilitation-only arm, predictive utility was minimal (largest importance

0.023). Together with the complexity and SEM results, these FC findings converge on a biphasic

account linking early default-mode–cerebellar features to short-term benefit and later precentral-

SMA-centered action-mode engagement to sustained improvement.

Discussion

In this randomized clinical trial, adding daily clinician-guided meditation to standard rehabilitation

produced progressive symptom improvement across eight months, alongside an early, modest

cognitive gain that plateaued. By estimating time-specific average treatment effects, we extend the

prior clinical report of this trial by clarifying the magnitude and timing of benefit, and by linking

clinical change to individual differences in response and phase-specific neuroimaging signatures.

A key behavioral insight is that symptoms and cognition improved on partly separable tracks. Early

cognitive gains predicted later symptom improvement but did not mediate it, suggesting that

cognition may function as an early marker of capacity to benefit (e.g., learning, engagement,

adherence, or strategy acquisition) rather than as the primary causal pathway by which meditation

reduces symptoms. Clinically, this implies that measuring early cognitive change could help identify

patients likely to accrue longer-term symptomatic benefit, while also cautioning against assuming

that strengthening cognition alone will automatically drive subsequent symptom relief.

We also observed meaningful treatment-response heterogeneity within the meditation arm.

Although average effects were robust, a subgroup continued to improve after 3 months while others

showed attenuated gains. Baseline correlates, including marital status and lower antipsychotic

burden, are pragmatic and clinically interpretable, consistent with roles for social support and

medication-related functional constraints in sustaining practice and recovery. Because subgroup sizes

were modest, these predictors should be treated as candidate stratification features for prospective

testing rather than definitive selection rules.

The longitudinal neuroimaging results provide a coherent, time-resolved account of brain–behavior

relationships during treatment. We found evidence for a biphasic pattern in which baseline default-

mode–cerebellar temporal complexity related to earlier symptom improvement, whereas mid-

treatment action-mode-sensorimotor–executive complexity, and especially precentral–SMA

connectivity, tracked longer-term gains. Importantly, these imaging measures should be interpreted

as predictive and associative markers of clinical trajectory rather than as direct proof that meditation

induces a specific network change that causes improvement. Nevertheless, the convergence

between complexity-based findings (lagged SEM results) and functional connectivity prediction

models supports a staged account of recovery in which early benefit aligns with default-mode–

cerebellar dynamics while later benefit aligns with increased engagement of systems supporting
goal-directed action and executive control.

These findings have practical implications for phase-aware personalization. Baseline default-mode–
cerebellar features could help anticipate early response, while early shifts in action-mode-
sensorimotor–executive coupling may indicate whether patients are likely to maintain improvement
over longer periods. In principle, this could motivate adaptive strategies (e.g., reinforcing movement-
linked or goal-directed practice elements for those showing early action-mode engagement, or
revising treatment plan when early neural change is absent), ideally within preregistered adaptive
trials that explicitly test such decision rules.

The recurrent involvement of the posterior cerebellum across complexity and connectivity markers is
notable. Cerebello–thalamo–cortical loops are increasingly implicated in predictive/feedforward
control and in calibrating cortical dynamics during skill learning and switching[30]. In this context,
baseline default-mode–cerebellar features may index capacity for early clinical change, whereas
subsequent action-mode-sensorimotor–executive engagement may reflect consolidation of
externally oriented control during sustained practice. A modest association between AMN–SEC
features and symptom change in the rehabilitation arm is plausible given its emphasis on manual,
goal-directed activities that recruit sensorimotor–executive systems. What appears distinctive in the
meditation arm is the temporal ordering: early default-mode–cerebellar features forecasting short-
term improvement, followed by mid-treatment action-mode-sensorimotor–executive features
forecasting longer-term gains. This sequence was not evident with rehabilitation alone, supporting
the phase-specific model rather than a nonspecific activity effect.

Several limitations temper interpretation. The sample was single-center, Asian male-only, limiting
generalizability across sex, culture, and clinical settings. Some analyses, particularly the imaging
screens and prediction models, were exploratory and pipeline-sensitive[31], and require out-of-
sample validation with harmonized preprocessing and prespecified analytic choices. Finally, while
longitudinal models support temporal precedence, they do not establish mechanistic causality[32];
future work combining preregistered phase-specific hypotheses with causal perturbations (e.g.,
targeted behavioral manipulation or neuromodulation) would be needed to test whether modulating
these circuits changes clinical outcome.

Conclusion

Daily clinician-guided meditation, delivered in a ward-compatible format as an
adjunct to rehabilitation, was associated with robust, time-increasing symptom

improvement and early cognitive gains in chronic schizophrenia. Clinical trajectories

were heterogeneous, and longitudinal neuroimaging identified phase-specific

predictive markers consistent with a biphasic shift from default-mode–cerebellar

involvement to action-mode-sensorimotor–executive engagement. These results

motivate personalized, stage-sensitive meditation delivery and call for preregistered

multisite studies to validate the proposed markers and treatment-adaptation

strategies.

Acknowledgement

The authors would like to thank all the participants, doctors, nurses and students who were involved in the implementation of this clinical trial. The authors also would like to thank Dr. Pedro Antonio Valdés-Sosa and Dr. María Luisa Bringas-Vega for their feedbacks and insights in improving this manuscript. The authors thank Dr. Hu Chuan-Peng for his suggestions during the 2025 COSN hackathon in Shenzhen, China.

Statement of Ethics

The trial protocol was approved by the Ethics Committee of Shanghai First Minzheng Mental Health Center (reference No. YJZXLL2017005). The study was registered at the WHO Primary Registry – Chinese Clinical Trials Registry: ChiCTR1800014913. All participants provided written informed consent at enrolment.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Funding Sources

This research was supported by the National Key R&D Program of China (No. 2017YFC0909200); the Shanghai Mental Health Center Key Project (No. 23zd01), Shanghai Jiao Tong University Joint Art and Science Funding (No. 14JCRZ05); Shanghai Jiao Tong University Medical and Engineering Joint Grant (No. YG2016ZD06); Shanghai Key Laboratory of Psychotic Disorders Open Grant (No. 17-K04), and the Integrated Innovation Research Team of Shanghai Mental Health Center.

The funder had no role in the design, data collection, data analysis, and reporting of this study. All authors were independent from the funders.

Author Contributions

Dr. Qing Wang had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Study conceptualization and design: Qing Wang (lead), Lizhao Du (supporting), and Donghong Cui (supporting). Acquisition, analysis, or interpretation of data: Qing Wang (lead), Lizhao Du (supporting), Jialing Sheng (supporting), Qiao Wang (supporting), Yuan Shi (supporting), Ting Xue (supporting), Zhong Sun (supporting), Yingying Tang (supporting), and Donghong Cui (supporting). Drafting of the manuscript: Qing Wang (lead), Lizhao Du (supporting), Qiao Wang (supporting), and Donghong Cui (supporting). Critical revision of the manuscript for important intellectual content: Qing Wang (lead), Lizhao Du (supporting), Qiao Wang (supporting),

and Donghong Cui (supporting). Statistical analysis: Qing Wang (lead), and Qiao Wang (supporting).

Obtained funding: Donghong Cui (lead), Qing Wang (supporting), Jialing Sheng (supporting), and

Zhong Sun (supporting). Administrative, technical, or material support: Qing Wang (lead), Jialing

Sheng (supporting), Zhong Sun (supporting), Yingying Tang (supporting), and Donghong Cui (lead).

Study supervision: Qing Wang (lead), Yingying Tang (supporting), and Donghong Cui (lead).

Data Availability Statement

The raw data of this study are not publicly available since they contain information that could

compromise the privacy of study participants. The data for statistical analysis and figures in this

paper are shared together with the codes via GitHub: https://github.com/Vincent-

wq/repo_sz_meditation. Reasonable requests for patient level data should be made to the

corresponding author and will be considered by the trial management group.

References

Sparacio A, IJzerman H, Ropovik I, Giorgini F, Spiessens C, Uchino BN, et al. Self-administered mindfulness interventions reduce stress in a large, randomized controlled multi-site study. Nat Hum Behav. 2024 June;1–10.

van Agteren J, Iasiello M, Lo L, Bartholomaeus J, Kopsaftis Z, Carey M, et al. A systematic review and meta-analysis of psychological interventions to improve mental wellbeing. Nat Hum Behav. 2021 May;5(5):631–52.

Ehmann S, Sezer I, Treves IN, Gabrieli JDE, Sacchet MD. Mindfulness, cognition, and long-term meditators: Toward a science of advanced meditation. Imaging Neuroscience. 2025 July;3:IMAG.a.82.

Kwok JYY, Chan LML, Lai CA, Ho PWL, Choi ZY, Auyeung M, et al. Effects of Meditation and Yoga on Anxiety, Depression and Chronic Inflammation in Patients with Parkinson's Disease: A Randomized Clinical Trial. Psychother Psychosom. 2025 Feb;94(2):101–18.

Cherkin DC, Sherman KJ, Balderson BH, Cook AJ, Anderson ML, Hawkes RJ, et al. Effect of Mindfulness-Based Stress Reduction vs Cognitive Behavioral Therapy or Usual Care on Back Pain and Functional Limitations in Adults With Chronic Low Back Pain: A Randomized Clinical Trial. JAMA. 2016 Mar;315(12):1240–9.

Wells RE, O'Connell N, Pierce CR, Estave P, Penzien DB, Loder E, et al. Effectiveness of Mindfulness Meditation vs Headache Education for Adults With Migraine: A Randomized Clinical Trial. JAMA Intern Med. 2021 Mar;181(3):317–28.

Kuyken W, Warren FC, Taylor RS, Whalley B, Crane C, Bondolfi G, et al. Efficacy of Mindfulness-Based Cognitive Therapy in Prevention of Depressive Relapse: An Individual Patient Data Meta-analysis From Randomized Trials. JAMA Psychiatry. 2016 June;73(6):565–74.

Polusny MA, Erbes CR, Thuras P, Moran A, Lamberty GJ, Collins RC, et al. Mindfulness-Based Stress Reduction for Posttraumatic Stress Disorder Among Veterans: A Randomized Clinical Trial. JAMA. 2015 Aug;314(5):456–65.

Goyal M, Singh S, Sibinga EMS, Gould NF, Rowland-Seymour A, Sharma R, et al. Meditation Programs for Psychological Stress and Well-being: A Systematic Review and Meta-analysis. JAMA Intern Med. 2014 Mar;174(3):357.

Liu Y-C, Li I-L, Hsiao F-H. Effectiveness of mindfulness-based intervention on psychotic symptoms for patients with schizophrenia: A meta-analysis of randomized controlled trials. Journal of Advanced Nursing. 2021;77(6):2565–80.

Xue T, Sheng J, Gao H, Gu Y, Dai J, Yang X, et al. Eight-month intensive meditation-based intervention improves refractory hallucinations and delusions and quality of life in male inpatients with schizophrenia: a randomized controlled trial. Psychiatry Clin Neurosci. 2024 Apr;78(4):248–58.

Fox KCR, Dixon ML, Nijeboer S, Girn M, Floman JL, Lifshitz M, et al. Functional neuroanatomy of meditation: A review and meta-analysis of 78 functional neuroimaging investigations. Neuroscience & Biobehavioral Reviews. 2016 June;65:208–28.

Menon V. 20 years of the default mode network: A review and synthesis. Neuron. 2023 Aug;111(16):2469–87.

Garrison KA, Zeffiro TA, Scheinost D, Constable RT, Brewer JA. Meditation leads to reduced default mode network activity beyond an active task. Cogn Affect Behav Neurosci. 2015 Sept;15(3):712–20.

Taren AA, Gianaros PJ, Greco CM, Lindsay EK, Fairgrieve A, Brown KW, et al. Mindfulness Meditation Training and Executive Control Network Resting State Functional Connectivity: A Randomized Controlled Trial. Psychosom Med. 2017 July;79(6):674–83.

Bremer B. Mindfulness meditation increases default mode, salience, and central executive network connectivity. Scientific Reports. 2022

He BJ. Scale-Free Properties of the Functional Magnetic Resonance Imaging Signal during Rest and Task. J Neurosci. 2011 Sept;31(39):13786–95.

Sokunbi MO, Gradin VB, Waiter GD, Cameron GG, Ahearn TS, Murray AD, et al. Nonlinear Complexity Analysis of Brain fMRI Signals in Schizophrenia. PLOS ONE. 2014 May;9(5):e95146.

Riegner G, Dean J, Wager TD, Zeidan F. Mindfulness Meditation and Placebo Modulate Distinct Multivariate Neural Signatures to Reduce Pain. Biological Psychiatry. 2025 Jan;97(1):81–8.

Dosenbach NUF, Raichle ME, Gordon EM. The brain's action-mode network. Nat Rev Neurosci. 2025 Mar;26(3):158–68.

Obermeier M, Mayr A, Schennach-Wolff R, Seemuller F, Moller H-J, Riedel M. Should the PANSS Be Rescaled? Schizophrenia Bulletin. 2010 May;36(3):455–60.

Ganz M, Poldrack RA. Data sharing in neuroimaging: experiences from the BIDS project. Nat Rev Neurosci. 2023 Dec;24(12):729–30.

Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, et al. fMRIPrep: a robust preprocessing pipeline for functional MRI. Nat Methods. 2019 Jan;16(1):111–6.

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc Natl Acad Sci USA. 2005 July;102(27):9673–8.

Varoquaux G, Sadaghiani S, Pinel P, Kleinschmidt A, Poline JB, Thirion B. A group model for stable multi-subject ICA on fMRI datasets. NeuroImage. 2010 May;51(1):288–99.

Friston KJ. Functional and Effective Connectivity: A Review. Brain Connectivity. 2011 Jan;1(1):13–36.

Rolls ET, Joliot M, Tzourio-Mazoyer N. Implementation of a new parcellation of the orbitofrontal cortex in the automated anatomical labeling atlas. NeuroImage. 2015 Nov;122:1–5.

Breiman L. Random Forests. Machine Learning. 2001 Oct;45(1):5–32.

Proust-Lima C, Philipps V, Liquet B. Estimation of Extended Mixed Models Using Latent Classes and Latent Processes: The *R* Package **lcmm**. J Stat Soft. 2017;78(2). DOI: 10.18637/jss.v078.i02

Lányi O, Zahemszky D, Wenning AS, Engh MA, Molnár Z, Horváth AA, et al. Cerebello-Thalamo-Cortical Dysconnectivity in Schizophrenia Spectrum Disorders: A Resting-State Functional Magnetic Resonance Imaging Meta-Analysis. BPS: CNNI. 2025 June;0(0). DOI: 10.1016/j.bpsc.2025.05.017

Wang Q, Aljassar M, Bhagwat N, Zeighami Y, Evans AC, Dagher A, et al. Reproducibility of cerebellar involvement as quantified by consensus structural MRI biomarkers in advanced essential tremor. Sci Rep. 2023 Jan;13(1):581.

Wang Q, Wang Q, Zhang R-Y. Claim causality with clarity. Psychoradiology. 2023 Mar;3:kkad007.

Figure Legends

Fig. 1. Trial design and time-specific clinical effects of clinician-guided meditation.

(a) Overview of the parallel-group randomized clinical trial and assessment schedule (baseline, 3 months, 8 months). (b) Time-specific average treatment effects (ATEs) of meditation (meditation + rehabilitation vs rehabilitation alone) on PANSS total percentage decrease rate (PDR). (c) ATEs on RBANS percentage increase rate (RIR), showing early improvement with plateauing thereafter. (d) Cross-lagged/mediation model testing temporal coupling between cognition and symptoms: early cognitive improvement predicts later symptom improvement but does not mediate the treatment effect on 8-month PDR (standardized coefficients shown in the figure).

Fig. 2. Heterogeneity of clinical response to meditation and baseline predictors.

(a) Latent-class trajectory model of PDR in the meditation arm identifying high- and normal-responder subgroups, shown alongside the rehabilitation-only group for context. (b) Standardized mean differences (SMDs) in baseline characteristics comparing high vs normal responders. (c) Permutation-based inference (10,000 permutations) for elastic-net logistic regression distinguishing high responders from normal responders; marital status and lower chlorpromazine-equivalent dose (CPZ) emerge as robust predictors (coefficients and p values shown in the figure).

Fig. 3. Biphasic network–symptom associations across treatment and time.

(a) Spatial map of the action-mode network–sensorimotor–executive component (AMN–SEC) derived from group ICA, with correlations between its composite complexity index (CCI) and clinical improvement (PDR and PANSS subscale decrease rates) by group and timepoint. (b) Spatial map of the default mode network–cerebellar-integrative component (DMN–CIC) with corresponding CCI–outcome correlations. (c) Two-wave structural equation model (SEM) testing phase-specific contributions: baseline DMN–CIC complexity predicts 3-month symptom improvement, whereas 3-month AMN–SEC complexity predicts additional 8-month gains (paths and standardized coefficients shown in the figure).

Abbreviations: AMN–SEC, action-mode network–sensorimotor–executive component; CCI, composite complexity index; DMN–CIC, default mode network–cerebellar integrative component; PDR, PANSS decrease rate; PPDR/PNDR/PGDR, PANSS positive/negative/general subscale decrease rates.

Fig. 4. Functional connectivity features predicting short- and long-term symptom improvement under meditation.

Elastic-net models (permutation-validated) predicting PDR in the meditation arm using AAL2 atlas functional connectivity (FC). (a) Prediction of 3-month PDR from baseline FC; highlighted edges (importance threshold as indicated) primarily involve DMN–CIC–related couplings. (b) Prediction of 8-month PDR from baseline FC; highlighted edges emphasize AMN–SEC–related couplings. (c) Prediction of 8-month PDR from early FC change (ΔFC, baseline to 3 months); strongest predictive features include precentral–SMA coupling. Only edges meeting the prespecified importance thresholds are summarized and visualized (thresholds stated in the figure).

Abbreviations: ΔFC, change in functional connectivity; FC, functional connectivity; PDR, PANSS decrease rate; SMA, supplementary motor area.