Modelling Trajectories of Antimicrobial Resistance in Human Populations

Oraya Srimokla[1*], Sean Cavany[1], James Petrie[2], Mark Pritchard[1], Anna Nebykova[1], Gisela Aguilar[1], Christiane Dolecek[1], and Ben Cooper[1]

[1]Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine

University of Oxford

[2]Independent Researcher

[*]Corresponding author: oraya.srimokla@ndm.ox.ac.uk

Abstract

Antimicrobial resistance (AMR) is recognised as a global health threat but temporal trends in the prevalence of resistance are typically obscured by patchy and heterogeneous data, particularly in lower and middle-income settings.

Previous approaches to estimating country-level resistance trajectories in such settings have used an ensemble approach, combining multiple flexible nonlinear algorithmic mean functions embedded within a Gaussian process framework without explicitly accounting for heterogeneities between data sources. We build on this work, but aim to overcome important limitations by explicitly accounting for variation between data sources and using a semi-mechanistic approach based on a logistic growth model with time-varying parameters. We take as an example *Acinetobacter baumannii*, a globally significant cause of antibiotic-resistant healthcare-associated infections and use data from 18 different sources to estimate trajectories in the prevalence of resistance in 32 countries in Asia between 2000 and 2022.

Working within a Bayesian framework, we compare six variants of the underlying model which vary in how data source heterogeneities are accounted for and in how the growth model parameters are assumed to vary by country and time and to depend on covariates and spatial relationships. We also consider a stacked ensemble of these six models. We assess out of sample model accuracy using the Leave-One-Out

Cross-Validation Information Criterion.

For four out of seven pathogen-antibiotic combinations considered, the stacked model had the highest out of sample prediction accuracy. For the other pathogen-antibiotic combinations, models that accounted for heterogeneity between data sources always performed better than models that did not. This framework improves upon existing approaches by offering a biologically plausible structure that can quantify the effects of covariates on resistance dynamics, effectively handle heterogeneous data sources and can be readily extended to incorporate more complex biologically motivated growth processes. This methodology provides a potentially valuable foundation for better understanding AMR resistance dynamics and evaluating public health interventions.

Author Summary

Understanding antimicrobial resistance trajectories and their drivers across countries and over time is important for quantifying the changing health burden of antimicrobial resistance, exploring the potential impact of interventions, and for setting control priorities and informing interventions. However, it is challenging because available surveillance data is sparse and comes from sources that differ in both microbiological practices (e.g. who is sampled and when) and the patient groups and facility types that are represented. We developed a way to model country-level resistance trajectories that accounts for these differences between data sources, represents changes in resistance using a biologically-informed logistic growth model, and that allows resistance trajectories for individual countries to be estimated by borrowing strength from other countries, accounting for spatial dependencies. We applied our method to 18 sources of *Acinetobacter baumannii* resistance data from 32 countries in Asia from 2000 to 2022, considering six different model variants as well as an ensemble approach using Bayesian stacking.

Our results show that accounting for variability between data sources improves how accurately models estimate the proportion of drug-resistant isolates for out-of-sample data. Models that accounted for differences between data sources outperformed other models that did not across all seven pathogen-antibiotic combinations. Combining model variants into a single ensemble model produced the most accurate resistance forecasts when predicting held-out data from the most recent four years, as measured by the mean absolute error. For countries with sparse or absent resistance data, the spatial models generated plausible resistance estimates that reflected trajectories observed in neighbouring countries. Our approach provides a potentially valuable pathway for better understanding resistance trajectories and assessing the potential impacts of different public health interventions.

1 Introduction

Antimicrobial resistance (AMR) threatens global health by compromising infection treatment and straining health systems worldwide [1, 2, 3, 4]. AMR occurs when microorganisms such as bacteria, viruses, parasites, and fungi develop resistance against antimicrobial medications, such as antibiotics, making infections more difficult to treat [3]. In response to the growing threat of AMR, international collaborations have been established to coordinate global action. The 2015 Global Action Plan (GAP) on AMR [5] led to the institution of AMR surveillance systems such as the WHO Global Antimicrobial Resistance and Use Surveillance System (GLASS) [6], the publication of national targets, and the creation of national action plans [7]. The 2024 UN General Assembly political declaration on AMR further strengthened these commitments [8]. Despite these efforts, significant challenges remain in understanding AMR trajectories [9].

Mathematical models have become essential tools for understanding AMR dynamics and can be broadly classified as phenomenological or mechanistic. Phenomenological models describe relationships in data without making assumptions about the underlying biology or transmission mechanisms [10]. For example, Browne et al. used a spatiotemporal stacked ensemble approach to describe the relationship between a set of covariates and resistance prevalence for *S.* Typhi and *S.* Paratyphi A in 75 endemic countries from 1990 to 2019 [11]. While strengths of this approach include its flexibility and apparent good performance for disease systems close to equilibrium [12], a limitation is that in settings which may be far from equilibrium resistance prevalence estimates may be produced that lack biological plausibility and are inconsistent with background domain-specific knowledge. Furthermore, because this approach does not account for intrinsic dynamics or represent processes driving non-equilibrium changes in resistance, it does not lend itself to the exploration of counterfactual scenarios or long-term forecasting. In contrast, mechanistic models aim to explicitly describe transmission dynamics through simplified representations of known processes and include model parameters with biologically meaningful interpretations [10]. For example, Kachalov et al. developed a compartmental model to represent the spread of carbapenem-resistant *Klebsiella pneumoniae* and extended-spectrum beta-lactamase-producing (ESBL) *Klebsiella pneumoniae* in community and hospital settings in Europe [13], while similar modelling approaches have been used to evaluate the effect of antimicrobial stewardship interventions [14, 15, 16, 17]. Both modelling approaches face substantial challenges when applied to global AMR data. Resistance data typically originate from various data sources, including clinical settings, national surveillance programs, and site-specific studies, each with distinct sampling protocols, reporting practices, and data processing standards [16, 18, 19, 3, 20]. Moreover, microbiological sampling practices can vary substantially between settings and are thought to have a significant impact on the directly

observed prevalence of resistance [21]. Additional challenges include sparse data coverage, particularly in low and middle-income countries, bias in available data towards healthcare-seeking populations, and low blood culture utilisation rates [16, 22, 23, 24, 21].

The approach we propose here can be thought of as a semi-mechanistic model. This represents a hybrid approach, sitting between fully mechanistic models, where the underlying dynamical model is a simplified representation of known processes, and phenomenological models, where nothing about the data generation process is assumed. Our mechanistic component builds on work of Emons et al. [25], who considered European AMR data for 887 pathogen-drug combinations. While they observed that for many pathogen-drug combinations resistance remained stable over two decades, for many others resistance trajectories could be well-described by a scaled logistic growth model with an initial exponential increase in resistance followed by stabilisation at an intermediate level. This scaled logistic growth model represents the solution to a simple mechanistic model for resistance where the dynamics of drug-resistant phenotypes are decoupled from those of drug-sensitive phenotypes. We model the relationship between the parameters of this underlying mechanistic model and covariates phenomenologically, without imposing assumptions about mechanisms or even directions of effect.

Our modelling framework also builds on the approach of Bhatt et al. who used a stacked ensemble approach to combine multiple nonlinear models for the mean function of a Gaussian process regression applied to *Plasmodium falciparum* prevalence data in sub-Saharan Africa [12, 26]. However, while Bhatt et al. used a variety of functions to directly map covariates to observed prevalence, in contrast we map covariates to parameters of a dynamic process at a particular place and time; the system's intrinsic dynamics then determine predicted resistance prevalence at any given country and time.

In this paper we first describe the example data we are considering. We then show how the logistic model emerges as a solution to a simple mechanistic model of resistance transmission, and describe how this basic model structure can be extended across six model variants to incorporate country and source-level variation, spatial correlation, and time-varying covariates that potentially impact resistance dynamics. Following this, we describe model fitting within a Bayesian framework, model performance evaluation using leave-one-out cross-validation, and the application of Bayesian model stacking to generate an ensemble model optimised for out-of-sample predictions. Finally, we present results comparing model performance across seven *Acinetobacter baumannii*-antibiotic combinations in Asia from 2000 to 2022, and discuss the implications of our findings for AMR surveillance and the potential for future public health intervention evaluation.

2 Methods

2.1 Data

Resistance data for *Acinetobacter baumannii* was obtained from the Global Research on Antimicrobial Resistance 2024 global burden study [4] which makes use of around 1.3 million clinical isolates for 113 countries globally from 1991 to 2022, combining data from surveillance programs, individual testing facilities, and systematic reviews. We extracted resistance data for seven classes of antibiotics: aminoglycosides, antipseudomonal penicillins, beta-lactams/beta-lactamase inhibitor combinations, carbapenems, fluoroquinolones, fourth-generation cephalosporins, and third-generation cephalosporins. For this analysis, we focus on countries in Asia and the period from 2000 to 2022, for which we retrieved data from 333936 clinical isolates across 18 data sources, 32 countries, and 7 pathogen-antibiotic combinations. See Section A in S1 Text for for the total isolates per year from 2000 to 2022 across Asia.

Covariate data were selected from previous studies identifying possible associations with resistance in *Acinetobacter baumannii* [27, 28, 29, 30, 31]. Covariate data included in the models are time-dependent and country-level gridded mean temperature data from the Copernicus Climate Change Service [32], estimates of total human antibiotic consumption (defined daily doses per 1000 inhabitants per day (DDD per 1000)) and the fraction of out-of-pocket healthcare expenditures from the 2024 Global Burden of Resistance study [4, 33]. All covariates included were available at year and country level.

2.2 Model Structure

Our underlying model of resistance dynamics is a logistic growth process and we consider variants of this basic model with parameters that vary in space and time as a function of covariates. There are a number of possible parameterisations of this model. In this paper, we focus on a parameterisation where the logistic curve emerges as the solution to a two-compartment ordinary differential equation, the susceptible-infectious-susceptible (SIS) transmission model. This SIS model can be expressed as

$$\frac{dX}{dt} = \frac{-\beta}{N}XI + \gamma * I \tag{1}$$

$$\frac{dI}{dt} = \frac{\beta}{N}XI - \gamma * I \tag{2}$$

where X represents the number susceptible to infection and I the number currently infectious.

Mechanistically, we might think of this model as a simple approach for representing carriage dynamics for variants of a particular bacterial species that are resistant to a particular antibiotic when the dynamics of drug-resistant and drug-sensitive strains are decoupled from each other (see Section B in S1 Text), an assumption that has some empirical support for a number of important nosocomial bacterial pathogens [34].

Under this assumption, the proportion of the population colonised with resistant strains, P_t, follows logistic growth dynamics [25], described by the Verhulst equation:

$$P_t = \frac{K}{1 + (\frac{K - P_0}{P_0})e^{-rt}} \tag{3}$$

where K is the carrying capacity, r is the growth rate, and P_0 is the initial proportion colonised. We parameterise this equation in terms of β and γ, rather than the standard carrying capacity and growth rate. In this parameterisation, the initial exponential growth rate, r can be defined as $\beta - \gamma$ and the carrying capacity as $(1 - \gamma/\beta)$ or, equivalently, $1 - 1/R_0$, where R_0 is the basic reproduction number (see Section B in S1 Text).

Note that, P_t cannot be directly equated to the observed proportion of clinical infections of a particular pathogen that are resistant. This latter quantity depends on additional factors including the proportion of people colonised with drug-sensitive strains, the risk of infection given colonisation (which may differ for resistant and sensitive strains), and the relative chances of drug-resistant and drug-sensitive infections being detected and reported, all which may vary with data. To account for these factors we introduce a source-specific scaling factor, η_s, later described below.

We extend this simple model to allow parameters governing the growth process to vary between countries, source, and over time to capture temporal changes in resistance dynamics. For models incorporating covariates, parameters can vary by country and year (see Table 1). For models without covariates, each country has country-specific but temporally constant values for β and γ and produces a singular growth curve per country. Six model variants are explored with different combinations of country and data source-level variation, spatial correlation effects, and covariates (see Table 2).

The proportion of the population colonised with resistant strains for the most general model (SP+S+X) at year t for country c and source s can be defined by the following piecewise logistic function:

$$P_{c,s,t_i} = \left(\frac{1 - \frac{\gamma_c}{\beta_{c,t_i}}}{1 + \left(\frac{1 - \frac{\gamma_c}{\beta_{c,t_i}}}{P_{c,s,t_{i-1}}} - 1 \right) e^{-(\beta_{c,t_i} - \gamma_c)(t_i - t_{i-1})}} \right) * \eta_s \tag{4}$$

where $P_{c,s,t_{i-1}}$ is the country- and source-specific resistance prevalence from the previous observation at index $i - 1$, η_s is a source-specific multiplicative scaling factor, and consecutive times t_{i-1} and t_i can be

unequally spaced. The initial condition, P_{c,s,t_0} is defined as:

$$P_{c,s,t_0} = \text{logit}^{-1}(P_0 + u_{P_0,c}) * \eta_s \tag{5}$$

where P_0 is the population-level initial resistance value in the logit space, and $u_{P_0,c}$ is the spatial random effect for the initial prevalence of resistance for country c.

For this model, we represent β_{c,t_i} as a function of time- and country-specific covariates:

$$\log(\beta_{c,t_i}) = B + u_{\beta,c} + \sum_{k=1}^{K} \theta_k \cdot X_{k,c,t_i} \tag{6}$$

where B is the population-level mean value, X_{k,c,t_i} are the K normalised covariates for country c at year t_i, $u_{\beta,c}$ represents the spatial effect for β_c for country c, and θ_k are the estimated covariate coefficients (see Section C and Section D in S1 Text). This structure ensures that β_{c,t_i} remains positive while allowing time-varying dynamics through the addition of covariates. For simplicity, we assume here that γ_c remains constant:

$$\log(\gamma_c) = G + u_{\gamma,c} \tag{7}$$

where G is the population-level mean value, and $u_{\gamma,c}$ is the spatial effect for γ_c for country c (see Table 1).

Spatial effects are incorporated through Gaussian process regression to capture spatial autocorrelation between neighbouring countries [35, 36]. To establish relationships between geographically close sites, we use a radial basis kernel function to define the covariance between countries according to the scaled distance (see Section D in S1 Text):

$$K_{i,j} = \zeta_\epsilon^2 * exp(-\rho^2 * D_{i,j}^2) \tag{8}$$

where $D_{i,j}$ is the scaled Euclidean distance from geographical midpoints between countries i and j, ζ_ϵ^2 is the maximum covariance between sites for model parameter ϵ, and ρ^2 is the rate covariance declines with distance. This covariance function is used to estimate spatially-correlated random effects for each country and is used to define $\beta_{c,t}$, γ_c, P_{c,s,t_0} [35, 36].

Simpler nested model variants are obtained by constraining components of the general model (SP+S+X). For model variants that retain spatial random effects, removing covariates gives a spatially-varying but temporally constant β_c and source-level scaling (SP+S). Further removing source-level scaling ($\eta_s = 1$)

gives a model with only spatial random effects (SP).

Country-level random effects allow β and γ to vary by country c with hierarchical priors (see Table 1 and Section E in S1 Text). The spatial structure can alternatively be replaced by country-level random effects by removing spatial effect terms $u_{\beta,c}$ and $u_{\beta,c}$ and population-level means B and G are instead drawn from a shared population-level distribution:

$$B_c \sim \text{Normal}(\mu_\beta, \sigma_\beta^2) \tag{9}$$

$$G_c \sim \text{Normal}(\mu_\gamma, \sigma_\gamma^2) \tag{10}$$

where μ_β and μ_γ are the population-level means and σ_β^2 and σ_γ^2 are the between-country variances (see Section E in S1 Text). Three model variants are obtained with country-level random effects: country-level effects with only source-level scaling (C+S), country-level effects with only covariates (C+X), and the simplest variant, country-level effects with no covariates and source-level scaling (C) (see Table 1 and Section C in S1 Text) .

Observations of resistant isolates are assumed to follow a binomial distribution:

$$y_{c,s,t} \sim \text{Binomial}(\text{N}_{c,s,t}, P_{c,s,t}) \tag{11}$$

where N_t is the total number of isolates tested for a given country-source combination at time t, and $P_{c,s,t}$ is the predicted proportion of clinical isolates from sources which are resistant.

Countries with Missing Data: For countries without resistance data, the Gaussian process framework enables spatial imputation using the posterior estimates of the maximum covariance between sites ζ^2, and the rate at which covariance declines with distance ρ^2 to calculate the covariance function for countries with unobserved resistance data. New values are predicted following a multivariate normal predictive distribution:

$$p(\mathbf{f}_*|\mathbf{f}) \sim \mathcal{N}(\mathbf{K}_*^T \mathbf{K}^{-1} \mathbf{f}, \mathbf{K}_{**} - \mathbf{K}_*^T \mathbf{K}^{-1} \mathbf{K}_*) \tag{12}$$

where f is the observed spatial effects for countries with resistance data, f_* is the spatial effects for countries with no resistance data, K is the covariance matrix of countries with resistance data, K_* is the covariance matrix with a combination of countries with and without resistance data, and K_{**} is the covariance matrix of countries with no resistance data [37, 38].

Ensemble modelling: To combine predictions across model variants, we apply Bayesian model stacking

Table 1. Definitions of model parameters for six model variants. Parameters are defined based on logistic growth dynamics where $P_{c,t}$ represents the proportion of resistant isolates at time t and country c. Here Model C: Model with country-level random effects, C+X: Country-level random effects model and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with source-level scaling, SP+S+X: Spatial model with source-level scaling and covariates.

Model	Rate of Resistance Coefficient (β)	Loss of Resistance Rate (γ)	Initial Prevalence (P_0)	Source Effect (η_s)
Model C	β_c (Country-specific)	γ_c (Country-specific)	$P_{0,c}$ (Country-specific)	None ($\eta_s = 1$)
Model C + X	$\beta_{c,t} = \exp(\log \beta_c + \sum \theta_k \cdot X_{k,c,t})$ (country-specific+time-varying covariates)	γ_c	$P_{0,c}$	None ($\eta_s = 1$)
Model C + S	β_c	γ_c	$P_{0,c}$	$P_{c,s,t} = P_{c,t} \times \eta_s$
Model SP	$\beta_c = \exp(\log \beta + k_{\beta,c})$ (spatial effect k_β)	$\gamma_c = \exp(\log \gamma + k_{\gamma,c})$ (spatial effect k_γ)	$P_{0,c} = \text{logit}^{-1}(P_{0\_logit} + k_{P_0,c})$ (spatial effect k_{P_0})	None ($\eta_s = 1$)
Model SP + S	$\beta_c = \exp(\log \beta + k_{\beta,c})$	$\gamma_c = \exp(\log \gamma + k_{\gamma,c})$	$P_{0,c} = \text{logit}^{-1}(P_{0\_logit} + k_{P_0,c})$	$P_{c,s,t} = P_{c,t} \times \eta_s$
Model SP + S + X	$\beta_{c,t} = \exp(\log \beta + k_{\beta,c} + \sum \theta_k \cdot X_{k,c,t})$ (spatial effect k_β + time-varying covariates)	$\gamma_c = \exp(\log \gamma + k_{\gamma,c})$	$P_{0,c} = \text{logit}^{-1}(P_{0\_logit} + k_{P_0,c})$	$P_{c,s,t} = P_{c,t} \times \eta_s$

using the loo (v2.8.0) package in R [39]. Model stacking generates ensemble predictions by finding the combination of models that maximises the leave-one-out (LOO) predictive density and the expected log pointwise predictive density (ELPD) of the ensemble predictions [12, 26, 40]. LOO predictive densities are obtained using the Pareto-smoothed importance sampling leave-one-out cross-validation (LOO-CV), a method for estimating pointwise out-of-sample prediction accuracy from a fitted Bayesian model [39]. The optimal model weights for each pathogen-antibiotic combination were obtained using the loo_model_weights() function, and used to determine the contribution of each model's predictive distribution to the ensemble [39, 26].

Table 2. Model variants and the model components that make up each model. Here Model C: Model with country-level random effects, C+X: Country-level random effects model and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with source-level scaling, SP+S+X: Spatial model with source-level scaling and covariates.

	Model C	Model C + X	Model C + S	Model SP	Model SP + S	Model SP + S + X
Country-level Random Effects	✓	✓	✓			
Covariates		✓				✓
Data source-level Random Effects			✓		✓	✓
Spatial Random Effects				✓	✓	✓

2.3 Model Fitting and Evaluation

Models were implemented in the Stan programming language, and model fitting and Hamiltonian Monte Carlo (HMC) sampling performed via the RStan interface (version 2.32.6).

Prior distributions for the models were selected to be weakly informative or informative when domain knowledge suggested biologically plausible parameter ranges (see Section C to F in S1 Text). We conducted prior and posterior predictive checks. For prior predictive checks, data were simulated from prior distributions to assess if the priors produce reasonable model outputs [41]. After fitting the model to empirical data, posterior predictive checks were done by simulating new replicated data sets based on the fitted model [41].

For each of the models, we assessed convergence using standard MCMC diagnostics including traceplots and Rhat statistics (with Rhat < 1.01). We additionally examined the number of divergent transitions, maximum tree depth, and pairs plots to identify issues with convergence or model parameterisation. Parameter recoverability was assessed by fitting the models to simulated data with known parameter values.

The models were then fitted with empirical data. For each of the six model variants, posterior parameter estimates, observation-level predictions, and country-specific growth curves for the years 2000 to 2022 were extracted from fitted model objects.

The performance of the six individual model variants (see Table 2) for each pathogen-antibiotic combination were compared using the Leave-One-Out Cross-Validation Information Criterion (LOOIC) obtained from applying the LOO-CV method via the loo (v2.8.0) package in R [42]. The LOOIC value was computed from the ELPD, with lower LOOIC values indicating better expected out-of-sample predictive performance.

To evaluate the models' performance for forecasting future resistance trends we fitted models to a training dataset and evaluated the predictions on a test set created by leaving out the most recent four years of data for each country. Predictive accuracy for this assessment was quantified using the mean absolute error (MAE) between observed and predicted values.

3 Results

After fitting model variants to all countries in Asia from 2000 to 2022 we found that all six model variants estimated qualitatively similar temporal trajectories for resistance for each country, though for countries with sparse data in the early 2000s, spatial models produced trajectories more influenced by neighbouring countries (see Fig 1 and Section G in S1 of Text). In most cases (192 of 214 antibiotic-country combinations) increasing resistance trends were estimated, though in a few cases (22 of 214 antibiotic-country combinations) declining trends were seen, most notably in Japan for all seven antibiotic combinations. Using models without covariates (C, C+S, SP, SP+S), we estimated the long-term saturation levels of resistance for each country and found that the majority of countries showed intermediate saturation levels rather than resistance fixation (see Section H in S1 of Text). For models with covariates, we observe year-to-year fluctuations in temporal resistance trajectories as a result of time-varying covariates (see Section G in S1 of Text). Estimated covariate coefficients for models incorporating covariates are additionally extracted (see Section I in S1 Text). As expected, uncertainty in estimated resistance proportions varied substantially by country and time, with high levels of uncertainty typically seen at the beginning of the time series where the data were most sparse. There was also substantial variation in levels of uncertainty in the estimated prevalence of resistance between the different models, with the spatial models (SP, SP+S, SP+S+X) typically giving narrower predictive intervals for country-level resistance proportions when data was sparse, a consequence of borrowing information from nearby countries.

Comparing out-of-sample predictive performance using leave-one-out cross-validation, in all cases, models that included random effects showed substantially lower LOOIC values (indicating better performance) than models that did not (Table 5). Comparing models that included covariates with corresponding models that did not (C+X v C and SP+S v SP+S+C) we found that including covariates consistently improved predictive performance for five out of the seven antibiotic classes considered (aminoglycosides, antipseudomonal penicillins, carbapenems, fluoroquinolones and third-generation cephalosporins), and in some cases led to substantial improvements. The model incorporating both country-level and source-level scaling (C+S) produced the lowest LOOIC value across five of seven antibiotic categories, indicating the best predictive performance (see Table 5). For the remaining two antibiotic categories (aminoglycosides and antipseudomonal penicillins), the model combining spatial effects and source-level scaling with covariates (SP+S+X) had the lowest LOOIC. Model C with only country-level random effects consistently had the second highest LOOIC, while the base spatial model (SP) had the highest LOOIC values for six of seven antibiotic categories. These results suggest that country-level effects and spatial effects alone are not able to capture heterogeneity in the

Fig. 1. Individual model fits using β, γ, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* and carbapenems model training for Asia. Posterior predictions for the proportion of clinical isolates that are resistant to carbapenems for five countries and six model variants. See Table 2 for model definitions. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

resistance data. In contrast, spatial models incorporating source-level scaling and/or covariates had relatively good predictive performance, ranking in the top three across all antibiotic combinations.

An ensemble modelling approach using Bayesian model stacking has potential to improve predictive performance. The model stacking weights are shown in Table 3. The model incorporating both country-level and source-level scaling (C+S) was assigned the highest weight for five out of seven antibiotic categories, indicating optimal prediction accuracy if the sampling is heavily weighted for this model. For the remaining two antibiotics categories, aminoglycosides and antipseudomonal penicillins, the highest weights were assigned to the spatial model with source-level scaling (SP+S) and the country-level effects model with covariates, respectively. The model with only country-level random effects (C), and/or the model with only spatial effects (SP) generally received lower weightings across different antibiotic categories. Examples of stacked model predictions for nine countries using these stacking weights are shown in Fig 2.

Antibiotic Class/Subclass	Model Stacking Weights		

	Model C	Model C+X	Model C+S	Model SP	Model SP+S	Model SP+S+X
Aminoglycosides	0.16	0.12	0.04	0.10	**0.32**	0.26
Antipseudomonal penicillins	0.03	**0.34**	0.12	0.02	0.26	0.23
Beta-lactams/beta-lactamase inhibitor combinations	0.08	< 0.001	**0.63**	0.13	0.06	0.10
Carbapenems	< 0.001	0.29	**0.48**	0.16	0.05	0.02
Fluoroquinolones	0.02	0.18	**0.44**	0.02	0.28	0.06
Fourth-generation cephalosporins	< 0.001	0.06	**0.54**	0.23	< 0.001	0.17
Third-generation cephalosporins	0.02	0.26	**0.37**	0.02	< 0.001	0.33

Table 3. Comparison of fitted models weights for each antibiotic category using model stacking. The models with the highest weights are shown in bold. See Table 2 for model definitions.

For models incorporating source-level random effects, the MAE was calculated for each country-source-time combination and for all other models, it was calculated for each country-time combination. The MAE values for all pathogen-antibiotic combinations are shown in Table 4. The stacked model had the lowest MAE value for four out of seven antibiotic categories, while the spatial model with source-level random effects and covariates (SP+S+X) had the lowest for two and the model with country-level random effects and source-level scaling (C+S) had the lowest MAE for a single antibiotic category. Similar to the LOOIC model comparison, models only incorporating country-level random effects (C) or only spatial effects (SP) tended to have lower predictive accuracy. The spatial model (SP) was found to have the largest MAE values for four of seven antibiotic categories.

Antibiotic Class/Subclass	Mean Absolute Error						
	Model C	Model C+X	Model C+S	Model SP	Model SP+S	Model SP+S+X	Stacked Model
Aminoglycosides	0.1684	0.1672	0.1525	0.1646	0.1513	**0.1511**	0.1532

Antipseudomonal penicillins	0.1284	0.1277	0.1222	0.1294	0.1257	0.1236	**0.1210**
Beta-lactams/beta-lactamase inhibitor combinations	0.1861	0.1854	0.1872	0.1848	0.1890	**0.1847**	0.1951
Carbapenems	0.1406	0.1403	0.1362	0.1423	0.1392	0.1374	**0.1361**
Fluoroquinolones	0.1187	0.1188	0.1215	0.1250	0.1215	0.1203	**0.1177**
Fourth-generation cephalosporins	0.1288	0.1302	0.1231	0.1273	0.1219	0.1221	**0.1216**
Third-generation cephalosporins	0.1333	0.1340	**0.1259**	0.1429	0.1367	0.1370	0.1267

Table 4. Comparison of mean absolute error (MAE) calculated for four years of held-out data across countries, years, and sources (for source-specific models). The MAE was calculated for a held-out test set containing the most recent four years of data for each country. The best model for each antibiotic category is shown in bold. Lower MAE values indicate more accurate model predictions. See Table 2 for model definitions.

Analysis using simulated data with the country-level random effects model with a β_c, γ_c, and $P_{0,c}$ parameterisation confirmed that the models were able to recover the true values of the parameters; posterior predictive checks further confirmed the model was able to capture the mean and standard deviation of the observed data (see Section J in S1 Text). The convergence of all six model variants was visually checked for all seven pathogen-antibiotic combinations using trace plots and pairs plots (see Section K in S1 Text). All plots exhibit convergence of all four chains in the trace plot, appropriate Rhat values, and do not show model divergences.

4 Discussion

We developed a Bayesian modelling framework that is able to account for time-varying parameters, country and source-level variation, covariates, and spatial correlation effects to estimate country-level antimicrobial resistance trajectories. We compared six model variants applied to seven *Acinetobacter baumannii*-antibiotic combinations across Asia from 2000 to 2022 and explored the applications of model stacking to generate ensemble predictions. We found that models incorporating source-level scaling consistently outperformed other models, as shown by the lowest LOOIC values. We accounted for spatial correlation effects in some of

(a)

Fig. 2. Stacked model predictions using $\beta\ \gamma$, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* and carbapenems for Asia from 2000 to 2022. Shows stacked model estimates for nine countries. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

Antibiotic Class/Subclass	Model Comparison using LOOIC differences from best model					
	Model C	Model C+X	Model C+S	Model SP	Model SP+S	Model SP+S+X
Aminoglycosides	431.3	430.4	38.3	466.9	4.9	**0**
Antipseudomonal penicillins	200.8	136.6	49.2	221.0	71.8	**0**
Beta-lactams/beta-lactamase inhibitor combinations	196.0	210.0	**0**	230.0	55.4	47.3
Carbapenems	378.2	329.9	**0**	432.2	100.9	95.8
Fluoroquinolones	663.6	567.7	**0**	722.9	27.7	13.3
Fourth-generation cephalosporins	131.9	132.7	**0**	130.8	31.7	30.7
Third-generation cephalosporins	312.2	221.9	**0**	334.3	56.6	9.5

Table 5. Comparison of fitted models for each antibiotic class using the leave-one-out cross-validation information criterion (LOOIC). Relative differences in LOOIC calculated from the best model (shown in bold with 0). Lower values indicate better predictive performance. See Table 2 for model definitions.

our models and observed an improvement in model performance when including covariates in the spatial and source-level scaling model (SP+S). We found that in most cases the ensemble approach, combining estimates from multiple component models, led to improved predictive accuracy. This framework is able to capture resistance dynamics of AMR across different countries, and enables estimates for countries where data are sparse by borrowing information from other countries.

While our modelling approach is broadly phenomenological in the sense that we aim to capture the sigmoidal growth process that changes in AMR prevalence typically follow without making explicit mechanistic assumptions, we note that the models with source-level scaling can be given a semi-mechanistic interpretation as the solution of a deterministic SIS model for the resistant strain, assuming no interaction of drug-resistant and drug-sensitive strain dynamics and a constant prevalence of drug-sensitive strains. This simple model might be considered an appropriate mechanistic representation of the transmission process when there is minimal niche overlap between drug-resistant and drug-sensitive strains [43].

The primary motivation for this parameterisation, however, is the presence of an analytical solution to this system (the logistic equation). This leads to computational efficiency making it an appealing choice compared to alternative models of sigmoidal growth when used as the basis for complex models applied to multicountry data. The underlying modelling framework we propose, however, can readily be extended to systems corresponding to different mechanistic assumptions. We derived our model in terms of β and γ as this parameterisation showed improved model convergence when compared alternative parameterisations. Within this framework we included time-varying parameters based on non-linear mappings between covariates that were thought likely to influence resistance dynamics such as antibiotic usage and hospital resourcing levels while accounting for spatial correlation and heterogeneity between data sources.

The framework presented is well-suited to capturing resistance dynamics where, as observed by Emons et al. across European pathogen-drug combinations, trajectories can reflect a mix of non-equilibrium and shifting equilibrium frequencies in response to changing antibiotic use. This framework has some advantages over alternative approaches that map covariates directly to observed prevalence data thus failing to account for intrinsic resistance dynamics that result from the communicable nature of the pathogens.

Our model comparison revealed that model structures capable of accounting for source-level heterogeneity (which also corresponds to the subset of models amenable to a mechanistic interpretation) had improved predictive accuracy. We found that models incorporating source-level scaling with the country-level effects model or the spatial model had the best performance across all antibiotic categories, shown by the lowest LOOIC. AMR data originate from various sources which have differences in patient care, patient characteristics, local resistance prevalence (within country variation), and microbiological culture practices (sampling protocols and laboratory methods). Accounting for source-level variability implicitly accounts for these differences. Our final model stacking weights similarly reflect the better performance of these models, as the highest weight is assigned to models with data source-level scaling for six out of seven antibiotic categories.

Incorporating covariates generally improved model performance across both models with only country-level effects (C) and the spatial effects model (SP). For the model with only country-level effects (C), adding covariates (C+X) reduced the LOOIC values for five out of seven antibiotic categories. We observed a similar pattern when adding covariates to our spatial models. When comparing the spatial model (SP), the spatial model with data source-level scaling (SP+S), and the spatial model with data source-level scaling and covariates (SP+S+X), we see a decrease in LOOIC values across all antibiotic categories when covariates are added to the spatial model with data-source level effects.

When comparing the base spatial model (SP) to the country-level effects model (C), the addition of spatial correlation effects did not consistently reduce LOOIC values across all antibiotic categories. Similarly,

combining spatial effects with source-level scaling (SP+S) did not always improve predictive performance relative to the country effects and data source-level scaling (C+S) model. Nevertheless, incorporating a spatial correlation structure provides a useful feature that enables the estimation of resistance trends for countries without observed resistance data. Through the Gaussian process structure of the spatial model, we can estimate trends for data-sparse countries by borrowing information from their neighbouring countries. Our current implementation uses geographic distance to define the spatial correlation structure and assumes neighbouring countries share similar resistance patterns. However, alternative spatial definitions could be explored within this framework. For example, human movement patterns through travel and migration can be more mechanistically relevant when considering a spatial structure in terms of resistance dynamics [17].

This study has several limitations. Our model parameterisation is derived from a two-compartment system that simplifies the dynamics of drug-resistant and drug-sensitive strains and is not intended to be given a literal mechanistic interpretation. More complex formulations that include hospital and community compartments and separate antibiotic use by hospital and community use and by antibiotic class might be expected to lead to more meaningful mechanistic insights as well as having the potential for better predictive performance. Also, while our framework currently allows for time-varying parameters driven by covariates, both evolutionary and ecological processes and unmeasured covariates could lead to changes in the dynamical system that the current framework does not capture. To address these there may be additional benefits in allowing model parameters to change over time for reasons other than covariate change, for example through the incorporation of a random walk process into the model. Finally, further extensions would be needed to account for potential biases in resistance data resulting from microbiological culture practices [21] and changing resistance breakpoints.

In this work, we developed a Bayesian semi-mechanistic modelling framework designed to address the challenge of estimating AMR dynamics across large spatial scales where data sources are often scarce and heterogeneous. This framework accounts for data source heterogeneity, covariate data, intrinsic resistance dynamics and spatial correlation structures. We found that the best-fitting models were those which both accounted for data source heterogeneity and which lent themselves to a biological interpretation. Using a model stacking approach consistently led to improved predictive performance. This methodology provides a framework for using diverse data sources to understand the dynamics of global resistance trends and offers the potential for extension to more explicitly mechanistic formulation which would provide a foundation for evaluating potential public health interventions.

Acknowledgements

I would like to thank the GRAM team at IHME and Oxford and all our GRAM data collaborators for building the foundation of this project and making this work possible. I would also like to thank the UK Department of Health and Social Care's Fleming Fund, and the Wellcome Trust for funding this project. I would like to acknowledge the NeoOBS Study Collaborators, Global Antibiotic Research and Development Partnership, Geneva, Switzerland. I acknowledge Cardiff University in relation to use of data from the BARNARDS Study.

Data Availability

The GRAM Project is committed to making data available to external users as widely as possible. Data collected in Oxford that are approved for public release can be accessed via the Antimicrobial Resistance (AMR) data repository hosted by the Infectious Diseases Data Observatory (IDDO) at https://www.iddo.org/antimicrobial-resistance. Data collected by the Institute for Health Metrics and Evaluation (IHME) can be accessed through the Global Health Data Exchange (GHDx) at https://ghdx.healthdata.org/. For data not publicly available through these repositories, access is subject to the terms and conditions of the original data contributors. Researchers wishing to request access to these data should contact the corresponding author. Requests will be reviewed in consultation with data owners, and access will be granted where permitted under the applicable data sharing agreements.

All code used in this study is publicly available on GitHub at https://github.com/orayasrimokla/methods_modelling_trajectories_AMR_manucript

Funding

This work was funded by the UK Department of Health and Social Care Fleming Fund and the Wellcome Trust.

References

1. Global Antimicrobial Resistance and Use Surveillance System (GLASS) report: antibiotic use data for 2022. Available from: https://www.who.int/publications/i/item/9789240108127 [Accessed on: 2025 May 6]

2. Prestinaci F, Pezzotti P, and Pantosti A. Antimicrobial resistance: a global multifaceted phenomenon. Pathogens and Global Health 2015 Oct; 109:309–18. DOI: 10.1179/2047773215Y.0000000030. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4768623/

3. Ahmed SK, Hussein S, Qurbani K, Ibrahim RH, Fareeq A, Mahmood KA, and Mohamed MG. Antimicrobial resistance: Impacts, challenges, and future prospects. Journal of Medicine, Surgery, and Public Health 2024 Apr; 2:100081. DOI: 10.1016/j.glmedi.2024.100081. Available from: https://www.sciencedirect.com/science/article/pii/S2949916X24000343

4. Naghavi M et al. Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050. The Lancet 2024 Sep; 404:1199–226. DOI: 10.1016/S0140-6736(24)01867-1. Available from: https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)01867-1/fulltext

5. Global action plan on antimicrobial resistance. Available from: https://www.who.int/publications/i/item/9789241509763 [Accessed on: 2025 May 6]

6. Global Antimicrobial Resistance and Use Surveillance System (GLASS). Available from: https://www.who.int/initiatives/glass [Accessed on: 2025 May 6]

7. Target. Global Database for Tracking Antimicrobial Resistance (AMR) Country Self- Assessment Survey (TrACSS). Available from: http://amrcountryprogress.org/ [Accessed on: 2025 May 6]

8. UN General Assembly High-Level Meeting on antimicrobial resistance 2024. Available from: https://www.who.int/news-room/events/detail/2024/09/26/default-calendar/un-general-assembly-high-level-meeting-on-antimicrobial-resistance-2024 [Accessed on: 2025 Nov 19]

9. Limmathurotsakul D, Dunachie S, Fukuda K, Feasey NA, Okeke IN, Holmes AH, Moore CE, Dolecek C, Doorn HRv, Shetty N, Lopez AD, and Peacock SJ. Improving the estimation of the global burden of antimicrobial resistant infections. The Lancet Infectious Diseases 2019 Nov; 19:e392–e398. DOI:

10.1016/S1473-3099(19)30276-2. Available from: https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(19)30276-2/fulltext

10. Porgo TV, Norris SL, Salanti G, Johnson LF, Simpson JA, Low N, Egger M, and Althaus CL. The use of mathematical modeling studies for evidence synthesis and guideline development: A glossary. Research Synthesis Methods 2019 Mar; 10:125–33. DOI: 10.1002/jrsm.1333. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6491984/

11. Browne AJ, Chipeta MG, Fell FJ, Haines-Woodhouse G, Kashef Hamadani BH, Kumaran EAP, Robles Aguilar G, McManigal B, Andrews JR, Ashley EA, Audi A, Baker S, Banda HC, Basnyat B, Bigogo G, Ngoun C, Chansamouth V, Chunga A, Clemens JD, Davong V, Dougan G, Dunachie SJ, Feasey NA, Garrett DO, Gordon MA, Hasan R, Haselbeck AH, Henry NJ, Heyderman RS, Holm M, Jeon HJ, Karkey A, Khanam F, Luby SP, Malik FR, Marks F, Mayxay M, Meiring JE, Moore CE, Munywoki PK, Musicha P, Newton PN, Pak G, Phommasone K, Pokharel S, Pollard AJ, Qadri F, Qamar FN, Rattanavong S, Reiner B, Roberts T, Saha S, Saha S, Shakoor S, Shakya M, Simpson AJ, Stanaway J, Turner C, Turner P, Verani JR, Vongsouvath M, Day NPJ, Naghavi M, Hay SI, Sartorius B, and Dolecek C. Estimating the subnational prevalence of antimicrobial resistant Salmonella enterica serovars Typhi and Paratyphi A infections in 75 endemic countries, 1990–2019: a modelling study. The Lancet Global Health 2024 Mar; 12:e406–e418. DOI: 10.1016/S2214-109X(23)00585-5. Available from: https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(23)00585-5/fulltext

12. Bhatt S, Cameron E, Flaxman SR, Weiss DJ, Smith DL, and Gething PW. Improved prediction accuracy for disease risk mapping using Gaussian process stacked generalization. Journal of The Royal Society Interface 2017 Sep. DOI: 10.1098/rsif.2017.0520. Available from: https://royalsocietypublishing.org/doi/10.1098/rsif.2017.0520

13. Kachalov VN, Nguyen H, Balakrishna S, Salazar-Vizcaya L, Sommerstein R, Kuster SP, Hauser A, Wiesch PAz, Klein E, and Kouyos RD. Identifying the drivers of multidrug-resistant Klebsiella pneumoniae at a European level. PLOS Computational Biology 2021 Jan; 17:e1008446. DOI: 10.1371/journal.pcbi.1008446. Available from: https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008446

14. Durazzi F, Pezzani MD, Arieti F, Simonetti O, Canziani LM, Carrara E, Barbato L, Onorati F, Remondini D, and Tacconelli E. Modelling antimicrobial resistance transmission to guide personalized antimicrobial stewardship interventions and infection control policies in healthcare setting: a pilot study.

Scientific Reports 2023 Sep; 13:15803. DOI: 10.1038/s41598-023-42511-5. Available from: https://www.nature.com/articles/s41598-023-42511-5

15. Changruenngam S, Modchang C, and Bicout DJ. Modelling of the transmission dynamics of carbapenem-resistant Klebsiella pneumoniae in hospitals and design of control strategies. Scientific Reports 2022 Mar; 12:3805. DOI: 10.1038/s41598-022-07728-w. Available from: https://www.nature.com/articles/s41598-022-07728-w

16. Niewiadomska AM, Jayabalasingham B, Seidman JC, Willem L, Grenfell B, Spiro D, and Viboud C. Population-level mathematical modeling of antimicrobial resistance: a systematic review. BMC Medicine 2019 Apr; 17:81. DOI: 10.1186/s12916-019-1314-9. Available from: https://doi.org/10.1186/s12916-019-1314-9

17. Rahbe E, Valdano E, Smith DR, Guillemot D, Cauchemez S, Glaser P, and Opatowski L. Global dynamics of antibiotic resistance: a modeling study of the spread of endemic and emerging resistance in E. coli. 2025 Oct. DOI: 10.1101/2025.09.25.25335519. Available from: https://www.medrxiv.org/content/10.1101/2025.09.25.25335519v1 [Accessed on: 2026 Mar 19]

18. Nateghi Haredasht F, Amrollahi F, Maddali MV, Marshall N, Ma SP, Cooper LN, Johnson AO, Wei Z, Medford RJ, Kanjilal S, Banaei N, Deresinski S, Goldstein MK, Asch SM, Chang A, and Chen JH. Antibiotic Resistance Microbiology Dataset (ARMD): A Resource for Antimicrobial Resistance from EHRs. Scientific Data 2025 Jul; 12:1299. DOI: 10.1038/s41597-025-05649-7. Available from: https://www.nature.com/articles/s41597-025-05649-7

19. Nazir A, Nazir A, Zuhair V, Aman S, Sadiq SUR, Hasan AH, Tariq M, Rehman LU, Mustapha MJ, and Bulimbe DB. The Global Challenge of Antimicrobial Resistance: Mechanisms, Case Studies, and Mitigation Approaches. Health Science Reports 2025 Jul; 8:e71077. DOI: 10.1002/hsr2.71077. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC12284435/

20. Holt KE, Carey ME, Chandler C, Cross JH, Dyson ZA, Furnham N, Glover RE, Virgo M, and Knight GM. Tools and challenges in the use of routine clinical data for antimicrobial resistance surveillance. npj Antimicrobials and Resistance 2025 May; 3:37. DOI: 10.1038/s44259-025-00105-3. Available from: https://www.nature.com/articles/s44259-025-00105-3

21. Lim C, Hantrakun V, Teerawattanasook N, Srisamang P, Teparrukkul P, Sumpradit N, Turner P, Day NP, Cooper BS, Peacock SJ, and Limmathurotsakul D. Impact of low blood culture usage on rates of antimicrobial resistance. The Journal of Infection 2021 Mar; 82:355–62. DOI: 10.1016/j.jinf.2020.10.040. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC7994019/

22. Frost I, Kapoor G, Craig J, Liu D, and Laxminarayan R. Status, challenges and gaps in antimicrobial resistance surveillance around the world. Journal of Global Antimicrobial Resistance 2021 Jun; 25:222–6. DOI: 10.1016/j.jgar.2021.03.016. Available from: https://www.sciencedirect.com/science/article/pii/S2213716521000825

23. Ashley EA, Shetty N, Patel J, Doorn R van, Limmathurotsakul D, Feasey NA, Okeke IN, and Peacock SJ. Harnessing alternative sources of antimicrobial resistance data to support surveillance in low-resource settings. Journal of Antimicrobial Chemotherapy 2019 Mar; 74:541–6. DOI: 10.1093/jac/dky487. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC6406030/

24. Dunachie SJ, Day NP, and Dolecek C. The challenges of estimating the human global burden of disease of antimicrobial resistant bacteria. Current Opinion in Microbiology 2020 Oct; 57:95–101. DOI: 10.1016/j.mib.2020.09.013. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC7763986/

25. Emons M, Blanquart F, and Lehtinen S. The evolution of antibiotic resistance in Europe, 1998–2019. PLOS Pathogens 2025 Apr; 21:e1012945. DOI: 10.1371/journal.ppat.1012945. Available from: https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1012945

26. Yao Y, Vehtari A, Simpson D, and Gelman A. Using Stacking to Average Bayesian Predictive Distributions (with Discussion). Bayesian Analysis 2018 Sep; 13:917–1007. DOI: 10.1214/17-BA1091. Available from: https://projecteuclid.org/journals/bayesian-analysis/volume-13/issue-3/Using-Stacking-to-Average-Bayesian-Predictive-Distributions-with-Discussion/10.1214/17-BA1091.full

27. Howard A, O'Donoghue M, Feeney A, and Sleator RD. Acinetobacter baumannii. Virulence 2012 May; 3:243–50. DOI: 10.4161/viru.19700. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3442836/

28. Kim YA, Kim JJ, Won DJ, and Lee K. Seasonal and Temperature-Associated Increase in Community-Onset Acinetobacter baumannii Complex Colonization or Infection. Annals of Laboratory Medicine 2018 May; 38:266–70. DOI: 10.3343/alm.2018.38.3.266. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5820073/

29. Filho SPF, Pereira MS, Nobre Rodrigues JL, Guimarães RB, Ribeiro da Cunha A, Corrente JE, Campos Pignatari AC, and Fortaleza CMCB. Seasonality and weather dependance of Acinetobacter baumannii complex bloodstream infections in different climates in Brazil. PLoS ONE 2021 Aug; 16:e0255593.

DOI: `10.1371/journal.pone.0255593`. Available from: `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8376013/`

30. Li W, Liu C, Ho HC, Shi L, Zeng Y, Yang X, Huang Q, Pei Y, Huang C, and Yang L. Association between antibiotic resistance and increasing ambient temperature in China: an ecological study with nationwide panel data. The Lancet Regional Health – Western Pacific 2023 Jan; 30. DOI: `10.1016/j.lanwpc.2022.100628`. Available from: `https://www.thelancet.com/journals/lanwpc/article/PIIS2666-6065(22)00243-7/fulltext`

31. Kritsotakis EI and Groves-Kozhageldiyeva A. A systematic review of the global seasonality of infections caused by *Acinetobacter* species in hospitalized patients. Clinical Microbiology and Infection 2020 May; 26:553–62. DOI: `10.1016/j.cmi.2019.09.020`. Available from: `https://www.sciencedirect.com/science/article/pii/S1198743X19305051`

32. Temperature and precipitation gridded data for global and regional domains derived from in-situ and satellite observations. Available from: `https://cds.climate.copernicus.eu/datasets/insitu-gridded-observations-global-and-regional?tab=overview` [Accessed on: 2025 Apr 6]

33. Browne AJ, Chipeta MG, Haines-Woodhouse G, Kumaran EPA, Hamadani BHK, Zaraa S, Henry NJ, Deshpande A, Reiner RC, Day NPJ, Lopez AD, Dunachie S, Moore CE, Stergachis A, Hay SI, and Dolecek C. Global antibiotic consumption and usage in humans, 2000–18: a spatial modelling study. The Lancet Planetary Health 2021 Dec; 5:e893–e904. DOI: `10.1016/S2542-5196(21)00280-1`. Available from: `https://www.sciencedirect.com/science/article/pii/S2542519621002801`

34. Ammerlaan HSM, Harbarth S, Buiting AGM, Crook DW, Fitzpatrick F, Hanberger H, Herwaldt LA, Keulen PHJ van, Kluytmans JAJW, Kola A, Kuchenbecker RS, Lingaas E, Meessen N, Morris-Downes MM, Pottinger JM, Rohner P, Santos RP dos, Seifert H, Wisplinghoff H, Ziesing S, Walker AS, and Bonten MJM. Secular Trends in Nosocomial Bloodstream Infections: Antibiotic-Resistant Bacteria Increase the Total Burden of Infection. Clinical Infectious Diseases 2013 Mar; 56:798–805. DOI: `10.1093/cid/cis1006`. Available from: `https://doi.org/10.1093/cid/cis1006`

35. McElreath R. Statistical Rethinking: A Bayesian Course with Examples in R and STAN. 2nd ed. New York: Chapman and Hall/CRC, 2020 Mar. DOI: `10.1201/9780429029608`

36. Stewart PS. An occupancy model with a Gaussian process for spatial autocorrelation in Stan. 2022 May. Available from: `https://peter-stewart.github.io/blog/gaussian-process-occupancy-tutorial/` [Accessed on: 2025 Jul 23]

37. Gaussian Processes. Available from: https://mc-stan.org/docs/stan-users-guide/gaussian-processes.html [Accessed on: 2025 Nov 19]

38. Wang J. An Intuitive Tutorial to Gaussian Process Regression. Computing in Science & Engineering 2023 Jul; 25. arXiv:2009.10862 [stat]:4–11. DOI: 10.1109/MCSE.2023.3342149. Available from: http://arxiv.org/abs/2009.10862

39. Vehtari A, Gabry J, Magnusson M, Yao Y, Bürkner PC, Paananen T, and Gelman A. loo: Efficient Leave-One-Out Cross-Validation and WAIC for Bayesian Models. 2015 Jun. DOI: 10.32614/CRAN.package.loo. Available from: https://CRAN.R-project.org/package=loo [Accessed on: 2025 Nov 19]

40. Model averaging/weighting via stacking or pseudo-BMA weighting — loo_model_weights. Available from: https://mc-stan.org/loo/reference/loo_model_weights.html [Accessed on: 2025 Nov 19]

41. Posterior and Prior Predictive Checks. Available from: https://mc-stan.org/docs/stan-users-guide/posterior-predictive-checks.html [Accessed on: 2025 Nov 19]

42. Efficient Leave-One-Out Cross-Validation and WAIC for Bayesian Models. Available from: https://mc-stan.org/loo/ [Accessed on: 2026 Mar 7]

43. Letten AD, Hall AR, and Levine JM. Using ecological coexistence theory to understand antibiotic resistance and microbial competition. Nature Ecology & Evolution 2021 Apr; 5:431–41. DOI: 10.1038/s41559-020-01385-w. Available from: https://www.nature.com/articles/s41559-020-01385-w

44. Shigemoto K. Various Logistic Curves in SIS and SIR Models. European Journal of Mathematics and Statistics 2023 Jan; 4. arXiv:2211.04186 [physics]:1–6. DOI: 10.24018/ejmath.2023.4.1.185. Available from: http://arxiv.org/abs/2211.04186

45. Bacaër N. Verhulst and the logistic equation (1838). *A Short History of Mathematical Population Dynamics*. Ed. by Bacaër N. London: Springer, 2011 :35–9. DOI: 10.1007/978-0-85729-115-8_6. Available from: https://doi.org/10.1007/978-0-85729-115-8_6 [Accessed on: 2025 Apr 6]

46. Besag J. Spatial Interaction and the Statistical Analysis of Lattice Systems. Journal of the Royal Statistical Society: Series B (Methodological) 1974; 36. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.2517-6161.1974.tb00999.x:192–225. DOI: 10.1111/j.2517-6161.1974.tb00999.x. Available from: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1974.tb00999.x

47. Besag J, York J, and Mollié A. Bayesian image restoration, with two applications in spatial statistics. Annals of the Institute of Statistical Mathematics 1991 Mar; 43:1–20. DOI: 10.1007/BF00116466. Available from: https://doi.org/10.1007/BF00116466

48. Riebler A, Sørbye SH, Simpson D, and Rue H. An intuitive Bayesian spatial model for disease mapping that accounts for scaling. Statistical Methods in Medical Research 2016 Aug; 25:1145–65. DOI: 10.1177/0962280216660421. Available from: https://doi.org/10.1177/0962280216660421

49. Morris M, Wheeler-Martin K, Simpson D, Mooney SJ, Gelman A, and DiMaggio C. Bayesian hierarchical spatial models: Implementing the Besag York Mollié model in stan. Spatial and Spatio-temporal Epidemiology 2019 Nov; 31:100301. DOI: 10.1016/j.sste.2019.100301. Available from: https://www.sciencedirect.com/science/article/pii/S1877584518301175

50. Spatial Models in Stan: Intrinsic Auto-Regressive Models for Areal Data. Available from: https://mc-stan.org/learn-stan/case-studies/icar_stan.html [Accessed on: 2025 May 5]

Supporting Information

S1 Text. Supporting Model Outputs

Combined file of supporting text and figures.

A   Data Summary

Figures to summarise data.

Fig. 3. Total number of isolates for global *Acinetobacter baumannii* data. The total number of isolates for 113 countries from 1991 to 2022 are shown across 7 antibiotic categories.

Fig. 4. Total number of isolates for *Acinetobacter baumannii* data across Asia. The total number of isolates for 32 countries in Asia from 2000 to 2022 are shown across 7 antibiotic categories.

Fig. 5. Total number of isolates for *Acinetobacter baumannii* data across Asia per year. The total number of isolates for 32 countries in Asia from 2000 to 2022 are shown across 7 antibiotic categories.

B   Model Derivation

A two-compartment ODE model can be used to represent the general dynamics of colonisation with drug-resistant strains. This framework assumes that colonisation with drug-sensitive strains remains approximately constant over time, and so drug-sensitive colonisation is not explicitly modelled. The differential equations describing the dynamics of this two-compartment model are shown in Equation 13 and Equation 14. Here, the X represents uncolonised individuals susceptible to infection and I represents individuals currently infectious.

$$\frac{dX}{dt} = \frac{-\beta}{N}XI + \gamma I \tag{13}$$

$$\frac{dI}{dt} = \frac{\beta}{N}XI - \gamma I \tag{14}$$

We assume a constancy of the population N to obtain a proportional value of the total population colonised with drug-resistant strains and can rewrite Equation 14 in terms of I, β, γ under this assumption.

$$\frac{dX}{dt} + \frac{dI}{dt} = 0 \tag{15}$$

$$\frac{dI}{dt} = \frac{\beta}{N}(N - I)I - \gamma I \tag{16}$$

$$= \beta(1 - \frac{I}{N})I - \gamma I \tag{17}$$

$$= \beta I - \frac{I^2}{N}\beta - \gamma I \tag{18}$$

$$= (\beta - \gamma)I - \frac{I^2}{N}\beta \tag{19}$$

An analytical solution to the two-compartment model is the Verhulst equation, also known as the logistic growth function (see Equation 22). We use the logistic growth function as an analytical solution to this two-compartment model [44]. This equation can be expressed in terms of the initial proportion of resistance P_0, a maximum limiting value for the population K, and the growth rate r (see in Equation 21) [45].

$$\frac{dP}{dt} = rP(1 - \frac{P}{K}) \tag{20}$$

$$= rP - \frac{r}{K}P^2 \tag{21}$$

$$P(t) = \frac{K}{1 + (\frac{K - P_0}{P_0})e^{-rt}} \tag{22}$$

We restructure Equation 19 in terms of Equation 21 to find the growth rate r and carrying capacity K in terms of β and γ.

$$r = \beta - \gamma \tag{23}$$

$$K = \frac{\beta - \gamma}{\beta}N \tag{24}$$

This allows us to express Equation 22 in terms of β and γ, as shown in Equation 25.

$$P(t) = \frac{1 - \frac{\gamma}{\beta}}{1 + (\frac{1 - \frac{\gamma}{\beta}}{P_0} - 1)e^{-(\beta - \gamma)t}} \tag{25}$$

618   Although the ODE system is defined when $\beta = \gamma$, a limitation in our closed form solution describing the

619 proportion of the population colonised with resistant strains (Equation 4) is that β and γ are not equal since

620 this equation would be undefined.

621   When looking at the ODE system, when $\beta = \gamma$, we find that,

$$P(t) = \frac{N \cdot P_0}{N + \beta \cdot P_0 \cdot t} \tag{26}$$

which is a different functional form from Equation 4 when $\beta = \gamma$. While a conditional branch could switch

between the solutions in Stan, conditioning on unknown parameters might cause a gradient discontinuity

that could compromise HMC sampling. In practice, this limitation is unlikely to affect our results as the

singularity is smoothly approached from either side, and when dealing with floating-point numbers the

probability of drawing a value of beta exactly equal to gamma is quite low.

The solution for $\beta \neq \gamma$

$$P_{\beta \neq \gamma}(t) = \frac{1 - \dfrac{\gamma}{\beta}}{1 + \left(\dfrac{1 - \frac{\gamma}{\beta}}{P_0} - 1 \right) e^{-(\beta - \gamma)t}}, \tag{27}$$

628 converges to the solution for when $\beta = \gamma$,

$$P_{\beta = \gamma}(t) = \frac{P_0}{1 + \beta P_0 t} \tag{28}$$

629   We evaluate the limit of the curve $P_{\beta \neq \gamma}(t)$ as $t * (\beta - \gamma) \rightarrow 0$. Using the Taylor expansion $e^{-\epsilon} =$

630 $1 - \epsilon + O(\epsilon^2)$, and noting that $O(\epsilon^2)$ goes to 0 faster as $\epsilon \rightarrow 0$, we substitute $e^{-(\beta - \gamma)t} \approx 1 - (\beta - \gamma)t$ to get:

$$P_{\beta \neq \gamma}(t) \approx \frac{1 - \dfrac{\gamma}{\beta}}{1 + \left(\dfrac{1 - \frac{\gamma}{\beta}}{P_0} - 1 \right)(1 - t * (\beta - \gamma))} \tag{29}$$

631   Expanding the denominator gives:

$$= \frac{1 - \dfrac{\gamma}{\beta}}{\dfrac{1 - \frac{\gamma}{\beta}}{P_0} - \dfrac{t * (\beta - \gamma)(1 - \frac{\gamma}{\beta})}{P_0} + (\beta - \gamma) * t} \tag{30}$$

632   Dividing both numerator and denominator by $(1 - \dfrac{\gamma}{\beta})$ gives:

$$= \frac{1}{\dfrac{1}{P_0} - \dfrac{t * (\beta - \gamma)}{P_0} + \dfrac{(\beta - \gamma) * t}{1 - \frac{\gamma}{\beta}}}. \tag{31}$$

Multiplying numerator and denominator by P_0:

$$P_{\beta \neq \gamma}(t) = \frac{P_0}{1 - (\beta - \gamma) t + \dfrac{P_0 (\beta - \gamma) t}{1 - \frac{\gamma}{\beta}}} \tag{32}$$

Substituting $1 - \dfrac{\gamma}{\beta} = \dfrac{\beta - \gamma}{\beta}$ into the last term of the denominator:

$$= \frac{P_0}{1 - (\beta - \gamma) t + \beta P_0 t}. \tag{33}$$

Taking the limit $(\beta - \gamma) t \to 0$, we get:

$$\lim_{(\beta - \gamma) t \to 0} P_{\beta \neq \gamma}(t) = \frac{P_0}{1 + \beta P_0 t} = P_{\beta = \gamma}(t) \tag{34}$$

C   Covariate Models

For the models with covariates, the β prior is set to explore the logarithmic space while the priors for P_0 and γ remain the same as the country-level effects model.

In Equation 35, θ_k is the coefficient of the covariate k and is set to have a weakly informative normal prior distribution. The covariate values are then multiplied by their estimated coefficient value, summed over the total number of covariates, and then added to their estimated country-level β_c term. The final value is then exponentially transformed.

$$\theta_k \sim \text{Normal}(0, 1) \tag{35}$$

$$log_\beta \sim \text{Normal}(0, 1) \tag{36}$$

$$\sigma_{log_\beta} \sim \text{Exponential}(0.5) \tag{37}$$

$$log_{\beta_c} \sim \text{Normal}(log_\beta, \sigma_{log_\beta}) \tag{38}$$

$$log(\beta_{c,t}) = \beta_c + \sum_{k=1}^{K} \theta_k \cdot X_{k,c,t} \tag{39}$$

D Spatial Effects

To incorporate spatial effects into the model, an initial approach was taken using the Besag-York-Mollié (BYM) model and extract the intrinsic conditional autoregressive component to smooth estimates based on neighbouring countries and to account for spatial interactions based on pairs of countries [46, 47, 48]. We incorporate these spatial models following the implementation of BYM model in Stan [49, 50]. We add in parameter ϕ to account to encode the neighbour relationships in terms of the pairwise difference formula as shown in Equation 40. The pairwise difference formula is calculated between i and j if they are neighbours and under the condition that i cannot equal j. A soft sum-to-zero constraint is also applied to ϕ to soft-centers the mean close to zero and is shown in Equation 41 [49]. A precision term, τ_{ϕ}, is added to estimate the standard deviation of ϕ, as shown in Equation 43. The spatial component, $\sigma_{\phi} * \phi_{c,i}$, is added related to I_0 and β as we expect neighbouring regions to have similar initial resistance rates and influence on transmission. When testing models with spatial effects, we only include source and spatial effects as we now relate regions to their neighbours.

$$p(\phi) \propto \exp\left(-\frac{1}{2} \sum_{i \sim j} (\phi_i - \phi_j)^2\right) \tag{40}$$

$$\sum(\phi) \sim \mathcal{N}(0, 0.001 \cdot C) \tag{41}$$

$$\tau_{\phi} \sim \text{Gamma}(1, 0.1) \tag{42}$$

$$\sigma_{\phi} = \text{inv}(\sqrt{\tau_{\phi}}) \tag{43}$$

$$I_{0,s\_spatial,i} = \text{logit}^{-1}(I_{0\_logit,s,i} + \sigma_{\phi} * \phi_{c,i}) \tag{44}$$

$$\beta_{s\_spatial,i} = \beta_{s,i} + \sigma_{\phi} * \phi_{c,i} + \sum_{j=1}^{n} \theta_j \cdot X_{j,i,c} \tag{45}$$

$$\gamma_{s\_spatial,i} = \gamma_{s,i} \tag{46}$$

After testing spatial effects in terms of the BYM model, an alternate method was explored in terms of Gaussian process regression to include spatial autocorrelation to reduce model run times and improve model convergence [35, 36].

To create a relationship between sites that are closer together, a kernel function can be used to define the covariance between countries based on distance. The radial basis kernel functions is shown as Equation 47 where D is the distance between countries, ζ^2 is the maximum covariance between sites, and ρ^2 is the rate that covariance declines. This relationship is then used to estimate the a spatial intercept term for each country following methods from previous examples [35, 36].

$$K_{i,j} = \zeta^2 * exp(-\rho^2 * D_{i,j}^2) \tag{47}$$

We incorporate the spatial effects intercept term, k, for P_0, β, and γ. For P_0, the spatial term is added to the population-level term and transformed to the inverse logit space as shown in Equation 48. For both *beta* and *gamma*, we add the spatial term in the log space and then exponentiated the sum of the spatial term and the population level parameter. The final β and γ terms used in the final probability equation is shown in Equations 49 and 50.

$$P_{0,c} = \text{logit}^{-1}(P_{0\_logit} + k_{P_0,c}) \tag{48}$$

$$\beta_c = exp(log_\beta + k_{\beta,c}) \tag{49}$$

$$\gamma_c = exp(log_\gamma + k_{\gamma,c}) \tag{50}$$

E   Country-level Priors

The prior distributions are shown. Here, we have two options we explored for the initial resistance P_0 prior. Option 1: We select a weakly informative normal prior distribution for P_{0_logit} since we do not have specific information on this parameter on a logit scale and having a hierarchical structure. Option 2: We select a beta(2,2) prior distribution for the initial resistance value for each country with no hierarchical structure. The

first option was preferred for fitting models starting at the initial time point for each country. The second option was preferred for fitting models all from the same year and starting at previous years where data is sparse since it is computationally more efficient. For this analysis, we use the second option for the initial resistance prior as we are generating predictions from 2000 onwards and data is sparse. For β and γ we select an exponential prior distribution with a value of three since the value for these parameters have a higher probability density closer to 0 and should be positive. $\sigma_{P_{0\_\text{logit}}}$, σ_β, and σ_γ all have an exponential prior distribution with a value of 0.5 as we expect this value to be positive and expect to have a slightly higher probability density closer to 0 since we do not expect major differences between countries for these parameters.

Prior distributions for the hierarchical model:

$$\text{Option 1 for } P_0 \text{ (logit):} \tag{51}$$

$$P_{0\_\text{logit}} \sim \text{Normal}(0, 1) \tag{52}$$

$$\sigma_{P_{0\_\text{logit}}} \sim \text{Exponential}(0.5) \tag{53}$$

$$P_{0\_\text{logit},c} \sim \text{Normal}(P_{0\_\text{logit}}, \sigma_{P_{0\_\text{logit}}}) \tag{54}$$

$$\tag{55}$$

$$\text{Option 2 for } P_0: \tag{56}$$

$$ProbP_{0,c} \sim \text{beta}(2, 2) \tag{57}$$

$$\tag{58}$$

$$\beta \sim \text{Exponential}(3) \tag{59}$$

$$\sigma_\beta \sim \text{Exponential}(0.5) \tag{60}$$

$$\beta_c \sim \text{Normal}(\beta, \sigma_\beta) \tag{61}$$

$$\tag{62}$$

$$\gamma \sim \text{Exponential}(3) \tag{63}$$

$$\sigma_\gamma \sim \text{Exponential}(0.5) \tag{64}$$

$$\gamma_c \sim \text{Normal}(\gamma, \sigma_\gamma) \tag{65}$$

After defining these priors, we can define the parameter values used as inputs into Equation **??**

$$P_{0,c} = \text{logit}^{-1}(P_{0\_\text{logit},c}) \text{ or } ProbP_{0,c} \tag{66}$$

$$\beta_c = \beta_c \tag{67}$$

$$\gamma_c = \gamma_c \tag{68}$$

F Data source-level Scaling

The prior distributions for the source-level parameters are shown. The models incorporating source-level scaling extend the country-level model and use the same priors as shown in the previous section for β_c, γ_c, and $P_{0,c}$. Here, the source-dependent multiplicative factor η_s is set to be bounded between 0 and 1 to ensure positive constraints and with a beta distribution density concentrated near 1. The structure of this model shows that the underlying curve for each country should be similar and the variation between sources should be relatively small when scaling the underlying curve.

$$\eta_s \sim \text{beta}(5, 1) \tag{69}$$

$$\tag{70}$$

G Resistance Trajectories

Resistance trajectories for each country across all pathogen-antibiotic combinations for Asia from 2000 to 2022.

H Long-term Saturation Levels

Long-term saturation levels estimated for each country for all pathogen-antibiotic combinations for Asia from 2000 to 2022.

I Covariate Model Predictions

The estimated covariate coefficients values are shown for models that incorporate covariates. These models include country-level random effects and covariates (C+X), and the spatial model with Gaussian process regression, source-level scaling, and covariates (SP+S+X). The covariates included are mean temperature,

Fig. 6. Individual model fits using $\beta\ \gamma$, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to aminoglycosides model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to aminoglycosides for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

(a)

(b)

(c)

(d)

(e)

Fig. 7. Individual model fits using β γ, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to antipseudomonal penicillins model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to antipseudomonal penicillins for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

(a)

(b)

(c)

(d)

(e)

Fig. 8. Individual model fits using $\beta\ \gamma$, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to beta-lactams/beta-lactamase inhibitor combinations model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to beta-lactams/beta-lactamase inhibitor combinations for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

(a)

(b)

(c)

(d)

(e)

Fig. 9. Individual model fits using β γ, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to carbapenems model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to carbapenems for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

Fig. 10. Individual model fits using β γ, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to fluoroquinolones model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to fluoroquinolones for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

(a)

(b)

(c)

(d)

(e)

Fig. 11. Individual model fits using β γ, P_0 parameterisation of the logistic growth function for resistance of *Acinetobacter baumannii* to fourth-generation cephalosporins model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to fourth-generation cephalosporins for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

Fig. 12. Individual model fits using β γ, P_0 parameterisation of the logistic growth function for resistance of _Acinetobacter baumannii_ to third-generation cephalosporins model training for Asia. a) to e) shows the posterior predictions for the proportion of clinical isolates that are resistant to third-generation cephalosporins for all countries and six model variants. Model C: Model with country-level random effects, C+X: Country-level random effects and covariates, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling, SP+S+X: Spatial model with Gaussian process regression, source-level scaling, and covariates. Solid lines represent the mean posterior prediction and the shaded regions show the 95% credible intervals with the different colours corresponding to each model type. The points show the observed resistance data used for training the models.

Fig. 13. Long-term saturation resistance levels of *Acinetobacter baumannii* to antibiotic combinations for models trained for Asia. a) to g) shows stabilisation values for aminoglycosides, antipseudomonal penicillins, beta-lactams/beta-lactamase inhibitor combinations, carbapenems, fluoroquinolones, fourth-generation cephalosporins, and third-generation cephalosporins, respectively. Points show posterior mean estimates with 95% credible intervals. Model C: Model with country-level random effects, C+S: Country-level random effects and source-level scaling, SP: Spatial model with Gaussian process regression, SP+S: Spatial model with Gaussian process regression and source-level scaling.

702 antibiotic consumption in terms of defined daily dose per 1000 inhabitant per day (DDD/1000), and out-

703 of-pocket healthcare expenditures reported per year. Alternative covariates can be included in the model

704 structure for extensions of this framework.

Antibiotic Class/Subclass	Average Covariate Coefficient Estimated [95% CrI]					
	Model C+X			Model SP+S+X		
	Temperature	DDD/1000	OOP_hexp	Temperature	DDD/1000	OOP_hexp
Aminoglycosides	0.143 [-0.031, 0.309]	0.210 [0.107, 0.322]	0.009 [-0.117, 0.132]	-0.074 [-0.204, 0.127]	0.251 [0.165, 0.337]	-0.028 [-0.133, 0.077]
Antipseudomonal penicillins	-0.082 [-0.224, 0.055]	0.365 [0.284, 0.446]	0.013 [-0.073, 0.095]	-0.010 [-0.159, 0.139]	0.484 [0.391, 0.577]	0.060 [-0.040, 0.159]
Beta lactam/beta-lactamase inhibitor combinations	-0.165 [-0.291, -0.040]	0.057 [-0.088, 0.214]	0.072 [-0.050, 0.190]	-0.147 [-0.267, -0.028]	0.245 [0.131, 0.358]	0.125 [0.012, 0.234]
Carbapenems	-0.199 [-0.300, -0.106]	0.330 [0.249, 0.403]	0.137 [-0.001, 0.273]	-0.192 [-0.287, -0.099]	0.211 [0.112, 0.322]	0.179 [0.062, 0.282]
Fluoroquinolones	-0.099 [-0.242, 0.042]	0.511 [0.443, 0.581]	0.186 [0.099, 0.274]	-0.044 [-0.195, 0.113]	0.187 [0.094, 0.284]	0.046 [-0.065, 0.156]
Fourth-generation cephalosporins	-0.128 [-0.210, -0.048]	0.123 [0.020, 0.215]	0.170 [0.067, 0.272]	-0.153 [-0.248, -0.059]	0.170 [0.071, 0.268]	0.189 [0.076, 0.298]
Third-generation cephalosporins	-0.412 [-0.504, -0.326]	0.142 [0.073, 0.214]	0.011 [-0.075, 0.107]	-0.353 [-0.442, -0.263]	0.254 [0.166, 0.342]	0.056 [-0.055, 0.177]

Table 6. Average coefficient value of covariates based on antibiotic class for mean temperature, antibiotic consumption in terms of defined daily dose per 1000 inhabitant per day (DDD/1000), and the out-of-pocket healthcare expenditures reported per year. The models incorporating covariates are country-level random effects and covariates (C+X), and the spatial model with Gaussian process regression, source-level scaling, and covariates (SP+S+X).

J Posteriors Densities from Simulated Data Model Fits

The models are fitted using simulated data with known model parameters to confirm parameter estimates are recoverable. The parameter posterior densities resulting from fitting simulated data with the country-level random effects model with a β, γ, I_0 (initial proportion colonised with drug-resistant strains P_0 in our model) parameterisation are shown. The resulting posterior densities are shown with the corresponding true parameter value used to generate data.

K Model Trace and Pair Plots

Model convergence was checked through trace and pair plots generated after fitting models to *Acinetobacter baumannii* resistance data for each pathogen-antibiotic combination. Example pair and trace plots for carbapenems are shown.

Fig. 14. Posterior densities from data simulation using β γ parameterisation of the logistic growth function a) shows the population level β and γ and b) shows the country-specific posterior densities recovered using the true parameter values shown by the dashed red line.

Fig. 15. Pairs plot of β **and** γ **the using** β γ**,** P_0 **parameterisation of the logistic growth function for resistance of** *Acinetobacter baumannii* **and carbapenems model trained for Asia from 2000 to 2022.** a) Shows pairs plots of country-level random effects b) pairs plot for base spatial-effects model

Fig. 16. Trace plots of β **and** γ **the using** β γ**,** P_0 **parameterisation of the logistic growth function for resistance of** *Acinetobacter baumannii* **and carbapenems model trained for Asia from 2000 to 2022.** a) Shows trace plots of the country-level random effects model b) trace plot for the spatial-effects model