Title: Proof of concept of meniscus reconstruction and tear simulation via subtractive segmentation and bi-planar MRI integration

Liqi Ng[1,2 #], Peng Zhang[1#], Maojiang Lyu[1,3 #], Xin Mu[1#], Jianle Liang[4], Peng Chen[1], Tian You[1], Lu Bai[1], Wanqi Ding[5], Yu Zhou[2,9], Mingjing Zhang[2], Swastina Nath Varma[2], Yuluan Wu[7], Weiji Wang[1], Yangchidung Lin[6], Xiaofeng Zhang[3], Yongcan Huang[8], Chaozong Liu[2], Yi Sun[1 *], Xintao Zhang[1*]

[1] Dept of Sports Medicine, Peking University Shenzhen Hospital, Shenzhen, China;

[2] Institute of Orthopaedic and Musculoskeletal Science, Royal National Orthopaedic Hospital, University College London, London, UK;

[3] Orthopaedics Department, Shanghai Sixth People's Hospital of Jinshan Branch, Shanghai, China;

[4] Dept of Radiology, Peking University Shenzhen Hospital, Shenzhen, China;

[5] College of Life Sciences and Oceanography, Shenzhen University, Shenzhen, China;

[6] Warshel Institute for Computational Biology, School of Medicine, The Chinese University of Hong Kong, Shenzhen, China;

[7] Dept of Bone and Joint Surgery, Peking University Shenzhen Hospital, Shenzhen, China;

[8] Dept of Spine Surgery, Peking University Shenzhen Hospital, Shenzhen, China;

[9] Postdoctoral Research Workstation, Orthopedic Hospital, Chongqing University of Chinese Medicine, Chongqing Orthopedic Hospital of Traditional Chinese Medicine, Chongqing, China;

Authors' info and contribution

Liqi Ng: liqi.ng.17@ucl.ac.uk; ORCID: 0000-0002-7049-0012; Data analysis, figure preparation, and manuscript writing

Peng Zhang: Drzhangpengvip@163.com; ORCID: 0009-0004-1367-3228; Data collection

Maojiang Lyu: doclvmaojiang@163.com; ORCID: 0009-0005-3035-6731; Data collection

Xin Mu: xin＿mu@sina.com; ORCID: 0009-0007-2109-4906; Data collection

Jianle Liang: 124860517@qq.com; ORCID: 0009-0004-8251-5688; Data collection

Peng Chen: doc＿chenpeng@163.com; ORCID: 0000-0002-0807-7809; Data collection and manuscript preparation

Tian You: 17914@163.com; ORCID: 0000-0001-7027-7230; Manuscript preparation

Lu Bai: boowboow@163.com; ORCID: 0000-0003-4360-9601; Manuscript preparation

Wanqi Ding: d17875662878@163.com; ORCID: 0009-0007-7276-9416; Manuscript

preparation

Yu Zhou: zyhenry123@163.com; ORCID: 0000-0002-3065-4625; Data analysis

Mingjing Zhang: mingjing.zhang.22@ucl.ac.uk; ORCID: 0009-0003-9017-3940; Data

analysis

Swastina Nath Varma: t.varma@ucl.ac.uk; ORCID: 0000-0002-9449-8772; Data analysis

Yuluan Wu: 13501476877@163.com; ORCID: 0009-0005-7735-7264; Review manuscript

Weiji Wang: vikey3638@163.com; ORCID: 0009-0000-4886-7496; Review manuscript

Yangchidung Lin: yangchidung@cuhk.edu.cn; ORCID: 0000-0003-2436-4602; Review

manuscript

Xiaofeng Zhang: zhangxiaofeng322@163.com; ORCID: 0009-0008-6931-5082; Review

manuscript

Yongcan Huang: y.c.huang@connect.hku.hk; ORCID: 0000-0001-8548-8233; Review

manuscript

Chaozong Liu: chaozong.liu@ucl.ac.uk; ORCID: 0000-0002-9854-4043; Review manuscript

Yi Sun: ericsunhku@outlook.com; ORCID: 0000-0002-9728-2205; Conception and design,

data acquisition and interpretation, drafting and final approval of the manuscript, and

obtaining the funds.

Xintao Zhang: zhangxintao@sina.com; ORCID: 0000-0001-9926-3756; Conception and

design, coordinating the collaboration, drafting and final approval of the manuscript, and

obtaining the funds.

Abstract (150 words)

Objective: To develop and validate a high-fidelity magnetic resonance imaging (MRI)-based

three-dimensional (3D) reconstruction of the meniscus for improved tears characterization

and quantification.

Methods: A subtractive segmentation algorithm for clinical MRI with bi-planar (coronal and

sagittal) model integration was employed to reconstruct 3D meniscal models from patients

with various tear patterns. Geometric accuracy was validated against physical measurements

and histological findings (Safranin-O/Fast green staining) with surgical specimens from

arthritic patients. Tear morphology, surgical outcome simulation, and longitudinal healing

efficiency were quantitatively assessed.

Results: The reconstruction models of menisci from osteoarthritis patients demonstrated high

morphometric fidelity, accurately replicating native meniscal dimensions (e.g., length, width,

rim height) and revealing subtle internal deformities resembling lipid or myeloid deposits.

The model enabled 3D visualization and quantification of tear patterns, including discoid,

horizontal, longitudinal, flap, and complex tears, and captured topographic changes such as

erosions in gout. Premorbid architecture of a discoid medial meniscus with complex tears was

virtually rebuilt, facilitating the distinction of reparable regions from zones requiring

meniscectomy. Two-year follow-up reconstructions revealed progressive reduction in tear

diastasis and surface-initiated healing after suture repair.

Conclusion: We demonstrate for the first time that MRI-based meniscal reconstruction

improves the detection, characterization, and management of meniscal tears, providing

detailed morphometric and intrasubstance measurements beyond conventional imaging and

arthroscopy. This approach supports the prediction of surgical outcomes and postoperative

monitoring, underscoring its substantial potential for personalized management of meniscal

injuries and osteoarthritis.

Keywords: Meniscal tears; Three-dimensional reconstruction; Magnetic resonance imaging;

Subtractive segmentation; Bi-planar integration

Introduction

The meniscus is a crescent-shaped fibrocartilaginous tissue situated between the femoral and tibial condyles within the knee joint capsule. It serves crucial functions in absorbing compressive and shear forces during movement, particularly in high-impact athletic activities, thereby contributing to cartilage lubrication and preserving joint mobility [1]. Meniscal injury is a common orthopedic condition, affecting approximately 0.6% of the general population, with prevalence exceeding 50% in patients with anterior cruciate ligament (ACL) tears [2,3] and 75% in those with osteoarthritis (OA)[4]. Tears disrupt meniscus architecture, resulting in nerve infiltration and cartilage erosion, and ultimately leading to the development of arthritis and deprivation of sports capacity [5]. Treatment options[6] (e.g., suturing or meniscectomy) and healing outcomes[7] heavily depend on the tear geometry, including its location, depth, and rim width, as well as the quality of the surrounding tissue. Therefore, precise visualization and simulation of meniscus tears are crucial for accurate diagnosis and effective surgical planning.

Compared to high-resolution intraoperative arthroscopy, two-dimensional (2D) magnetic resonance imaging (MRI) offers a non-invasive cross-sectional evaluation of meniscus tears, thereby minimizing the risks of infection, surgical uncertainties, and iatrogenic damage[8-10]. However, complex tear configurations, such as radial, horizontal, or bucket-handle tears, are often underestimated due to their limited through-plane spatial resolution [11]. 2D MRI demonstrates lower accuracy and poor agreement with arthroscopy in detecting ramp lesions at the medial menisco-synovial junction[12], particularly in the context of concomitant ACL injuries [11], despite reports suggesting MRI may achieve over 85% diagnostic specificity after optimization [13]. Furthermore, over 65% of lateral meniscus posterior horn tears are missed on ordinary MRIs [14]. Interobserver reliability remains suboptimal due to experiences, cognitive fatigue, and time pressures [15], even among seasoned clinicians [16]. Besides, the transplantation of size-mismatched allografts disrupts contact mechanics that predispose to secondary injury [17]. Three-dimensional (3D) MRI with signal-to-noise ratio enhancement [18], though demonstrating superior sensitivity in detecting ligament and meniscal tears, remains substantially underutilized in the clinic. Thus, meniscus reconstruction derived from 2D MRI is strongly desired for accurate tear evaluation and the fabrication of anatomically shaped implants [19]. Meniscus engineering created with various scaffolds significantly increases the demand for high-index volumetric models with sufficient topographic details [20].

3D reconstruction based on clinical MRI enables a reliable and unbiased evaluation of

pathological changes; for instance, 3D vascular volumetric reconstruction facilitates

hepatobiliary surgical planning [21]. Kruger et al. and Li et al. manually extracted the meniscal

signals from sagittal plane images for automated segmentation and 3D modeling [22], yet

interference from neighboring structures was considerable, resulting in an incomplete

representation of the architecture. Generating a new algorithmic logic to pinpoint the

meniscus compartment is thus demanded. Moreover, adding coronal or axial MRI imaging

planes may reduce the inter-slice signal loss and capture abundant geometric details, although

this remains unexplored. Overall, the purpose of this study was to develop a concrete MRI-

based meniscus simulation and evaluate its performance in tear detection across several types

and repair assessment via a two-year longitudinal follow-up case study.

Materials & Methods

MRI image acquisition

The MRI images were collected from 32 patients under ethics approval from Peking

University Shenzhen Hospital (PUSZH R2022-170-1). Patients suffered from discomfort or

disability related to the knee joint and received meniscus suture or segment removal

surgeries. Cases where the patients' pre-operative X-rays indicated open epiphysis or post-

operative MRI re-evaluation was missing, were not involved, and thus, 10 cases (tear

simulation: 9; follow-up study: 1) were presented in this study (**Supplementary Table 1**). All

data were collected prospectively between 2020 and 2024.

All MRI scans were conducted on a stretched knee with 3.0 T spectra (Siemens Healthineers,

Erlangen, Germany) using the standard protocols. MRI scans were obtained with multiple

series at axial, sagittal, and coronal planes, including T1-weighted images with a short

repetition time (TR) and time to echo (TE), and proton density (PD) weighted images with

long TR and short TE sequence. In brief, sequence parameters per plane were: repetition time

3050 ms, echo time 42 ms, flip angle 150°, pixel bandwidth 170 Hz, matrix 320 × 320, 30

slices, Field of View (FOV) of 16cm × 16cm, slice thickness 3 mm, and interslice gap of

3mm. Proton density-weighted images with turbo-spin-echo (TSE) were selected for the

reconstruction. The skeletal age was determined according to a completely closed growth

plate of the femoral and tibial knee epiphyses. Experienced musculoskeletal radiologists

assessed meniscus quality, and two orthopaedic surgeons evaluated tears or extrusion via a

'bow-tie configuration' located between the knee joint[23].

MRI image process and 3D reconstruction

The MRI scan data were exported and saved as Digital Imaging and Communications in

Medicine (DICOM) format, and further inserted into Materialise Interactive Medica Control

System (Mimics) 21.0 software (Materialise, Leuven, Belgium) for next-step analysis. Via

thresholding, two separate masks were created for semi-automatic subtractive processing: the

first mask (M_1) encompassed all tissues and spacing in the overall image, while the second

mask (M_2) highlighted denser tissues (e.g., knee joint bones and tissues surrounding the

meniscus) (**Fig. 1**). After creating both masks, the denser mask was subtracted from the

overall mask, isolating the meniscus ($M_{cropped}$) with clear distinction from adjacent

compartments while retaining minor residual noise, which was subsequently removed

manually (M_{clean}). 3D reconstruction was performed in 3-matic Research 13.0 (Materialise,

Leuven, Belgium) for both coronal and sagittal planes, with minimized wrapping and

smoothing. These planes were merged via bi-planar integration to achieve high-resolution

complementary modelling.

Algorithm

(1) Computation of meniscal datasets

The meniscal segmentation is obtained through the following computational procedure:

$$M_{meniscus(x,y,z)}$$

$$= \{[(\Theta[I(x,y,z); I_{min}, I_{max}] \cdot \Gamma[I(x,y,z); I_{min} + I_\Delta, I_{max}])$$

$$\times \Omega(x,y,z; X,Y,Z)] - \Psi(x,y,z)\}^{\delta(x,y,z)}$$

Here, $M_{meniscus(x, y, z)}$ denotes the definitive binarized segmentation output at spatial coordinates

(x, y, z). The operators Θ and Γ implement dual-thresholding based on intensity bounds [I_{min},

I_{max}], incorporating adaptive intensity offset (I_Δ). The function $\Omega_{(x, y, z; X, Y, Z)}$ formalizes a

region of interest (ROI) via a bounding box with coordinates (X, Y, Z). $\Psi_{(x, y, z)}$ designates the

expert-curated exclusion mask for noise reduction. $\delta_{(x, y, z)}$ enables a voxel-wise confidence

measure for precision-weighted validation.

(2) Coregistration conducted with a rigid transformation:

$$V_{aligned} = R \cdot V_{original} + t$$

$V_{aligned}$ constitutes the transformed vertex coordinates. R is the rotation matrix, t is the

translation vector, and $V_{original}$ represents the original vertex set derived from an individual

plane.

(3) Volumetric fusion is performed via weighted integration:

$$V_{merged} = \alpha\, V_{sagittal} + (1 - \alpha) V_{coronal}$$

Here, V_{merged} denotes the coherent integration of 3D reconstruction, while the coefficient α

balances the contributions from the sagittal ($V_{sagittal}$) and coronal ($V_{coronal}$) components.

Data processing and statistics

The genuine meniscus harvested during total knee arthroplasty (TKA) surgery was first

compared with the 3D digital model for topography assessment. The Meniscus model was

aligned as previously described [24], and size measurement was performed with statistical

shape analysis based on the point correspondence model. Three positioning points along the

inner circumference (A: the most anterior location; P: the most posterior location; C: the

valley point, **Supplementary Fig. 1**) were determined as direct contact with the tibial

plateau, and their planes were superimposed for appropriate alignment. The root mean square

error (RMSE) was determined by the landmark locations between the original meniscus and

the digital model. Four dimensions of total length (L_T), gap length (L_G), anterior (W_A), and

posterior width (W_P) were determined. The gap ratio ($R_G = L_G/L_T$) and the length-width ratio

($R_{LW} = L_T/W_P$) were described to elaborate on the coverage of the tibia plateau cartilage and

the semi-circular shape of the meniscal tissue. Moreover, the cross-sectional geometry was

determined by the maximum height (H), width (W), and slope (H0/W0). These parameters

were introduced in **Supplementary Figure S1**.

Various types of tears were subjected to reconstruction and quantified based on their sizes.

Time-course meniscus reconstruction was carried out in one patient prior to knee arthroplasty

and meniscus repair surgery, and with three scans within the two-year visit. The standard

rehabilitation program was employed. The scans were performed using the same set-up and

re-segmented by the same reader to minimize the intra-observer differences. All size

measurements were conducted by three persons (with over 2 years of experience in MRI

imaging analysis) to avoid inter-observer variability. Data were analyzed using IBM SPSS 25

software and presented as means ± standard deviations (SD). The differences between the

original meniscus tissue and its digital model were compared using an unpaired t-test, with p

< 0.05 considered significant.

Results

Digital model construction

We initially evaluated the robustness of 3D reconstruction in recapitulating the geometric structure of the native menisci from two elderly osteoarthritis patients: a Male diagnosed with intra-capsule lipid-filled cyst formation (**Fig. 2a–d)** or a Female with classified Type III meniscal tears (**Fig. 2e–h**). Diagnostic MRI images captured in coronal and sagittal planes distinguished meniscal signals from surrounding bone, tendon, and muscle (**Fig. 2a&e**); however, both the size and topography of the meniscus remained barely discernible. In contrast, digital simulations generated vivid depictions of the menisci, enabling intuitive multi-angle assessment and quantitative analysis (**Fig. 2b&f**). Parameters critical to overall morphology, including total length (L_T), gap length (L_G), posterior (W_P), and anterior width (W_A), showed close alignment between reconstruction models and genuine tissues (**Fig. 2i**), indicating high fidelity of the digital models in replicating gross geometry. Cross-sectional measurements revealed topographic continuity and internal structures that are not observable in 2D MRI. The average rim heights of the medial menisci from two patients were 7.97 ± 0.85 mm and 4.20 ± 0.75 mm. Meniscal width (W0) gradually decreased from the valley point toward the anterior and posterior ends, reflecting its fan-shaped crescent morphology (**Fig. 2b&c**). Conversely, the other injured meniscus exhibited a narrowed width near the valley point, correlating with region-specific focal defects in native tissues (**Fig. 2f&g**). The curvature metric H0/W0 was significantly higher in the mid-body zone of the injured meniscus (**Fig. 2f&j**), but apparently lower in the discoid meniscus (**Fig. 2b, j**). Thus, our digital models, developed through subtractive segmentation with biplanar MRI integration, could accurately recapitulate meniscal geometry with minimal assay variation. Notably, the reconstructions captured authentic internal deformations in the male patient's meniscus, enabling in-depth observation of focal intrasubstance lesions (**Fig. 2d**). These spherical structures exhibited the absence of fibrochondrocytes and were nearly, if not at all, negative to safranin-O/Fast green staining, indicating a non-calcified composition [25], likely chondral lipid [26] or amyloid deposition [27]. Interestingly, this male OA patient was diagnosed with lipid-filled cyst formation in the tibia plateau capsule, implying a systemically disordered lipogenesis.

Tear simulation and quantification

We further investigated whether various types of meniscal tears could be quantitatively simulated. Clinically, tears are morphologically categorized as horizontal, longitudinal, radial, or complex types, and specifically described by delamination or displaced flap at advanced stages, or differentiated by location, such as anterior/posterior horns or red/white zones. First,

we modeled a lateral discoid meniscus (**Fig. 3a**), which, according to our *menis*MRI,

exhibited exceptionally clear full-thickness meniscal signals on spanning coronal and sagittal

planes. 3D reconstruction showed the characteristic pan-like morphology vividly. It detected

a subtle horizontal tear with delamination at the anterior horn (**Fig. 3a**, red asterisk in Gross

image), measuring approximately 2.42×0.15 mm. Cross-sectional analysis described the

internally existing structural derangement that was arthroscopically confirmed as chondral

lipid deposition during the meniscoplasty surgery (**Fig. 3a**, red area highlighted in the Cross-

sectional image). Proceeding to a female patient in her 50s, diagnosed with complex meniscal

tears, our simulation revealed comprehensive pathological deformation (**Fig. 3b**, arrow heads

in gross images), comprising two horizontal peripheral tears: one in the antero-mid meniscal

body of the medial meniscus (length: 10.5mm; width: 2.3±0.4mm); and another in the

posterior horn of the lateral meniscus (length: 13.4mm; width: 1.59±0.24mm). The

longitudinal tear in the postero-mid meniscal body, together with a horizontal tear, resulted in

full-thickness dissociation from the main body. Moreover, our model clearly demonstrated a

flap tear in the anterior horn (length: 16.2 mm; width: 2.3 mm), which was overlooked during

arthroscopy. Overall, our simulation predicts significant risks of the detachment of the flap

segments and dissociation of the longitudinal tear fragments, warranting urgent surgical

remediation. To be noted, cross-sectional simulation revealed intra-meniscal deformities of

the lateral meniscus, potentially indicating its degeneration and susceptibility to delamination.

Besides, flap tears could occasionally be mistaken for bucket-handle tears. In a case where

conventional 2D MRI diagnosed delamination and complex tearing in the left lateral

meniscus, our constructed digital model was able to recapture in detail 1) an extensive

delamination widespreading from the anterior to posterior horns (**Fig. 3c**, red asterisk in

angled gross view); and 2) concomitant longitudinal and radial tears in the meniscal body

configuring a flap tear that had been mistaken by a "bucket-handle tear" in 2D MRI. In

addition, a severe horizontal tear (length: 14mm; width: 2.0±0.3mm) in the medial meniscus

was defined, revealing a high risk of delamination that was overlooked during the initial ex

vivo diagnosis (**Fig. 3c**, arrowhead in gross images).

Our digital platform enabled the reconstruction of native meniscus geometry prior to tear

formation and facilitated in silico prediction of surgical outcomes. As in the case mentioned

above, we rebuilt the premorbid architecture of torn meniscus in a quantitative way (**Fig. 4a**),

demonstrating its Watanabe type II incomplete discoid morphology, and thereby proposing

operative recommendations as 1) arthroscopic partial meniscectomy for the unstable flap tear

component and 2) suture repair for delamination injuries at the anterior horn-body junction.

Moreover, in another case involving right medial meniscal delamination combined with flap

tear, our model accurately replicated both the complex tears at the meniscal body and

postero-mid meniscal delamination (length: 18.4mm; width: 3.7±0.9mm) (**Fig. 3d**, red

asterisks in Gross image). Though partial meniscectomy with radiofrequency-assisted

meniscal contouring had been performed and modelled postoperatively (**Fig. 4b**), we

simulated posterior horn suture repair as a viable alternative to preserve its height

(4.85±1.68mm vs 2.71±0.61mm) (**Fig. 4b**). Additionally, a focal longitudinal anterior horn

defect (length: 20mm; width: 3.0±0.5mm), likely secondary to flap tear displacement, was

identified in the lateral meniscus despite unremarkable 2D MRI and arthroscopic examination

(**Fig. 3d,** arrow heads in Gross image). Through computational processing of slice-layer

images with contour completion (e.g., A1-3 in Fig. 4c), we reconstructed the presumed

displaced flap tear component, restoring native meniscal anatomy (**Fig. 4c**).

Last but not least, high-resolution reconstruction was able to precisely recapitulate the surface

topography. In a gout-diagnosed case with intra-articular urate crystal deposition, arthroscopy

confirmed erosion originating from tophi, which was accurately mirrored by our digital

model (**Fig. 3e**, arrowheads in arthroscopic images), including several high-definition erosion

sites (diameter: 1.1±0.3mm) at the medial meniscus posterior horn. Notably, the lateral

meniscus exhibited full-thickness erosion (diameter: 3.13mm) coalescing with a horizontal

tear (length: 16.7mm; width: 2.7±0.8mm) in its peripheral third, indicating high propensity

for complex tear progression, delamination, and partial meniscal extrusion.

Meniscal healing evaluation

Given limited accuracy (48%-76%) and interobserver disagreement for predicting meniscal

tear reparability using 2D MRI [28], we then questioned the reliance of meniscal reconstruction

in a patient with two-year follow-up (**Fig. 5**). Preoperative simulation identified a

17.4×2.0mm longitudinal tear at the antero-mid body, demonstrating clearly the disruption of

the meniscus continuity (**Fig. 5b**, pre-op). The patient underwent meniscus-preserving

surgery with two vertical mattress sutures followed by standardized rehabilitation. Serial

MRI-based modeling revealed progressive healing: gap dimensions reduced to 11.5×1.0mm

at 3 months postoperatively (**Fig. 5b**, 3m post-op) and complete gap resolution after 2 years

(**Fig. 5b**, 2y post-op), suggesting effective tear coaptation and fibrocartilaginous regeneration.

In agreement with previous findings, a tear line was observable on the surface of the repaired

meniscus [29]. It should be noted that bridging was initiated on the surface, preceding internal

fusion. Furthermore, the 2-year post-operative simulation revealed internal structural

irregularities at the repair site, consistent with scar tissue formation, implying a potential

predisposition to degenerative changes and secondary injury, which encourages long-term

computational monitoring of the meniscus health in this patient.

<u>Discussion</u>

This study, for the first time, adopts a subtractive segmentation approach and bi-planar

integration to achieve high-resolution MRI-based 3D reconstruction of the meniscus,

enabling virtual resection of its anatomy and thereby facilitating the recapture of tears being

overlooked or obscured on 2D MRI. This model enables intuitive observation and

straightforward assessment of diverse tear types, facilitating precise tear measurement and

enhancing surgical planning and outcome prediction. Moreover, it encourages adaptive post-

operative rehabilitation based on time-course monitoring of tear healing. Although not the

focus of this study, it is important to emphasize that size-matched meniscal implantation is

essential to prevent secondary injury, underscoring the necessity of 3D models with adequate

geometric accuracy. Lastly, such a model can be integrated into augmented reality platforms

for intraoperative navigation, thereby improving surgical precision and efficacy.

Subtractive segmentation technique enabled the construction of highly clear 3D meniscal

models (*menis*MRI), surpassing the clarity achievable with conventional MRI. A proton

density (PD)-weighted turbo spin-echo (TSE) sequence was selected to reduce scan time

while maintaining diagnostic capability, as this protocol is widely established for evaluating

joint injuries and pathologies, including the detection of high signal intensity indicative of

meniscal tears [30]. Previously reported techniques that rely entirely on manual segmentation

are prone to omitting subtle anatomical features. In contrast, the subtractive approach retains

these details by automatically eroding extraneous tissues, requiring only minimal manual

removal of residual floating debris.

Studies have indicated that sagittal plane MRI images are most suitable for segmenting the

meniscus due to reduced partial volume effects at the medial and lateral ends, thereby

minimizing segmentation inaccuracies [8,10,22]. Here, we showed that 3D reconstruction relying

on a single plane is insufficient. Instead, integrating and merging data from at least two

orthogonal planes, specifically the coronal and sagittal views as implemented in this study,

proves optimal for accurate 3D model generation (Figure 1). Moreover, a slice thickness of

less than 2 mm is recommended for clear meniscal segmentation [31], whereas the standard

clinical protocol typically uses approximately 3 mm. Such a bi-planar approach helps

mitigate interslice signal loss. It enables the construction of two complementary models,

thereby improving continuity and reducing spatial omissions, particularly along peripheral

and curved regions of the meniscus.

Numerous studies have sought to improve the diagnostic specificity and sensitivity for tears

such as bucket handle tears by combining the MRI signs of absence of the bow tie, double

posterior cruciate ligament, and disproportional posterior horn [32], or for discoid lateral

meniscus, based on its tibial plateau coverage and the slope angle of the medial tibial

eminence[10], to name just a few. Conventional MRI lacks efficiency in detecting subtle

degenerative tears, such as ramp lesions or specific longitudinal and radial tears located in the

posterior horn [4,11,12]. Demographic factors are also used adjunctively; for instance, flap tears

are typically older (30–50 years) than those with bucket-handle tears (15–35 years). In

contrast, our reconstruction permits unbiased multi-angled observation of meniscal

architecture, thereby facilitating comprehensive differentiation of tear patterns and severity.

Moreover, the digital model enables the remodeling of the torn meniscus to virtually simulate

its native geometry and surgical outcomes. Given that meniscectomy increases OA incidence[5]

with inferior functional outcomes compared to repair [29], we reconstructed a discoid meniscus

with complex tearing. Clearly, we identified regions amenable to partial meniscectomy and

suture repair, guiding clinical practice (**Fig. 3c and 4a**). However, in another case where our

simulation suggested anterior horn suturing to preserve its rim height, meniscectomy was

ultimately employed (**Fig. 3d and 4b**). Above all, we demonstrated in a gout model that

meniscal surface topography can be accurately captured despite smoothening artifacts

typically introduced during reconstruction. This detailed evaluation of surface morphology

may advance our understanding of joint lubrication and the probability of cartilage damage.

Degenerative tears manifest as internal meniscal deformities and complex tears of the

posterior horn, arising from chronic shear and hoop stresses [33], which contribute to pain[34] and

represent a prognostic indicator of knee OA[4]. Our digital model effectively mimics and

vividly visualizes these focal internal deformities. In contrast to meniscal calcifications,

commonly found in aged and arthritic menisci and mediated by osteogenic differentiation via

ANKH and mTOR/ATF4 signaling [35], the internal defects identified here (Fig. 2) more

closely resemble chondral lipid [26] or amyloid deposits [26,36], which is evidenced by their

acellular nature and negative staining to Fast green. Arthroscopic observation of yellowish fat

tissue may support this notion. Further histological validation using Oil Red and Congo Red

staining is warranted. The accumulation of chondral lipid and amyloid within menisci is

strongly associated with OA progression[26,36]. In addition, meniscal histology revealed

pronounced tissue fissuring and disorganization of collagen bundles, underlying

microstructural disruption near the defects. Remarkably, follow-up reconstruction of repaired

menisci reveals reduction of tear diastasis without second-look arthroscopy, characterized by

the formation of a bridging structure that may indicate the osteochondral junction and

corresponds to tree branch-like MRI signs previously reported as a hallmark of ongoing

meniscus healing[37]. Interestingly, this reparative process appeared to commence peripherally

before progressing inward, which may be associated with the profound vascularity at the

meniscus margins.

This study has several limitations. Firstly, the sample size of this study is small, and cases

involving meniscal extrusion resulting from discrete root disruption were not included in this

model. In addition, larger cohorts will facilitate the development of a comprehensive tear-

classification scoring system with AI-assisted approaches. Last but not least, our

reconstruction relying on T1 MRI lacks the capacity to delineate the meniscal vascularity,

biochemical composition, and mechanics under physiological conditions. The breakdown and

disorientation of the collagen meshwork arising from the tears compromise meniscal

mechanics[38], which is measurable by non-invasive diffusion tensor imaging combined with

fiber tractography (DTI-FT). Dynamic stress imaging (DSI) recapitulates meniscus

mechanics under loads. Above all, pre-existing endoligamentous vessels within the peripheral

meniscus (10% - 30%)[39] are critical for meniscus nourishment and tear healing[40]. Hence,

integrating Contrast-Enhanced MRI, DTI-FT, and DSI, our reconstruction may present

anatomically composited and mechanically informed advanced functional 3D models.

Conclusions

This study is the first to employ a subtractive segmentation technique for semi-automated

identification of the meniscus on clinical MRI, combined with bi-planar model integration to

reconstruct its three-dimensional architecture. Our results suggest that this virtual

reconstruction accurately recapitulates the meniscal morphology, enabling detailed 3D

visualization and morphometric analysis of tear patterns and severity, as well as simulation of repair efficiency. This study presents a valuable strategy for personalized surgical planning and the development of biomimetic meniscal implants.

Author's contributions:

Liqi Ng: Conception and design, data acquisition and interpretation, producing the figures, drafting of the article

Peng Zhang: Provision of study materials and tissue preparation, data collection, and analysis.

Maojiang Lyu: Data collection and interpretation, statistical analysis.

Xin Mu: Data collection and interpretation.

Jianle Liang: MRI imaging and acquisition of study materials.

Peng Chen: Data collection, producing the figures.

Tian You: Critical revision of the article.

Lu Bai: Critical revision of the article.

Wanqi Ding: Data collection.

Yu Zhou: Interpretation of results.

Mingjing Zhang: Critical revision of the article.

Swastina Nath Varma: Critical revision of the article.

Yuluan Wu: Critical revision of the article.

Weiji Wang: Crtical revision of the article.

Yangchidung Lin: Conception and design, critical revision of article.

Xiaofeng Zhang: Critical revision of the article.

Yongcan Huang: Critical revision of the article.

Chaozong Liu: Conception and design, interpretation of results, critical revision of article.

Yi Sun: Conception and design, data acquisition and interpretation, drafting and final approval of the manuscript, and obtaining the funds.

Xintao Zhang: Conception and design, coordinating the collaboration, drafting and final approval of the manuscript, and obtaining the funds.

Conflict of interest statement

All authors had no conflict of interest that could inappropriately influence the bias of the presented work.

Acknowledgement:

This work is jointly supported by Guangdong Basic and Applied Basic Research Foundation (2024A1515010104&2025A1515012557), Shenzhen Science and Technology Program (JCYJ20240813115906009 & JCYJ20220530160218040), Shenzhen High-level Hospital Construction Fund and Peking University Shenzhen Hospital Scientific Research Fund (KYQD2022226, KYQD2023345 and KYQD2025456), Innovation and Development Initiative for Orthopaedic Drug Research, China Association of Chinese Medicine (GSKQNJJ-2023-011).

References

1. Fox, A.J., Wanivenhaus, F., Burge, A.J., Warren, R.F. & Rodeo, S.A. The human meniscus: a review of anatomy, function, injury, and advances in treatment. *Clinical Anatomy* **28**, 269-287 (2015).

2. Choi, N.-H., Yang, B.-S., Lee, D.-M. & Lee, C.-R. Association of Increased Lateral Femoral Condylar Ratio With Lateral Meniscus Posterior Root Tear in Noncontact ACL Injury. *Orthopaedic Journal of Sports Medicine* **12**, 23259671231224023 (2024).

3. Keyhani, S., Esmailiejah, A.A., Mirhoseini, M.S., Hosseininejad, S.-M. & Ghanbari, N. The prevalence, zone, and type of the meniscus tear in patients with anterior cruciate ligament (ACL) injury; does delayed ACL reconstruction affects the meniscal injury? *Archives of Bone and Joint Surgery* **8**, 432 (2020).

4. Ward, R.J., Driban, J.B., MacKay, J.W., McAlindon, T.E., Lu, B., Eaton, C.B., Lo, G.H., Barbe, M.F. & Harkey, M.S. Meniscal degeneration is prognostic of destabilzing meniscal tear and accelerated knee osteoarthritis: Data from the Osteoarthritis Initiative. *J Orthop Res* **41**, 2418-2423 (2023).

5. Li, L., Yang, L., Zhang, K., Zhu, L., Wang, X. & Jiang, Q. Three-dimensional finite-element analysis of aggravating medial meniscus tears on knee osteoarthritis. *Journal of orthopaedic translation* **20**, 47-55 (2020).

6. Anderson, A.F., Irrgang, J.J., Dunn, W., Beaufils, P., Cohen, M., Cole, B.J., Coolican, M., Ferretti, M., Glenn Jr, R.E. & Johnson, R. Interobserver reliability of the International Society of Arthroscopy, Knee Surgery and Orthopaedic Sports Medicine (ISAKOS) classification of meniscal tears. *The American journal of sports medicine* **39**, 926-932 (2011).

7. Wright, R.W., Huston, L.J. & Haas, A.K. Ten-Year Outcomes of Second-Generation, All-
Inside Meniscal Repair in the Setting of ACL Reconstruction. *J Bone Joint Surg Am* **105**,
908-914 (2023).

8. El-Hagrasy, A.M.A., Theckayil, A.J., Khan, M.A., Khan, H.N. & Butt, A.J. Magnetic
Resonance Imaging Is an Effective First-Line Noninvasive Tool for Meniscal Tear
Detection: A Retrospective Comparative Analysis With Knee Arthroscopy. *Arthrosc*
Sports Med Rehabil **7**, 101065 (2025).

9. D'Ambrosi, R., Di Maria, F., Ursino, C., Ursino, N., Di Feo, F., Formica, M. &
Kambhampati, S.B. Magnetic resonance imaging shows low sensitivity but good
specificity in detecting ramp lesions in children and adolescents with ACL injury: A
systematic review. *J ISAKOS* **9**, 371-377 (2024).

10. You, M., Li, J., Zhang, X., Chen, G. & Li, J. Diagnostic Efficiency of MRI in Child and
Adolescent Lateral Discoid Meniscus. *Orthop Surg* **17**, 900-908 (2025).

11. Kim, S.H., Lee, H.J., Jang, Y.H., Chun, K.J. & Park, Y.B. Diagnostic Accuracy of
Magnetic Resonance Imaging in the Detection of Type and Location of Meniscus Tears:
Comparison with Arthroscopic Findings. *J Clin Med* **10**(2021).

12. Vo, T.T., Nguyen, D.T., Dinh Le, N.A., Van Nguyen, K.H., Vuu, H.K., Le, T.A. & Le
Phan, N.B. Evaluation of meniscal injury on magnetic resonance imaging and knee
arthroscopy in patient with anterior cruciate ligament tear. *SICOT J* **10**, 56 (2024).

13. Escoda Menendez, S., Garcia Gonzalez, P., Meana Moris, A.R., Del Valle Soto, M. &
Maestro Fernandez, A. Evaluation of the reliability and accuracy of MRI for the diagnosis
of meniscal ramp lesions. *Acta Radiol* **65**, 1382-1389 (2024).

14. Kim, S.H., Lee, H.-J., Jang, Y.-H., Chun, K.-J & Park, Y.-B. Diagnostic accuracy of
magnetic resonance imaging in the detection of type and location of meniscus tears:
comparison with arthroscopic findings. *Journal of clinical medicine* **10**, 606 (2021).

15. Benjaminse, A., Webster, K.E., Kimp, A., Meijer, M. & Gokeler, A. Revised approach to
the role of fatigue in anterior cruciate ligament injury prevention: a systematic review
with meta-analyses. *Sports medicine* **49**, 565-586 (2019).

16. Strawbridge, J.C., Schroeder, G.G., Garcia-Mansilla, I., Singla, A., Levine, B.D.,
Motamedi, K., Jones, K.J. & Kremen, T.J., Jr. The Reliability of 3-T Magnetic Resonance
Imaging to Identify Arthroscopic Features of Meniscal Tears and Its Utility to Predict
Meniscal Tear Reparability. *Am J Sports Med* **49**, 3887-3897 (2021).

17. Bilgen, B., Jayasuriya, C.T. & Owens, B.D. Current concepts in meniscus tissue
engineering and repair. *Advanced healthcare materials* **7**, 1701407 (2018).

18. Song, B., Tan, W., Xu, Y., Yu, T., Li, W., Chen, Z., Yang, R., Hou, J. & Zhou, Y. 3D-MRI
combined with signal-to-noise ratio measurement can improve the diagnostic accuracy
and sensitivity in evaluating meniscal healing status after meniscal repair. *Knee Surg*
Sports Traumatol Arthrosc **27**, 177-188 (2019).

19. Li, H., Li, P., Yang, Z., Gao, C., Fu, L., Liao, Z., Zhao, T., Cao, F., Chen, W. & Peng, Y.
Meniscal regenerative scaffolds based on biopolymers and polymers: recent status and
applications. *Frontiers in Cell and Developmental Biology* **9**, 661802 (2021).

20. Niu, W., Guo, W., Han, S., Zhu, Y., Liu, S. & Guo, Q. Cell-based strategies for meniscus
tissue engineering. *Stem cells international* **2016**(2016).

21. Tongdee, R., Narra, V., Oliveira, E., Chapman, W., Elsayes, K. & Brown, J. Utility of 3D
magnetic resonance imaging in preoperative evaluation of hepatobiliary diseases. *HPB* **8**,
311-317 (2006).

22. Kruger, N., McNally, E., Al-Ali, S., Rout, R., Rees, J.L. & Price, A.J. Three-dimensional
reconstructed magnetic resonance scans: Accuracy in identifying and defining knee
meniscal tears. *World Journal of Orthopedics* **7**, 731 (2016).

23. Helms, C.A. The meniscus: recent advances in MR imaging of the knee. *American*
journal of roentgenology **179**, 1115-1122 (2002).

24. Vrancken, A., Crijns, S., Ploegmakers, M., O'kane, C., van Tienen, T.G., Janssen, D.,
Buma, P. & Verdonschot, N. 3D geometry analysis of the medial meniscus–a statistical
shape modeling approach. *Journal of anatomy* **225**, 395-402 (2014).

25. Hellberg, I., Karjalainen, V.P., Finnila, M.A.J., Jonsson, E., Turkiewicz, A., Onnerfjord,
P., Hughes, V., Tjornstrand, J., Englund, M. & Saarakkala, S. 3D analysis and grading of
calcifications from ex vivo human meniscus. *Osteoarthritis Cartilage* **31**, 482-492
(2023).

26. Park, S., Park, H.J. & Kim, J.N. Usefulness of the fast spin-echo multi-point Dixon image
of the knee joint on 3.0 T MRI for evaluation of meniscal tear. *Acta Radiol*,
2841851251347780 (2025).

27. Yanagisawa, A., Ueda, M., Sueyoshi, T., Nakamura, E., Tasaki, M., Suenaga, G.,
Motokawa, H., Toyoshima, R., Kinoshita, Y., Misumi, Y., Yamashita, T., Sakaguchi, M.,
Westermark, P., Mizuta, H. & Ando, Y. Knee osteoarthritis associated with different kinds
of amyloid deposits and the impact of aging on type of amyloid. *Amyloid* **23**, 26-32
(2016).

28. Holland, C.T., Tse, S., Bateni, C.P., Chen, D. & Lee, C.A. MRI-Based Prediction of
Meniscal Tear Repairability Demonstrates Limited Accuracy and Reliability. *J Clin Med*
14(2025).

29. Ahmed, I., Dhaif, F., Khatri, C., Parsons, N., Hutchinson, C., Staniszewska, S., Price, A.
& Metcalfe, A. The meniscal tear outcome (METRO) review: A systematic review
summarising the clinical course and outcomes of patients with a meniscal tear. *Knee* **38**,
117-131 (2022).

30. Lefevre, N., Naouri, J.F., Herman, S., Gerometta, A., Klouche, S. & Bohu, Y. A current
review of the meniscus imaging: proposition of a useful tool for its radiologic analysis.
Radiology research and practice **2016**(2016).

31. Shen, X., Zuo, J., Li, Z., Xiao, J. & Liu, T. Morphological analysis of normal meniscus on
magnetic resonance imaging (MRI)-based three-dimensional reconstruction models in
healthy Chinese adults. *Medical Science Monitor: International Medical Journal of*
Experimental and Clinical Research **26**, e927101-927101 (2020).

32. Pakdemirli, E., Cesur, T. & Bozkurt, I. MRI in the Diagnosis of Bucket Handle Tears:
What Is the Current Situation? *Cureus* **15**, e43324 (2023).

33. Howell, R., Kumar, N.S., Patel, N. & Tom, J. Degenerative meniscus: Pathogenesis,
diagnosis, and treatment options. *World J Orthop* **5**, 597-602 (2014).

34. Tsai, L., Matzkin, E., Jones, M.H., Miller, R.E. & Katz, J.N. Potential sources of pain in
symptomatic degenerative meniscal tear: A narrative review. *Osteoarthr Cartil Open* **7**,
100616 (2025).

35. Shakya, B.R., Karjalainen, V.P., Hellberg, I., Finnila, M.A.J., Elkhouly, K., Sjogren, A.,
Turkiewicz, A., Onnerfjord, P., Hughes, V., Tjornstrand, J., Englund, M. & Saarakkala, S.
Prevalence and classification of meniscal calcifications in the human knee. *Osteoarthritis*
Cartilage **32**, 1443-1451 (2024).

36. Solomon, A., Murphy, C.L., Kestler, D., Coriu, D., Weiss, D.T., Makovitzky, J. &
Westermark, P. Amyloid contained in the knee joint meniscus is formed from
apolipoprotein A-I. *Arthritis Rheum* **54**, 3545-3550 (2006).

37. Schwach, M., Grange, S., Klasan, A., Putnis, S., Philippot, R. & Neri, T. MRI Criteria for
Healing at 1 Year After Repair of a Traumatic Meniscal Tear. *Am J Sports Med* **51**, 3693-
3700 (2023).

38. Ewing, J.W. *Articular cartilage and knee joint function: basic science and arthroscopy*,
(Lippincott Williams & Wilkins, 1990).

39. Harner, C.D., Janaushek, M.A., Kanamori, A., Yagi, M., Vogrin, T.M. & Woo, S.L.

Biomechanical analysis of a double-bundle posterior cruciate ligament reconstruction.

The American journal of sports medicine **28**, 144-151 (2000).

40. Dratsch, T., Zäske, C., Siedek, F., Rauen, P., Hokamp, N.G., Sonnabend, K., Maintz, D.,

Bratke, G. & Iuga, A. Reconstruction of 3D knee MRI using deep learning and

compressed sensing: a validation study on healthy volunteers. *European Radiology*

Experimental **8**, 47 (2024).

Figures and tables

Figure 1. Workflow for meniscal semi-automatic subtractive segmentation and 3D reconstruction via bi-planar integration. An initial global mask was created from clinical MRI scans at both coronal and sagittal plans, and further subtracted from a denser segmentation to extract a meniscus-specific subtractive mask. Meniscal signals were enhanced in *menis*MRI to facilitate 3D reconstruction. Digital representation of the meniscus was obtained by merging two complementary 3D models derived from the two planes.

 Figure 2. Meniscus reconstruction and fit analysis. (a&e) 2D MRI, (b&f) reconstructed 3D
digital model, (c&g) gross morphology and (d&h) Safranin O/Fast green staining of the
menisci from osteoarthritis knee joints . Comparative views of exterior and interior geometry
were shown. Star: meniscal defect site; A: anterior, P: posterior. Scale bar: 500μm. (i&j)
Geometry quantification based on the genuine menisci (Gen-) and their digital models (Dig-).
W_A: anterior width, W_P: posterior width, L_G: Gap length, L_T: total length, W_0: inner width,
H_0: inner height.

Figure 3. Meniscal tears simulation. (a) Right lateral discoid meniscus; (b) Right lateral
meniscus with complex tears; (c) Left lateral discoid meniscus with complex tears and
delamination; (d) Right medial meniscus with delamination; (e) meniscus with tophi. For
each tear type, images for its clinical MRI (2D MRI), meniscus-enhanced MRI (*menisMRI*),
corresponding cross-sectional view from the 3D model, and the gross reconstruction were
presented. Scale bar: 500μm. Triangles indicate tears that were obscure or overlooked in
clinical MRI or during arthroscopy; Red stars indicate tears that were verified by clinical
MRI or arthroscopy.

Figure 4. Robustness of meniscal 3D reconstruction for virtual simulation and

prediction of meniscal changes. (a) Simulation of the premorbid structure in a meniscus

with complex tears (as described in Fig. 3c). Red dot line indicates the flap tear fragment. (b)

Surgical outcome prediction for the meniscus is shown in Fig. 3d. Black stars indicate flap

tear removed by meniscectomy. Triangles mark delaminated areas, with morphology

following suture repair or meniscectomy reconstructed. (c) Predictive reconstruction of a

meniscal defect via site-specific cross-sectional supplementation.

Figure 5. Follow-up reconstruction and evaluation of meniscal healing. (a) Representative MRI images of the meniscus with a longitudinal tear preoperatively, and at 3 months and 2years post-suture surgery. (b) Menisci reconstruction and cross-sectional views at the same time points. Cross-sections views through the suture site illustrate the healing progression.

729 **Supplementary Figure 1. Schematic illustration of meniscus geometry measurement. (a)**

730 Gross and cross-sectional measurements. (b) Parameters measurement summarization.

b

P_H	Posterior half point
P	Posterior inner point
C	Central
A	Anterior inner point
A_H	Anterior half point
W_P	Posterior width
W_A	Anterior width
L_G	Gap length
L_T	Total length
W	Width
W_0	Inner width
H	Height
H_0	Inner height
B	Bulge

Gender	Age Group	Side	Where analyzed in this study	MRI Diagnosis	Arthroscopy	Surgical intervention
M	71-75	R	Fig. 2	Osteoarthritis; Intra-capsule lipid-filled cyst formation		TKA
F	66-70	R		Osteoarthritis; Meniscus tear (Type III) in anterior/ posterior horn		
F	51-55	R	Fig. 3a	Lateral discoid meniscus tear		Partial meniscectomy
M	51-55	R	Fig. 3b	Type III posterior horn and body tears (Medial); lateral meniscus tear	Posterior horn tear (Medial); Longitudinal tear in white-zone (Lateral)	Partial Meniscectomy
F	41-45	L	Fig. 3c&4a	Lateral discoid meniscus tear	Discoid meniscus, intra-substance deposition of yellowish fat-like tissue, posterior horn complex tear with delamination (Lateral)	Meniscectomy and suture repair
M	31-35	R	Fig. 3d&4b-c	Posterior horn tears with cyst (Medial)	Mid-body to posterior horn horizontal tear with flap, parameniscal cyst (Medial)	Partial meniscectomy

M	36-40	R	Fig. 3e	Anterior horn tear (Medial)	Anterior horn flap tear (Medial); Gouty erosion (Lateral)	Suture
M	31-35	L	Fig. 5	Medial meniscus tear	Bucket-handle tear (Medial)	Suture repair