Title Page

Title: Individual-specific resting-state networks predict language dominance in drug-resistant epilepsy

Authors:

Mervyn Lim Jun Rui, MBBS, MPH, FRCS[1,2,3], Zhang Shaoshi, PhD[1], Shreya Pande, B.Eng[1], Xue Aihuiping, PhD[1], Ru Kong, PhD[1], Kareem Zaghloul, MD, PhD[2], Sara Inati, MD[2], Thomas Yeo Boon Thye, PhD[1]

1. Computational Brain Imaging Group, Yong Loo Lin School of Medicine, National University of Singapore
2. National Institute of Neurological Disorders and Stroke, National Institute of Health
3. Division of Neurosurgery, Department of Surgery, National University Hospital

Corresponding author: Thomas Yeo Boon Thye, Computational Brain Imaging Group, Yong Loo Lin School of Medicine, National University of Singapore, Tahir Foundation Building (MD1), 12 Science Drive 2, #13-05C, National University of Singapore, Singapore 117549. Phone: +65 97391162, Email: thomas.yeo@nus.edu.sg.

Manuscript word count: 2743

Key Points

Question: Can individual-specific network topography from resting-state functional magnetic resonance imaging (fMRI) predict task-based language dominance in drug-resistant epilepsy?

Findings: In this multi-centre case-control study of 91 participants with drug-resistant epilepsy and 40 healthy controls, individual-specific networks outperformed group-average networks and generalized well to an independent cohort. Language network topography differed significantly across left (mean lateralization index (LI) = 0.165 ± 0.106), bilateral (LI = 0.056 ± 0.074), and right (LI = 0.023 ± 0.055) dominance groups (p = 0.002).

Meaning: Resting-state fMRI can estimate high-quality individual-specific cortical networks that predict language dominance, providing a non-invasive tool for presurgical evaluation.

Abstract

Importance: Identifying language dominance is a crucial step in epilepsy surgery planning. We applied a precision functional brain mapping approach to estimate individual-specific cortical resting-state networks in drug-resistant epilepsy and predict language dominance.

Objective: To determine whether individual-specific cortical network topography can predict task-based language dominance in drug-resistant epilepsy.

Design: Retrospective case-control study conducted between January 2024 and August 2025.

Setting: Multicentre population-based study including healthy participants from the Human Connectome Project, and participants with drug-resistant epilepsy from the National Institutes of Health (NIH) and the University of Iowa.

Participants: Eligible participants had drug-resistant epilepsy defined by International League Against Epilepsy criteria and were undergoing pre-surgical evaluation. All participants underwent neuroimaging, with a subset receiving concurrent intracranial electrical stimulation during fMRI.

Main Outcomes and Measures: Individual-specific cortical network topography and prediction of task functional magnetic resonance imaging language dominance.

Results: Ninety-one participants with drug-resistant epilepsy were included: 61 (67.0%) temporal lobe epilepsy, 29 (31.9%) extra-temporal lobe epilepsy, and 1 (1.1%) undetermined seizure onset zone. The mean age was 33.0 ± 11.4 years and 50 (54.9%) were male. There were 40 healthy participants with a mean age of 29.0 ± 4.0 years, and 16 (40.0%) were male.

We developed a multi-session hierarchical Bayesian model (MS-HBM) trained on NIH data to estimate individual-specific networks in drug-resistant epilepsy. MS-HBM trained on epilepsy data outperformed group-average networks or MS-HBM trained on healthy participants and generalized well to an independent dataset. During concurrent intracranial electrical stimulation, cortical activation and deactivation aligned more closely to individual-specific networks than group-average networks. Individual-specific language network topography significantly differed across left (mean lateralization index (LI) = 0.165 ± 0.106; area-under-the-curve (AUC) = 0.82), bilateral (LI = 0.056 ± 0.074; AUC = 0.72), and right (LI = 0.023 ± 0.055; AUC = 0.83) language dominance groups (p = 0.002).

Conclusions and Relevance: Our model is publicly available (github link), which may be used to predict language dominance from approximately 10 minutes of resting-state fMRI. This provides a practical, non-invasive tool for presurgical evaluation of drug-resistant epilepsy.

Introduction

Identifying language dominance is a crucial step in epilepsy surgery planning. Current guidelines recommend that task functional magnetic resonance imaging (fMRI) may be considered as a non-invasive alternative to the intracarotid amobarbital test[1-3]. However, the spatial pattern and strength of task-evoked activation are highly dependent on patient compliance and the specific language paradigm used[4,5]. Resting-state fMRI avoids this dependency on task performance, but prior studies showed that resting-state functional connectivity predicts task fMRI language dominance only modestly[6-9].

We previously developed an approach to derive large-scale resting-state networks of the cerebral cortex[10] as well as a multi-session hierarchical Bayesian model (MS-HBM) for estimating individual-specific networks with limited quantities of data – MS-HBM networks obtained using only 10 minutes of data were comparable to other approaches requiring approximately 50 minutes[11]. Large-scale functional networks vary across individuals[12-14] and differ between healthy and pathological brains[15,16]. Moreover, individual-specific differences in network topography were able to predict inter-individual variation in cognition and personality[11]. Despite significant progress in estimating individual cortical networks in healthy individuals, their properties and behavioural relevance remain poorly characterized in drug-resistant epilepsy.

Here, we applied our approach for deriving group-average[10] and individual-specific cortical networks[11] to participants with drug-resistant temporal lobe and extra-temporal lobe epilepsy undergoing presurgical evaluation at the National Institute of Neurological Disorders and Stroke, National Institutes of Health (NIH dataset)[6]. We showed that individual-specific networks using MS-HBM trained on drug-resistant epilepsy were of higher quality than group-average networks and MS-HBM trained on healthy participants.

We further validated and tested generalizability of the MS-HBM model in an independent drug-resistant epilepsy dataset[17], showing that during concurrent intracranial electrical stimulation with fMRI, cortical activation and deactivation better conformed to individual-specific network boundaries than to group-average boundaries. Finally, we showed that individual-specific networks predict task-based language dominance in individual patients.

Our model is publicly available (github link), which may be used to estimate high-quality individual-specific cortical networks and predict language dominance in a new individual with drug-resistant epilepsy using approximately 10 minutes of resting-state fMRI data.

Methods

Participants and Datasets

In this study, we included one open-source dataset of healthy participants and two independent datasets of participants with drug-resistant epilepsy undergoing pre-surgical evaluation. Human Connectome Project (HCP) participants[11] had a mean age of 29.0 ± 4.0 years, 16 (40.0%) were male, and 12 (30.0%) self-identified as non-White and/or Hispanic. NIH epilepsy participants[6] had a mean age of 31.9 ± 11.2 years and 33 (50.8%) were male. Of these, 42 (64.6%) had temporal lobe epilepsy and 23 (35.4%) had extra-temporal lobe epilepsy (Table S1). Electrical-stimulation fMRI (esfmri) dataset[17] participants had a mean age of 35.8 ± 11.5 years, and 17 (65.4%) were male. Of these, 19 (73.1%) had temporal lobe epilepsy, 6 (23.1%) had extra-temporal lobe epilepsy, and 1 (3.8%) had undetermined seizure onset zone (Table S2).

All datasets included structural and resting-state fMRI scans. In the esfmri dataset, post-operative fMRI was acquired with concurrent intracranial electrical stimulation delivered through stereoencephalography electrodes and interleaved with echo planar imaging volume acquisition[17]. Alternating blocks of approximately 30 seconds of no stimulation or stimulation

was delivered during the fMRI scan. No cognitive task was given to the participant, and no behavioural effects were evoked during the stimulation period.

Further details regarding the demographics, data collection, and pre-processing are provided in the supplementary methods[6,17-19]. After pre-processing, there were 40 HCP participants with 4 runs each[11]. Of the 48 NIH participants with two or more runs, 34 (70.8%) participants were used for the training set, and 14 (29.2%) participants were used for the test set. For the esfmri dataset, there were 11 participants that had at least two valid pre-operative runs and 17 participants that had at least two valid post-operative runs (Figure 1A).

Derivation of group-average networks and training of MS-HBM for estimating individual-specific networks

Our approach has been described previously[10,11]. At each cortical region (81,924 vertices), Pearson's correlation between the fMRI time series and 1175 regions of interests were binarized by keeping the top 10% of correlations. The connectivity profiles were then clustered using a mixture of von Mises-Fisher distributions to derive the group-average 15-networks[10,20]. In addition to computing group-average networks for the 40 HCP participants and 34 NIH participants, we also compared our results to group-average networks derived from an independent dataset of 15 intensively-sampled healthy participants from Du et al[18].

We trained MS-HBM models on either the 40 HCP participants (using the HCP group-average[11] or Du group-average[18]) or 34 NIH participants to estimate model parameters, including the inter-subject functional connectivity variability, intra-subject functional connectivity variability, spatial smoothness prior, and the inter-subject spatial variability prior[11]. Individual-specific networks were visually confirmed using model-free seed-based functional connectivity. Inter-subject similarity in network topography across the 15-networks

was quantified using the Dice similarity coefficient across all 57 NIH participants with at least one valid run.

Validation and generalization of MS-HBM trained on drug-resistant epilepsy

To evaluate the quality of individual-specific networks estimated from MS-HBM trained on drug-resistant epilepsy or healthy participants, we computed two previously established metrics[11,14,21,22]: the weighted resting-state connectional homogeneity and weighted task (electrical-stimulation) functional inhomogeneity. These metrics encode the principle that if an individual-specific network captured the system-level organization of the individual's cortex, each network should exhibit homogeneous connectivity and function.

We first evaluated the resting-state homogeneity of each network on the NIH test set of 14 participants that were not used for training MS-HBM. The generalizability of the MS-HBM model was then tested using pre-operative (11 participants) and post-operative (17 participants) esfmri data, as well as task inhomogeneity during electrical stimulation in the post-operative data. For estimating individual-specific networks in the post-operative data, we used only blocks with no stimulation after discarding the first 4 frames of each block to remove residual effects of the haemodynamic response from the previous block[23].

To ensure that runs used for estimating individual-specific networks were independent from runs used for computing evaluation metrics, we used a leave-one-run-out cross-validation method. For each participant with n valid runs, individual-specific networks were estimated using n-1 runs, and evaluation metrics were computed on the left-out run. This was repeated for all runs, and an average result was obtained for each participant. Statistical tests were performed using two-tailed paired t-tests between each pair of networks (dof = number of participants – 1). Statistical significance was taken to be q<0.05 after false discovery rate correction.

The average resting-state homogeneity and task inhomogeneity were compared across six networks: (1) HCP group-average networks; (2) individual-specific networks estimated using HCP MS-HBM; (3) Du group-average networks; (4) individual-specific networks estimated using Du MS-HBM; (5) NIH group-average networks; and (6) individual-specific networks estimated using NIH MS-HBM. Resting-state homogeneity was computed by averaging the Pearson's correlations between each vertex's time course with the average time course of its network. The correlations were then averaged across all 15 networks while accounting for network size[11,21,22]. Task inhomogeneity was computed as the average standard deviation of cortical activation z-scores within each network from unthresholded generalized linear models of intracranial electrical stimulation[19]. The standard deviation was then averaged across all 15 networks while accounting for network size[11,21,22]. A higher standard deviation indicated higher task inhomogeneity. Further details on the computation of the generalized linear models are provided in the supplementary methods.

To visualise the effects of intracranial electrical stimulation on cortical networks, group-average network boundaries and individual-specific network boundaries were overlayed on the electrical-stimulation z-score activation maps. We hypothesised that electrical-stimulation evoked activity would align more closely to individual-specific network boundaries than group-average network boundaries.

Predicting language dominance using individual-specific network topography

All patients underwent language task fMRI using an auditory description decision task as described previously[5,6]. Task-based language dominance was determined by a trained epileptologist (SI) through visual inspection of activation maps in the frontal and temporal regions at three thresholds (p < .01, p < .001, top 10% of activations) and confirmed at a

multi-disciplinary epilepsy conference. There were a total of 43 NIH participants with valid task language dominance and at least one valid resting-state fMRI run.

In several participants with drug-resistant epilepsy, regions typically associated with the language network exhibited connectivity patterns characteristic of the default mode network, suggesting network reorganization. To capture this variability, we computed a resting-state laterality index (LI) based on the network topography of both the Language A and Language/ Default B networks:

$$\text{Resting-state LI} = \frac{LH - RH}{LH + RH},$$

where LH and RH represent the number of vertices in the left and right hemispheres, respectively, assigned to the individual-specific Language A and Language/ Default B networks. Resting-state LI values for participants with left, bilateral, or right task-based language dominance were compared using one-way ANOVA, with $p < 0.05$ considered statistically significant. Finally, receiver operating curves and area-under-the-curve statistics were computed to evaluate the predictive power of resting-state LI for each language dominance group.

Results

Individual-specific resting-state networks in drug-resistant epilepsy capture inter-individual variability and outperform group-level networks

We estimated large-scale resting-state networks in drug-resistant epilepsy despite structural and functional abnormalities. Networks were categorized into 9 canonical groups (Visual, Somatomotor, Auditory, Limbic, Control, Dorsal Attention, Salience/Ventral Attention,

Language, and Default), which were broadly consistent with prior literature[10,11,18,24]. Group-level networks in healthy participants were more spatially distributed than those in drug-resistant epilepsy (Figure 1B). For example, association networks such as the control network were subdivided into three networks in HCP data[11], but were represented by a single, less distributed network in drug-resistant epilepsy. Instead, two spatially focal limbic networks emerged in orbitofrontal and parahippocampal regions. These differences were not explained by motion as the average relative root mean square framewise displacement was 0.059 ± 0.023 vs. 0.076 ± 0.015 and the average voxel-wise differentiated signal variance (DVARS)[25] was 23.0 ± 4.6 vs. 58.4 ± 6.8 for the NIH dataset compared to the HCP dataset respectively.

Using MS-HBM, individual-specific networks captured inter-individual variability while preserving canonical topographical features of healthy participants. However, in some participants, inferior frontal language regions exhibited stronger connectivity with middle temporal and temporopolar areas typically associated with the default mode network in healthy individuals, suggesting network reorganization (Figure 2A vs. 2B). Language connectivity was visually confirmed using identical seeds in the inferior frontal and lateral temporal regions across participants (Figure 2C). Finally, we observed that somatomotor and visual networks exhibited lower inter-subject variability (DICE similarity coefficient ranging from 0.66 – 0.71) than association networks such as the salience/ ventral attention, dorsal attention, control, and language networks (DICE similarity coefficient ranging from 0.35 – 0.50; Figure 2D).

Next, we compared the quality of group-level networks estimated from the HCP data, Du atlas, and NIH data with the individual-specific networks estimated in 14 NIH subjects using the three MS-HBM models. Individual-specific networks estimated using NIH MS-HBM

performed significantly better than all group-average networks and MS-HBM trained on healthy participants, even after correcting for multiple comparisons (Figure 3A).

MS-HBM trained on drug-resistant epilepsy generalized to an independent dataset and showed the lowest electrical-stimulation inhomogeneity

MS-HBM trained on healthy participants might perform worse than NIH MS-HBM because of site and scanner differences. Therefore, we tested if NIH MS-HBM was generalizable to an independent dataset of drug-resistant epilepsy. Individual-specific networks estimated using NIH MS-HBM performed significantly better in resting-state homogeneity than all group-average networks and MS-HBM trained on healthy participants, even after correcting for multiple comparisons (Figure 3B and 3C).

In addition, individual-specific networks estimated using NIH MS-HBM performed significantly better than group-average networks for electrical-stimulation inhomogeneity, even after correcting for multiple comparisons (Figure 4A). We observed that during concurrent intracranial electrical stimulation and fMRI scans, cortical evoked activity aligned better to individual-specific network boundaries than group-average network boundaries (Figure 4B). Taken together, these results suggest that MS-HBM trained using drug-resistant epilepsy can estimate high-quality, generalizable individual-specific networks that captured system-level cortical organization better than group-average networks.

Individual-specific language network topography predicts task-based language dominance

Having shown that individual-specific networks estimated from NIH MS-HBM captured inter-individual variability in cortical resting-state networks, we next tested if language network

topography were able to predict task-based language dominance. Resting-state LI differed

significantly across left (mean LI = 0.165 ± 0.106), bilateral (mean LI = 0.056 ± 0.074), and

right (mean LI = 0.023 ± 0.055) task-based language dominance (p = 0.002). Receiver

operating curve analyses showed area-under-the-curve statistics of 0.82 for left, 0.72 for

bilateral, and 0.83 for right language dominance versus other participants. Conversely,

individual-specific networks estimated using MS-HBM trained on healthy participants were

not able to differentiate between task-based language dominance (HCP MS-HBM: p = 0.150;

Du MS-HBM: p = 0.303).

Discussion

Precision functional mapping techniques have been shown to reliably estimate the intrinsic

cortical organisation of an individual, but typically required long scan durations of an hour or

more fMRI data[12,13,26] and have not been applied to drug-resistant epilepsy. Using MS-HBM,

we showed that despite structural abnormalities and differences in functional connectivity,

individual-specific networks may be reliably estimated in drug-resistant epilepsy even with

only approximately 10 minutes of single-session resting-state fMRI data. MS-HBM trained on

drug-resistant epilepsy captured the organization of an individuals' cerebral cortex better

than group-average networks or MS-HBM trained on healthy participants. Furthermore, NIH

MS-HBM generalized to an external dataset of drug-resistant epilepsy despite site, scanner,

and pre-processing differences.

Although cortical networks in drug-resistant epilepsy broadly resembled canonical resting-

state networks identified in healthy participants, we observed that spatial topography was

more focal and individual-level differences emerged that were not typically observed in

healthy participants. These differences could not be explained by motion. In particular,

inferior frontal language regions exhibited stronger connectivity with regions typically

associated with the default mode network in healthy individuals, which may be due to the long-term effects of drug-resistant epilepsy on functional connectivity[27-32]. Existing data showed that alterations in functional connectivity appear to be related to the duration and severity of epilepsy, and were associated with cognitive impairment, suggesting progressive cortical reorganization in the setting of recurrent seizures over time[33,34]. Thus, further research using precision functional mapping may provide deeper insight into the effects of chronic epileptogenic activity on reorganization of the resting-state cortical networks[35].

These differences in the resting-state language network in drug-resistant epilepsy may potentially explain why previous research in resting-state laterality using seed-based connectivity were limited in predicting task-based language dominance[6-9]. Here, we showed that by accounting for both the Language A and the Language/ Default B networks, individual-specific network topography was able to predict task-based language dominance. Future studies should validate this resting-state laterality index against the intracarotid amobarbital test[3] or cortical electrical stimulation[36] and assess generalizability across other drug-resistant epilepsy cohorts. By characterizing functional organization at the individual level, we hypothesize that individual-specific networks may also predict the lateralization of other cognitive functions[33,37] as well as post-operative functional outcomes[38-40]. Thus, mapping individual-specific cortical networks from brief resting-state scans may offer a practical, non-invasive approach for presurgical epilepsy evaluation in the future[41].

Methodological Considerations

We used healthy participants from the Human Connectome Project and two drug-resistant epilepsy datasets to show that MS-HBM trained on drug-resistant epilepsy outperformed group-average networks and MS-HBM trained on healthy participants despite site, scanner, and pre-processing differences. We studied participants with temporal lobe and extra-temporal lobe epilepsy undergoing presurgical evaluation, and further validation of the

networks in other types of drug-resistant epilepsy (e.g. idiopathic generalized epilepsy) is required.

The esfmri dataset contained unique data that allowed us to validate the quality of individual-specific networks using evoked activity during concurrent intracranial electrical stimulation. However, it must be noted that the esfmri dataset did not contain true post-operative resting-state fMRI data and we concatenated blocks of fMRI data when no stimulation was applied to simulate resting-state fMRI scans[17]. When comparing functional connectivity within the same participants across pre-operative and post-operative data, we identified differences that may reflect either (1) the effects of neurosurgery; (2) distortion in the fMRI signal due to the implanted electrodes; or (3) scanner differences. Thus, we tested the quality of individual-specific networks estimated using both the pre-operative and post-operative data and showed that individual-specific networks estimated by NIH MS-HBM performed the best regardless of the data used.

Conclusion

We present a method for estimating reliable, high-quality individual-specific cortical networks in drug-resistant epilepsy and showed that individual-specific language network topography predicts task-based language dominance. Our model is publicly available (github link), which may be used to estimate individual-specific cortical networks and predict language dominance in a new individual with drug-resistant epilepsy given approximately 10 minutes of single-session resting-state fMRI data.

Funding

References

1.   Szaflarski JP, Gloss D, Binder JR, et al. Practice guideline summary: Use of fMRI in the presurgical evaluation of patients with epilepsy: Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. Jan 24 2017;88(4):395-402. doi:10.1212/wnl.0000000000003532

2.   Lado FA, Ahrens SM, Riker E, et al. Guidelines for Specialized Epilepsy Centers: Executive Summary of the Report of the National Association of Epilepsy Centers Guideline Panel. *Neurology*. Feb 2024;102(4):e208087. doi:10.1212/WNL.0000000000208087

3.   Binder JR, Swanson SJ, Hammeke TA, et al. Determination of language dominance using functional MRI: a comparison with the Wada test. *Neurology*. Apr 1996;46(4):978-84. doi:10.1212/wnl.46.4.978

4.   Binder JR, Swanson SJ, Hammeke TA, Sabsevitz DS. A comparison of five fMRI protocols for mapping speech comprehension systems. *Epilepsia*. Dec 2008;49(12):1980-97. doi:10.1111/j.1528-1167.2008.01683.x

5.   Gaillard WD, Berl MM, Moore EN, et al. Atypical language in lesional and nonlesional complex partial epilepsy. *Neurology*. Oct 30 2007;69(18):1761-71. doi:10.1212/01.wnl.0000289650.48830.1a

6.   Rolinski R, You X, Gonzalez‐Castillo J, et al. Language lateralization from task‐based and resting state functional MRI in patients with epilepsy. *Human Brain Mapping*. 2020;41(11):3133-3146. doi:10.1002/hbm.25003

7.   Doucet GE, Pustina D, Skidmore C, Sharan A, Sperling MR, Tracy JI. Resting-state functional connectivity predicts the strength of hemispheric lateralization for language processing in temporal lobe epilepsy and normals. *Human Brain Mapping*. 2015;36(1):288-303. doi:10.1002/hbm.22628

8.   Pur DR, Eagleson R, Lo M, Jurkiewicz MT, Andrade A, de Ribaupierre S. Presurgical brain mapping of the language network in pediatric patients with epilepsy using resting-state fMRI. *J Neurosurg Pediatr*. Jan 8 2021;27(3):259-268. doi:10.3171/2020.8.PEDS20517

9.   Phillips NL, Shatil AS, Go C, Robertson A, Widjaja E. Resting-State Functional MRI for Determining Language Lateralization in Children with Drug-Resistant Epilepsy. *AJNR Am J Neuroradiol*. Jul 2021;42(7):1299-1304. doi:10.3174/ajnr.A7110

10.   Yeo BT, Krienen FM, Sepulcre J, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*. Sep 2011;106(3):1125-65. doi:10.1152/jn.00338.2011

11.   Kong R, Li J, Orban C, et al. Spatial Topography of Individual-Specific Cortical Networks Predicts Human Cognition, Personality, and Emotion. *Cereb Cortex*. Jun 1 2019;29(6):2533-2551. doi:10.1093/cercor/bhy123

12.   Laumann TO, Gordon EM, Adeyemo B, et al. Functional System and Areal Organization of a Highly Sampled Individual Human Brain. *Neuron*. Aug 5 2015;87(3):657-70. doi:10.1016/j.neuron.2015.06.037

13.   Gordon EM, Laumann TO, Gilmore AW, et al. Precision Functional Mapping of Individual Human Brains. *Neuron*. Aug 16 2017;95(4):791-807 e7. doi:10.1016/j.neuron.2017.07.011

14.   Gordon EM, Laumann TO, Adeyemo B, Petersen SE. Individual Variability of the System-Level Organization of the Human Brain. *Cerebral Cortex*. 2015;doi:10.1093/cercor/bhv239

15.   Laumann TO, Ortega M, Hoyt CR, et al. Brain network reorganisation in an adolescent after bilateral perinatal strokes. *Lancet Neurol*. Apr 2021;20(4):255-256. doi:10.1016/S1474-4422(21)00062-4

16.   Lynch CJ, Elbau IG, Ng T, et al. Frontostriatal salience network expansion in individuals in depression. *Nature*. 2024;doi:10.1038/s41586-024-07805-2

17.   Thompson WH, Nair R, Oya H, et al. A data resource from concurrent intracranial stimulation and functional MRI of the human brain. *Sci Data*. Aug 5 2020;7(1):258. doi:10.1038/s41597-020-00595-y

18.   Du J, DiNicola LM, Angeli PA, et al. Organization of the human cerebral cortex estimated within individuals: networks, global topography, and function. *Journal of neurophysiology*. 2024;131(6):1014-1082. doi:10.1152/jn.00308.2023

19.   Oya H, Howard MA, Magnotta VA, et al. Mapping effective connectivity in the human brain with concurrent intracranial electrical stimulation and BOLD-fMRI. *J Neurosci Methods*. Feb 1 2017;277:101-112. doi:10.1016/j.jneumeth.2016.12.014

20.   Lashkari D, Vul E, Kanwisher N, Golland P. Discovering structure in the space of fMRI selectivity profiles. *Neuroimage*. Apr 15 2010;50(3):1085-98. doi:10.1016/j.neuroimage.2009.12.106

21.   Schaefer A, Kong R, Gordon EM, et al. Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*. 2018;28(9):3095-3114. doi:10.1093/cercor/bhx179

22.   Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex*. Jan 2016;26(1):288-303. doi:10.1093/cercor/bhu239

23.   Pedersen M, Zalesky A. Intracranial brain stimulation modulates fMRI-based network switching. *Neurobiol Dis*. Aug 2021;156:105401. doi:10.1016/j.nbd.2021.105401

24.   Buckner RL, Krienen FM, Yeo BTT. Opportunities and limitations of intrinsic functional connectivity MRI. *Nature Neuroscience*. 2013;16(7):832-837. doi:10.1038/nn.3423

25.   Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. Jan 1 2014;84:320-41. doi:10.1016/j.neuroimage.2013.08.048

26.   Braga RM, Buckner RL. Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. *Neuron*. 2017;95(2):457-471.e5. doi:10.1016/j.neuron.2017.06.038

27.   Laufs H, Hamandi K, Salek‑Haddadi A, Kleinschmidt AK, Duncan JS, Lemieux L. Temporal lobe interictal epileptic discharges affect cerebral activity in "default mode" brain regions. *Human Brain Mapping*. 2006;28(10):1023-1032. doi:10.1002/hbm.20323

28.   Sporns O, Luo C, Qiu C, et al. Disrupted Functional Brain Connectivity in Partial Epilepsy: A Resting-State fMRI Study. *PloS one*. 2012;7(1)doi:10.1371/journal.pone.0028196

29.   Bettus G, Guedj E, Joyeux F, et al. Decreased basal fMRI functional connectivity in epileptogenic networks and contralateral compensatory mechanisms. *Human Brain Mapping*. 2008;30(5):1580-1591. doi:10.1002/hbm.20625

30.   Pittau F, Grova C, Moeller F, Dubeau F, Gotman J. Patterns of altered functional connectivity in mesial temporal lobe epilepsy. *Epilepsia*. 2012;53(6):1013-1023. doi:10.1111/j.1528-1167.2012.03464.x

31.   Valdes-Sosa PA, Liao W, Zhang Z, et al. Altered Functional Connectivity and Small-World in Mesial Temporal Lobe Epilepsy. *PloS one*. 2010;5(1)doi:10.1371/journal.pone.0008525

32.   Li R, Wang H, Wang L, et al. Shared and distinct global signal topography disturbances in subcortical and cortical networks in human epilepsy. *Human Brain Mapping*. 2020;42(2):412-426. doi:10.1002/hbm.25231

33.   Englot DJ, Konrad PE, Morgan VL. Regional and global connectivity disturbances in focal epilepsy, related neurocognitive sequelae, and potential mechanistic underpinnings. *Epilepsia*. 2016;57(10):1546-1557. doi:10.1111/epi.13510

34.   van Diessen E, Diederen SJH, Braun KPJ, Jansen FE, Stam CJ. Functional and structural brain networks in epilepsy: What have we learned? *Epilepsia*. 2013;54(11):1855-1865. doi:10.1111/epi.12350

35.   Robert S, Granovetter MC, Patterson C, Behrmann M. Hemispheric functional organization, as revealed by naturalistic neuroimaging, in pediatric epilepsy patients with cortical resections. *Proc Natl Acad Sci U S A*. Jul 9 2024;121(28):e2317458121. doi:10.1073/pnas.2317458121

36.   Austermuehle A, Cocjin J, Reynolds R, et al. Language functional MRI and direct cortical stimulation in epilepsy preoperative planning. *Ann Neurol*. Apr 2017;81(4):526-537. doi:10.1002/ana.24899

37.   Ankeeta A, Zhang Q, Javidi SS, Modi S, Sperling MR, Tracy JI. Individualized Functional and Structural Language Lateralities in Temporal Lobe Epilepsy and Their Impact on Memory. *Hum Brain Mapp*. Jul 2025;46(10):e70250. doi:10.1002/hbm.70250

38.   Akbarian B, Sainburg LE, Janson A, et al. Association Between Postsurgical Functional Connectivity and Seizure Outcome in Patients With Temporal Lobe Epilepsy. *Neurology*. 2024;103(7)doi:10.1212/wnl.0000000000209816

39.   Bettus G, Bartolomei F, Confort-Gouny S, et al. Role of resting state functional connectivity MRI in presurgical investigation of mesial temporal lobe epilepsy. *Journal of Neurology, Neurosurgery & Psychiatry*. 2010;81(10):1147-1154. doi:10.1136/jnnp.2009.191460

40.   Boerwinkle VL, Mohanty D, Foldes ST, et al. Correlating Resting-State Functional Magnetic Resonance Imaging Connectivity by Independent Component Analysis-Based Epileptogenic Zones with Intracranial Electroencephalogram Localized Seizure Onset Zones and Surgical Outcomes in Prospective Pediatric Intractable Epilepsy Study. *Brain Connect*. Sep 2017;7(7):424-442. doi:10.1089/brain.2016.0479

41.   Boerwinkle VL, Mirea L, Gaillard WD, et al. Resting-state functional MRI connectivity impact on epilepsy surgery plan and surgical candidacy: prospective clinical work. *J Neurosurg Pediatr*. Mar 20 2020:1-8. doi:10.3171/2020.1.PEDS19695

42.   Van Essen DC, Ugurbil K, Auerbach E, et al. The Human Connectome Project: a data acquisition perspective. *Neuroimage*. Oct 1 2012;62(4):2222-31. doi:10.1016/j.neuroimage.2012.02.018

43.   Smith SM, Beckmann CF, Andersson J, et al. Resting-state fMRI in the Human Connectome Project. *Neuroimage*. Oct 15 2013;80:144-68. doi:10.1016/j.neuroimage.2013.05.039

44.   Holmes AJ, Hollinshead MO, O'Keefe TM, et al. Brain Genomics Superstruct Project initial data release with structural, functional, and behavioral measures. *Sci Data*. 2015;2:150031. doi:10.1038/sdata.2015.31

45.   Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage*. Feb 1999;9(2):179-94. doi:10.1006/nimg.1998.0395

46.   Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage*. Oct 2002;17(2):825-41. doi:10.1016/s1053-8119(02)91132-8

47.   Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *NeuroImage*. 2003;20(2):870-888. doi:10.1016/s1053-8119(03)00336-7

48.   DuPre E, Salo T, Ahmed Z, et al. TE-dependent analysis of multi-echo fMRI with tedana. *Journal of Open Source Software*. 2021;6(66)doi:10.21105/joss.03669

49.   Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based registration. *Neuroimage*. Oct 15 2009;48(1):63-72. doi:10.1016/j.neuroimage.2009.06.060

50.   Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*. 2004;23 Suppl 1:S208-19. doi:10.1016/j.neuroimage.2004.07.051

51.   Esteban O, Markiewicz CJ, Blair RW, et al. fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature Methods*. 2018;16(1):111-116. doi:10.1038/s41592-018-0235-4

52.   Gorgolewski K, Burns CD, Madison C, et al. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Front Neuroinform*. 2011;5:13. doi:10.3389/fninf.2011.00013

53.   Woolrich MW, Behrens TE, Smith SM. Constrained linear basis sets for HRF modelling using Variational Bayes. *Neuroimage*. Apr 2004;21(4):1748-61. doi:10.1016/j.neuroimage.2003.12.024

Figure Legends and Tables

A) Overview of analysis and population flowcharts

1. Model development

65 NIH participants drug-resistant epilepsy

17 participants <2 rs-fMRI runs

48 participants ≥2 rs-fMRI runs

Model Training: 34 (70.8%)

Model Testing: 14 (29.2%)

2. External validity

26 esfmri participants drug-resistant epilepsy

pre-operative data: 11 participants ≥2 rs-fMRI runs

post-operative data: 17 participants ≥2 es-fMRI runs

3. Language laterality

65 NIH participants drug-resistant epilepsy

8 participants no rs-fMRI

57 participants ≥1 rs-fMRI run

14 participants no task-fMRI

43 participants Language test

B) Model training for estimating individual-specific networks

HCP group-average Du group-average NIH group-average

Visual A	Dorsal Attn A
Visual B	Dorsal Attn B
Somatomotor A	Sal/Vent Attn A
Somatomotor B	Sal/Vent Attn B
Auditory	Language A
Limbic A	Lang/Def B
Limbic B	Default A
Control	

Train MS-HBM (HCP 40 participants)

Train MS-HBM (HCP 40 participants)

Train MS-HBM (NIH 34 participants)

Estimate Individual-specific Networks

Figure 1: Study overview and population flow chart. (A) We developed a MS-HBM model for estimating individual-specific networks in drug-resistant epilepsy, validated and tested the model in an independent dataset, and showed that the topography of individual-specific language networks was able to predict task-based language dominance. (B) Group-average networks from healthy participants of the HCP dataset, the Du et al atlas, or drug-resistant epilepsy participants from the NIH dataset were used to train different MS-HBM models. 15-

network labels were shown for NIH group-average networks. Networks in drug-resistant epilepsy appeared more focal compared to healthy participants

Figure 2: Individual-specific resting-state cortical networks in drug-resistant epilepsy. (A-B) Individual-specific networks of two representative NIH participants estimated using NIH MS-HBM. (C) Language network boundaries of both participants showing functional connectivity from identical seeds placed in the inferior frontal and lateral temporal region to different language networks in the two subjects. (D) Dice similarity coefficient for the 15-networks across all 57 NIH participants. Higher dice coefficients (yellow) indicate greater inter-subject topographic similarity, while lower dice coefficients (dark red) indicate higher variability. Inter-subject variability in drug-resistant epilepsy is higher in association than sensory networks.

A) NIH test set (14 participants)

		HCP		Du		NIH	
		avg	ind	avg	ind	avg	ind
HCP	avg		p<0.001	p=0.010	p<0.001	p<0.001	p<0.001
HCP	ind	p<0.001		p<0.001	p=0.004	p=0.871	p<0.001
Du	avg	p=0.010	p<0.001		p<0.001	p<0.001	p<0.001
Du	ind	p<0.001	p=0.004	p<0.001		p=0.794	p<0.001
NIH	avg	p<0.001	p=0.871	p<0.001	p=0.794		p=0.003
NIH	ind	p<0.001	p<0.001	p<0.001	p<0.001	p=0.003	

B) esfmri pre-operative data (11 participants)

		HCP		Du		NIH	
		avg	ind	avg	ind	avg	ind
HCP	avg		p<0.001	p=0.002	p<0.001	p<0.001	p<0.001
HCP	ind	p<0.001		p<0.001	p=0.808	p=0.001	p<0.001
Du	avg	p=0.002	p<0.001		p<0.001	p<0.001	p<0.001
Du	ind	p<0.001	p=0.808	p<0.001		p=0.001	p<0.001
NIH	avg	p<0.001	p=0.001	p<0.001	p=0.001		p<0.001
NIH	ind	p<0.001	p<0.001	p<0.001	p<0.001	p<0.001	

C) esfmri post-operative data (17 participants)

		HCP		Du		NIH	
		avg	ind	avg	ind	avg	ind
HCP	avg		p<0.001	p<0.001	p<0.001	p<0.001	p<0.001
HCP	ind	p<0.001		p<0.001	p=0.114	p=0.370	p<0.001
Du	avg	p<0.001	p<0.001		p<0.001	p<0.001	p<0.001
Du	ind	p<0.001	p=0.114	p<0.001		p=0.286	p<0.001
NIH	avg	p<0.001	p=0.370	p<0.001	p=0.286		p<0.001
NIH	ind	p<0.001	p<0.001	p<0.001	p<0.001	p<0.001	

Figure 3: Resting-state homogeneity of group-average and individual-specific networks.

Individual-specific networks estimated using NIH MS-HBM outperformed group-average

networks or MS-HBM trained on healthy participants. Results were consistent across the (A) NIH test set, (B) esfmri pre-operative data, and (C) esfmri post-operative data. Six different networks were evaluated: (1) group-average and (2) individual-specific networks estimated from the HCP data; (3) group-average and (4) individual-specific networks estimated from the Du atlas; (5) group-average and (6) individual-specific networks estimated from the NIH data. Left: boxplots of median resting-state homogeneity. Right: t-test p-values comparing pairs of approaches. Green indicates that the approach on the row performed significantly better than the column, red indicates worse, and grey indicates non-significant differences, after correcting for multiple comparisons using false discovery rate of q < 0.05.

A) esfmri post-operative data (17 participants)

		HCP		Du		NIH	
		avg	ind	avg	ind	avg	ind
HCP	avg		p<0.001	p=0.132	p=0.007	p=0.012	p=0.004
HCP	ind	p<0.001		p=0.026	p=0.948	p=0.362	p=0.420
Du	avg	p=0.132	p=0.026		p=0.004	p=0.050	p=0.005
Du	ind	p=0.007	p=0.948	p=0.004		p=0.351	p=0.337
NIH	avg	p=0.012	p=0.362	p=0.050	p=0.351		p=0.032
NIH	ind	p=0.004	p=0.420	p=0.005	p=0.337	p=0.032	

B) sub-339 run-01 electrical stimulation evoked activity

Figure 4: Electrical-stimulation inhomogeneity and evoked cortical activation map. Individual-specific networks estimated using NIH MS-HBM performed the best in electrical-stimulation inhomogeneity and network boundaries aligned better to electrical-stimulation evoked activity compared to group-average networks. (A) Left shows boxplots of median electrical stimulation inhomogeneity across the six networks: (1) group-average and (2) individual-specific networks estimated from the HCP data; (3) group-average and (4) individual-specific networks estimated from the Du atlas; (5) group-average and (6) individual-specific networks estimated from the NIH data. Right: t-test p-values comparing pairs of approaches. Green indicates that the approach on the row performed significantly better than the column, red indicates worse, and grey indicates non-significant differences, after correcting for multiple comparisons using false discovery rate of q < 0.05. (B) Generalised linear model of electrical-stimulation during fMRI showing z-score normalised beta-coefficients overlayed with the NIH group-average network boundaries on the left and the participant's individual-specific network boundaries on the right. White arrowheads show evoked activity that aligned better with individual-specific networks than group-average networks.

Figure 5: Individual-specific language network topography predicts task-based language dominance. (A) Boxplots of language laterality index for participants with left, bilateral, or

right language dominance derived from task fMRI, comparing individual-specific networks

estimated from HCP MS-HBM, Du MS-HBM, or NIH MS-HBM. (B) Receiver operating

curves and area-under-the-curve statistic for predicting language dominance using

individual-specific networks estimated by NIH MS-HBM.

Supplementary Methods

Participants and Datasets

The HCP dataset consisted of 40 participants from the Human Connectome Project (HCP)

S900 release. The mean age was 29.0 ± 4.0 years, there were 16 (40.0%) male participants,

and 12 (30.0%) participants self-reported as non-White and/or Hispanic race. Each subject

had one 0.7mm isotropic T1 structural scan and four runs of 2mm isotropic resting-state

fMRI with a TR of 0.72s and lasting 14min 33s each. All imaging data were collected on a

custom-made Siemens 3 T Skyra scanner using a multiband sequence. Details of the data

collection may be found elsewhere[1,2].

The NIH dataset consisted of 65 participants with drug-resistant epilepsy undergoing

presurgical evaluation at the NIH epilepsy center[3] (Table S1). The mean age of participants

was 31.9 ± 11.2 years, and there were 33 (50.8%) male participants. There were 42 (64.6%)

participants with temporal lobe epilepsy, and 23 (35.4%) participants with extra-temporal

lobe epilepsy. All participants gave informed written consent under a research protocol

approved by the NIH Combined Neurosciences Institutional Review Board[3]. Each participant

had one 0.8 x 0.75 x 0.75mm T1 and T2 structural scans, up to four runs of 3mm isotropic

resting-state fMRI with a TR of 2.5s and lasting 6min 25s each, and a 3mm isotropic auditory

description decision language task fMRI run with a TR of 2.0s and lasting 5min. All imaging

data were collected on a 3.0 Tesla scanner at the NIH Nuclear Magnetic Resonance Center

using a multiband sequence on either a Siemens Skyra scanner or a GE Discovery scanner.

Details of the data collection may be found elsewhere[3].

The esfmri dataset consisted of 26 participants with drug-resistant epilepsy undergoing

presurgical evaluation at the University of Iowa[4] (Table S2). The mean age of participants

was 35.8 ± 11.5 years, and there were 17 (65.4%) male participants. There were 19 (73.1%)

participants with temporal lobe epilepsy, 6 (23.1%) participants with extra-temporal lobe

epilepsy, and 1 (3.8%) participant where the seizure onset zone was not determined. Intracranial electrodes were implanted to localize the epileptogenic zone, and were determined by clinical criteria. All participants gave informed written consent under a research protocol approved by the University of Iowa Institutional Review Board[4]. Each participant had one 1.0 x 1.0 x 0.8mm T1 structural scan, up to five runs of 3.4 x 3.4 x 4.0mm pre-operative resting state fMRI with a TR of 2.26s lasting 4.8min each, and up to eleven runs of 3mm isotropic electrical-stimulation fMRI with a TR of 3.0s lasting approximately 10min each. For each post-operative fMRI scan, bipolar intracranial electrical stimulation using depth electrodes was interleaved between echo planar imaging volume acquisition during a 100ms period where there was no scanner radiofrequency or gradient switching[4]. Stimulation was blocked with alternating durations of approximately 30 seconds of STIM-ON and STIM-OFF each. No cognitive task was given to the participant, and no behavioural effects were evoked during the stimulation period. Details of the data collection may be found elsewhere[4,5].

No.	Age	Sex	Hand	Seizure Onset Zone	Task Laterality	MRI	Valid Runs
1	26-30	F	R	Right Mesial Temporal Sclerosis	L	Siemens Skyra	2
2	20-25	F	R	Right Parietal	U	GE Discovery MR750	2
3	36-40	M	R	Left > Right Temporal	B	GE Discovery MR750	1
4	30-35	M	R	Right > Left Medial Temporal	L	Siemens Skyra	2
5	20-25	F	R	Left Mesial Temporal Sclerosis	U	GE Discovery MR750	1
6	36-40	M	R	Right Mesial Temporal Sclerosis	L	Siemens Skyra	0
7	30-35	F	R	Right Temporoparietal	L	GE Discovery MR750	2
8	26-30	F	R	Right Temporal	B	GE Discovery MR750	2
9	30-35	F	R	Right Mesial Temporal Sclerosis	L	Siemens Skyra	3
10	20-25	M	L	Left Mesial Temporal Sclerosis	L	GE Discovery MR750	2

11	36-40	F	R	Left Temporal	R	GE Discovery MR750	2
12	56-60	F	R	Right Frontal	L	GE Discovery MR750	2
13	16-20	M	R	Right Frontal	L	GE Discovery MR750	1
14	36-40	F	R	Left Frontal/ Insula	U	GE Discovery MR750	1
15	20-25	M	R	Right Frontal	U	GE Discovery MR750	2
16	30-35	M	R	Left Temporal Periventricular Nodular Heterotopia	B	Siemens Skyra	3
17	36-40	M	L	Left > Right Frontal	U	GE Discovery MR750	1
18	36-40	F	L	Left Mesial Temporal Sclerosis	R	GE Discovery MR750	2
19	20-25	M	R	Left Frontal/ Insula	L	GE Discovery MR750	2
20	36-40	M	R	Right Frontal	U	GE Discovery MR750	2
21	30-35	F	R	Left Mesial Temporal Sclerosis	L	Siemens Skyra	1
22	16-20	F	R	Left Frontotemporal	R	GE Discovery MR750	2
23	51-55	M	R	Left Temporal	L	Siemens Skyra	3
24	26-30	F	R	Right Temporal	R	GE Discovery MR750	2
25	30-35	F	R	Left Temporal	L	Siemens Skyra	2
26	16-20	F	R	Right Frontal	L	GE Discovery MR750	2
27	36-40	F	R	Left Temporal	B	GE Discovery MR750	0
28	56-60	F	R	Left Mesial Temporal Sclerosis	L	Siemens Skyra	3
29	26-30	M	B	Bilateral Frontal	L	GE Discovery MR750	2
30	30-35	F	R	Left > Right Temporal	L	Siemens Skyra	3
31	26-30	F	L	R Insula > Frontal	R	GE Discovery MR750	2
32	20-25	F	B	Right Polymicrogyria, Bilateral Periventricular Nodular Heterotopia	B	GE Discovery MR750	2
33	51-55	M	R	Left Frontal Schizencephaly	U	GE Discovery MR750	0
34	26-30	F	R	Left Mesial Temporal Sclerosis	B	Siemens Skyra	3
35	41-45	M	R	Right Frontal	L	Siemens Skyra	3

36	26-30	F	R	Right Mesial Temporal Sclerosis	U	Siemens Skyra	1
37	30-35	M	R	Right Parietal	L	Siemens Skyra	3
38	41-45	F	L	Right Frontal	B	Siemens Skyra	3
39	16-20	F	R	Right Temporal	U	Siemens Skyra	0
40	56-60	M	R	Left Temporal	B	Siemens Skyra	1
41	20-25	M	R	Left Temporal	L	Siemens Skyra	3
42	36-40	F	R	Right Temporal	U	Siemens Skyra	3
43	20-25	M	L	Right Temporal Arteriovenous Malformation	L	Siemens Skyra	3
44	16-20	M	L	Left Parietal	B	Siemens Skyra	3
45	20-25	M	R	Left Frontal	L	Siemens Skyra	2
46	26-30	M	R	Left Temporal Low Grade Glioma	L	Siemens Skyra	3
47	20-25	F	R	Left Frontal	L	Siemens Skyra	3
48	26-30	M	R	Right Parieto-occipital	L	Siemens Skyra	0
49	30-35	M	R	Left Mesial Temporal Sclerosis	U	Siemens Skyra	3
50	20-25	M	R	Right Frontal	L	Siemens Skyra	4
51	30-35	M	L	Left Temporal	U	Siemens Skyra	3
52	20-25	F	R	Right Temporal	L	Siemens Skyra	2
53	26-30	F	R	Left Temporal	U	Siemens Skyra	3
54	56-60	M	R	Right Temporal	L	Siemens Skyra	1
55	6-10	M	R	Left Frontal	L	Siemens Skyra	3
56	26-30	M	R	Bilateral Temporal	U	Siemens Skyra	3
57	36-40	M	R	Left Temporal	U	Siemens Skyra	3
58	20-25	M	R	Right Temporal	L	Siemens Skyra	3
59	51-55	F	R	Right Temporal	L	Siemens Skyra	3
60	30-35	F	R	Right Temporal	R	Siemens Skyra	0
61	20-25	F	R	Right Parietal/ Temporal	L	Siemens Skyra	0
62	26-30	M	R	Left Temporal	U	Siemens Skyra	3
63	26-30	M	R	Right Temporal	L	Siemens Skyra	2
64	51-55	F	R	Left Temporal	L	Siemens Skyra	0
65	30-35	M	R	Left Temporal	B	Siemens Skyra	2

Table S1: Demographic, MRI, and epilepsy characteristics of participants from the NIH dataset.

No.	Age	Sex	Hand	Seizure Onset Zone	pre-op MRI	pre-op Runs	post-op MRI	post-op Runs
1	46-50	F	L	Left mesial temporal lobe, Left frontal lobe	Siemens Trio	0	Siemens Trio	5
2	31-35	M	R	Right anterior frontal lobe	Siemens Trio	0	Siemens Trio	2
3	46-50	F	R	Left mesial temporal lobe	Siemens Trio	0	Siemens Skyra	1
4	31-35	F	R	Right mesial temporal lobe, Left mesial temporal encephalomalacia	Siemens Trio	1	Siemens Skyra	4
5	26-30	M	R	Left insula	Siemens Trio	1	Siemens Skyra	1
6	26-30	F	R	Bilateral mesial temporal lobe	Siemens Trio	5	Siemens Skyra	0
7	31-35	F	R	Right mesial temporal lobe	Siemens Trio	4	Siemens Skyra	7
8	46-50	F	R	Right hippocampus	Siemens Trio	0	Siemens Skyra	0
9	41-45	M	L	Left occipital lobe	Siemens Trio	5	Siemens Skyra	4
10	31-35	M	R	Left mesial temporal lobe	Siemens Trio	1	Siemens Skyra	5
11	36-40	M	L	Right temporal pole, Left temporal base	Siemens Trio	5	Siemens Skyra	10
12	31-35	M	R	Bilateral mesial temporal lobe	Siemens Trio	3	Siemens Skyra	2
13	41-45	M	R	Not determined	None	-	Siemens Skyra	6
14	31-35	M	B	Left frontal cystic mass	GE Discovery	2	Siemens Skyra	0
15	36-40	M	R	Left mesial temporal lobe	None	-	Siemens Skyra	1
16	26-30	M	R	Right mesial temporal lobe	None	-	Siemens Skyra	8
17	31-35	M	R	Left temporal pole	GE Discovery	2	Siemens Skyra	0
18	46-50	F	R	Right mesial temporal lobe	GE Discovery	3	Siemens Skyra	2
19	36-40	M	R	Right mesial temporal lobe, Right frontal pole	GE Discovery	2	Siemens Skyra	0

20	21-25	M	R	Right amygdala	None	-	Siemens Skyra	7
21	11-15	M	R	Left superior frontal gyrus cavernoma	None	-	Siemens Skyra	3
22	21-25	F	R	Right mesial temporal lobe, Possible right frontal base	GE Discovery	3	Siemens Skyra	6
23	56-60	M	R	Left mesial temporal lobe	GE Discovery	1	Siemens Skyra	2
24	56-60	F	R	Left mesial temporal lobe	None	-	Siemens Skyra	0
25	16-20	M	R	Left frontal encephalomalacia	GE Discovery	4	Siemens Skyra	3
26	21-25	M	L	Right mesial temporal lobe	GE Discovery	5	Siemens Skyra	2

Table S2: Demographic, MRI, and epilepsy characteristics of participants from the esfmri dataset.

Pre-processing of Imaging Data

The HCP dataset was released with data already pre-processed (smoothed using a 2mm full-width half-maximum kernel, denoised with ICA-FIX and aligned with MSMAll) and were detailed elsewhere[6,7]. The preprocessed fMRI data was projected onto the FreeSurfer fsaverage6 surface space. Additional nuisance regression, censoring, and spatial smoothing was computed using the surface-based pipeline of Yeo et al[6].

Pre-processing of the NIH dataset followed the surface-based pipeline of Yeo et al[8,9]. Structural data were processed using recon-all (FreeSurfer 7.1.1)[10]. Resting-state fMRI data were processed with the following: (i) removal of first 4 frames, (ii) slice time correction using FSL[11], (iii) motion correction and outlier detection using rigid body translation and rotation[11], (iv) susceptibility-induced spatial distortion correction[12], (v) multi-echo denoising[13], (vi) boundary-based registration[14], (vii) regression of the global, white matter, ventricular signals,

6 motion parameters, and their temporal derivatives, (viii) censored frames were interpolated with the Lomb-Scargle periodogram[15], (ix) bandpass filtering of 0.009Hz – 0.08Hz, and lastly, (x) the data was projected onto FreeSurfer fsaverage6 surface space and smoothed using a 6mm full-width half-maximum kernel. Outlier detection using framewise displacement (FD) and voxel-wise differentiated signal variance (DVARS) were computed using fsl_motion_outliers[16]. Frames that exceeded a threshold of 0.2mm FD or 50 DVARS as well as one frame before and two frames after were censored. Segments of data lasting less than five contiguous frames were also censored[17]. After pre-processing, 8 participants did not have more than one valid resting-state run, 9 participants had one run, 23 participants had two runs, 24 participants had three runs, and one participant has four runs.

The esfmri dataset was released with data pre-processed using fMRIPrep 1.5.1rc1[18] (based on Nipype 1.2.0[19]) and was described in detail elsewhere[4]. Frames that exceeded a threshold of 0.2mm FD or 1.2 standardized derivative of the root-mean-square variance were censored. Runs that had more than 50% motion outliers were discarded. Linear regression was used to regress out 6 motion parameters, the whole brain, white matter, and cerebrospinal fluid regressors and their derivatives. The fMRI time-series were then resampled to the FreeSurfer fsaverage6 space and spatially smoothed using a 6mm full-width half-maximum kernel (FreeSurfer 5.3.0)[10]. After pre-processing, there were 11 participants that had at least two valid pre-operative runs and 17 participants that had at least two valid post-operative runs.

Derivation of group-average networks and training of MS-HBM for estimating individual-specific networks

Our approach has been described previously[6,8]. Briefly, the pre-processed fMRI data were projected onto fsaverage6 surface meshes that consisted of 40,962 vertices per hemisphere. The connectivity profile of a cortical region (vertex) was defined to be its functional coupling

to 1175 regions of interest (ROIs), which consisted of single vertices uniformly distributed across the fsaverage6 surface meshes. For each run, the Pearson's correlation between the fMRI time series at each spatial location (81,924 vertices) and the 1175 ROIs were computed. The 81,924 x 1175 correlation matrix was then binarized by keeping the top 10% of correlations. To obtain group-average networks, each vertex's connectivity profiles were averaged across all resting-state fMRI runs of all participants. The averaged connectivity profiles were then clustered using a mixture of von Mises-Fisher distributions to obtain the 15-network group-average for each dataset[8,20].

We developed group-average networks and trained MS-HBM using either 40 participants from the Human Connectome Project (HCP dataset)[6], or 34 participants with drug-resistant epilepsy from the National Institutes of Health (NIH dataset). In addition, we used a group-average atlas of 15 densely sampled healthy participants from Du et al[7] to train MS-HBM on the 40 participants from the HCP dataset. Model parameters estimated included the inter-subject functional connectivity variability, intra-subject functional connectivity variability, spatial smoothness prior, and the inter-subject spatial variability prior[6]. Using each of the three MS-HBM models, individual-specific networks were estimated for each participant with drug-resistant epilepsy. For all individual-specific networks estimated, the weight of the group spatial prior was taken to be 80, and the weight of the smoothness prior was taken to be 10[6].

Generalized linear models of intracranial electrical stimulation during fMRI scans

Generalized linear models were used to quantify the effects of electrical stimulation on cortical activity. A boxcar function of the stimulus duration (varies from 50-90 ms) for each post-operative run was computed[4]. The hemodynamic response function was then modelled by convolving the boxcar function with an optimised set of three gamma basis functions generated using FSL Linear Optimal Basis Sets[21] to account for any non-canonical

haemodynamic response function shapes caused by the stimulation (parameters used were m1 = 0-2s; m2 = 3-6s; m3 = 3-8s; m4 = 3-8s; c = 0-0.3). The amplitude of all basis functions were normalized to peak values of 1 by dividing by the maximum amplitude before computing the haemodynamic response function.

The generalized linear models were computed using freesurfer's mri_glmfit on the fsaverage6 data and corrected at the vertex-level for the lag one autoregression correlation matrix averaged across all valid runs. The vertex-level unthresholded beta-coefficients of the contrast consisting of all three haemodynamic response function shapes were then Z-score normalised by subtracting the mean and dividing by the standard deviation across all beta-coefficients of both hemispheres.

References

1.   Van Essen DC, Ugurbil K, Auerbach E, et al. The Human Connectome Project: a data acquisition perspective. *Neuroimage*. Oct 1 2012;62(4):2222-31. doi:10.1016/j.neuroimage.2012.02.018

2.   Smith SM, Beckmann CF, Andersson J, et al. Resting-state fMRI in the Human Connectome Project. *Neuroimage*. Oct 15 2013;80:144-68. doi:10.1016/j.neuroimage.2013.05.039

3.   Rolinski R, You X, Gonzalez‐Castillo J, et al. Language lateralization from task‐based and resting state functional MRI in patients with epilepsy. *Human Brain Mapping*. 2020;41(11):3133-3146. doi:10.1002/hbm.25003

4.   Thompson WH, Nair R, Oya H, et al. A data resource from concurrent intracranial stimulation and functional MRI of the human brain. *Sci Data*. Aug 5 2020;7(1):258. doi:10.1038/s41597-020-00595-y

5.   Oya H, Howard MA, Magnotta VA, et al. Mapping effective connectivity in the human brain with concurrent intracranial electrical stimulation and BOLD-fMRI. *J Neurosci Methods*. Feb 1 2017;277:101-112. doi:10.1016/j.jneumeth.2016.12.014

6.   Kong R, Li J, Orban C, et al. Spatial Topography of Individual-Specific Cortical Networks Predicts Human Cognition, Personality, and Emotion. *Cereb Cortex*. Jun 1 2019;29(6):2533-2551. doi:10.1093/cercor/bhy123

7.   Du J, DiNicola LM, Angeli PA, et al. Organization of the human cerebral cortex estimated within individuals: networks, global topography, and function. *Journal of neurophysiology*. 2024;131(6):1014-1082. doi:10.1152/jn.00308.2023

8.   Yeo BT, Krienen FM, Sepulcre J, et al. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol*. Sep 2011;106(3):1125-65. doi:10.1152/jn.00338.2011

9.   Holmes AJ, Hollinshead MO, O'Keefe TM, et al. Brain Genomics Superstruct Project initial data release with structural, functional, and behavioral measures. *Sci Data*. 2015;2:150031. doi:10.1038/sdata.2015.31

10.   Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage*. Feb 1999;9(2):179-94. doi:10.1006/nimg.1998.0395

11.   Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage*. Oct 2002;17(2):825-41. doi:10.1016/s1053-8119(02)91132-8

12.   Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *NeuroImage*. 2003;20(2):870-888. doi:10.1016/s1053-8119(03)00336-7

13.   DuPre E, Salo T, Ahmed Z, et al. TE-dependent analysis of multi-echo fMRI with tedana. *Journal of Open Source Software*. 2021;6(66)doi:10.21105/joss.03669

14.   Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based registration. *Neuroimage*. Oct 15 2009;48(1):63-72. doi:10.1016/j.neuroimage.2009.06.060

15.   Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. Jan 1 2014;84:320-41. doi:10.1016/j.neuroimage.2013.08.048

16.   Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*. 2004;23 Suppl 1:S208-19. doi:10.1016/j.neuroimage.2004.07.051

17.   Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cereb Cortex*. Jan 2016;26(1):288-303. doi:10.1093/cercor/bhu239

18.   Esteban O, Markiewicz CJ, Blair RW, et al. fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature Methods*. 2018;16(1):111-116. doi:10.1038/s41592-018-0235-4

19.   Gorgolewski K, Burns CD, Madison C, et al. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Front Neuroinform*. 2011;5:13. doi:10.3389/fninf.2011.00013

20.   Lashkari D, Vul E, Kanwisher N, Golland P. Discovering structure in the space of fMRI selectivity profiles. *Neuroimage*. Apr 15 2010;50(3):1085-98. doi:10.1016/j.neuroimage.2009.12.106

21.   Woolrich MW, Behrens TE, Smith SM. Constrained linear basis sets for HRF modelling using Variational Bayes. *Neuroimage*. Apr 2004;21(4):1748-61. doi:10.1016/j.neuroimage.2003.12.024