Delayed introduction and susceptible variation drive spatial asynchrony in pertussis epidemics

Sang Woo Park[1,*]

1 School of Biological Sciences, Seoul National University, Seoul, Korea

*Corresponding author: sangwoopark@snu.ac.kr

Abstract

Infectious disease outbreaks typically exhibit a large-scale, spatial synchrony, reflecting coupling through shared environmental forcing and human mobility. During South Korea's unprecedented large pertussis outbreak in 2024—2025, however, asynchronous epidemic patterns were observed throughout the country: some regions experienced two epidemic waves, while others regions experienced only the first or second wave. Using high-resolution surveillance data from 252 municipalities and a simple transmission model, we show that heterogeneity in introduction timing and local susceptibility can drive such fine-scale asynchrony even when transmission is spatially homogeneous. These results redefine epidemic synchrony as a condition contingent not only on shared transmission dynamics, but also on variation in initial epidemic conditions.

Introduction

Spatiotemporal variation in epidemic dynamics offers a powerful means for understanding factors that determine the timing and magnitude of epidemic waves across populations [1, 2, 3, 4]. Ecological theory predicts that spatial coupling—via shared environmental forcing and human mobility—can synchronize epidemic dynamics between neighboring locations, sometimes manifesting as travelling waves, where epidemic waves propagate through space [5, 6, 7]. Consistent with this theory, large-scale spatial synchrony has been observed across diverse pathogens, including measles [7], dengue [8], influenza [9], RSV [4], and SARS-CoV-2 [10], highlighting the ubiquity of spatial coupling in shaping epidemic dynamics.

Between 2024 and 2025, an unprecedented large pertussis outbreak occurred in Korea, resulting in more than 50,000 cases by early 2025 (Figure 1) [11]. This outbreak marked a dramatic increase in the disease compared to 292 cases reported in 2023 and 21 cases reported in 2022, despite high vaccine coverage (91.1% of identified 2024 cases had a known vaccination history) [12, 11]. The outbreak consisted of two waves that peaked around late July and December 2024 (Figure 1A) and exhibited large spatial heterogeneity in timing (Figure 1B,C) and magnitude (Figure 1D). For example, both Seoul, the capital city located in the northwest region, and Busan, the second most populated city located at the southeast end, experienced two waves, whereas Daegu, the fourth most populated city located < 100km away from Busan, only experienced the second wave (Figure 1C; see Figure S1 for epidemic dynamics in other regions). This pronounced asynchrony deviates from our expectations based on spatial coupling, raising questions about mechanisms driving this fine-scale spatiotemporal heterogeneity in pertussis epidemic [13].

To address this question, we combine a detailed spatiotemporal surveillance data from 252 municipalities in Korea (Figure 1) with mathematical modeling approaches. We begin by showing that asynchrony in pertussis epidemics, captured by heterogeneous epidemic timing, is associated with differences in introduction delays. In contrast, the analysis of transmission dynamics reveals synchrony in underlying transmission patterns. Integrating these findings into a mathematical model illustrates that the apparent asynchrony can be explained by differences in initial epidemic conditions (i.e., introduction delays and initial susceptibility).

Asynchrony in pertussis epidemics associated with introduction delays

First, to characterize spatial heterogeneity in pertussis waves across Korea, we quantify the center of gravity (Figure 1E), which represents the mean timing of cases [1]. Notably, we find a sharp contrast between early (orange) and late (light blue) waves between many neighboring municipalities (Figure 1E), suggesting spatial asynchrony in pertussis dynamics. To systematically evaluate this apparent asynchrony, we estimated the spatial synchrony function, which captures changes in correlation

Figure 1: **Spatiotempordal dynamics of pertussis outbreak in Korea, 2024—2025.** (A) Total number of reported cases in Korea. (B) Number of reported cases per 100,000 individuals by municipality, ordered by latitude. (C) Number of reported cases per 100,000 individuals in four most populated cities in Korea. (D) Map of pertussis cases per 100,000 individuals between January 2024 and November 2025. (E) Map of center of gravity, which represents the mean timing of cases. (F) Spatial synchrony in logged cases as a function of distance. The red solid line and shaded regions represent the median estimate and the corresponding 95% confidence interval. The dashed horizontal line represents the mean correlation. (G) Relationship between center of gravity and the timing of introduction, defined as the first week when the number of reported cases is greater than 1 in 100,000. The red solid line and shaded regions represent the linear regression fit and the corresponding 95% confidence interval.

in epidemic dynamics as a function of distance [7]. The spatial synchrony function confirms a sharp reduction in cross-correlation with distance (Figure 1F). For example, within 100km of distance, the cross-correlation drops below 0.25. By comparison, pertussis epidemics in the US maintained cross-correlations above 0.2 even at 1,000 km [14], highlighting the fine-scale asynchrony in pertussis epidemics in Korea. Interestingly, the variation in the center of gravity significantly correlates with the introduction timing ($\rho = 0.53$; 95%CI: 0.44—0.62; Figure 1G), suggesting that differences in introduction timing may contribute to the apparent spatial asynchrony.

Figure 2: **Spatial variatoin in the estimates of effective reproduction number.** (A) Each black line represents the estimated effective reproduction number for a given municipality with more than 400 total cases. The red solid line and shaded regions represent the median and 95% quantiles of effective reproduction number estimates. (B) Relationship between correlation coefficient for $\mathcal{R}(t)$ and correlation coefficient for reported cases across all pairwise municipality combinations with more than 400 total cases. (C) Spatial synchrony in $\mathcal{R}(t)$ (red) versus logged cases (black) as a function of distance. The solid line and shaded regions represent the median estimate and the corresponding 95% confidence interval. The dashed horizontal line represents the mean correlation. (D) Estimated multiplicative effects on $\mathcal{R}(t)$ from the regression analysis. The blue shaded violin represents the distribution of intercept effect, where a separate intercept term is estimated for each municipality. The red lines represent the estimated susceptible depletion effect for each municipality. The orange line and shaded regions represent the estimated changes in transmission from a smooth term and the corresponding 95% confidence interval.

Synchrony in temporal changes in pertussis transmission

We then ask whether the observed epidemic asynchrony can be explained by the differences in transmission patterns. To do so, we quantify the effective reproduction number $\mathcal{R}(t)$, which measures the average number of new infections caused by a single infected individual [15, 16]. For this analysis, we focus on municipalities with

more than 400 total cases reported because $\mathcal{R}(t)$ estimates can be unstable when the number of infections are low.

Notably, $\mathcal{R}(t)$ estimates appear to be largely homogeneous across different regions, despite the apparent heterogeneity in the observed epidemic dynamics (Figure 2A). Comparisons of correlation coefficients in $\mathcal{R}(t)$ versus correlation coefficients in reported cases between all pairwise municipality combinations confirm this observation (Figure 2B). In 830 out of 861 municipality combinations, correlation in $\mathcal{R}(t)$ is higher than correlation in cases. This is reflected in the spatial synchrony function for $\mathcal{R}(t)$ (Figure 2C), where the mean cross-correlation for $\mathcal{R}(t)$ ($\rho = 0.71$) is considerably higher than that for cases ($\rho = 0.30$).

Figure 3: **Fine-scale spatiotemporal variation in pertussis epidemic can be explained with a simple SEIR model.** (A) Comparisons between the observed and predicted epidemic dynamics across municipalities with more than 400 total cases, ordered by latitude.. (B) The bar plot represents the distribution of R squared values for model fits. The vertical dashed line represents the median. (C) Relationship between the estimated and predicted center of gravity. The solid line indicated the one-to-one relationship. (D) Relationship between the initial susceptible $S(0)$ and infected $I(0)$ fraction with center of gravity. Points represent the estimates for each municipality. (E) Impact of initial susceptibility on pertussis epidemic dynamics. (F) Estimated time-varying pertussis transmission rate. The red line and shaded regions represent the posterior median and the corresponding 95% credible interval.

To better understand factors driving changes in $\mathcal{R}(t)$, we perform a regression

analysis using a generalized additive model with three covariates [17, 18, 19]: (1) a municipality-level fixed effect allowing each municipality to have its own intercept; (2) a cumulative-cases term capturing the effect of susceptible depletion; and (3) a single smooth term shared across municipalities, which allows for temporal changes in pertussis transmission over time, such as seasonal forcing. First, the smooth term suggests a strong transmission reduction around August and February, which coincide with the timing of school breaks (Figure 2D). Second, the model estimates a significant effect of effect of susceptible depletion (Figure 2D; $p < 0.001$). Notably, this simple regression model explains 84% of variance in $\mathcal{R}(t)$, suggesting that the observed temporal fluctuations in transmission can be largely explained by school-related seasonality and the gradual depletion of susceptible individuals. Taken together, the analysis of center of gravity and $\mathcal{R}(t)$ suggest that differences in introduction timing and susceptible host dynamics are likely main drivers for spatial asynchrony in pertussis epidemics in Korea, rather than variation in seasonal transmission patterns.

Mathematical modeling of fine-scale spatio-temporal pertussis epidemic

To test whether variation in introduction timing and susceptible dynamics alone can explain the observed epidemic patterns, we extended a standard susceptible-exposed-infected-removed (SEIR) model [20] to jointly estimated a single, time-varying transmission term, which is shared across all regions. We also allowed spatial variation in initial susceptible $S(0)$ and infected $I(0)$ fraction as well as the reporting rate ρ. Variation in $S(0)$ captures differences in population-level susceptibility, whereas variation in $I(0)$ effectively captures differences in introduction timing. For this analysis, we also focus on municipalities with more than 400 total cases. The model is fitted using a Bayesian inference software rstan [21].

Our simple transmission model that assume shared transmission rate can capture fine-scale spatiotemporal variation of pertussis outbreak in Korea (Figure 3A) with a median R squared value of 0.86 (95% quantile: 0.71—0.91). Furthermore, we find a near-perfect correlation between the predicted and observed center of gravity (Figure 3C). So what factors determine the timing of epidemic, and therefore the center of gravity? We address this question by simulating our model between April 2024 and April 2025 across plausible ranges of $S(0)$ and $I(0)$.

First, increasing $I(0)$ leads to an earlier epidemic because it permits earlier introduction of the infection to the population (Figure 3D). This is consistent with our earlier observations (Figure 1F). Similarly, increasing $S(0)$ also results in an earlier epidemic. This is because high levels of susceptibility allows for a large first wave, which prevents the second wave through the overcompensatory susceptible depletion (Figure 3E). Finally, we estimate that the large reduction in transmission around late July 2024 and late January 2025 allowed the occurrence of two distinct epidemic waves (Figure 3F). this estimated timing of transmission reduction is consistent with

our estimates from the $\mathcal{R}(t)$ analysis and coincide with the timing of school holidays in Korea.

We validate our model by taking the estimated transmission term and fitting the remaining parameters (i.e., initial infected, initial susceptible, and reporting probability) to data from the remaining 210 municipalities with less than 400 total cases (Figure S2). While model predictions appear consistent with the observed patterns (Figure S2A), we estimate a lower value for the median R squared: 0.58 (95% quantile: 0.04–0.87; Figure S2B). Nonetheless, our analysis suggests that the majority of asynchrony in pertussis epidemics can be explained via the differences in introduction timing and initial susceptibility.

Summary

Understanding mechanisms that drive spatiotemporal variation in epidemic dynamics is a major aim for infectious disease dynamics and control [1, 2, 3, 4, 7, 8, 9] The large pertussis outbreaks in Korea that occurred between 2024 and 2025 presents a unique perspective on a fine-scale spatiotemporal asynchrony heterogeneity in epidemic dynamics: some municipalities experienced two epidemic waves while other places experienced only the first or second without a clear patterns of synchrony. We propose that this variation can be explained by the differences in introduction delays and initial susceptibility, where high levels of susceptibility can cause a large first wave and prevent the second wave through susceptible depletion. Incorporating this idea into a simple transmission model by allowing for a joint estimation of a single transmission rate with variation in initial epidemic conditions confirms our hypothesis. Together, these findings demonstrate that differences in initial epidemic conditions— rather than variation in transmission—can generate fine-scale asynchrony, offering a general framework for understanding the spatiotemporal dynamics of re-emerging infections.

Caveats and future directions

Even though our model explains the observed dynamics reasonably well, there are several limitations to our work. First, our model is purely deterministic and does not account for stochasticity, which is likely an important factor for characterizing the establishment of infections after the initial invasion and the subsequent dynamics, especially in small populations [22, 23, 24]. Second, we estimated that a variation in susceptibility can explain the observed epidemic patterns, but we were unable to identify the mechanism that explains this variation. For example, we hypothesized that past vaccine coverage would explain variation in susceptibility given the role of waning immunity in driving pertussis re-emergence [25, 26, 27], but we did not find a significant correlation (Figure S3).

Our parameter estimates must be interpreted with care as our model relies on several simplifying assumptions. For example, we assumed the same values of basic reproduction number $\mathcal{R}_0$ across all municipalities. Therefore, any regional variation in $\mathcal{R}_0$, driven by differences in contact rates and age distributions would be captured in our estimate of the initial susceptible fraction. In other words, our estimated susceptible fraction should be interpreted as the effective susceptibility that also captures behavioral and sociological factors, beyond immunological differences. We also did not consider an age structure in our model, but age-structured contact patterns play an important role in shaping pertussis transmission [20]. For example, [28] recently suggested transmission among elderly may contributed to the persistence of pertussis in Korea. Despite these limitations, our work provides a quantitative foundation and empirical evidence for understanding epidemic asynchrony.

Conclusion

The continued emergence of new pathogens and re-emergence of vaccine preventable infections, such as pertussis, highlights the importance of identifying hotspots of future outbreaks and developing targeted intervention strategies [29, 30, 31]. The apparent asynchrony in pertussis epidemic in Korea, driven by the differences in introduction timing and population-level susceptibility, underlines the importance of detailed pathogen surveillance for early detection and strategic allocation of prevention and control resources.. Immunological surveillance platforms will be particularly useful for evaluating potential risk of pathogen re-emergence in the future [32, 33]. A detailed understanding of spatial variation in population-level immunity will be crucial for predicting spatiotemporal dynamics of future outbreaks.

Materials and Methods

Data

The weekly surveillance data on pertussis cases across 252 municipalities (1st week of 2024 to the 44th week of 2025) were obtained from a publicly available website, the Infectious Disease Portal, by the Korea Disease Control and Prevention Agency [34]. Population size data as of September 2025 were obtained from publicly available website by the Ministry of the Interior and Safety [35]. Vaccine coverage data were obtained from a publicly available website by the Korea Disease Control and Prevention Agency [36]. Longitude and latitude data for each municipality were obtained from a publicly available GitHub repository [37]. Finally, shape files used for constructing maps were obtained from a publicly available website [38].

Center of gravity

We quantified the center of gravity to characterize variation in the mean timing of epidemic. Typically, the center of gravity is calculated based on annual incidence for recurrent epidemics. Since we are focused on analyzing a single outbreak, we use the entire time series of reported cases C_t to computer the center of gravity for each municipality:

$$\text{Center of gravity} = \frac{\sum_{t=1}^{t_{\max}} C_t \times t}{\sum_{t=1}^{t_{\max}} C_t}. \tag{1}$$

We then calculate the correlation coefficient between the center of gravity and the timing of introduction, defined as the first week when the number of reported cases is greater than 1 in 100,000.

Spatial synchrony of reported cases

We characterized the spatial synchrony of pertussis epidemic, which captures changes in pairwise correlation as a function of distance [7]. We used logged values of reported cases between May 2024 and March 2025, excluding municipalities that had no reported cases ($n = 250$). Spatial synchrony was calculated using the `ncf` package in R [39]. We used 1,000 bootstraps to generate the median estimate and the corresponding 95% confidence intervals.

Effective reproduction number

We calculated the effective reproduction number $\mathcal{R}(t)$ to characterize changes in pertussis transmission over time. First, we took the case time series C_t and fitted a generalized additive model assuming a Poisson error to smooth the time series [18]. Then, we estimated $\mathcal{R}(t)$ using the method of [16]:

$$\mathcal{R}(t) = \frac{i(t)}{\sum_{k=1}^{n} i(t-k)g(k)}, \tag{2}$$

where $i(t)$ is the smoothed incidence and n is the maximum length of generation interval in weeks (assumed to be 5 weeks). The generation-interval distribution $g(k)$ is assumed to follow a gamma distribution with a mean of 9.47 days and a standard deviation of 6.22 days. The reproduction number estimates can be unrealistically high when the number of infections is close to zero. Therefore, we only use estimates between May 2024 and March 2025 throughout the paper. We truncated all $\mathcal{R}(t)$ estimates exceeding 5 to a maximum value of 5.

First, we compared correlation coefficients between $\mathcal{R}(t)$ estimates and those between logged cases across all pairwise municipality combinations (Figure 2B in the main text). For this analysis, we only use data from municipalities with more than 400 total cases. Then, we compared the spatial synchrony for $\mathcal{R}(t)$ estimates and

228 synchrony for logged cases (Figure 2C in the main text). For this analysis, we used
229 $\mathcal{R}(t)$ estimates from all municipalities without any exclusion.

230   Finally, we fitted a generalized additive model to logged values of $\mathcal{R}(t)$ to quantify
231 the effect of susceptible depletion and temporal variation in intrinsic transmission
232 [17, 18, 19]:

$$\log(\mathcal{R}_{t,m}) = \alpha_m + \beta d_{t,m} + s(t), \tag{3}$$

233 where $\mathcal{R}_{t,m}$ represents $\mathcal{R}(t)$ estimate at time t in municipality m; α_m represents the
234 municipality-specific intercept term; β represents the effect of susceptible depletion;
235 $d_{t,m}$ represents the cumulative cases at time t in municipality m; and $s(t)$ represents
236 the smooth term capturing temporal variation in transmission. For this analysis, we
237 only use $\mathcal{R}(t)$ estimates from municipalities with more than 400 total cases.

Transmission model

Finally, we fitted a simple transmission model using Bayesian inference to test
whether variation in introduction timing and susceptible dynamics alone can ex-
plain the observed epidemic patterns. Specifically, we extended the SEIR model,
which is commonly used for pertussis transmission [20], to allow for a joint estima-
tion of a single, time-varying transmission term that is shared across all regions and
a separate estimation of initial conditions and reporting rates. To do so, we first
discretized the SEIR model following the approach of [40]:

$$\beta(t) = \mathcal{R}_0(1 - \exp(-\gamma\Delta t))\delta(t) \tag{4}$$

$$\mathrm{FOI}(t) = \frac{\beta(t)I_m(t-\Delta t)}{N_m} \tag{5}$$

$$\Delta S_m(t) = [1 - \exp(-\mathrm{FOI}(t)\Delta t)]\, S_m(t-\Delta t) \tag{6}$$

$$\Delta E_m(t) = [1 - \exp(-\sigma\Delta t)]\, E_m(t-\Delta t) \tag{7}$$

$$\Delta I_m(t) = [1 - \exp(-\gamma\Delta t)]\, I_m(t-\Delta t) \tag{8}$$

$$S_m(t) = S(t-\Delta t) - \Delta S_m(t) \tag{9}$$

$$E_m(t) = E(t-\Delta t) - \Delta E_m(t) + \Delta S_m(t) \tag{10}$$

$$I_m(t) = I(t-\Delta t) - \Delta I_m(t) + \Delta E_m(t) \tag{11}$$

where $S_m(t)$, $E_m(t)$, and $I_m(t)$ represent the number of susceptible, exposed, and
infectious individuals in municipality m at time t, respectively; N_m represents the
population size of municipality m; Δt represents the simulation time step, which
assumed to be 1 week; $\beta(t)$ represents the shared time-varying transmission rate;
$\mathcal{R}_0$ represents the shared basic reproduction number, assumed to equal 17 [41]; $\delta(t)$
represents the normalized time-varying transmission rate; σ represents the rate at
which individuals develop infectiousness, which is assumed to be $\sigma = -\log(1 - 7/8)$
such that the mean latent period is 8 days; and γ represents the rate at which
individuals recover, which is assumed to be $\sigma = -\log(1 - 7/15)$ such that the mean
latent period is 15 days.

Here, we modeled changes in transmission using the $\delta(t)$ term, which is given a normal prior around 1:

$$\delta(t) \sim \text{Normal}(1, 0.2). \tag{12}$$

To allow for smooth variation in transmission, we also incorporated a random walk prior:

$$\delta(t) \sim \text{Normal}(\delta(t - \Delta t), \sigma_\delta), \tag{13}$$
$$\sigma_\delta \sim \text{Half-Normal}(0, 0.1), \tag{14}$$

where a half-normal prior on σ_δ constrains the smoothness of $\delta(t)$.

The municipality-specific observation process is modeled based on a Poisson distribution:

$$\text{cases}_{t,m} \sim \text{Poisson}(\rho_m \Delta S_m(t)) \tag{15}$$
$$\rho_m \sim \text{Uniform}(0, 1) \tag{16}$$

where $\text{cases}_{t,m}$ represents the reported cases at time t in municipality m; ρ_m represents the municipality-specific reporting rate; and $\Delta S_m(t)$ represents the expected number of new infections between time $t - \Delta t$ and t.

Finally, we estimate a municipality-specific initial conditions by imposing the following priors:

$$s_m(0) \sim \text{Uniform}(0, 1), \tag{17}$$
$$i_m(0) \sim \text{Half-Normal}(0, 0.001), \tag{18}$$

where $s_m(0)$ represents the initial fraction susceptible in municipality m, and $i_m(0)$ represents the initial fraction infected in municipality m. Then, the initial conditions for the model is specified as follows:

$$S_m(0) = N_m s_m(0), \tag{19}$$
$$E_m(0) = N_m i_m(0), \tag{20}$$
$$I_m(0) = N_m i_m(0). \tag{21}$$

We simultaneously fit this model time series data between May 2024 and March 2025 from all municipalities with more than 400 total cases (a total of 42 municipalities) using a Bayesian inference software rstan [21].

To assess the model fit, we compute R squared values for each municipality. This is done by calculating the squared value of the correlation coefficient between logged values of the observed cases+1 and a posterior median of the logged predictions $\rho_m \Delta S_m(t)$.

Finally, we evaluate the impact of initial conditions on pertussis epidemic dynamics by varying $s(0)$ between 0.13 and 0.23 and $i(0)$ between 7×10^{-6} and 1.5×10^{-3}. For each simulation, we quantify the center of gravity.

Transmission model validation

We validate our model by fitting the model to remaining municipalities (i.e., those with less than 400 total cases). In doing so, we assume that changes in transmission rate $\delta(t)$ is known and only estimate the initial conditions, $s_m(0)$ and $i_m(0)$, as well as the reporting rate ρ. For this validation, we simply used the optimization function in rstan [21] rather than performing a full Bayesian inference. Then, we compute R squared values for each municipality in the same way as before.

Relationshiip between estimated initial susceptible and vaccine coverage

We assessed the relationshiip between estimated initial susceptible and vaccine coverage using linear regression. Specifically, assuming that the waning of vaccinal immunity is the main cause for pertussis re-emergence, we compared DTaP (Diphtheria-Tetanus-Pertussis) vaccine coverage for 3 years old as of 2015 and the estimated initial susceptible in each municipality and performed a linear regression. We used vaccine coverage as of 2015 because this was the oldest data that were publicly available. We considered using estimates from municipalities with > 400 cases, since the parameter estimates are expected to be more reliable, but were unable to find any clear signature between vaccine coverage and estimated $s(0)$.

Acknowledgements

S.W.P. was supported by the New Faculty Startup Fund from Seoul National University.

Author contributions

S.W.P. conceived of the study, conducted the analysis, wrote the manuscript, and oversaw the work.

Competing interests

The authors declare no competing interests.

Data availability

All data and code used in this paper are available on https://github.com/parklab-snu/asynchrony-pertussis.

Supplementary Figures

Figure S1: **Spatiotempordal dynamics of pertussis outbreak in Korea across first-level administrative divisions, 2024—2025.** Time series are arranged based on their approximate locations.

Figure S2: **Validation of our transmission model using time series data from municipalities with less than 400 total cases.** (A) Comparisons between the observed and predicted epidemic dynamics across municipalities with less than 400 total cases, ordered by latitude. (B) The bar plot represents the distribution of R squared values for model fits. The vertical dashed line represents the median.

Figure S3: **Relationship between estimated initial susceptible and vaccine coverage for (A) municipalities with > 400 cases and (B) for all municipalities.** Each point represents the estimate initial susceptible and vaccine coverage value for each municipality. Red lines and shaded regions represent the linear regression fit and corresponding 95% confidence intervals.

References

[1] Virginia E Pitzer, Cécile Viboud, Wladimir J Alonso, Tanya Wilcox, C Jessica Metcalf, Claudia A Steiner, Amber K Haynes, and Bryan T Grenfell. Environmental drivers of the spatiotemporal dynamics of respiratory syncytial virus in the United States. *PLoS pathogens*, 11(1):e1004591, 2015.

[2] Benjamin D Dalziel, Stephen Kissler, Julia R Gog, Cecile Viboud, Ottar N Bjørnstad, C Jessica E Metcalf, and Bryan T Grenfell. Urbanization and humidity shape the intensity of influenza epidemics in US cities. *Science*, 362(6410):75–79, 2018.

[3] Margarita Pons-Salort, M Steven Oberste, Mark A Pallansch, Glen R Abedi, Saki Takahashi, Bryan T Grenfell, and Nicholas C Grassly. The seasonality of nonpolio enteroviruses in the united states: Patterns and drivers. *Proceedings of the National Academy of Sciences*, 115(12):3078–3083, 2018.

[4] Rachel E Baker, Ayesha S Mahmud, Caroline E Wagner, Wenchang Yang, Virginia E Pitzer, Cecile Viboud, Gabriel A Vecchi, C Jessica E Metcalf, and Bryan T Grenfell. Epidemic dynamics of respiratory syncytial virus in current and future climates. *Nature communications*, 10(1):5512, 2019.

[5] Michael P Hassell, Hugh N Comins, and Robert M Mayt. Spatial structure and chaos in insect population dynamics. *Nature*, 353(6341):255–258, 1991.

[6] JA Sherratt. Periodic travelling waves in cyclic predator–prey systems. *Ecology Letters*, 4(1):30–37, 2001.

[7] Bryan T Grenfell, Ottar N Bjørnstad, and Jens Kappey. Travelling waves and spatial hierarchies in measles epidemics. *Nature*, 414(6865):716–723, 2001.

[8] Derek AT Cummings, Rafael A Irizarry, Norden E Huang, Timothy P Endy, Ananda Nisalak, Kumnuan Ungchusak, and Donald S Burke. Travelling waves in the occurrence of dengue haemorrhagic fever in thailand. *Nature*, 427(6972):344–347, 2004.

[9] Cécile Viboud, Ottar N Bjørnstad, David L Smith, Lone Simonsen, Mark A Miller, and Bryan T Grenfell. Synchrony, waves, and spatial hierarchies in the spread of influenza. *science*, 312(5772):447–451, 2006.

[10] Robert S Paton, Christopher E Overton, and Thomas Ward. The rapid replacement of the SARS-CoV-2 Delta variant by Omicron (B. 1.1. 529) in England. *Science Translational Medicine*, 14(652):eabo5395, 2022.

[11] Hye-Jin Kim, Young Joon Park, Dongkeun Kim, Jihee Lee, Yun Kyoung Kim, Soonryu Seo, Jin Seon Yang, Yeoeun Yun, Eunbyeol Wang, Subin Park, Seo Yeon Ko, Jin Lee, Jeeyeon Shin, Wookeon Lee, and Seonhee Ahn. Pertussis surveillance and occurrence report in the Republic of Korea: from 2024 to the first half of 2025. *Public Health Weekly Report*, 18(43):1631–1651, 11 2025.

[12] Hyun Mi Kang, Taek-Jin Lee, Su Eun Park, and Soo-Han Choi. Pertussis in the post-COVID-19 era: resurgence, diagnosis, and management. *Infection & Chemotherapy*, 57(1):13, 2024.

[13] Scott M Duke-Sylvester, Luca Bolzoni, and Leslie A Real. Strong seasonality produces spatial asynchrony in the outbreak of infectious diseases. *Journal of the Royal Society Interface*, 8(59):817–825, 2011.

[14] Marc Choisy and Pejman Rohani. Changing spatial epidemiology of pertussis in continental USA. *Proceedings of the Royal Society B: Biological Sciences*, 279(1747):4574–4581, 2012.

[15] Jacco Wallinga and Marc Lipsitch. How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings of the Royal Society B: Biological Sciences*, 274(1609):599–604, 2007.

[16] Anne Cori, Neil M Ferguson, Christophe Fraser, and Simon Cauchemez. A new framework and software to estimate time-varying reproduction numbers during epidemics. *American journal of epidemiology*, 178(9):1505–1512, 2013.

[17] Dennis E te Beest, Michiel van Boven, Mariëtte Hooiveld, Carline van den Dool, and Jacco Wallinga. Driving factors of influenza transmission in the netherlands. *American journal of epidemiology*, 178(9):1469–1477, 2013.

[18] Simon N Wood. *Generalized additive models: an introduction with R*. Chapman and Hall/CRC, 2017.

[19] Stephen M Kissler, Christine Tedijanto, Edward Goldstein, Yonatan H Grad, and Marc Lipsitch. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science*, 368(6493):860–868, 2020.

[20] Pejman Rohani, Xue Zhong, and Aaron A King. Contact network structure explains the changing epidemiology of pertussis. *Science*, 330(6006):982–985, 2010.

[21] Bob Carpenter, Andrew Gelman, Matthew D Hoffman, Daniel Lee, Ben Goodrich, Michael Betancourt, Marcus Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. Stan: A probabilistic programming language. *Journal of statistical software*, 76:1–32, 2017.

[22] Bärbel F Finkenstädt, Ottar N Bjørnstad, and Bryan T Grenfell. A stochastic model for extinction and recurrence of epidemics: estimation and inference for measles outbreaks. *Biostatistics*, 3(4):493–510, 2002.

[23] James O Lloyd-Smith, Sebastian J Schreiber, P Ekkehard Kopp, and Wayne M Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, 2005.

[24] FMG Magpantay, M Domenech De Cellés, P Rohani, and AA King. Pertussis immunity and epidemiology: mode and duration of vaccine-induced immunity. *Parasitology*, 143(7):835–849, 2016.

[25] Helen J Wearing and Pejman Rohani. Estimating the duration of pertussis immunity using epidemiological signatures. *PLoS pathogens*, 5(10):e1000647, 2009.

[26] Tina Tan, Tine Dalby, Kevin Forsyth, Scott A Halperin, Ulrich Heininger, Daniela Hozbor, Stanley Plotkin, Rolando Ulloa-Gutierrez, and Carl Heinz Wirsing Von König. Pertussis across the globe: recent epidemiologic trends from 2000 to 2013. *The Pediatric infectious disease journal*, 34(9):e222–e232, 2015.

[27] Matthieu Domenech de Cellès, Felicia MG Magpantay, Aaron A King, and Pejman Rohani. The impact of past vaccination coverage and immunity on pertussis resurgence. *Science translational medicine*, 10(434):eaaj1748, 2018.

[28] Seonghui Cho, Dong Wook Kim, Chiara Achangwa, Junseo Oh, and Sukhyun Ryu. Pertussis in the elderly: Plausible amplifiers of persistent community transmission of pertussis in South Korea. *Journal of Infection*, 89(3), 2024.

[29] David L Heymann and Guénaël R Rodier. Hot spots in a wired world: Who surveillance of emerging and re-emerging infectious diseases. *The Lancet infectious diseases*, 1(5):345–353, 2001.

[30] Mahmoud M Naguib, Patrik Ellström, Josef D Järhult, Åke Lundkvist, and Björn Olsen. Towards pandemic preparedness beyond COVID-19. *The Lancet Microbe*, 1(5):e185–e186, 2020.

[31] Hai Nguyen-Tran, Sang Woo Park, Kevin Messacar, Samuel R Dominguez, Matthew R Vogt, Sallie Permar, Perdita Permaul, Michelle Hernandez, Daniel C Douek, Adrian B McDermott, et al. Enterovirus D68: a test case for the use of immunological surveillance to develop tools to mitigate the pandemic potential of emerging pathogens. *The Lancet Microbe*, 3(2):e83–e85, 2022.

[32] Michael J Mina, C Jessica E Metcalf, Adrian B McDermott, Daniel C Douek, Jeremy Farrar, and Bryan T Grenfell. A global lmmunological observatory to meet a time of pandemics. *Elife*, 9:e58989, 2020.

[33] Hai Nguyen-Tran, Sang Woo Park, Matthew R Vogt, Perdita Permaul, Alicen B Spaulding, Michelle L Hernandez, Jennifer A Bohl, Sucheta Godbole, Tracy J Ruckwardt, Peter W Krug, et al. Dynamics of endemic virus re-emergence in children in the USA following the COVID-19 pandemic (2022–23): A prospective, multicentre, longitudinal, immunoepidemiological surveillance study. *The Lancet Infectious Diseases*.

[34] Korea Disease Control and Prevention Agency. Notifiable Infectious Diseases. 2025. `https://dportal.kdca.go.kr/pot/is/rginEDW.do`. Accessed Nov 2, 2025.

[35] Ministry of the Interior and Safety. Resident Population and Household Statistics by Administrative District. 2025. `https://jumin.mois.go.kr/`. Accessed Nov 2, 2025.

[36] Korea Disease Control and Prevention Agency. 2015 National Immunization Coverage Survey. 2025. `https://nip.kdca.go.kr/irhp/infm/goNatnVcntStatView.do`. Accessed Nov 2, 2025.

[37] cubensys. Korea_district. 2018. `https://github.com/cubensys/Korea_District`. Accessed Nov 2, 2025.

[38] Hyung-jun Kim. Download the latest administrative divisions (SHP) of the Republic of Korea, GIS Developer. 2023. `http://www.gisdeveloper.co.kr/?p=2332`. Accessed Nov 2, 2025.

[39] Ottar N. Bjornstad. *ncf: Spatial Covariance Functions*, 2022. R package version 1.3-2.

[40] Daihai He, Edward L Ionides, and Aaron A King. Plug-and-play inference for disease dynamics: measles in large and small populations as a case study. *Journal of the Royal Society Interface*, 7(43):271–283, 2010.

[41] Roy M Anderson and Robert M May. *Infectious diseases of humans: dynamics and control.* Oxford university press, 1991.