Genotypic analysis of *Plasmodium falciparum* malaria parasites in a clinical trial of RTS,S vaccination in combination with seasonal malaria chemoprevention

Emily LaVerriere[1,2], Katrina Kelley[1,2], Alassane Dicko[3], Issaka Sagara[3], Jean-Bosco Ouedraogo[4], Issaka Zongo[4], Halidou Tinto[4], Daniel Chandramohan[5], Brian Greenwood[5], Daniel E. Neafsey*[1,2]

Affiliations:

1. Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA, USA.

2. Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA.

3. The Malaria Research and Training Center, University of Science, Technology and Techniques of Bamako, Bamako, Mali.

4. Institut des Sciences et Techniques, Institut de Recherche en Sciences de la Santé, Bobo-Dioulasso, Burkina Faso.

5. Department of Disease Control, London School of Hygiene & Tropical Medicine, London, UK.

* Corresponding author. Email: neafsey@hsph.harvard.edu

Abstract

Both vaccination and seasonal chemoprevention are effective interventions against malaria, but both morbidity and mortality due to malaria remain high, particularly in sub-Saharan Africa. A recent clinical trial showed that the combination of these interventions provided substantial protection against malaria over a five-year period. We performed amplicon-based genotyping of highly polymorphic parasite antigens within 1,530 samples from this clinical trial.

We evaluated the complexity of infection within these samples, finding that participants who

received both interventions had less complex infections (fewer genetically distinct parasite

strains) than participants who received either intervention given alone. We also evaluated the

prevalence of parasites matching the vaccine construct CSP, at the level of haplotypes,

epitopes, and individual amino acid changes. We found no significant differences in the

prevalence of vaccine-matching alleles between vaccinated and unvaccinated groups. For

context, this study genotyped about half as many non-vaccinated participants as the original

phase 3 trial for RTS,S. While a comparable number of vaccinated individuals were genotyped

in both studies, the infections in this study had less complex infections, leading to an overall

lower number of parasite strains to analyze. These results suggest that the combination of

therapies provides protection from infection, in addition to clinical disease, and they highlight the

importance of continued molecular surveillance as malaria interventions are deployed.

Introduction

Malaria remains a persistent global health threat, causing an estimated 597,000 deaths

per year, primarily in children under five years of age (1). The burden of malaria is greatest in

sub-Saharan Africa, where in many regions disease transmission is seasonal in nature due to

rainfall-associated changes in the population size of *Anopheles spp.* mosquitoes that transmit

the disease (2). While immunity to clinical disease has been demonstrated to accrue with age,

acquisition of asymptomatic infections is age-independent, and occurs with similar timing across

age groups following the commencement of the rainy season (3,4).

Effective interventions for malaria in seasonal transmission settings are therefore timed

to coincide with increased transmission, and include seasonal chemoprevention (SMC) and,

more recently, seasonal vaccination. Seasonal chemoprevention usually involves multiple,

regularly administered doses of sulfadioxine-pyrmethamine and amodiaquine during the

expected transmission season (5). Multiple vaccine candidates and schedules have been tested

(6,7), and the RTS,S/AS01$_E$ and R21 vaccines are currently recommended for use by the WHO

in malaria-endemic areas, each with a recommended three-dose primary schedule, followed by

a booster dose (8,9). The timing of administration of the booster dose is usually age based but

countries with highly seasonal malaria transmission may consider giving the booster dose prior

to the malaria transmission season (9). Both SMC and vaccination provide protection against

disease that wanes over time. A recent clinical trial compared the combination of SMC and

seasonal vaccination with the RTS,S/AS01$_E$ vaccine (subsequently termed RTS,S) to each

intervention on its own in areas with strongly seasonal malaria transmission (10,11). In this trial,

the combination of SMC with RTS,S significantly reduced the incidence of uncomplicated

disease, severe disease, and death compared to either intervention on its own.

In this study, we performed genotyping of highly polymorphic parasite antigens in

participant samples obtained from children in each of the three arms of the combined SMC and

vaccination clinical trial (RTS,S alone, SMC alone, and the combination) in order to understand

the nature of protection of the individual and combined interventions against infection. Our

objectives were to determine whether the vaccine provided differential protection against

parasites that had an exact match at the circumsporozoite protein (CSP) antigen to the RTS,S

vaccine construct, and to estimate the number of genomically distinct parasite lineages within

each sample as a measure of protection against infection. Previous large-scale genotyping

studies have examined the parasites present in infections from multiple RTS,S clinical trials

(12,13). The initial genotyping analysis of the phase 3 trial found increased vaccine efficacy

against parasite strains matching the CSP genotype used within the vaccine. More recent work

analyzing a phase 2b clinical trial comparing different vaccine dosing regimens found that all

vaccine regimens tested blocked some parasite infections, and that complexity of infection was

reduced in participants in the RTS,S arm compared to the control arm.

In this study, we generated parasite genotyping data from samples meeting the clinical

case definition for malaria from all three arms of the combined SMC and vaccination clinical trial

(10,11), allowing us to profile the number of clones in each. While the original study

demonstrated a clear effect of the combined interventions on clinical disease, these newly-

generated genotyping data demonstrate evidence of combined efficacy against infection in a

non-strain-specific manner.

Methods

<u>Samples / original trial</u>

We processed samples from the clinical trial NCT04319380, registered on

ClinicalTrials.gov, as previously described (10,11). This study was a double-blind, individually-

randomized, controlled, non-inferiority and superiority phase 3 trial done at two sites: the

Bougouni district and neighboring areas in Mali and Houndé district, Burkina Faso. The original

trial protocol was reviewed and approved by the ethics committees of the London School of

Hygiene & Tropical Medicine, UK; the University of Science, Techniques, and Technologies,

Bamako, Mali; the ethics committee for health research, Burkina Faso; and the regulatory

authorities in Burkina Faso and Mali.

Briefly, 6,861 children 5 to 17 months of age were enrolled and randomly assigned to

receive either sulfadoxine-pyrimethamine and amodiaquine (the chemoprevention-alone group,

2,287 children), RTS,S/AS01$_E$ (the vaccine-alone group, 2,288 children), or both therapies (the

combination group, 2,286 children). The chemoprevention-alone group received tetanus or

tetanus-diphtheria toxoid vaccine in place of RTS,S/AS01$_E$, and the vaccine-alone group also

received sulfadoxine-pyrimethamine and amodiaquine placebos. Participants each received five

doses of RTS,S/AS01$_E$ or placebo vaccination: three in April-June 2017, followed by annual

doses in 2018 and 2019. Participants each received four courses of sulfadoxine-pyrimethamine

and amodiaquine or placebo per year. All participants received a long-lasting insecticide-treated

bed net when enrolled into the study.

Uncomplicated malaria was the primary outcome of the original trial, defined as fever

(either measured at least 37.5□ or fever within the previous 48 hours) and *P. falciparum*

parasitemia of at least 5,000 parasites per mm^3. Participants presenting to trial health facilities

with suspected malaria were tested with rapid diagnostic tests, and those with positive tests

received artemether-lumefantrine and gave a blood sample for microscopic confirmation.

The initial trial lasted three years (10), and participants who did not have severe

reactions to the interventions (febrile convulsions after vaccination) were invited to re-enroll in

the extension study for the subsequent two years (11). In 2020, 5098 children consented to re-

enroll in the extension study; these participants retained the same treatment groups as before,

resulting in 1,683, 1,705, and 1,710 participants in the extension study combination group,

vaccine-only group, and chemoprevention-only groups, respectively. Participants no longer

received SMC when they reached the age of five years, in line with national guidelines, and

seasonal vaccination was no longer given after they reached this age. Thus, participants who

enrolled when aged 5-11 months received two additional doses of RTS,S/AS01$_E$ or placebo

vaccination in June 2020 and June 2021, as well as four cycles of sulfadoxine-pyrimethamine

and amodiaquine or placebo per year in 2020 and 2021, whilst those enrolled when aged 12-17

months received only one additional year of SMC and seasonal vaccination in 2020.

A 1:2 random sample from all samples which came from infections meeting the clinical

definition of malaria in the original trial (symptomatic disease with ≥5,000 parasites/µL identified

via microscopy) was selected for genotyping across intervention and country for each year

(2017, 2019, 2021). This selection resulted in 730 samples from the vaccine-only group, 572

samples from the chemoprevention-only group, and 228 samples from the combination group

(**Table 1**). Sequencing was attempted on all selected samples.

Table 1.

Country	Burkina Faso			Mali		
Year	RTS,S	SMC	RTS,S+SMC	RTS,S	SMC	RTS,S+SMC
2017	114	50	12	85	76	10
2019	204	182	61	65	82	37
2021	154	71	58	108	111	50

Number of samples per group, year, and country. All samples were from infections meeting the

clinical definition of malaria in the original trial (symptomatic disease with ≥5,000 parasites/µL

identified via microscopy).

Sequencing

We extracted genomic DNA from the dried blood spot samples and prepared them for

sequencing using the 4CAST amplicon panel, as previously described (14), with no changes in

protocol from the original publication. This amplicon panel targets four highly polymorphic

regions of the *P. falciparum* genome, within the *csp, ama1, sera2*, and *trap* genes. Particularly

relevant for this study, the amplicon within *csp* covers the polymorphic C-terminal region that is

part of the RTS,S vaccine construct. The coverage of this region of *csp* allows for direct

identification of parasites matching or mis-matching the vaccine strain in this region. The high

diversity captured within this amplicon panel also allows estimation of complexity of infection

(COI; number of genotypically distinct strains within an infection) more accurately than a single

amplicon alone.

We sequenced the majority of samples (n=1,542) on two lanes of Illumina NovaSeq

instruments, with 768 samples per lane. We sequenced a second batch of samples (n=192) on

an Illumina MiSeq instrument. All sequencing runs were paired-end, 2x250 bp reads. Samples

were processed in 96-well plates, with each plate containing extraction and PCR controls that

were carried through sequencing and analysis. Data from these samples were submitted to the

NCBI Sequencing Read Archive (http://www.ncbi.nih.gov/sra) under accession PRJNA1345094.

 <u>Analysis</u>

We processed the data through a previously described pipeline (14), designed for

analyzing deeply sequenced multiplexed amplicon sequencing data and parsing it into

haplotypes. We set a threshold for the minimum number of reads per haplotype per sample at

200 reads (for samples sequenced on NovaSeq) or 20 reads (for samples sequenced on

MiSeq). We also required a haplotype to have a within-sample frequency of at least 1% as

estimated by sequencing reads, and we removed any singleton haplotypes from the full dataset.

We examined the positive and negative control wells present on every 96-well plate; control

wells were required to have the included strain (Dd2) detected, and negative controls were

required to have no haplotypes with reads above 200 (for NovaSeq samples) or 20 (for MiSeq

samples). The negative control in one 96-well plate did not pass these checks, so all samples

on that plate were removed from the final dataset.

One of the two NovaSeq lanes had several sequencing artifacts. Specifically, three

nucleotide changes within the CSP amplicon were detected in some samples (genomic

coordinates 221383, 221403, and 221412, all changed to C). We inspected this region in

previously generated amplicon sequencing datasets of parasites from similar regions, and none

of these three changes appeared (3,15). We also looked within the Pf3k dataset of whole-

genome data (16) and did not see these nucleotide changes in any samples. Finally, we

selected eight samples where these nucleotide changes were present, re-amplified this region

with the CSP primers, and confirmed with Sanger sequencing that the nucleotide changes were

not present in the samples. We masked these three positions in all downstream analysis.

We performed all subsequent analyses on denoised haplotype calls in R 4.2.0, with the

packages Biostrings, DescTools, grantham, here, RColorBrewer, stats, and tidyverse (17–23).

We estimated COI as the maximum number of distinct haplotypes identified at an individual

4CAST locus (CSP, AMA1, SERA2, TRAP) for an individual sample. All statistical tests were

two-sided, unless otherwise noted. We tested for significance using the "kruskal.test",

"fisher.test", and "p.adjust" functions from the stats package, as well as the "BinomDiffCI"

function from the DescTools package.

Results

Figure 1.

All panels visualize the proportion of samples at estimated complexity of infection (COI) level,

from 1 (monoclonal) to 5+ (highly polyclonal). A) Samples are stratified by study arm. Samples

are more likely to have lower COI in the combination-therapy group (Kruskal-Wallis test, $x^2(2)$ =

9.11, P = 0.011; Dunn test with Holm correction, both vs. SMC only: adjusted P = 0.0079). B)

Samples stratified by country. COI tended to be higher in samples in Burkina Faso than in

samples from Mali (Kruskal-Wallis test, $x^2(1)$ = 22.35, P = 1.41 e^{-5}; mean COI in Burkina Faso =

2.3, mean COI in Mali = 1.8). C) Samples stratified by study arm and country. Samples are

significantly less complex in the combination group in Burkina Faso (Kruskal-Wallis test, $x^2(2)$ =

6.84, P = 0.03; Dunn test with Holm correction, both vs. SMC only: adjusted P = 0.027).

Samples are less complex in the combination group in Mali, though not significantly (Kruskal-

Wallis test, $x^2(2)$ = 3.76, P = 0.15). D) Samples stratified by time period and country. COI

distributions were consistent over time in Burkina Faso (Kruskal-Wallis test, $x^2(2)$ = 0.22, P =

0.90), but differed in later years in Mali (Kruskal-Wallis test, $x^2(2) = 8.85$, P = 0.011; Dunn test
with Holm correction, 2017-2018 vs. 2021-2022, adjusted P = 0.012).

<u>Infections are less complex in individuals who received RTS,S/AS01$_E$ with SMC</u>

We generated high-quality *P. falciparum* sequencing data from 1,530 samples in total, all

of which came from infections meeting the clinical definition of malaria in the original trial:

symptomatic disease with ≥5000 parasites/µL identified via microscopy. Concordant with the

original study (10), the number of clinical disease cases was much lower in the individuals who

received both SMC and RTS,S, leading to an uneven number of samples to sequence between

each study arm. Thus, we sequenced 228 samples from the RTS,S with SMC arm, compared to

730 samples from the RTS,S arm and 572 samples from the SMC arm (**Table 1**).

We first leveraged the genotyping data to identify complex (polyclonal) infections. We

estimated the complexity of infection, or the number of genetically distinct parasites identified in

each sample. We found that monoclonal infections were more common in individuals who

received both RTS,S and SMC than in those who received SMC only (**Figure 1A**; Kruskal-

Wallis test, $x^2(2) = 9.11$, P = 0.011; Dunn test with Holm correction, both vs. SMC only: adjusted

P = 0.0079).

We also compared COI distributions by country (**Figure 1B**). Both countries had similar

proportions of monoclonal infections - approximately half of all sequenced infections were

monoclonal (48.5% in Burkina Faso (n=393), 52.3% in Mali (n=390). However, among the

infections with more than one clone detected, COI tended to be higher in individuals in Burkina

Faso than in individuals from Mali (Kruskal-Wallis test, $x^2(1) = 22.35$, P = 1.41 e^{-5}; mean COI in

Burkina Faso = 2.3, mean COI in Mali = 1.8). When stratified by both study arm and country

(**Figure 1C**), COI distributions follow a similar trend as in **Figure 1A**. In Burkina Faso, the

pattern of monoclonal infections being more common in the RTS,S + SMC study arm was

observed again (Kruskal-Wallis test, $x^2(2) = 6.84$, P = 0.03; Dunn test with Holm correction, both

vs. SMC only: adjusted P = 0.027). Monoclonal infections remained more common in RTS,S +

SMC individuals in Mali as well, though the trend did not reach the threshold of statistical

significance (Kruskal-Wallis test, $x^2(2) = 3.76$, P = 0.15).

COI distributions remained consistent over time in Burkina Faso (**Figure 1D**; Kruskal-

Wallis test, $x^2(2) = 0.22$, P = 0.90). In Mali, monoclonal infections became less frequent in later

years (Kruskal-Wallis test, $x^2(2) = 8.85$, P = 0.011; Dunn test with Holm correction, 2017-2018

vs. 2021-2022, adjusted P = 0.012).

Figure 2.

A) Frequency of the top ten most common amino acid CSP haplotypes (each with a distinct
color), including the 3D7-matching haplotype (second most common, colored in black). All less
common haplotypes are grouped together and colored in grey. 3D7-matching haplotype
prevalence did not differ between vaccinated and unvaccinated participants (Fisher's exact test,
one-sided, P = 0.386). B) Grantham distance for each haplotype compared to the 3D7
haplotype is on the x-axis (distance is a sum of the Grantham distance from each non-3D7-
matching amino acid per haplotype). The difference in prevalence of the haplotype between
vaccinated and unvaccinated groups is on the y-axis. The lines spanning each point plot the
95% confidence interval around the difference in prevalences.
232
<u>Parasites matching the vaccine strain appear at similar proportions in all groups</u>

We compared the prevalence of CSP haplotypes matching the 3D7 vaccine strain

between participants who received RTS,S and those who did not (**Figure 2a**). Prevalence of

amino acid haplotypes matching 3D7 did not differ between vaccinated participants (127 3D7-

matching infections, prevalence = 14.4%) and unvaccinated participants (81 3D7-matching

infections, prevalence = 15.1%; Fisher's exact test, one-sided, P = 0.386). Additionally, we

found no difference when considering the frequency of 3D7-matching haplotypes in place of

prevalence; 3D7-matching haplotypes in vaccinated participants had a prevalence of 8.73%

among all haplotypes, compared to a prevalence of 8.56% among non-vaccinated participants

(Fisher's exact test, one-sided, P = 0.579). Full lists of nucleotide and amino acid haplotypes

observed in participants stratified by vaccination status are shown in **Supplemental Tables 1-2**.

We also considered how similar the common haplotypes were to 3D7, by summing the

Grantham distance of each mismatched amino-acid in a given haplotype (**Figure 2b**). None of

the most common haplotypes were very similar to 3D7 (sum of Grantham distances ranged

from 319 to 533 per haplotype).

Table 2.

Locus	3D7-matching allele prevalence			P-value
	RTSS+	RTSS-	Difference	
CSP	0.147	0.152	-0.005	0.386
Th2R	0.157	0.162	-0.005	0.641
Th3R	0.214	0.223	-0.009	0.609
DV10	0.177	0.173	0.004	0.778

Prevalence of 3D7-matching CSP alleles in vaccinated (RTSS+) and unvaccinated (RTSS-)
populations. The Locus column describes the allele being compared, including the full amino
acid haplotype (CSP) and the amino acid haplotypes for each of the three CSP epitopes (Th2R,
Th3R, and DV10). P-values are from Fisher's exact test, one-sided.

We also evaluated whether evidence of allele-specific protection could be identified at

the three epitopes previously described for CSP: DV10, Th2R, and Th3R, rather than at the full

haplotype level (**Table 2**). We compared the prevalence of amino acid haplotypes spanning

each of these epitope regions individually, and we again found no difference associated with

vaccination status forTh2R, Th3R, or DV10 (Fisher's exact tests, one-sided; P = 0.641, 0.609,

0.778, respectively). The incidence of each epitope haplotype per study arm is described in

Supplemental Tables 3-5.

Finally, we investigated individual amino acid positions within these epitopes to look for

evidence of 3D7-specific protection. Specifically, we examined amino acid positions 299, 301,

317, 354, 356, 359, 361, which all vary in these populations, consistent with previous deep

sequencing studies of CSP (13,15). We examined each position on its own, and we compared

the distributions of amino acids observed at that position across each of the study arms. The

prevalence of the 3D7-matching allele at position 321 was higher in vaccinated participants (n =

281, prevalence = 0.319) compared to unvaccinated participants (n = 151, prevalence = 0.283.

While the initial hypothesis testing was significant (Fisher's exact test, one-sided, P = 0.01),

after adjusting for multiple hypothesis testing, the result is on the edge of significance (adjusted

P = 0.22, after Benjamini-Hochberg correction). None of the other distributions varied

significantly from each other, even before correcting for multiple hypothesis testing (Fisher's

exact test applied to each residue, prevalences and p-values in **Supplemental Table 6**, raw

number of appearances of each residue in **Supplemental Table 7**).

Discussion

In this study, we generated amplicon sequencing data to explore parasite diversity within

symptomatic infections in a clinical trial which compared the combination of SMC with seasonal

vaccination with RTS,S with either intervention given alone (10). Genotyping data has the

capacity to both confirm and, in this case, recapitulate the efficacy of the interventions in the

trial. While the original study found an increase in protection against symptomatic disease in

those who received both SMC and vaccination (10,11), we found an additional benefit of the

combined interventions. Participants who received both interventions were also more likely to

exhibit lower complexity infections than those receiving only SMC or RTS,S alone. This

suggests that participants who received both interventions were more effective at blocking a

subset of infections, whether polyclonal infections were founded via multiple infectious mosquito

bites or single polyclonal infectious bites. This finding suggests that the combined

chemoprevention and vaccination not only protects against clinical disease, but also has

additive protection against parasite infection itself, which can occur regardless of clinical

symptoms. This protection may manifest as lower complexity infections, as previously seen in

individuals vaccinated with RTS,S (12).

The genotyping data generated also allowed us to look for differential protection as a

result of vaccination, which was observed in the original phase 3 trial of the RTS,S/AS01$_E$

vaccine (13). We did not observe vaccination-related differences in the prevalence of 3D7-

matching alleles when considering full haplotypes or epitopes. Two factors may have resulted in

a greater temporal lag and a smaller impact of vaccination on the observed breakthrough

infections genotyped here compared to the phase 3 trial. First, the administration of SMC in the

combination group may have delayed the time to first infection within each season of

transmission, thus leading to first infections occurring when vaccine efficacy has started to

decrease. Second, the samples genotyped in this study were a random selection of all

symptomatic cases (stratified by country and year), compared to the phase 3 trial which

genotyped the first post-vaccination infection per individual.

We did observe increased prevalence of the 3D7-matching CSP amino acid 321N in

vaccinated individuals, although this difference did not reach statistical significance after

correcting for multiple hypothesis testing. However, we cannot rule out that a smaller magnitude

difference in protection could be present below our limit of detection. With a minimum of 500

samples per group, estimated prevalence of the 3D7-matching haplotype of 0.15 in the

unvaccinated population, and a significance threshold of 0.05, this study had only 80% power to

detect a difference of ~0.05 in prevalence between groups. Any smaller differences in

prevalence would require higher sample size. As vaccination rates increase throughout malaria-

endemic regions, continued surveillance of malaria infections in vaccinated and unvaccinated

populations will be necessary for detecting any signals of vaccine escape, and larger studies will

be better powered to estimate the magnitude of any allele-specific protective efficacy that is

present.

Conclusion

Combining seasonal malaria chemoprevention with vaccination protected against clinical

disease in a previous clinical trial. In the study reported in this paper, we found that the

combination of SMC and seasonal vaccination also led to reduced levels of polyclonality within

samples from participants with clinical disease, indicating increased protection against infection

itself relative to participants receiving only one intervention. Infection polyclonality was similar

between the RTS,S-only and SMC-only study, suggesting comparable protection against

infection provided by each intervention given alone, as was noted in analysis of clinical

infections. These genotyping results underscore the additive value of vaccination and SMC for

protecting children against malaria in seasonal transmission settings. In the RTS,S-only and

RTS,S+SMC study arms we found no evidence of significantly greater protection against

parasites with CSP C-terminus sequences matching the 3D7 vaccine strain, suggesting that

such protection is modest in magnitude if present. Nevertheless, we advocate continued

surveillance for evidence of vaccine escape as RTS,S and R21 are more widely deployed in

Africa, when power to detect allele-specific protection may be enhanced.

Acknowledgements

We thank Sanni Ali and Paul Snell for their assistance with sample selection and coordination.

The overall trial on which the study described in this paper and the laboratory analyses

described was funded by the UK Joint Global Health Trials (Department of Health and Social

Care (DHSC), the Foreign, Commonwealth & Development Office (FCDO), the Global

Challenges Research Fund (GCRF), the Medical Research Council (MRC) and the Wellcome

Trust (WT)) (grant MR/V005642/1). This UK funded award was part of the EDCTP2 programme

supported by the European Union. The trial was also funded by PATH's Malaria Vaccine

Initiative (CVIA 083-MAL99) through a grant to PATH (INV-007217) from the Bill and Melinda

Gates Foundation. Research grants were made to the London School of Hygiene & Tropical

Medicine, United Kingdom, with sub-contracts to the Malaria Research and Training Center

(MRTC), Mali, the Institut de Recherche en Sciences de la Santé (IRSS) and to the University of

Oxford.

References

1. World Health Organization. World malaria report 2024: addressing inequity in the global malaria response. 2024.

2. Cairns M, Roca-Feltrer A, Garske T, Wilson AL, Diallo D, Milligan PJ, et al. Estimating the potential public health impact of seasonal malaria chemoprevention in African children. Nat Commun. 2012 June 6;3:881.

3. LaVerriere E, Johnson ZM, Shieh M, Nziza N, Alter G, Buckee CO, et al. Marked heterogeneity in malaria infection rate in a Malian longitudinal cohort. Nat Commun. 2025 July 15;16(1):6512.

4. Tran TM, Li S, Doumbo S, Doumtabe D, Huang CY, Dia S, et al. An Intensive Longitudinal Cohort Study of Malian Children and Adults Reveals No Evidence of Acquired Immunity to Plasmodium falciparum Infection. Clin Infect Dis. 2013 July 1;57(1):40–7.

5. ACCESS-SMC Partnership. Effectiveness of seasonal malaria chemoprevention at scale in west and central Africa: an observational study. Lancet Lond Engl. 2020 Dec 5;396(10265):1829–40.

6. Datoo MS, Natama HM, Somé A, Bellamy D, Traoré O, Rouamba T, et al. Efficacy and immunogenicity of R21/Matrix-M vaccine against clinical malaria after 2 years' follow-up in children in Burkina Faso: a phase 1/2b randomised controlled trial. Lancet Infect Dis. 2022 Dec;22(12):1728–36.

7. RTS,S Clinical Trials Partnership. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial. Lancet Lond Engl. 2015 July 4;386(9988):31–45.

8. University of Oxford. University of Oxford. 2023 [cited 2025 Sept 20]. Oxford R21/Matrix-M™ malaria vaccine receives WHO recommendation for use paving the way for global roll-out | University of Oxford. Available from: https://www.ox.ac.uk/news/2023-10-02-oxford-r21matrix-m-malaria-vaccine-receives-who-recommendation-use-paving-way-global

9. World Health Organization. WHO recommends groundbreaking malaria vaccine for children at risk [Internet]. 2021 [cited 2025 Sept 20]. Available from: https://www.who.int/news/item/06-10-2021-who-recommends-groundbreaking-malaria-vaccine-for-children-at-risk

10. Chandramohan D, Zongo I, Sagara I, Cairns M, Yerbanga RS, Diarra M, et al. Seasonal Malaria Vaccination with or without Seasonal Malaria Chemoprevention. N Engl J Med. 2021 Sept 9;385(11):1005–17.

11. Dicko A, Ouedraogo JB, Zongo I, Sagara I, Cairns M, Yerbanga RS, et al. Seasonal vaccination with RTS,S/AS01E vaccine with or without seasonal malaria chemoprevention in children up to the age of 5 years in Burkina Faso and Mali: a double-blind, randomised, controlled, phase 3 trial. Lancet Infect Dis. 2024 Jan;24(1):75–86.

12. Juraska M, Early AM, Li L, Schaffner SF, Lievens M, Khorgade A, et al. Genotypic analysis of RTS,S/AS01E malaria vaccine efficacy against parasite infection as a function of

dosage regimen and baseline malaria infection status in children aged 5-17 months in
Ghana and Kenya: a longitudinal phase 2b randomised controlled trial. Lancet Infect Dis.
2024 May 6;S1473-3099(24)00179-8.

13. Neafsey DE, Juraska M, Bedford T, Benkeser D, Valim C, Griggs A, et al. Genetic
Diversity and Protective Efficacy of the RTS,S/AS01 Malaria Vaccine. N Engl J Med. 2015
Nov 19;373(21):2025–37.

14. LaVerriere E, Schwabl P, Carrasquilla M, Taylor AR, Johnson ZM, Shieh M, et al.
Design and implementation of multiplexed amplicon sequencing panels to serve genomic
epidemiology of infectious disease: A malaria case study. Mol Ecol Resour.
2022;22(6):2285–303.

15. Early AM, Lievens M, MacInnis BL, Ockenhouse CF, Volkman SK, Adjei S, et al. Host-
mediated selection impacts the diversity of Plasmodium falciparum antigens within
infections. Nat Commun. 2018 11;9(1):1381.

16. MalariaGEN Plasmodium falciparum Community Project. MalariaGEN Plasmodium
falciparum Community Project. https://www.malariagen.net/resource/34; 2016.

17. Magno R, Duarte I, Maia AT. grantham: Calculate the Grantham Distance [Internet].
2025. Available from: https://CRAN.R-project.org/package=grantham

18. Müller K. here: A simpler way to find your files [Internet]. 2020. Available from:
https://CRAN.R-project.org/package=here

19. Neuwirth E. RColorBrewer: ColorBrewer palettes [Internet]. 2022. Available from:
https://CRAN.R-project.org/package=RColorBrewer

20. Pagès H, Aboyoun P, Gentleman R, DebRoy S. Biostrings: Efficient manipulation of
biological strings [Internet]. 2025. Available from:
https://bioconductor.org/packages/Biostrings

21. R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria: R Foundation for Statistical Computing; 2025. Available from:
https://www.R-project.org/

22. Signorell A. DescTools: Tools for Descriptive Statistics [Internet]. 2025. Available from:
https://CRAN.R-project.org/package=DescTools

23. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to
the {tidyverse}. J Open Source Softw. 2019;4(43):1686.