Cross-Program Secondary Analyses and Public Health Innovation: The RADx Data Hub as a Resource for Integrated COVID-19 Research

[1*]Peter W. Rose, [2*]Thais Rivas, [3*]Nader Mehri, [4*]Oswaldo Alonso Lozoya, [5*]Liesl Jeffers-Francis, [6#]Keriayn N. Smith, [7]Marquea D. King, [8]Mete U. Akdogan, [6]Jasmine Snipe, [6]Brandy Farlow, [7]Michael A. Keller, [8]Mark A. Musen, [6]Stephanie Suber, [6]Ashok Krishnamurthy

[*] Equal contributions

[1] San Diego Supercomputer Center, University of California San Diego, La Jolla, CA 92093, USA (https://orcid.org/0000-0001-9981-9750)

[2] Department of Health and Human Sciences, University of North Carolina at Greensboro, Greensboro, NC 27405, USA

[3] School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27517, USA (https://orcid.org/0000-0002-6820-8022)

[4] RTI International, Durham, NC, 27713, USA (https://orcid.org/0000-0001-6779-7382)

[5] Biology Department, College of Science and Technology, North Carolina Agricultural and Technical State University, Greensboro NC 27411 (https://orcid.org/0000-0001-7725-6336)

[6] Renaissance Computing Institute (RENCI), University of North Carolina at Chapel Hill, Chapel Hill, NC 27517, USA (https://orcid.org/0000-0002-4351-2765; https://orcid.org/0009-0002-8157-1421, https://orcid.org/0000-0002-3215-7622)

[7] Booz Allen Hamilton Inc., McLean, VA 22102, USA (https://orcid.org/0009-0004-6688-4491)

[8] Stanford Center for Biomedical Informatics Research, Stanford University, Palo Alto, CA 94304, USA (https://orcid.org/0000-0003-1520-1481, https://orcid.org/0000-0003-3325-793X)

[#]Correspondence: kns@email.unc.edu

Word count = 6955

Abstract

The COVID-19 pandemic highlighted the need for timely, standardized research data to address future health challenges. From the pandemic, there has been a plethora of diverse data, including patient demographics, clinical outcomes, biospecimen information, and technology development frameworks. Yet, much of these data have been fragmented and without clear provenance and standardization. To overcome these limitations, the National Institutes of Health

(NIH) established the RADx Data Hub (Data Hub) as a centralized platform to promote Rapid

Acceleration of Diagnostics (RADx) data distribution under the Findable, Accessible,

Interoperable, and Reusable (FAIR) data principles. The Data Hub aggregated diverse COVID-

19 diagnostic and clinical datasets into a harmonized repository that supports data integration,

quality control, and secondary research. We independently tested Data Hub biomedical data

management and data analysis capabilities to investigate its utility in viral monitoring, health

disparity research, biomarker identification, and precarity modeling. We found that the Data

Hub's accessibility enabled efficient secondary research, including diagnostic and clinical data

reuse. This supports repository use in accelerating infectious disease research and studying

disease outcomes. Further, the Data Hub's centralized, FAIR-aligned structure exemplifies a

sustainable data stewardship model, extending beyond COVID-19 to addressing future public

health emergencies, while strengthening the capacity to do reproducible science within broader

biomedical research.

Introduction

The COVID-19 pandemic exposed critical research gaps that precluded the efficient data

collection, integration, and analysis needed to enable a rapid public health response. To assist,

the National Institutes of Health (NIH) launched the Rapid Acceleration of Diagnostics (RADx)

Initiative (1). This initiative, comprised of four initial programs: RADx Radical (RADx-rad), RADx

Underserved Populations (RADx-UP), RADx Tech, and RADx Digital Health Technologies

(RADx DHT), diverse yet complementary programs spanning experimental diagnostics, health

disparities research, technology development, and digital health solutions (1,2). Specifically,

RADx-rad focused on novel diagnostic approaches for improved COVID-19 detection, including

innovative assays and targets, while RADx-UP focused on understanding and addressing

COVID-19 health disparities (1). RADx Tech accelerated COVID-19 test development,

validation, and commercialization, and RAD DHT generated and integrated digital health data

from diverse sources (1,2).

The final program, the RADx Data Hub (Data Hub; radxdatahub.nih.gov), was formed to

centralize and standardize RADx data from more than 180 studies and 1700 data files (2),

aggregating vast and valuable COVID-19 data to enable secondary research and accelerate

diagnostic innovations and public health impacts (2). Within the Data Hub, structured

approaches were developed to validate, deidentify, and integrate data, supporting findable,

accessible, interoperable, and reusable (FAIR) data principles (2). For users, the Data Hub

provided harmonized data access within a secure environment equipped with appropriate

analytical tools, including statistical software and Jupyter notebooks, to facilitate secondary

intra- and cross-study RADx program data investigations (2).

The goal behind this organization and access was to eliminate common primary data collection

time and resource constraints faced by researchers, to facilitate enriched exploration of

research questions and foster efficient hypothesis testing to accelerate COVID-19 (and related)

knowledge generation. The Data Hub's harmonization framework categorized data elements

into tiers reflecting their degree of standardization and shared use across multiple studies (2).

This structure integrates diverse datasets and enables comparative analysis (including Tier 1,

universal data elements across all studies) to support comprehensive foundational information.

Additional tiered variables represent common, program-specific data elements, tailored to

certain study subsets or non-harmonized, study-specific variables (2). For example, metadata

on RADx-rad specimen types, analytes, detection approaches, and biological targets were

systematically catalogued across diagnostic sub-projects (2,3).

The goal of these Data Hub functionalities is to support broad secondary research to advance

understanding of COVID-19 and similar diseases. This includes population-level viral monitoring

through diagnostic and environmental surveillance data, tracking viral/variant emergence and

spread to inform public health responses, and insights into vulnerable community testing access

and infection outcome inequities. Further, biomarker identification is an important use case for

diagnostics and prognostics, with integrable datasets for researchers to identify biological

signatures applicable for disease detection and monitoring. Predictive modeling with

harmonized Hub datasets should also support statistical and machine learning approaches to

forecast outbreak dynamics, providing insights into disease trajectories that can strengthen

preparedness and response planning.

Here, we independently tested Data Hub capacity to address empirical research questions,

seeking to understand the utility of this and similar data repositories to enable secondary

research and data validation, both for COVID-19 and other infectious and non-infectious

diseases.

Materials and Methods

Broad utility testing

RADx-rad data were accessed through the dbGaP portal (https://dbgap.ncbi.nlm.nih.gov/home).

We identified 44 RADx-rad studies with available datasets in the Data Hub

(https://radxdatahub.nih.gov/) and imported them into a Data Hub workspace configured with

AWS SageMaker Studio. The workspace, provisioned with 100 GB of disk space and 16 GB of

memory, provided direct data access. Each study appeared in a folder labeled by its dbGaP

accession number, containing data and dictionary files (CSV) and metadata (JSON). The

metadata included key identifiers such as subproject names, which were used for aggregation

and cross-study analysis. Analyses were performed in Jupyter Notebooks within a reproducible

CONDA environment. The notebooks, environment specification, and RADx-rad data

106 dictionaries annotated with tier information (https://github.com/radxrad/common-data-elements)

107 are publicly available on GitHub (https://github.com/radxrad/radx-analyzer).

108 *Wastewater analyses*

109 For comparative wastewater studies, sampling frame and locations were: i) NC A&T

110 (Greensboro, NC): grab or time-composite wastewater samples collected from dormitory sewer

111 laterals and central campus interceptors on scheduled sampling days (2021-2023), with

112 sampling twice per week during spring and fall semesters and once per week during summer;

113 and ii) Miami University (Oxford, OH): wastewater samples from community-scale sewersheds

114 with sampling once per week, available via the RADx Infectious Disease (RADx-rad) program

115 and accessed through the Data Hub. Samples from NC A&T were transported on ice,

116 concentrated, extracted, and quantified for SARS-CoV-2 RNA copies/L by RT-qPCR using

117 standard assays (e.g., N1/N2), with appropriate calibration and controls (4). For NC A&T data

118 aggregation, per-date mean across dorms was computed (zeros treated as missing if flagged as

119 below limit of detection (LoD)). For Miami University data (source Data Hub) aggregation, per-

120 date mean across sites was computed from processed RADx data. All time series are shown on

121 a log10 scale to accommodate multi-order-of-magnitude variation.

122 Examined variant windows included dominant variant periods (Alpha, Delta, Omicron BA.1,

123 BA.5, XBB), annotated as light-grey bands using widely accepted date ranges for U.S.

124 circulation (5,6). For site comparability, NC A&T dorm-level catchments represented smaller,

125 higher-resolution populations, whereas Miami University sewersheds represented broader

126 community populations, affecting signal amplitude and variance.

127 *Health Disparity Analyses*

128 We accessed six Data Hub datasets focused on community interventions to increase COVID-19

129 testing and vaccination among Latino populations in California, Maryland, New York, and

Florida. Search filters included the terms "Latino" and "Latinx," "community intervention,"

"COVID-19," and "social determinants of health." Participants completed structured

questionnaires assessing demographic, socioeconomic, and health-related characteristics,

including education level, employment status, sex, age group, and health insurance coverage.

COVID-19 positivity was determined through self-reporting or verified diagnostic test results.

Additional items captured perceptions and concerns about the COVID-19 vaccine, including

safety, development speed, effectiveness, and trust in government or healthcare systems. The

measures used included: i) COVID-19 Positivity Rate: Defined as the proportion of participants

who reported a positive COVID-19 test result within each subgroup (education, employment,

insurance type, sex, and age); ii) Vaccine Concerns: Categorized as safety/side effects, rapid

development, doubts about effectiveness, distrust of government, distrust of healthcare, or

religious/moral concerns; and iii) Sociodemographic Variables: Education (less than high

school, high school, some college, bachelor's, graduate), employment status (full-time, part-

time, unemployed, homemaker, self-employed, student), and insurance type (uninsured, public,

private). For data management, all six source files were combined, and variable names were

aligned across datasets to enable cross-study comparison. Data cleaning procedures

addressed inconsistent entries, including "missing," "other," and "prefer not to answer"

responses. Qualitative data were extracted from interviews and focus groups within each study,

capturing detailed information on community outreach efforts, vaccine concerns, and testing and

vaccination barriers. Both quantitative and qualitative measures were used to evaluate uptake,

participation rates, and participant perceptions of vaccination and testing programs. Data were

last accessed on November 5, 2025. Authors did not have access to data that could identify

individual study participants.

Qualitative data were analyzed thematically to identify recurring barriers, facilitators, and

perceptions related to COVID-19 testing and vaccination. Quantitative measures of participation

and vaccination uptake were summarized descriptively to contextualize cross-site intervention

effectiveness. Mixed-methods data integration provided a comprehensive understanding of how

social determinants of health, such as education, employment, and insurance access, affected

COVID-19 outcomes within Latino communities.

Retrospective characterization of RNA-seq library complexity

Raw sequencing volumes: Based on reported information retrieved from the published study,

specimen RNA-seq data exhibited a broad range of total sequencing volumes, between 1.5M-

160M reads sequenced per sample (median: 32.4M, IQR: 24.7M-39.4M) and transcriptome

read coverage levels between 11.4x-46.1x (median: 29.8x, IQR: 26.1x-33.5x). These values

corresponded to rates of sequenced reads per 1x transcriptome coverage equivalent between

117K-3.72M reads across specimens (median: 1.05M, IQR: 843K-1.27M).

Sequencing saturation metrics: Further inspection of the gene expression matrix supplied to the

Data Hub revealed substantial changes in the amount and dispersion for the total counts of

reads aligned uniquely to transcripts, between 28.6K-37.9M counts per sample (median: 4.30M,

IQR: 2.10M-6.85M) which corresponds to 1.50-1.28K fold difference in volume-to-aligned read

rates (median: 5.88, IQR: 4.50-11.2). Normalized volume-to-aligned read rates represent a wide

range of sequencing saturation levels across specimens in the expression matrix from the

original study. Interestingly, these rates corresponded to estimates for effective aligned depth,

i.e., the relation between total numbers of aligned reads and the 1x transcriptome coverage

equivalent rate in a specimen, between 0.01x-15.8x (median: 4.78x, IQR: 2.42x-7.05x) implying

that the overall gene expression matrix obtained from the original study, after unique sequenced

reads alignment and deduplication, is also sparse and under-sampled in terms of number of

genes detected and gene read counts per specimen, respectively.

Parametric focusing for specimen selection from unbalanced and sparse RNA-seq data

To handle the original Data Hub study's sparse expression matrix (7), we selected a reduced

search space of parametrically segregated, variably expressed (facultative) genes, i.e., gene

knee plot. This approach shows improved performance under unsupervised clustering analysis

compared to approaches that include all genes regardless of their total aligned read tallies (8).

The SALSA analytical workflow is a statistically agnostic method for biomarker extraction from

single-cell RNA-seq data; it segregates single cell representations (barcodes) agnostically,

based on latent patterns of shared gene expression and is tailored to process unary expression

data from large combinations of individual low-complexity libraries sequenced beyond

saturation. Further, to mitigate disparate library complexity saturation in gene expression

analysis impacts, we adopted parametric sweeping of rank-ordered total read counts per

barcode, i.e., specimen knee plot, a technique from scRNA-seq data processing by which

barcodes from putative cell multiplets (with an "excess" volume of reads), cell singlets (with a

"sufficient" volume of reads), and fractional cell debris (with a "shallow" volume of reads) are

segregated by their total read counts under a probabilistic mixture parametric model (8).

Transcriptome readout efficiency aggregate metrics supported our parametric-based

segregation of rank-ordered specimens based on total aligned reads, identifying subsets of

sequenced specimens in which gene expression information, which flows from total reads

sequenced to net aligned reads and transcriptome equivalent coverage, may lose

correspondence. The pool of 142 sufficient specimens (2.6M-19M total aligned reads per

sample) kept for unsupervised classification and differential expression analysis showed

matching ranges for transcriptome readout efficiency metrics that tracked commensurately

along the bioinformatics workflow: volume-to-aligned fold differences between 1.50-19.5

(median: 5.01, IQR: 4.34-6.66) and effective aligned depth rates between 1.59x-15.8x (median:

6.37x, IQR: 4.54x-7.72x). In contrast, 6 excess specimens were discarded from further analysis

(21M-38M total aligned reads per sample), displaying narrow transcriptome readout efficiency

metric ranges, suggesting overwhelming representation of RNA from highly expressed genes:

volume-to-aligned fold differences between 3.60-4.47 (median: 3.99, IQR: 3.72-4.45) and

effective aligned depth rates between 8.32x-11.1x (median: 9.58x, IQR: 8.78x-10.46x). 69

shallow specimens (29K-2.5M total aligned reads per sample) were also discarded from further

analysis, as they presented wide and mismatched transcriptome readout efficiency metric

ranges by orders of magnitude, characteristic of data from oversaturated and clonal sequencing

of low-complexity libraries: volume-to-aligned fold differences between 3.74-1.28K (median:

16.1, IQR: 8.61-27.5) and effective aligned depth rates between 0.01x-5.95x (median: 1.65x,

IQR: 0.83x-2.46x).

Precarity modeling

RADx-UP promoted equitable COVID-19 testing, vaccination, and care access among

disproportionately affected communities; it was the source of Precarity modeling data. The Data

Hub's Study Explorer and Metadata Specification tools assisted in selecting nine RADx-UP

projects, applying filters for study population, design, and variable availability to identify datasets

containing standardized Common Data Elements (CDEs) relevant to household hardship and

vaccination. Together, these projects contributed de-identified data from 5,397 adults across

diverse settings including rural, urban, and Tribal communities, representing historically

underrepresented populations in biomedical research.

Six household challenge variables were selected (including limited access to healthcare,

unstable housing, food insecurity, lack of clean water, difficulty obtaining medications, and

limited transportation) and modeled as observed indicators in a CFA to derive a latent Precarity

construct. Weighted least squares mean and variance-adjusted estimation (WLSMV) was used

for ordinal data, and model fit was assessed using χ^2, CFI, TLI, and RMSEA indices. Factor

scores from the validated model were then used in a multivariable logistic regression predicting

COVID-19 vaccination uptake, adjusting for education, age, sex, and insurance coverage.

Analyses were conducted in R (v4.3) using the lavaan and glm packages.

Results

1) Utility of RADx Study Data

1.1. Description of RADx-rad Studies used in Utility Testing

To test general Data Hub data utility, we analyzed 44 RADx-rad studies accessed through the

Data Hub and grouped into three research areas: i) wastewater-based surveillance, ii)

Multisystem Inflammatory Syndrome in Children (MIS-C) diagnosis, and iii) novel diagnostic

methods development. Collectively, these studies span community-level SARS-CoV-2

surveillance, clinical diagnostics, and point-of-care technology innovation.

Six subprojects focused on wastewater monitoring for population-level SARS-CoV-2 (and its

variants) detection. These studies evaluated viral RNA detection in environmental samples,

examined temporal trends, and benchmarked wastewater signals against case incidence data,

demonstrating wastewater surveillance as a scalable public health monitoring tool. Eight

Predicting Viral-Associated Inflammatory Disease Severity in Children with Laboratory

Diagnostics and Artificial Intelligence (PreVAIL-kIds) subprojects addressed MIS-C diagnostic

and biomarker discovery. These studies investigated biomarkers, clinical characteristics, and

outcomes prediction, providing insights into pediatric complications of SARS-CoV-2 infection in

vulnerable populations. The remaining 30 subprojects developed innovative diagnostic

technologies across six thematic areas (Table 1).

Table 1. Diagnostic Technologies

Technology	Description
SCENT (Electronic Noses)	VOC-based breath analysis for SARS-CoV-2 detection
Chemosensory Testing	At-home smell tests for rapid, low-cost screening

Exosome-Based Technologies	Microfluidic saliva assays for viral proteins and RNA, antibodies, and host exosome signatures
Novel Biosensing	Home-based diagnostics and breathalyzers for sensitive, rapid testing
Automatic Detection and Tracing	Electrochemical and mechanical sensors, touch-based detection, aerosolized virus monitoring, and privacy-preserving contact tracing
Multimodal Surveillance	Integrated systems such as smart masks, dialysis facility monitoring, and multimodal data streams for early detection and population surveillance

1.2. Analysis of Data Elements Across Studies

Each dataset included a data dictionary with standardized study variables and defined names,

types, and definitions, ensuring consistent representation and interpretation across studies. The

46 Tier 1 data elements, defined by NIH as describing patient-level metadata (demographics,

disability status, health history, mental health status, and COVID-related symptoms), were

harmonized across all RADx projects to enable cross-study analyses. Tier 2 elements, focused

on technology development, are harmonized across RADx-rad subprojects, while Tier 3 data

elements comprise project-specific variables that could not be further harmonized.

Both the relative proportions among Tiers 1-3 and the number of data elements within each tier

varied considerably across studies (Fig 1A-B). Although 46 Tier 1 data elements were defined,

not all were available or applicable to every study (Fig 1B). Tier 2 data elements showed

broader coverage across diagnostic technology studies, whereas Tier 3 data elements reflected

project-specific measurements. The PreVAIL-kIds (MIS-C) studies, in particular, exhibited

substantial diversity, using primarily Tier 3 data elements. This variation highlights the breadth

of RADx-rad data within the Data Hub and opportunities for further harmonization to enhance

cross-study comparability.

1.3. Analysis of Diagnostic Method Development Studies

RADx-rad diagnostic projects encompassed wide-ranging technologies, including

chromatographic, electrochemical, fluorescent, mechanical, spectroscopic, and olfactory

modalities (Fig 1C). This diversity reflects complementary signal generation strategies, designed

to optimize sensitivity and robustness across diagnostic contexts. Projects reported multiple

performance metrics categories (Fig 1D), including Binary classification metrics: sensitivity,

specificity, predictive values, percent agreement, and area under the ROC curve (AUC);

Analytical metrics: limit of LoD, limit of quantitation (LoQ), and limit of blank (LoB), used to

benchmark assay sensitivity; and technical metrics: cost per test and turnaround time,

addressing practical field deployment considerations.

Despite variable reporting, several platforms achieved >90% sensitivity and specificity, with low

LoDs enabling early infection detection. Cross-reactivity testing against influenza and seasonal

coronaviruses confirmed diagnostic specificity, while validation across Alpha, Beta, Delta,

Omicron, and other variants demonstrated robustness in dynamic epidemiological contexts (Fig

1E).

Diagnostic studies employed diverse specimen types, including saliva, breath, blood/serum, and

nasal swabs (Fig 1F). Saliva and breath were most common because they are non-invasive and

suitable for at-home testing. Analytes included viral spike proteins, proteases, and antibodies,

enabling detection across multiple stages of infection (Fig 1G). Biorecognition approaches

spanned antibodies, aptamers, enzyme-based reporters, and VOC recognition systems (Fig

1H).

2. Wastewater SARS-CoV-2 dynamics

2.1. Selection of Data Hub wastewater study and external comparator

To assess whether Data Hub-housed wastewater data may contextualize campus-level

surveillance, we paired (i) an external dataset from North Carolina Agricultural and Technical

State University (NC A&T) with (ii) a community-scale wastewater dataset from Miami

University, available in the Data Hub ("University of Miami SARS-CoV-2 wastewater

surveillance" study). The Miami University dataset was clearly documented, sampled repeatedly

over the COVID-19 period, and represented a broad, community-scale sewershed, suitable as a

reference for the higher-resolution NC A&T dorm/interceptor data. This pairing enabled testing

of cross-site comparability and the added value of Data Hub resources for campuses already

collecting local wastewater data.

2.2. Semester-linked wastewater dynamics across campuses

After harmonizing both datasets to daily means (NC A&T: per-date mean across dorms, zeros

treated as missing if below LoD; Miami: per-date mean across sites from the processed RADx

table), both campuses showed semester-linked trends with surges near term starts (Fig 2A),

consistent with prior findings that campus population shifts appear in wastewater signals (4).

Signal amplitude and sharpness differed by catchment scale (defined as the population and

sewer infrastructure feeding a given sampling point). Miami University (community-scale mean)

exhibited higher-amplitude, sharper peaks around campus returns, while NC A&T (dorm-level

mean) showed a lower baseline with episodic increases (4). These differences are expected,

because larger, community-scale sewersheds integrate more people and introductions, while

dormitory sewer laterals represent smaller, higher-resolution catchments with more stochastic

variation.

2.3. Alignment with SARS-CoV-2 variant periods

To interpret these peaks, we overlaid dominant U.S. variant periods (Alpha, Delta, Omicron

BA.1, BA.5, XBB) as contextual, light-grey bands (Fig 2A). The largest SARS-CoV-2 RNA

concentration increases (i.e., the highest peaks in the wastewater signal) at both campuses fell

within Delta and Omicron windows, matching clinical reports that Omicron lineages spread

faster and showed immune escape even with shorter clinical lead times (5,6). This concordance

indicates that Data Hub wastewater data can serve as a temporal reference for campus data

(Fig 2A).

2.4. Local vs. regional resolution from paired datasets

A key objective was to determine whether the paired datasets could distinguish local from

regional signals. NC A&T's dorm-level signals provided localized, actionable signals tied to

specific campus areas, supporting targeted communication or testing. In contrast, the Data Hub

series for Miami University provided broad, regional context and external validation, showing

whether campus increases reflected broader community rise. Together, these findings

decompose the wastewater signal into campus trends relative to the surrounding community (4).

2.5. Biosurveillance capacity from metagenomics

Finally, to determine whether the workflow could support endpoints beyond SARS-CoV-2, we

summarized NC A&T metagenomics results. Campus wastewater contained 361 bacterial, 6

fungal, and 5 protist taxa (Fig 2B), demonstrating that the same sampling and data-integration

workflow can support broader biosurveillance and environmental health monitoring (9). This

finding complements the comparative wastewater analysis and confirms that campus

wastewater also captures diverse microbial signatures, enabling future expansion to RSV,

influenza, and anti-microbial resistance (AMR) monitoring.

3. Analysis of COVID-19 in Diverse Populations

To assess the capacity to address COVID-19-related health disparities and evaluate

community-based intervention impacts on COVID-19 testing and vaccine uptake in Latino

populations, we used data from six NIH RADx studies across California, Maryland, New York,

and Florida. We combined quantitative and qualitative data, standardized variables, and

assessed structural and behavioral factors to evaluate engagement barriers and facilitators.

3.1. COVID-19 Positivity by Sex and Education Level in Latino Participants

COVID-19 positivity varied by both sex and education across Latino participants in the six RADx

studies (Fig 3A). Participants with less than a high school education had the highest positivity

rates (60% of women and 57% of men), indicating increased vulnerability among individuals

with limited access to stable employment, healthcare, and preventive resources (Fig 3A).

Positivity decreased with higher educational attainment. Among participants with a bachelor's

degree, 22% of women and 29% of men tested positive; among those with graduate degrees,

rates fell to 20% for women and rose to 47% for men. The unexpectedly higher rate among

graduate-educated men likely reflects a smaller sample size (low *n*) and potential occupational

heterogeneity.

Participants with associate's degrees or some college education showed moderate positivity

rates (28-39%), possibly influenced by job type and workforce participation. Men with "some

college" exhibited slightly higher rates (39%) than women (33%), possibly due to employment in

construction, manufacturing, transportation, and service roles where physical distancing and

remote work were limited. Women consistently showed lower positivity rates across most

educational levels, possibly relating to differences in occupational exposure.

3.2. COVID-19 Positivity by Age and Insurance Type

Participants aged 30-39 years had the highest positivity rates across insurance types, reaching

37.8% among those with private insurance, 34.1% among the uninsured, and 33.3% among

those with public insurance (Fig 3B). This pattern suggests that early- and mid-career adults

faced substantial exposure regardless of insurance coverage. Among participants aged 40-49,

positivity remained elevated across insurance groups (32.5-36.0%), then declined with age. In

the 50-59 group, positivity fell to 23.8-31.2%, with the highest among those on public insurance,

potentially reflecting underlying health vulnerabilities and increased testing access through

public programs.

Participants aged 60-69 showed a pronounced disparity: 29.5% positivity among those with

public insurance compared to 17.3% for the uninsured and 22.7% with private insurance,

suggesting differential testing access and chronic disease burden among older Medicare or

Medicaid beneficiaries. In the 70+ group, positivity was lowest among publicly insured

participants (16.7%) but increased to 27.8% for those with private insurance. Among young

adults (18-29), positivity was consistent across all insurance types (~25%), suggesting that

social behaviors and occupational exposure, rather than coverage, drove infection risks (Fig

3B).

Overall, insurance coverage did not uniformly protect against infection; exposure risks,

occupational context, and healthcare access jointly influenced outcomes. Age-stratified

disparities highlight the need for targeted outreach and prevention strategies, especially for

middle-aged Latino adults balancing high exposure risk and variable healthcare access. These

findings reflect the intersection of economic precarity and inadequate healthcare access, where

uninsured, working-age adults experienced delayed testing and reduced preventative care.

3.3. COVID-19 Positivity by Employment Status in Latino Participants

Participants who were employed part-time (37.6%) and self-employed (34.5%) had the highest

positivity rates (Fig 3C), possibly due to inconsistent workplace protections, lack of paid sick

leave, and dependence on multiple income sources that increase daily contact. Unemployed

participants (34.1%) showed comparably high rates, likely influenced by prior exposure in high-

contact jobs or ongoing socioeconomic vulnerability. Full-time workers (32.6%) had slightly

lower positivity, possibly due to more stable employment with structured safety protocols or

remote work options. Homemakers (19.4%) and students (17.1%) had the lowest infection

rates, consistent with reduced occupational exposure and greater ability to quarantine (Fig 3C).

These trends indicate that unstable employment, lack of paid sick leave, and absence of

workplace protections may heighten exposure among Latino populations employed in essential

sectors such as agriculture, food service, construction, and custodial labor.

3.4. Vaccine Concerns and Community Trust

Latino participant vaccine attitudes reflected safety concerns, misinformation, and institutional

mistrust (*Fig 3D*). Fear of side effects or doubts about vaccine safety, reported by 1,580

participants, highlighting enduring influence of misinformation and limited access to culturally

relevant health communication. Media influence was substantial; 1,100 participants indicated

that their views were shaped by media coverage, often inconsistent or sensationalized early in

the pandemic. Similarly, 1,050 participants were unsure of the vaccine's effectiveness, and 980

participants believed it had been developed too quickly, reflecting both scientific uncertainty and

communication gaps around vaccine trials and approval processes.

Trust and social concerns were also substantial (*Fig 3D*). Participants cited distrust of the

healthcare system or government (580 participants), reflecting longstanding inequities in

healthcare access and treatment that shape vaccine hesitancy in Latino communities. A smaller

subset of participants raised religious (350) or personal beliefs (200 personal freedom; 200

vaccine not needed) as factors influencing their decision, while 250 participants said they did

not know what the vaccine contained, suggesting that knowledge gaps could be addressed

through accessible education campaigns. Overall, these findings demonstrate that vaccine

hesitancy among Latino populations is multifactorial, influenced by safety and efficacy concerns

and institutional distrust.

4. Biomarker Discovery

410 *4.1. Putative biomarker extraction by secondary use of peripheral blood RNA-seq data from*

411 *pediatric MIS-C patients*

412 We utilized RNA-seq data from 217 whole blood specimens collected under the RADx-rad

413 program from SARS-CoV-2 patients presenting with varying MIS-C severity (7). This secondary

414 analysis had two complementary objectives: i) to identify sets of highly discriminant expressed

415 genes that characterize specimens among latent groupings of transcriptional similarity, and ii) to

416 assess whether those groups aligned with reported MIS-C severity (*Fig 4A*).

417 Specimen libraries showed variable effective transcriptome representation, producing a sparse,

418 under-sampled cross-study gene expression matrix (see *Materials and Methods*). The tallies of

419 aligned read counts per gene were also heavily unbalanced across individual specimens,

420 despite deriving from standardized alignment and bioinformatics workflows (7). We reasoned

421 that the undersampled expression matrix in our study was similar ultra-sparse scRNA-seq

422 cell×gene datasets, so we could identify specimens with poor transcriptome readout efficiency

423 to discard from our analysis (8). This strategy yielded 6 specimens with excess data volume,

424 142 with sufficient data volume, and 69 with shallow data volume (*Fig 4B*) (see *Materials and*

425 *Methods*).

426 Based on gene expression matrix distribution features reminiscent of single-cell RNA-seq data

427 footprints, we adopted the SALSA analytical workflow (see *Materials and Methods*) (8), which

428 treats specimen-level data similarly to single-cell approaches, and was previously established to

429 retrieve candidate biomarkers from sparse and untargeted RNA-seq data (8,10). This also

430 allowed us to reduce the analytical search space for unsupervised biomarker extraction by

431 focusing statistical analysis on agnostically determined facultative genes. We focused our

432 unsupervised clustering analysis to ~3,300 facultative genes (*Fig 4C*) and applied dimensional

433 reduction of gene expression patterns via unsupervised clustering. In all, we identified 9 major

specimen clusters (M1 through M9), driven by expression patterns across 1,085 differentially

expressed genes (DEGs) grouped into 6 distinct gene expression modules. Sorting data by

cluster and gene-module memberships confirmed that the 142 sufficient specimens formed a

sparse specimen×gene expression matrix (*Fig 4D*).

4.2. Clustering of specimens by severity

We examined whether major specimen clusters overlapped with distinct MIS-C severity

patterns. We mapped reported MIS-C severity levels onto their corresponding specimens and

observed that severe MIS-C mapped heterogeneously, but concentrated within a subset of

major clusters, especially specimens from M5, M6, and M7 (*Fig 4E*). Among the 1,085 DEGs,

we identified candidate expression biomarkers most closely associated with major clusters. This

analysis yielded a minimal assignment probability baseline of 6.37×10^{-5}, with 135 DEGs having

assignment probabilities greater than the baseline to at least one cluster. We excluded genes

with a maximum assignment probability across clusters less than a random assignment

threshold of 1 in 270, resulting in 100 top-scoring candidate biomarkers. These biomarkers

tracked almost exclusively with one of seven latent gene expression classes (LC1 through LC7),

each corresponding to one or two combined major specimen clusters. We used two-way

unsupervised hierarchical assignment score clustering, which helped identify a smaller subset of

20 top-scoring biomarkers with the sharpest latent class membership probabilities by visual

inspection (*Fig 4G*).

Specimens from patients with severe MIS-C predominantly mapped to major clusters M5

through M7 (*Fig 4F*). Established correspondence between major clusters and latent gene

expression classes (M5 → LC3, M6 → LC6, and M7 → LC4, as shown in *Fig 4G*), linked

specimen donors with an enrichment of expression for distinct biomarker genes characteristic of

specimen groupings. We performed gene set enrichment analysis on the 100 candidate

biomarkers, grouped by their distinct latent class associations, to assess i) whether biomarkers

linked to specimens from donors with severe MIS-C reflected previously reported gene

expression findings in SARS-CoV-2-infected human primary cells and tissues; and ii) whether

severe MIS-C-associated gene sets represented enriched responses from specific anatomical

structures, cell types, or biomarkers (see *Materials and Methods*).

4.3. Cluster Characterization

Using Enrichr (11) we found that specimens exhibited varying informative biomarker potential

levels (Table 2). The M5, M6, and M7 clusters had transcriptional indicators of immune/blood,

optical, or lung systems (Fig 4H-I).

Table 2. Biomarker potential of Candidates

Cluster	Top-Scoring Candidate Number (Biomarkers)	Top-Scoring Candidates (Biomarker candidates in Bold Font)	Justification
M5	19 (7)	***ACSL1**, ALPL, BCL2A1, CD177, **G0S2**, HIF1A, ICAM1, IFITM2, IFITM3, **IL1R2**, **IRS2**, MGAM, MMP9, **NFKBIA**, **PFKFB3**, S100A8, S100A12, SNORA73B, **TNFAIP3***	Gene set enrichment showed strong association with transcriptional signatures upregulated in lung, sclera, and natural killer (NK) cells infected with SARS-CoV-2 or sampled from patients with mild to severe COVID-19. Functional ontologies enriched for monocyte-like immune responses.
M7	49 (2)	*ANPEP, APOL6, ARHGEF1, CD53, CR1, CSF2RB, CTSS, FCGR2A, FGL2, FPR1, **GBP2**, GCA, ICAM3, IFITM1, **IGHG2**, IGHM, IGLC2, IRF1, ITGA2B, ITGAM, JAK3, LCP1, LILRA5, LRRK2, LYZ, MLLT6, MPIG6B, MT-CO2, NIBAN1, NLRC5, NOP53, PARP14, PGD, PIK3AP1, PLEC, PRF1, RNF213, S100A4, SERPINA1, SLC2A3, SLC11A1, SMCHD1, SPOCK2, STAT1, SYNE1, TAP1, TAPBP, TMC8, VNN2*	Gene set enrichment showed strong association with transcriptional signatures upregulated in lung, sclera, and natural killer (NK) cells infected with SARS-CoV-2 or sampled from patients with mild to severe COVID-19. Functional ontologies enriched for neutrophil-like immune responses.

M6	15 (4)	*ASCC2, BCL2L1, BPGM, C9orf78, FECH, KRT1, MARCHF8, MBNL3, **NCOA4**, PSMF1, **RPIA**, SESN3, **SNCA**, TNS1, **TRIM58***	Gene set enrichment showed association with differentially regulated gene expression patterns, at different stages of SARS-CoV-2 infection and measured in blood from human patients. Functional ontologies enriched for erythroid-like expression signatures.

Our secondary analysis shows that blood specimens from patients with severe MIS-C display

immune signatures at the gene expression level consistent with enhanced phagocytic activity

(as in M5 and M7 genes), suggesting active hemolytic reactions in a subset of patients (as in

M6 specimens). These features can be identified through first-tier differential expression

screening using at least three of thirteen measurable biomarker genes.

5. Structural Hardship, Educational Attainment, and COVID-19 Vaccination: A

Confirmatory Factor Analysis of Precarity

5.1. Study of Structural Barriers in Vaccine Uptake

We analyzed pooled data from nine RADx-UP studies accessed through the Data Hub to

evaluate how cumulative household hardship interacts with educational attainment to shape

COVID-19 vaccination uptake. Together, these projects contributed data from 5,397 adult

participants across diverse populations and settings, including urban underserved communities,

syringe-exchange clients, medically underserved adults, and caregivers of children with

intellectual and developmental disabilities. The Data Hub's standardized architecture enabled

consistent vaccination status, demographic, and hardship indicator measurement across

studies. The combined sample was demographically diverse (53% female, 47% male, with a

mean age of 49 years), with 45% reporting education beyond high school. Overall, 76% of

participants were vaccinated against COVID-19, though coverage varied across populations.

5.2. Household Precarity Factor

To quantify cross-study cumulative household hardship, we applied a Confirmatory Factor

Analysis (CFA) on six harmonized hardship indicators available through the Data Hub's

Common Data Elements (12–14). These indicators included limited healthcare access, unstable

housing, food insecurity, lack of clean water, difficulty obtaining medications, and limited

transportation. All six ordinal variables were measured consistently across studies due to Data

Hub harmonization, allowing them to be modeled jointly as observed indicators of a shared

latent construct. As shown in Fig 5A, most participants reported no challenges, though minor

and major challenges were common, particularly for food security, transportation, and access to

healthcare and medications. These distributions provided the empirical basis for modeling

cumulative hardship using a latent factor approach.

The CFA results showed that all six indicators loaded strongly on a single latent household

Precarity factor (standardized loadings λ = 0.72 to 0.90). Model fit was excellent (χ^2[9] = 48.2, *p*

< 0.001; CFI = 0.998; TLI = 0.996; RMSEA = 0.046). Higher factor scores reflected greater

cumulative hardship and varied substantially across studies, geographic regions, and

educational levels. This validated Precarity score provided a unified structural hardship measure

across the RADx-UP projects and served as the primary independent variable for evaluating its

relationship with COVID-19 vaccination uptake in downstream regression models (Fig 5B).

5.3. Vaccination Uptake Model

We estimated a multivariable logistic regression model to evaluate how household precarity and

educational attainment jointly predict COVID-19 vaccination uptake, adjusting for age, age-

squared, sex at birth, substance use risk, and health insurance status. Educational attainment

was represented using two indicator variables: "high school/GED" and "beyond high school",

with "less than high school" as the reference category."

Greater household precarity was associated with lower vaccination odds (OR = 0.83, $p < 0.10$),

indicating reduced uptake as structural hardship increased. Participants with education beyond

high school had substantially higher odds of vaccination (OR = 1.51, $p < 0.001$), while those

with only a high school/GED resembled the reference group (OR = 0.97). The interaction

between precarity and education revealed a key moderating effect. The Precarity × Beyond

High School term was statistically significant (OR = 0.56, $p < 0.001$), demonstrating that the

protective effect of higher education diminishes sharply as household hardship increases (Fig

5C).

Discussion

The Data Hub was designed to be an exemplar of an integrated, FAIR-compliant data platform,

demonstrating how centralized and harmonized repositories can accelerate research for future

public health challenges. We independently investigated diverse uses, and found that the Data

Hub, a central repository with an integrated analysis platform (2) seamlessly facilitated

secondary research. The program's harmonized datasets provide a foundation not only for the

current pandemic response but also for future-ready diagnostic and surveillance systems.

First, using hub-based RADx-rad data, we found that cutting-edge diagnostic methods can be

developed and validated across diverse specimen types and viral variants. The Data Hub's

harmonized datasets enabled systematic, side-by-side innovative technology comparisons,

demonstrating that RADx-rad transformed infectious disease diagnostics, enabling rapid,

accurate, and scalable detection. Moreover, the Data Hub's secure workspace provided

integrated tools for data querying, analysis, and machine learning, which facilitated

reproducibility and accelerated discovery.

Secondly, we found that wastewater-based epidemiology (WBE) provided early, unbiased

detection of SARS-CoV-2 trends, often anticipating clinical activity, particularly around semester

transitions (4). Comparing NC A&T's dorm-level campus data with a Miami University's Data

Hub-based community-scale showed that catchment scale differences (dorm vs. community)

explained amplitude/variance differences and underscored the complementarity of local and

regional surveillance layers: local data are better for action; regional data are better for context.

Consistent with clinical and epidemiologic studies, we confirmed that the Delta variant was

associated with higher clinical severity, while Omicron lineages (BA.1/BA.5/XBB) exhibited

greater transmissibility and immune escape, driving rapid campus surges despite lower per-

infection severity than Delta (5,6). Operationally, because faster growth and immune escape

can stress campus operations (absenteeism, isolation housing) even when severity is lower,

WBE can help to detect and compare these dynamics in near-real time for targeted mitigation

(4).

We noted that interpretation is affected by low-copy samples near the LoD, occasional temporal

gaps, evolving clinical testing volumes, plumbing/site differences, and incomplete

flow/population metadata. To improve cross-site comparability in future work, we recommend

integrating normalization using pepper mild mottle virus (PMMoV) (15), expanding surveillance

to RSV/Influenza/AMR targets, and adopting dashboards that link wastewater, clinical, and

mobility data for campus-level decision support (9). Importantly, this analysis also shows that

Data Hub wastewater studies can serve as a reference layer when a campus already has local

wastewater but wants to understand whether its patterns match a broader community signal.

Regarding health disparities, we found evidence supporting prior observations that the COVID-

19 pandemic disproportionately affected marginalized United States communities, with Latino

populations experiencing some of the highest rates of infection, hospitalization, and mortality.

By integrating findings from six RADx-UP community intervention datasets across multiple U.S.

states to examine how social determinants and community engagement strategies shaped

COVID-19 outcomes among Latino populations, we found that that lower educational

attainment, unstable employment, and lack of insurance were strongly associated with higher

infection rates. These findings support the hypothesis that structural and social determinants of

health contribute substantially to Latino community infection risks and show the enduring

influence of structural inequities on health.

Vaccine hesitancy was a persistent barrier, driven primarily by safety concerns, perceived

vaccine development speed, and government institution distrust. Prior work in 2020 indicated

that Latino communities experienced a 54% excess mortality compared with 12% among non-

Hispanic Whites, and infection rates were over three times higher than the general population

(16). Vaccination rates lagged behind, with only 32% of Latinos vaccinated by May 2021

compared with 43% of non-Hispanic Whites (17). These disparities were partially influenced by

social determinants of health, including limited healthcare access, employment instability,

educational inequities, and structural barriers that reduce prevention and care opportunities.

Strengthening vaccine confidence requires transparent communication, community-driven

outreach, and linguistically and culturally tailored messaging delivered by trusted local voices

(18). These results reinforce the need for health equity frameworks that address upstream

determinants such as employment, education, and healthcare access, while leveraging

community trust to deliver public health interventions.

Nevertheless, this health disparity analysis is descriptive and cross-sectional, limiting the ability

to infer causality or temporal relationships between socioeconomic factors and infection risk.

Self-reported data may introduce recall or reporting bias, particularly regarding COVID-19

testing and vaccination attitudes. The sample may also not represent all Latino subpopulations,

and state-level differences were not statistically adjusted. Despite these limitations, the findings

provide valuable insight into disparity patterns that warrant further investigation. Future research

should include longitudinal monitoring and multisite collaborations to assess community

partnership durability, social media-based outreach impacts, and promotoras-led intervention

scalability for other minority populations. Mixed-method approaches integrating behavioral,

social, and contextual data will be critical for developing sustainable, inclusive strategies that

strengthen community resilience beyond the COVID-19 pandemic and ensure more equitable

responses to future public health emergencies.

SARS-CoV-2 infections can elicit both COVID-19 in adults and MIS-C in children, but these two

complex diseases are often researched and managed separately, impeding effective disease

detection and monitoring. This has slowed the discovery of novel actionable biomarkers and

phenotypic traits applicable to both diseases. As such, our understanding of commonalities

between COVID and MIS-C molecular physiology remains limited. Using Data Hub-based multi-

modal research data from adults and children diagnosed with COVID and/or MIS-C, we

evaluated potential indicators of SARS-CoV-2 infection that may predict COVID-19 or MIS-C

severity starting from clinical or molecular patient data.

The PreVAIL kIds cohort, a translational research consortium of eight independently-run

NICHD-led clinical studies with roughly 10,000 pediatric participants (19), offers molecular

profiling data amenable to multi-modal integration and biomarker discovery. Using whole blood

RNA-seq data from one of these studies (7), we found strong associations between MIS-C

severity and highly enriched immune-related transcriptional profiles, linking clinical observations

to underlying physiological mechanisms. These results posit testable molecular biomarker

candidates to forecast MIS-C incidence, severity, and progression across the SARS-CoV-2

infection and host response timeline. By amalgamating physiological and clinical traits across

severity levels, our secondary analysis reveals new analytical strategies to project multisystemic

disease severity in patients presenting with early symptoms or clinical SARS-CoV-2 infection

diagnoses.

The biggest challenge in processing these datasets came from disparities between total

sequenced and aligned read counts, possibly due to differentially saturated sequencing

libraries. To address this, we adopted a statistical workflow tailored for sparse expression

datasets (i.e., single-cell RNA-seq analysis) that pre-emptively mitigated statistical bias,

prioritized robustly measured gene expression levels, and optimized latent association

discovery between observed clinical phenotypes and molecular data profiles in a streamlined

fashion.

Finally, we show how the centralization of data within the Data Hub can enable equity-focused

research by harmonizing complex, multi-site data for advanced analytic modeling. The Data

Hub's harmonized datasets, transparent metadata, and analytic infrastructure enabled

reproducible cross-cohort modeling. Using CFA, we derived a unified Precarity score that

quantifies cumulative household hardship, a key dimension of structural vulnerability affecting

COVID-19 vaccine uptake.

The results reveal a central paradox: while education is typically protective for health behaviors,

its benefit diminishes when structural barriers are high. Individuals with higher education but

greater precarity were substantially less likely to be vaccinated. This pattern supports a

polysocial risk framework, which recognizes that overlapping social risks including housing,

transportation, and healthcare access, operate collectively rather than in isolation. From a policy

perspective, these findings suggest that interventions addressing upstream structural hardships,

rather than focusing solely on education or awareness, may yield greater, more equitable gains

in vaccine coverage.

While the Data Hub was effective in harmonizing and centralizing RADx Data, investigators

should be aware of some challenges. These include data missingness, including incomplete or

inconsistent demographic/clinical information and residual variability in study design and patient

populations. While adopting metadata and FAIR-aligned ontologies partially alleviates these

issues, further standardization may be required based on investigators' study questions. Data

access and governance frameworks can also limit rapid data access and cross-institutional

analyses. Implementing interoperable governance models and harmonization policies across

NIH-supported repositories could reduce such bottlenecks. Moreover, future work linking RADx

data (acute COVID infection data) with initiatives such as REsearching COVID to Enhance

Recovery (RECOVER; long COVID) (20), may facilitate investigations into long-term sequelae

of SARS-CoV-2 infection, while integrations with All of Us (21) or NHLBI's BioData Catalyst (22)

could accelerate broader research on health disparities, biomarker trajectories, and social

determinants of health on infectious disease outcomes.

Acknowledgements

We thank the participants of RADx-funded studies, the RADx Data Hub data generators and

coordinators including RADx study investigators, and the (C)DCCs that provided the foundation

that enabled this work. We thank Marcos Martinez-Romero, Alissa Fujimoto, Nilesh Mistry,

Ashley Sier, Aubrie Weyhmiller, Aria Henry, and Ashley Evans, and Leah Mason, for technical

assistance and operational support in the execution of this work. We thank current and former

RADx Data Hub Partners for their efforts on the RADx Data Hub.

Author Contributions

P.W.R., T.R., N.M., O.A.L., L.J.F., and K.N.S. wrote and revised the manuscript. M.D.K.,

M.U.A., J.S., and B.F. provided critical content input and feedback, and reviewed the

manuscript. M.A.K., M.A.M., S.S., and A.K. provided support as principal investigators of the

RADx Data Hub.

Funding

656 This work was supported by the National Institutes of Health (NIH) under Other Transactions

657 Authority award 1OT2DB000009.

658 **Competing interests**

659 The authors declare no competing financial interests.

Figure Legends

Figure 1. **Inventory of study characteristics across NIH RADx-rad projects.** Overview of data element tiers, specimen types, analytes, detection methods, and biological targets reported across RADx-rad subprojects. A. Distribution of Tier 1-3 data elements across studies, grouped by subproject. Tier 1 elements represent patient-level metadata harmonized across RADx projects; Tier 2 elements represent technology-development variables harmonized across RADx-rad subprojects; Tier 3 elements are project-specific and not further harmonized. B. Breakdown of Tier 1 data elements by category, including demographics, disability status, health history, mental health, and symptoms. Although 46 harmonized Tier 1 elements were defined, not all were reported in every study. C. Types and frequencies of signal-detection methods employed across diagnostic subprojects. D. Metrics reported to evaluate diagnostic performance, including sensitivity, specificity, predictive values, percent agreement, and area under the ROC curve (AUC), as well as analytical and technical measures such as limit of detection (LoD) and turnaround time. E. SARS-CoV-2 variants tested across subprojects, including Alpha (B.1.1.7), Beta (B.1.351), Delta (B.1.617.2), Omicron (BA lineages), and other variants of concern, as well as common respiratory viruses (e.g., human coronaviruses and influenza) included for cross-reactivity testing. F. Specimen types analyzed across subprojects, including saliva, breath, blood/serum, and nasal swabs. G. Classes of analytes detected by RADx-rad diagnostic methods, including viral proteins, RNA, antibodies, and host biomarkers. H. Biorecognition mechanisms used across subprojects, such as antibodies, aptamers, enzyme-based reporters, and volatile-organic-compound (VOC) recognition systems.

Figure 2. **Wastewater SARS-CoV-2 dynamics with campus microbiome context.** A. Miami vs NC A&T wastewater RNA over time (2021-2023). NC A&T = orange, Miami = blue; y-axis = RNA copies/L (log10). Dashed vertical lines mark semester starts. Gray bands indicate

dominant variant waves (Alpha, Delta, Omicron BA.1, BA.5, XBB). NC A&T = mean across

dorms per date; Miami = mean per date across sites. B. NC A&T wastewater microbial diversity

(metagenomics). Unique taxa detected: Bacteria 361, Fungi 6, Protists 5.

Figure 3. **COVID-19 testing and vaccine concerns among Latino participants.** A. COVID-19

positivity rates stratified by sex (male, female) and education level (less than high school, high

school graduate, some college or higher). B. COVID-19 positivity rates stratified by age group

(18-29, 30-49, 50+) and insurance type (insured, uninsured). C. COVID-19 positivity rates

stratified by employment status (full-time, part-time, unemployed, essential worker). D. The most

commonly reported COVID-19 vaccine concerns, including safety, efficacy, development speed,

and institutional trust. Data were pooled from six RADx studies of Latino communities across

California, Maryland, New York, and Florida. Percentages reflect the proportion of participants

within each subgroup reporting positive COVID-19 tests or specific vaccine concerns.

Figure 4. **Identification of biomarkers of COVID-19/MIS-C severity in children by**

secondary analysis of sparse RNA-seq data from whole blood specimens. A. Schematic of

RNA-seq analytical strategy to coalesce unsupervised clustering of participant transcriptomes

with clinical metadata from participants in phs002781 study. B. Identification of subsample of

142 specimens to include in secondary analysis based on comparability in volume of reads

sufficiency. C. Identification of 3,349 facultative genes of interest to leverage for expression

pattern recognition by unsupervised clustering based on density of expression. D. Specimen ×

Gene heatmap of gene expression fold differences with respect to the global mean of all

specimens, ordered by patterns of expression similarity grouped into gene modules (x-axis) and

major sample clusters (y-axis: M1 through M9). E. Plot of 2D projection of the hierarchical

clustering dendrogram of specimens, with coloring to indicate major cluster assignments. F.

Overlay of non-parametric density contours of disease severity onto constellation plot of

specimens, with inset labeling to indicate major cluster assignments. G. Hierarchical clustering

of estimated item-response probabilities via latent class analyses for a subset of 20 top-scoring

candidate biomarkers (left, heatmap: yellow to red gradient) and all 9 major specimen clusters

(right, heatmap: white to black gradient) that correspond with 7 latent classes. H. Heatmap of

enrichment significance for gene sets in the "COVID-19 Related Gene Sets 2021" database

from Enrichr using 100 top-scoring candidate biomarkers corresponding with major specimen

clusters per latent class analysis. I. Bar graphs of top 5 enriched gene sets in the "HuBMAP

ASCT+B augmented with RNAseq Coexpression" database from Enrichr using subsets of

genes from 100 top-scoring biomarkers which correspond with major clusters of specimens M5-

M7 and overlap with study participants presenting severe COVID-19/MIS-C.

Figure 5. **The Paradox of Education: How Structural Barriers Shape Vaccine Uptake.**

Panels A-C show how structural hardship interacts with education to influence COVID-19

vaccination uptake. A. The prevalence of six household challenges, healthcare, housing, food,

water, medication, and transportation, that together form the foundation of the Precarity

construct. B. The Confirmatory Factor Analysis (CFA) model combines these indicators into a

single Precarity score (factor loadings = 0.72-0.90). C. Adjusted odds ratios demonstrating that

individuals with greater household precarity were less likely to be vaccinated, and that high

precarity substantially reduced the protective effect of higher education. Data are drawn from

5,397 participants across nine RADx-UP studies representing diverse underserved U.S.

populations.

References

1. Tromberg BJ, Schwetz TA, Pérez-Stable EJ, Hodes RJ, Woychik RP, Bright RA, et al.

Rapid Scaling Up of Covid-19 Diagnostic Testing in the United States - The NIH RADx

Initiative. N Engl J Med. 2020 Sep 10;383(11):1071–7.

2. Martínez-Romero M, Horridge M, Mistry N, Weyhmiller A, Yu JK, Fujimoto A, et al. A Cloud-

Based Platform for Harmonized COVID-19 Data: Design and Implementation of the Rapid

Acceleration of Diagnostics (RADx) Data Hub. JMIR Public Health Surveill. 2025 Aug

20;11:e72677.

3. Ledgerwood-Lee M, Hubenko A, Ray P, Theriault Y, Brickner H, Vazquez LF, et al. A User-

Centered Service Model for Accelerating COVID-19 Diagnostic Innovation: The RADx-rad

Dx Core Approach. IEEE Open J Eng Med Biol. 2025;6:472–9.

4. Bhatia S, Maswanganye TN, Jeje O, Winston D, Lamssali M, Deng D, et al. Wastewater

Speaks: Evaluating SARS-CoV-2 Surveillance, Sampling Methods, and Seasonal Infection

Trends on a University Campus. Microorganisms. 2025 Apr 17;13(4):924.

5. Lewnard JA, Hong VX, Patel MM, Kahn R, Lipsitch M, Tartof SY. Clinical outcomes

associated with SARS-CoV-2 Omicron (B.1.1.529) variant and BA.1/BA.1.1 or BA.2

subvariant infection in Southern California. Nat Med. 2022 Sep;28(9):1933–43.

6. Nyberg T, Ferguson NM, Nash SG, Webster HH, Flaxman S, Andrews N, et al. Comparative

analysis of the risks of hospitalisation and death associated with SARS-CoV-2 omicron

(B.1.1.529) and delta (B.1.617.2) variants in England: a cohort study. Lancet Lond Engl.

2022 Apr 2;399(10332):1303–12.

7. Loy CJ, Sotomayor-Gonzalez A, Servellita V, Nguyen J, Lenz J, Bhattacharya S, et al.
Nucleic acid biomarkers of immune response and cell and tissue damage in children with
COVID-19 and MIS-C. Cell Rep Med. 2023 Jun;4(6):101034.

8. Lozoya OA, McClelland KS, Papas BN, Li JL, Yao HHC. Patterns, Profiles, and Parsimony:
Dissecting Transcriptional Signatures From Minimal Single-Cell RNA-Seq Output With
SALSA. Front Genet. 2020 Oct 9;11:511286.

9. Kitajima M, Ahmed W, Bibby K, Carducci A, Gerba CP, Hamilton KA, et al. SARS-CoV-2 in
wastewater: State of the knowledge and research needs. Sci Total Environ. 2020 Oct
15;739:139076.

10. Lozoya OA, Papas BN, inventors; US Department of Health, assignee. Massively paralleled
multi-patient assay for pathogenic infection diagnosis and host physiology surveillance
using nucleic acid sequencing. United States patent application US 18/031,165. 2023 Nov
23.

11. Xie Z, Bailey A, Kuleshov MV, Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene Set
Knowledge Discovery with Enrichr. Curr Protoc. 2021 Mar;1(3):e90.

12. Kline RB. Principles and practice of structural equation modeling. Guilford publications;
2023 May 24.

13. Bollen KA. Structural equations with latent variables. John Wiley & Sons; 1989 May 12.

14. Schumacker RE, Lomax RG. A Beginner's Guide to Structural Equation Modeling. 4th ed.
Routledge; 2016.

15. Rainey AL, Liang S, Bisesi JH, Sabo-Attwood T, Maurelli AT. A multistate assessment of

population normalization factors for wastewater-based epidemiology of COVID-19. Sambri

V, editor. PLOS ONE. 2023 Apr 12;18(4):e0284370.

16. Riley AR, Chen YH, Matthay EC, Glymour MM, Torres JM, Fernandez A, et al. Excess

mortality among Latino people in California during the COVID-19 pandemic. SSM - Popul

Health. 2021 Sep;15:100860.

17. Bigelow BF, Saxton RE, Martínez DA, Flores-Miller A, Shin JM, Parent C, et al. High Uptake

and Series Completion of COVID-19 Vaccine at Community-Based Vaccination for Latinos

With Limited English Proficiency. J Public Health Manag Pract JPHMP. 2022 Dec

1;28(6):E789–94.

18. Buro AW, Roman Candelaria K, Bailey R, Luna F, Albizu-Jacob A, Stern M, et al.

Exploration of Multilevel Barriers and Strategies That Affected Early COVID-19 Vaccination

and Testing in Rural Latino Communities in Southwest Florida. Int J Environ Res Public

Health. 2022 Sep 18;19(18):11785.

19. Li GX, Gopchandani K, Brazer N, Tippett A, Choi C, Hsiao HM, et al. Clinical Features and

Outcomes of Pediatric and Adult Patients Hospitalized for Coronavirus Disease 2019: A

Comparison Across Age Strata. Open Forum Infect Dis. 2024 Jul 30;11(8):ofae443.

20. Reese JT, Blau H, Casiraghi E, Bergquist T, Loomba JJ, Callahan TJ, et al. Generalisable

long COVID subtypes: findings from the NIH N3C and RECOVER programmes.

eBioMedicine. 2023 Jan;87:104413.

21. The All of Us Research Program Investigators. The "All of Us" Research Program. N Engl J

Med. 2019 Aug 15;381(7):668–76.

22. Ahalt S, Avillach P, Boyles R, Bradford K, Cox S, Davis-Dusenbery B, et al. Building a

collaborative cloud platform to accelerate heart, lung, blood, and sleep research. J Am Med

Inform Assoc. 2023 Jun 20;30(7):1293–300.

A

COVID-19 Positivity Rates by Sex and Education Level in Latino Participants

B

COVID-19 Positivity Rates by Age and Insurance Type in Latino Participants

C

COVID-19 Positivity Rates by Employment Status in Latino Participants

D

Top COVID-19 Vaccine Concerns in Latino Participants

A

B

C