Reporting quality in psychological interventions. Implications for cumulative science and evidence-based practice: A scoping review.

Vives, J.[1], Lorente, S.[1], Casellas, A.[2], López-González, E.[3],
Marín-García, J.A.[4], Molla, C.[3], Rodrigo, M.F.[3], Viguer, P.[3], and Losilla, J.-M.[1]

[1] Universitat Autònoma de Barcelona; [2] Universitat de Vic-Universitat Central de Catalunya
[3] Universitat de València; [4] Universitat Politècnica de València

Abstract

Background: Poor quality in research reporting is a persistent issue that has hindered the advancement of scientific knowledge across all health science disciplines. Similarly, the progress of evidence-based practice has been hampered, as the replication of interventions is often impeded by the incomplete reporting of implementation details.

Objective: To map how reports of psychological interventions describe implementation at both the whole intervention level and the level of recipients' behaviours or actions (BeA), and to examine whether reporting quality varies by field, guideline use, or indicators of intervention complexity.

Methods. This scoping review followed PRISMA-ScR. Searches were conducted on PsycINFO (ProQuest) and Web of Science (Clarivate) for publications (2014-2024) describing the development and/or implementation of psychological interventions in clinical, public health, social/organizational or educational fields. A random sample of 200 publications per field was screened for inclusion. Following a two-stage selection process involving double screening and consensus resolution, 80 studies were included. Data were extracted using a Delphi-based instrument and analysed using descriptive statistics and χ^2 tests to compare different aspects of reporting across fields, guideline use, and complexity features.

Results. Only 11.3% reported development-only objectives, while most combined development with effectiveness (40%) and feasibility (48.8%). Reporting quality tools were seldom used (13.8%). Interventions typically targeted a single recipient category (87.5%); multilevel targeting was rare (6.3%). Implementation sequence was often replicable (73.8%), but structured aids were underused (tables/lists 38.8%; diagrams 16.3%). Costs were rarely reported (3.8%). At the BeA level, structural features were usually replicable (e.g., type/form of administration, periodicity, place, timing), whereas content/materials (36.3% and 47.5%), adaptation/customization (35%), and fidelity/adherence (27.5%) were frequently incomplete. Few significant between field differences emerged. Reporting tended to improve as the number of BeA increased.

Conclusions. Psychological intervention reports commonly omit critical replicability details, especially provider/evaluator prerequisites, content/materials, adaptation rules, fidelity, and costs, despite available guidance. Routine use of structured representations (e.g., flow diagrams, BeA tables) and established reporting tools (e.g., TIDieR, CONSORT-SPI) could close the most consequential gaps and facilitate replication and implementation at scale.

Registration: INPLASY 2025.7.0098.

Keywords: Psychological interventions; Reporting quality; Implementation; Fidelity; Complex interventions; Scoping review.

Introduction

Evidence-based practice is grounded in cumulative science, which is usually based on systematic reviews of the results of primary studies. However, the ability of a systematic review to synthesize evidence is often reduced, or even precluded, by deficiencies in the reporting of primary studies (Turner et al., 2012; Losilla et al., 2018). Poor quality in research reporting, whether due to a lack of completeness, transparency and/or precision, is a pressing problem that has hindered the advancement of scientific knowledge across all health science disciplines for decades (Glasziou et al., 2014; Rosengaard et al., 2024). Moreover, inadequate reporting of research not only may prevent its replication, but it is also a major source of research waste, which in areas such as biomedicine is estimated to account for 85% of research resources (Ioannidis et al., 2014; Glasziou & Chalmers, 2018; Almaqrami et al., 2020; Blanco et al., 2020).

The quality of reporting in primary studies focusing on an intervention, whether concerning its development, feasibility, efficacy, effectiveness or efficiency, is not expected to differ. In the case of interventions, however, another key implication arises in relation to evidence-based practice. If the characteristics related to the implementation of proposed psychological interventions are not reported with sufficient information, it is not possible to faithfully replicate them. Consequently, any practice involving the application of such interventions cannot truly be considered evidence based.

The emergence of reporting guidelines, such as CONSORT (Begg, 1996; Moher, Schulz & Altman, 2001; Schulz et al., 2010; Hopewell et al., 2025), STROBE (Vandenbroucke et al., 2007), PRISMA (Liberati et al., 2009; Page et al., 2021), SPIRIT (Chan et al., 2013, 2025) or TREND (Des Jarlais, Lyles & Crepaz, 2004), has led to an improvement in the quality of research reporting. Nevertheless, the impact of these guidelines and authors' adherence to them remain in need of improvement (Bastuji-Garin et al., 2013; Mannocci et al., 2014; Clayson et al., 2019; Orduña-Malea et al., 2023). In the case of interventions, the recent publication of guidelines to develop interventions, such as those focusing on complex interventions of Medical Research Council (MRC) (Craig et al., 2008; Tanner-Smith & Grant, 2018; Skivington et al., 2021) and the Agency for Healthcare Research and Quality (AHRQ) (Guise et al., 2017b), and the availability of intervention-specific reporting tools and guides such as CReDECI 2 (Möhler, Köpke & Meyer, 2015; Tanner-Smith & Grant, 2018), GUIDED (Duncan et al., 2020), TIDieR (Hoffmann et al., 2014), iCAT_SR (Lewin et al., 2017) and RIPI-f (Lopez-Alcalde et al., 2022), may have contributed to some improvement in the reporting quality of interventions.

Note that some of these guides target complex interventions and address how the factors contributing to their complexity should be reported to enable accurate replication and evaluation. Taking the factors that define the complexity of an intervention into account allows for a more thorough description of the characteristics of published interventions and to explore whether these factors, which can be challenging to report appropriately, are related to the reporting quality. Although several definitions of complex interventions attempt to distinguish them from non-complex ones, complexity is neither a binary characteristic nor a property of a single type of intervention (Petticrew, 2011; Lewin et al., 2017). Rather, any intervention can

be placed on a continuum ranging from a low degree to a high degree of complexity (Moore et al., 2017). While the debate surrounding the concept of 'complex intervention' is still ongoing, the most influential conceptualizations published in recent years, such as those of the MRC and the AHRQ aforementioned, agree that, in an intervention, complexity may arise, first and foremost, from the properties of the intervention itself, being the presence of multiple components, and the relations between them and the outcomes (especially moderations and mediations) the most relevant properties, along with the adaptability or customizability of the protocol.

Partitioning the intervention into components may facilitate the study of its efficacy and effectiveness (Petticrew et al., 2015; Skivington et al., 2021). However, none of the most influential definitions of complex interventions clearly defines what a component is. The only definition of component we are aware of is offered by iCAT_SR (Lewin et al., 2017), where a component is defined as a "discrete, active element of the intervention that could be implemented independently of other elements" (op. cit., p. 3). Based on this definition, each behavior or action (BeA) promoted by the intervention in its recipients could be considered the atomic constituent of an intervention. Each BeA may have specific implementation characteristics, which may at the same time have a verifiable effect on one or more outcomes, so that its contribution, whether direct, moderated or mediated, to the change pursued by the intervention can be better identified in a detailed and replicable report.

While the presence of multiple components and the relations that may derive from them are the first source of complexity, the second source arises from the system (e.g., an education or healthcare system, or a company) in which the intervention operates (Hawe, Shiell & Riley, 2009; Rutter et al., 2017). A system may be a multidimensional entity composed of multiple organization layers, each hosting independent yet interacting processes. When complexity is examined at the system level, an intervention can be considered an event expected to contribute to change at one or more levels and/or categories of recipients (e.g., users, professionals, managers, policymakers). Therefore, the number of levels and categories of recipients targeted by the intervention is another important factor contributing to its complexity.

Even though it is unreasonable to implement an intervention for which there is no empirical evidence of its efficacy in achieving the desired outcomes, it is also not justifiable to implement it if the report of their development does not contain the necessary information to do so in the precise manner intended by its authors. In order to improve the reporting of interventions, the MRC (Skivington et al., 2021) encourages the publication of their development separately from the assessment of its results (such as efficacy, effectiveness, efficiency or feasibility). It is to be seen the impact of this recommendation on the literature reporting interventions.

In this study, our aim was to examine the salient features of intervention reports and to explore whether some of these features are related to their quality. To ensure certain degree of homogeneity and according to our expertise, we focused on psychological interventions. According to Munn (2018), a scoping review represents the most appropriate type of systematized review to address such exploratory objectives. Specifically, our scoping review aimed to:

a) Describe the most common features of the intervention reports.

b) Analyze the replicability of the interventions.

c) Explore the presence of elements that characterize complex interventions and whether they
are related to the quality and replicability of intervention reports.

d) Analyze the relationship between the quality of the intervention report and its replicability
with: 1) the objective of the intervention report (devoted exclusively to the description of its
development and its implementation, or also including the evaluation of its feasibility, efficacy,
effectiveness or efficiency); 2) the research field to which the intervention belongs (clinical,
public health, social and organizational, or education); and 3) the use of reporting tools or
guidelines.

Method

This review was conducted following the Preferred Reporting Items for Systematic Reviews
and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) (Tricco et al., 2018). The
protocol was prospectively registered at INPLASY (Vives et al., 2025b).

In accordance with the principles of open science, all datasets and supplementary information
referenced in the following sections are publicly available in the CORA_RDR public repository
and can be accessed at https://doi.org/10.34810/data2792. The PRISMA-ScR checklist applied
can be consulted in S1_PRISMA-ScR.

Search strategy

The search was conducted in PsycInfo (Proquest) and Web of Science (Clarivate), with the aim
of collecting data from a representative sample of recently published studies focusing on
reporting on psychological interventions in the four application domains: clinical, public health,
social and organizational, and educational. The general search structure (("intervention*" OR
"treatment*" OR "therap*" OR "program*") NEAR/5 ("develop*" OR "protocol" OR
"feasibility" OR "effica*" OR "effectiv*")) was adapted to accommodate specific descriptors
of each of the four psychological research fields (see full search syntax in Appendix A).

Eligibility criteria

Eligible studies were those:

1. Describing the development or implementation of psychological interventions, whether they
also included empirical assessments of their feasibility, efficacy, effectiveness, or efficiency.

2. Providing information about the implementation process, even if parts of this process
referenced previous publications (e.g., a previous protocol, a previously published manual,
etc.).

3. Whose reported interventions belonged to the four research fields of clinical, public health,
social and organizational, or educational.

4. Published between 2014 and 2024, in order to focus this study on the recent research.

Selection process

A stratified random selection of 200 studies from each of the four target research fields was
made, followed by a random sequential selection of 21 studies, one of which was used to
perform the pilot test of data extraction, and the remaining 20 to extract the information of
interest for our objectives, including a total of 80 studies in the review. Two reviewers (JML,

JV) independently assessed the inclusion criteria based on the titles and abstracts of the studies
and made the final selection from the full text. Discrepancies during this selection process were
resolved by consensus between the two reviewers, with the participation of a third reviewer
specialist in the field of intervention (AC, SL, JM/MFR, CM/EL/PV) when necessary. The
degree of agreement between the raters was calculated using the Kappa index.

Data collection

References identified by the search strategy were entered into Zotero bibliographic software,
and duplicates were removed automatically.

Two authors (JML, JV) generated an initial list of items based on the review of: a) publications
focused on establishing the characteristics that determine the complexity of interventions (Craig
et al., 2008; Guise et al., 2017a; Rutter et al., 2017; Tanner-Smith & Grant, 2018; Skivington et
al., 2021); and b) tools and guides to assess the quality of intervention reports, with particular
attention to those that consider the complexity of interventions and incorporate guidelines or
elements in their proposals aimed at capturing the characteristics of the complexity of the
interventions. The tools that served as the basis for writing the items were SPIRIT (*Standard
Protocol Items: Recommendations for Interventional Trials*) (Chan et al., 2013; Hopewell et al.,
2022), iCAT-SR (*Complexity Assessment Tool for Systematic Reviews*) (Lewin et al., 2017),
Consort-SPI (Montgomery et al., 2018), CReDECI 2 (*Criteria for Reporting the Development
and Evaluation of Complex Interventions in healthcare*) (Möhler, Köpke & Meyer, 2015;
Tanner-Smith & Grant, 2018), RIPI-f (*Reporting Integrity of Psychological Interventions
delivered face-to-face*) (Lopez-Alcalde et al., 2022), StaRI (*Standards for Reporting
Implementation Studies*) (Pinnock et al., 2015, 2017), TIDieR (*Template for Intervention
Description and Replication checklist*) (Hoffmann et al., 2014) and TREND (*Transparent
Reporting of Evaluations with Non-Randomized designs*) (Des Jarlais, Lyles & Crepaz, 2004).

The same authors developed a second version of the items to be used for data extraction, this
time including response options. A two-round Delphi survey with all the research team (AC,
CM, EL, JM, JML, JV, MR, PV, SL) was conducted using this second version as the starting
point. The results of the Delphi allowed the development of a third version of the data extraction
spreadsheet. Then, all research team members participated in the pilot test, which involved
applying this third version to one published study from each of the four research areas. The
results of a pilot test, together with the research team's comments, led to modifications and/or
additions to items and response options that were deemed necessary, resulting in the final
version of the data extraction spreadsheet (see S3_Data_Extraction_Spreadsheet) with response
options protected by drop-down lists and checkboxes (see Fig. 1a and Fig. 1b).

Although most elements of the data extraction spreadsheet are relatively straightforward,
certain items warrant additional clarification. Items were categorized as green, which assessed
characteristics of the intervention not directly related to its implementation, or blue, which
evaluated whether the information reported on implementation was sufficient to enable accurate
and unambiguous replication. While both green and blue items shared response options such as
Yes and No, their interpretation differed: a Yes response for a green item indicated that the
statement was factually true (e.g., a Yes for Item G1a signified that at least one study objective
involved assessing the feasibility of the intervention), whereas a Yes response for a blue item
signified that the reported information was replicable (e.g., a Yes for Item G7 indicated that the
implementation sequence could be reproduced).

Figure 1a.
Data extraction. Report on the general characteristics of the intervention.

Item		Resp	Met
G0	Description	Specify	
G1	Objective/s of the study <u>in addition to the development</u> of the intervention program:	Resp	
	a. Evaluation of the **feasibility** of the intervention	Y/N	
	b. Evaluation of the expected effects (*efficacy, effectiveness*) of the intervention	Y/N	
G2	Previous publication of **intervention program** (completely o partially) or registration of the **intervention protocol**	R/P/C/N	
	a. Part of the intervention program is described in **non peer-reviewed documents** (e.g., unpublished papers, conference proceedings, grey literature)	Y/N	
	b. Part of the intervention program is described in **inaccessible documents** (e.g., broken links)	Y/N	
G3	Reporting **quality assessment tool** of the study	Y/N	Specify
		Resp	**Met**
G4	**Number of categories of recipients** (i.e., those whose intervention *effects* are tested; *individual level* -e.g. categories: patients, professionals, policy makers-, *group/team level* -e.g. categories: family, clinical staff, teaching staff, patient support groups-, or *system level* -e.g. categories: communities, health centers/hospitals, schools, organizations-)	O/M/L	☐
G5	**Facilitators or barriers** (i.e., characteristics or factors related to material resources, recipients, providers and/or context that may facilitate or hinder the implementation of the intervention)	Y/N	☐
G6	**Prerequisites for the <u>whole</u> intervention** (necessary to implement the whole intervention; should include a description of the previous tasks to be carried out by the providers, evaluators and/or recipients of the intervention, or specific characteristics of these or the context in which the intervention is implemented):	Resp	Met
	a. Providers (e.g., training, competencies/skills, irrelevant/unnecessary)	Y/D/I/N	☐
	b. Evaluators (e.g., training, competencies/skills, irrelevant/unnecessary)	Y/D/I/N	☐
	c. Recipients (e.g., training, competencies/skills, specific characteristics, irrelevant/unnecessary)	Y/D/I/N	☐
	d. From the context (*setting*) (e.g., social, cultural, political, economic, physical, geographical/spatial, digital, irrelevant/unnecessary)	Y/D/I/N	☐
G7	**Implementation sequence** (both the behaviors or actions -BeA- of the recipients and the tasks to be performed by providers and evaluators)	Y/I/N	☐
	a. Report includes a **diagram (flowchart or infographic)** of the intervention program	Y/N	☐
	b. Report includes **table/s (or list/s)** describing the structure of the intervention program	Y/N	☐
G8	**Intervention outcomes** (adding up the different outcomes of all BeA)	S/M	☐
G9	**Costs** (i.e., costs of the necessary resources, e.g., personnel, materials, equipment, for the implementation of the intervention)	Y/N	☐
G10	**General considerations** on the quality of the report or on problems for the replicability of the intervention	Specify	
	Extraction time:	Minutes	

211 Notes: Y: yes or reported and/or replicable; D: description only and not clearly stated; I: reported, but not replicable (not enough Information); N: no or not reported; R: protocol registered; P: intervention previously published (partially);
212 C: intervention previously published (completely); O: one category; M: multi-category; L: multi-level; S: single; M: multiple; BeA: behaviors and actions; Resp: response; Met: information included in the methods section.

214 Figure 1b.

215 *Data extraction. Report of each of the recipients' behaviors or actions (BeA).*

Item		BeA 1		[...]	BeA n	
B0	BeA **Description**	Specify			Specify	
			Met			Met
B1	**Levels and categories of recipients** (i.e., those whose intervention *outcomes* are collected)	Resp	☐		Resp	☐
	a. Individual level (e.g. categories: patients, professionals, policy makers)	N/O/M	Specify		N/O/M	Specify
	b. Group/team level (e.g. categories: family, clinical staff, teaching staff, patient support groups)	N/O/M	Specify		N/O/M	Specify
	c. System level (e.g. categories: communities, health centers/hospitals, schools, organizations)	N/O/M	Specify		N/O/M	Specify
		Resp	Met		Resp	Met
B2	**Facilitators or barriers** (i.e., characteristics or factors related to material resources, recipients, providers and/or context that may facilitate or hinder the implementation of the BeA).	Y/N	☐		Y/N	☐
B3	Outcomes and effects of a <u>particular</u> BeA:	Resp	Met		Resp	Met
	a. Number of outcomes (specific, but not exclusive, to particular BeA and measured on their recipients)	N/S/M	☐		N/S/M	☐
	b. Moderation or mediation effect on the outcomes of other BeA?	Y/N/U	☐		Y/N/U	☐
B4	**Adaptation/Customization or not** (i.e., protocol admits implementation flexibility)	Y/D/N	☐		Y/D/N	☐
B5	**Evaluation of the fidelity** of the implementation (including adherence)	Y/I/N	☐		Y/I/N	☐
B6	Prerequisites for a <u>particular</u> BeA:	Resp	Met		Resp	Met
	a. Providers (e.g., training, competencies, skills, qualification, profession, irrelevant/unnecessary)	Y/D/I/N	☐		Y/D/I/N	☐
	b. Evaluators (e.g., training, competencies, skills, irrelevant/unnecessary).	Y/D/I/N	☐		Y/D/I/N	☐
	c. Recipients (e.g., training, competencies, skills, personal characteristics, symptomatology, motivation, access to/or specific resources, irrelevant/unnecessary)	Y/D/I/N	☐		Y/D/I/N	☐
	d. From the **context** (*setting*) (e.g., social, cultural, political, economic, physical, geographical/spatial, digital, irrelevant)	Y/D/I/N	☐		Y/D/I/N	☐
B7	Implementation:	Resp	Met		Resp	Met
	a. Type of administration (i.e., remote, face-to-face, mixed, *tailored*).	Y/I/N	☐		Y/I/N	☐
	b. Form of administration (i.e., targeting a single individual/group/system, targeting 2 or more individuals/groups simultaneously, mixed, *tailored*)	Y/I/N	☐		Y/I/N	☐
	c. Supervision/support (i.e., directed [requires concurrent intervention of providers] with one-way interaction [e.g., supervision], directed with two-way interaction [e.g., on-demand support], autonomous [not require concurrent intervention of providers], mixed, *tailored*)	Y/I/N	☐		Y/I/N	☐
	d. Materials and support tools (e.g., audio, video, text/booklet/leaflet, computer hardware and software, mobile phone apps, gaming, pharmacological, etc., or irrelevant/unnecessary)	Y/I/N	☐		Y/I/N	☐
	e. Content and development of the sessions	Y/I/N	☐		Y/I/N	☐
	f. Periodicity (i.e., time unit; e.g., once, daily, weekly, monthly, *tailored*)	Y/I/N	☐		Y/I/N	☐
	g. Frequency per time unit (e.g., once, a certain number of times, *tailored*)	Y/I/N	☐		Y/I/N	☐
	h. Total frequency (e.g., specific number, *tailored*)	Y/I/N	☐		Y/I/N	☐
	i. Duration of sessions, dose or intensity (e.g., specific, *tailored*)	Y/I/N	☐		Y/I/N	☐
	j. Geographical, spatial, physical and/or digital place (e.g., home, specific/specialized place [e.g., school, clinic, gym, *outdoor*, etc.], *tailored*, irrelevant/unnecessary)	Y/I/N	☐		Y/I/N	☐
	k. Moment in time (e.g., specific, *tailored*, irrelevant)	Y/I/N	☐		Y/I/N	☐
	l. Social, cultural, political and/or economic setting, system, environment (e.g., specific, *tailored*, irrelevant)	Y/I/N	☐		Y/I/N	☐

216
Notes: Y: yes or reported and/or replicable; D: description only and not clearly stated; I: reported, but not replicable (not enough <u>I</u>nformation); N: no, none or not reported; O: one category; M: multi-category; L: multi-level; S: single; M: multiple; U: unclear; BeA: behaviors and actions; Resp: response; Met: information included in the methods section.

All items referring to information that should be reported in the Methods section included a
checkbox ("Met"), which was ticked if that information was indeed reported in that section.

Items were organized into two spreadsheets: one containing items that pertained to the whole
intervention, and another containing the items that affected each of the BeA of the intervention's
target recipients. Items G6 and B6 include a "D" response option ("description only and not
clearly stated"), which is absent in the other items. These items aim to capture information
about prerequisites related to providers, evaluators, recipients, and contextual factors. During
the pilot test, it was observed that some studies described the characteristics of the sample
(participants), the professionals who administered or measured the results, or the characteristics
of the intervention's implementation context, but did not specify whether this information
referred to prerequisites for implementing the intervention or was merely a description of how
the study was conducted. For these cases, response option "D" was considered.

The data extraction process was conducted similarly to the study selection process. Two
reviewers performed independently the data extraction. Discrepancies during this process were
resolved by consensus between the two reviewers, with the participation of a third reviewer
specialist in the intervention field (AC, SL, JM/MFR, CM/EL/PV) when necessary. Prior to
data extraction, the two raters identified and agreed on the BeA of each study according to the
definition provided in the introduction: any constituent of an intervention with specific
implementation characteristics that might also have a verifiable effect on one or more
intervention outcomes was considered a potential BeA.

Data analysis

A narrative synthesis of the results was carried out using descriptive statistical tables and bar
charts. The analyses were organized according to the specific objectives set for the scoping
review. Thus, a first descriptive analysis of the general characteristics of the interventions and
their BeA was carried out, and the possible differences between the four research fields were
explored. Secondly, possible differences in the quality of reporting (replicability) of items
related to the implementation of the intervention were evaluated due to: a) the research field of
the intervention; b) the use by the study authors of guidelines for writing the report; and c) the
characteristics of the intervention related to its complexity, such as the number of BeA, the
number of categories and levels of the intervention recipients, and the level of adaptability or
customization of the implementation reported by its authors (JAMOVI v.2.6.4).

The raw and curated data may be consulted in S4_Raw_Data and S5_Curated_Data,
respectively. Data extracted from the main variables is shown in Table 1.

Ethics statement

This scoping review does not require approval from an ethics committee, as no data was
collected directly from humans or animals.

Table 1.
Data extraction. Main variables.

ID	Field	G_1	G_2	G_3	G_5	G_7	G_9	nBeA	G_4	BeA_1a	BeA_1b	BeA_1c	G_6	BeA_6	G_8	BeA_3a	BeA_3b	BeA_4	BeA_5	BeA_7
Adnan, 2023	Soc	Fea; Eff	NP	No	No	Y	No	2	O	O	No	No	I	I	>1	0	No	No	N	I
Aharonovich, 2018	Clin	Fea; Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	0	No	No	N	I
Ahlqvist-Björkroth, 2017	Hea	Dev	NP	No	No	Y	No	1	L	M	O	No	Y	Y	>1	0	No	No	N	I
Akkermans, 2015	Soc	Eff	CP	No	No	Y	No	1	O	O	No	No	I	I	>1	0	No	NCS	N	I
Andruschko, 2018	Edu	Fea; Eff	NP	No	Yes	N	No	4	O	O	No	No	I	I	>1	0	No	Yes	N	I
Appiah, 2022	Hea	Fea	NP	No	Yes	Y	No	1	O	O	No	No	I	I	>1	0	No	No	N	I
Berman, 2016	Clin	Dev	NP	No	No	Y	No	1	O	O	No	No	N	N	>1	>1	No	NCS	N	Y
Bork, 2020	Edu	Eff	NP	No	No	Y	No	2	O	O	No	No	N	N	>1	0	No	No	N	I
Boudreaux, 2015	Hea	Fea; Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	NCS	Y	I
Caporale-Berkowitz, 2021	Soc	Eff	CP	No	No	Y	No	2	O	O	No	No	N	N	>1	0	No	NCS	N	I
Capps, 2024	Edu	Fea; Eff	NP	Yes	Yes	Y	No	4	M	M	No	No	I	Y	>1	>1	No	Yes	Y	Y
Christensen, 2014	Hea	Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	NCS	N	I
Chu, 2015	Edu	Fea; Eff	CP	No	No	Y	No	2	L	O	O	No	I	I	>1	0	No	Yes	Y	Y
Cohrs, 2019	Edu	Eff	NP	No	No	Y	No	2	O	O	No	No	I	I	>1	0	No	NCS	I	I
Coker, 2019	Soc	Dev	NP	No	No	N	No	1	O	O	No	No	N	N	>1	0	No	No	N	I
Drake, 2024	Hea	Eff	NP	No	No	Y	No	1	O	O	No	No	N	N	>1	0	No	NCS	N	I
Eklund, 2021	Edu	Fea; Eff	PP	No	No	Y	No	2	O	O	No	No	I	I	>1	0	No	NCS	I	I
Elderson-Van Duin, 2023	Edu	Eff	NP	No	No	Y	No	3	O	No	O	No	I	I	>1	0	Yes	Yes	N	I
Emerson, 2017	Edu	Fea; Eff	NP	No	Yes	I	No	3	O	O	No	No	I	I	>1	0	No	Yes	N	I
Erturk Kilic, 2023	Soc	Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	NCS	N	I
Facchini, 2016	Hea	Fea	NP	No	Yes	Y	Yes	5	M	O	O	No	Y	I	>1	0	Yes	Yes	N	Y
Faretta, 2016	Clin	Dev	CP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
Farrand, 2017	Clin	Dev	NP	No	No	Y	No	3	O	O	No	No	Y	Y	>1	>1	No	Yes	N	Y
Ferolino, 2024	Soc	Fea; Eff	PP	No	No	Y	No	4	M	O	O	No	I	I	>1	0	Unclear	No	N	I
Fletcher, 2024	Soc	Dev	NP	No	No	N	No	2	O	O	No	No	Y	Y	>1	0	No	No	N	I
Fung, 2018	Hea	Eff	NP	No	No	N	No	4	O	O	No	No	I	I	>1	0	No	Yes	N	I
Georgiou, 2021	Soc	Eff	NP	No	No	I	No	1	O	O	No	No	I	I	>1	0	No	No	N	I

ID	Field	G_1	G_2	G_3	G_5	G_7	G_9	nBeA	G_4	BeA_1a	BeA_1b	BeA_1c	G_6	BeA_6	G_8	BeA_3a	BeA_3b	BeA_4	BeA_5	BeA_7
Goldenthal, 2024	Edu	Fea	NP	No	Yes	Y	No	4	L	O	No	No	I	I	>1	0	Yes	No	N	I
Hettiarachchi, 2022	Edu	Eff	NP	No	Yes	Y	No	3	O	O	No	No	I	I	1	0	Unclear	Yes	N	I
Huang, 2016	Hea	Fea	PP	No	Yes	Y	No	3	O	O	No	No	I	I	>1	0	No	No	Y	I
Hussong, 2024	Hea	Fea; Eff	NP	No	No	Y	No	3	O	O	No	No	I	I	>1	0	Unclear	Yes	I	I
James, 2020	Soc	Fea	NP	No	No	Y	No	7	O	O	No	No	I	I	>1	0	Yes	Yes	N	I
Janka, 2022	Soc	Dev	Pr	No	No	Y	No	3	O	O	No	No	Y	Y	>1	0	No	No	Y	I
Jarero, 2016	Clin	Fea	PP	No	No	Y	No	8	O	O	No	No	I	I	1	0	Yes	Yes	N	I
Jiwani, 2024	Hea	Eff	NP	No	No	N	No	4	O	O	No	No	Y	Y	1	0	Unclear	Yes	N	I
Johnsen, 2024	Edu	Eff	Pr	No	No	Y	No	5	M	O	O	No	Y	Y	>1	0	Yes	Yes	I	I
Kahler, 2014	Hea	Fea; Eff	PP	No	No	Y	No	8	O	O	No	No	I	Y	>1	0	No	Yes	Y	I
Kelly, 2024	Soc	Dev	Pr	Yes	No	N	No	4	O	O	No	No	Y	Y	>1	>1	No	Yes	I	I
Kemer, 2022	Edu	Eff	NP	No	No	Y	No	2	O	O	No	No	Y	Y	>1	>1	No	No	N	Y
Kjellström, 2022	Soc	Eff	Pr	Yes	No	I	No	3	M	M	O	No	I	Y	>1	>1	No	No	N	I
Knaevelsrud, 2017	Hea	Fea; Eff	Pr	No	No	I	No	3	O	O	No	No	I	Y	>1	>1	No	No	Y	I
Kokubo, 2023	Soc	Eff	NP	No	No	I	No	4	O	O	O	No	I	Y	>1	>1	No	No	N	I
Komischke-Konnerup, 2023	Clin	Fea; Eff	NP	No	No	Y	No	3	O	O	No	No	I	I	>1	0	Yes	Yes	N	I
Kopelman-Rubin, 2019	Clin	Fea; Eff	NP	No	No	Y	No	4	O	O	No	No	Y	Y	>1	0	Yes	Yes	Y	I
Koponen, 2021	Edu	Fea; Eff	PP	No	No	I	No	5	O	O	No	No	I	Y	>1	>1	No	Yes	Y	I
Kraiss, 2018	Clin	Dev	Pr	No	No	Y	No	4	O	O	No	No	I	I	>1	0	Yes	No	N	I
Kulkarni, 2014	Edu	Eff	NP	No	No	Y	No	3	O	O	No	No	Y	Y	>1	>1	No	NCS	N	Y
Kuschner, 2017	Clin	Fea	NP	No	Yes	Y	No	4	L	O	O	No	I	Y	>1	0	No	No	N	Y
Kwan, 2023	Soc	Fea; Eff	NP	Yes	Yes	Y	No	2	O	O	No	No	I	Y	>1	>1	No	No	Y	I
La Greca, 2016	Hea	Fea; Eff	PP	No	Yes	I	No	2	O	O	O	No	I	Y	>1	>1	No	NCS	Y	I
Lovett, 2021	Edu	Fea; Eff	PP	No	Yes	Y	No	3	L	No	O	O	Y	Y	>1	>1	No	Yes	Y	I
Maj, 2024	Clin	Eff	Pr	No	Yes	Y	No	1	O	O	No	No	I	I	>1	>1	No	Yes	Y	I
Matsuno, 2021	Hea	Fea; Eff	NP	Yes	Yes	Y	No	3	O	O	No	No	I	Y	>1	>1	No	Yes	Y	I
Mesurado, 2019	Hea	Eff	NP	No	No	I	No	5	O	O	No	No	N	Y	>1	>1	No	No	N	I
Michelon, 2024	Hea	Fea; Eff	Pr	No	No	Y	No	3	O	O	No	No	I	I	>1	>1	No	Yes	I	I
Molek-Winiarska, 2024	Soc	Eff	NP	No	Yes	I	No	4	O	O	No	No	I	Y	>1	>1	No	NCS	N	I
Morgan, 2014	Clin	Fea; Eff	PP	Yes	Yes	I	No	6	O	O	No	No	I	Y	>1	>1	No	No	N	I

ID	Field	G_1	G_2	G_3	G_5	G_7	G_9	nBeA	G_4	BeA_1a	BeA_1b	BeA_1c	G_6	BeA_6	G_8	BeA_3a	BeA_3b	BeA_4	BeA_5	BeA_7
Motlova, 2021	Edu	Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
Newins, 2021	Soc	Fea; Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	Y	I
Nozawa, 2022	Edu	Fea; Eff	Pr	Yes	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
O'Garo, 2023	Clin	Eff	PP	No	Yes	I	No	2	O	O	No	No	I	Y	>1	>1	No	Yes	N	I
Oldershaw, 2024	Clin	Fea; Eff	NP	Yes	Yes	I	No	5	O	O	No	No	I	Y	>1	>1	No	Yes	Y	I
Olsson, 2024	Soc	Fea; Eff	Pr	Yes	Yes	Y	Yes	1	O	O	No	No	I	I	>1	>1	No	Yes	Y	I
Otsuka, 2015	Clin	Fea; Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
Painter, 2019	Clin	Eff	NP	No	No	Y	No	2	O	O	No	No	I	I	>1	>1	No	No	N	I
Pakarinen, 2018	Hea	Eff	Pr	Yes	No	Y	No	2	O	O	M	No	I	I	>1	>1	No	No	N	I
Passoni, 2018	Soc	Eff	PP	No	No	Y	No	3	O	O	No	No	I	I	>1	>1	No	No	N	I
Ramirez, 2021	Clin	Eff	NP	No	Yes	Y	No	3	O	O	No	No	I	I	>1	>1	No	No	N	I
Rodrigo, 2022	Clin	Eff	CP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
Saffaran, 2022	Hea	Fea; Eff	NP	No	No	Y	No	2	O	O	No	No	I	I	>1	>1	No	No	Y	I
Saulnier, 2024	Clin	Fea; Eff	Pr	No	No	Y	No	2	O	O	No	No	I	I	>1	>1	No	No	Y	I
Schneider, 2016	Soc	Fea; Eff	NP	No	No	I	Yes	1	O	O	No	No	I	I	>1	>1	No	No	Y	I
Sherwood, 2023	Hea	Fea; Eff	CP	No	No	Y	No	2	O	O	No	No	I	I	>1	>1	No	Yes	Y	Y
Tran, 2020	Edu	Fea; Eff	Pr	No	No	I	No	1	O	O	No	No	I	I	>1	>1	No	No	N	Y
Walker, 2014	Hea	Eff	NP	No	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	Y
Ward, 2020	Soc	Eff	NP	No	No	Y	No	2	O	O	No	No	I	I	>1	>1	No	No	N	I
Wasik, 2020	Edu	Fea; Eff	NP	No	No	Y	No	3	O	O	No	No	I	I	>1	>1	No	No	Y	I
Weigel, 2015	Edu	Eff	Pr	Yes	No	Y	No	1	O	O	No	No	I	I	>1	>1	No	No	N	I
Zajonz, 2024	Clin	Eff	NP	No	No	N	No	1	O	O	No	No	I	I	>1	0	No	No	N	I
Zhang, 2014	Clin	Eff	NP	No	No	Y	No	3	O	O	No	No	I	I	>1	0	No	Yes	N	I

257
Note 1. Table Columns. Field: Research field; G_1: Objectives; G_2: Previous publication of intervention program or protocol; G_3: Reporting quality assessment tool; G_5: Facilitators or barriers; G_7: Implementation
sequence; G_9: Costs; nBeA: Number of BeA; G_4: Number of categories of recipients; BeA_1a: Levels & categories of recipients: Individual level; BeA_1b: Levels & categories of recipients: Group/team level; BeA_1c:
Levels & categories of recipients: System level; G_6: Prerequisites of the whole intervention: Global; BeA_6: Prerequisites of the particular BeA: Global; G_8: Intervention outcomes; BeA_3a: Number of outcomes/effects;
BeA_3b: Moderation or mediation effect on the outcomes of other BeA; BeA_4: Adaptation/customization; BeA_5: Evaluation of the fidelity (including adherence); BeA_7 Implementation: Global

Note 2: Abbreviations: Clin: Clinical; Edu: Education; Hea:Health; Soc: Social; Eff: Effectiveness; Fea: Feasibility; Dev: Development only; CP: Intervention previously published (completely); NP: Without prior
publication or registration; PP: Intervention previously published (partially); Pr: Protocol registered; L: Multi-level: M: Multi-category; O: One-category; D: Description only, no prerequisite stated; I: Reported, but not
replicable (missing info); N: Not reported; Y: Reported and/or replicable; NCS: Not clearly stated

Results

Search results

Figure 2 shows the results of the search strategy, reported according to the PRISMA flowchart
(Page et al., 2021). Finally, 84 studies were selected, 4 were used for testing the extraction
procedure, and the remaining 80 were included in the review. The interrater agreement for
abstract and title selection was 100%, so no kappa coefficient needed to be calculated,
whereas the interrater agreement for full text selection was kappa k=0.78 (CI95%: 0.59-
0.96). After discussion to solve the discrepancies, the interrater agreement percentage for full
text selection was 100%.

Figure 2.
PRISMA flow diagram of the selection process.

Characteristics of the whole interventions

Table 2 and Figure 3 summarize the key characteristics of intervention programs in four
research fields (clinical, health, social, and education). The analysis included chi-square tests
to assess differences between fields. Below, we first describe the overall patterns observed for
each characteristic across all fields combined, followed by a comparison across the individual
fields.

Figure 3.
Characteristics of the whole interventions.

Across all fields, only 11.3% of studies focused exclusively on the development of the
intervention program. Most publications combined development with an additional objective,
most often the assessment of effectiveness (40%) or both feasibility and effectiveness (40%).
Feasibility evaluation was reported in just under half of the studies (48.8%), while the majority
(80%) assessed the effectiveness. Most studies (61.3%) did not previously publish the
intervention or register their protocol, and only 13.8% applied a reporting quality assessment
tool.

In terms of recipients, most interventions targeted a single category (87.5%), while a small
number addressed multiple levels (6.3%). Facilitators or barriers for the intervention were
reported in only 25% of cases. Finally, nearly all interventions (96.3%) included multiple
outcomes.

No statistically significant differences were found across research fields, though descriptive
trends emerged. Development-only objectives appeared in clinical and social research (20%
each) but were absent in education, while health studies rarely stopped at development (5%).
Combined evaluations of feasibility and effectiveness were most frequent in education (50%)
and least in social research (30%). Feasibility assessments dominated health (60%) and
education (55%) but were less common in social studies (35%). Effectiveness was almost
universally assessed in education (95%) and less so in clinical research (70%).
305

306   Table 2.

307   *Characteristics of the whole intervention by field.*

		All fields %	Clinical %	Health %	Social %	Education %	Between fields X² sig.
Objective/s of the study in addition to the development of the intervention program	Development only	11.3	20	5	20	0	0.438
	+Effectiveness	40	35	35	45	45	
	+Feasibility	8.7	10	15	5	5	
	+Feasibility+Effectiveness	40	35	45	30	50	
Evaluation of feasibility of the intervention	Yes	48.8	45	60	35	55	0.399
	No	51.2	55	40	65	45	
Evaluation of the effectiveness of the intervention	Yes	80	70	80	75	95	0.224
	No	20	30	20	25	5	
Previous publication of intervention program (completely o partially) or registration of the intervention protocol	Intervention previously published (completely)	7.5	10	5	10	5	0.998
	Intervention previously published (partially)	13.7	15	15	10	15	
	Protocol registered	17.5	15	15	20	20	
	Without prior publication or registration	61.3	60	65	60	60	
Reporting quality assessment tool applied	Yes	13.8	10	10	20	15	0.788
	No	86.2	90	90	80	85	
Number of categories of recipients of the whole intervention	One-category	87.5	95	90	90	75	0.368
	Multi-category	6.2	0	5	10	10	
	Multi-level	6.3	5	5	0	15	
Facilitators or barriers for the whole intervention	Yes	25	30	25	15	30	0.659
	No	75	70	75	85	70	
Intervention outcomes of the whole intervention	Single	3.8	5	5	0	5	0.792
	Multiple	96.2	95	95	100	95	

309 Reporting quality tools were rarely used, ranging from 10% of clinical and health to 20% of
310 social studies. Most interventions targeted a single recipient category, especially in clinical
311 (95%) and health (90%), whereas education showed greater diversity, with 15% of multi-level
312 interventions compared to 0-5% in the other fields. Finally, reporting of facilitators or barriers
313 was highest in education and clinical (30%) and lowest in social research (15%).

Characteristics of the intervention's recipients' behaviors or actions (BeA)

Table 3 and Figure 4 outline the characteristics related to the BeA, both at global level and for
each of the four specific research fields that were analyzed.

Figure 4.
Characteristics of the intervention recipients' behaviors or actions (BeA).

Across all fields, most interventions included more than one recipients' BeA. While 30%
addressed a single BeA and 21.3% two BeA, the largest proportion (37.5%) reported 3–4 BeA,
and a smaller proportion (11.3%) addressed 5–8 BeA. Most studies described recipients'
behaviors at a single level (86.3%), 97.6% considered the individual level, 15.1% the
group/team level, and only 1.3% the system level.

Facilitators or barriers for particular BeA were almost absent (1.3%). In terms of outcomes, just
over half of the interventions (55%) reported multiple outcomes or effects for BeA, while 45%
reported none. Interactive effects among BeA were infrequently explored: only 11.3% of studies
reported moderating or mediating effects of the BeA on the outcomes of other BeA, and 5%
were unclear.

Table 3.

Characteristics of the intervention's recipients' behaviors or actions (BeA) by field.

		All fields %	Clinical %	Health %	Social %	Education %	Between fields X² sig.
Number of BeA	One BeA	30	35	30	35	20	
	Two BeA	21.2	15	20	25	25	0.953
	3-4 BeA	37.5	35	35	35	45	
	5-8 BeA	11.3	15	15	5	10	
Number of levels of recipients of particular BeA	One level	86.2	95	80	85	85	0.572
	Two levels	13.8	5	20	15	15	
Levels & categories of recipients of particular BeA: Individual level	One-category	93.8	100	95	95	85	
	Multi-category	3.7	0	5	5	5	0.298
	No	2.5	0	0	0	10	
Levels & categories of recipients of particular BeA: Group/team level	One-category	13.7	5	15	15	20	
	Multi-category	1.3	0	5	0	0	0.534
	No	85	95	80	85	80	
Levels & categories of recipients of particular BeA: System level	One-category	1.3	0	0	0	5	0.386
	No	98.7	100	100	100	95	
Facilitators or barriers for particular BeA	Yes	1.2	0	0	0	5	0.386
	No	98.8	100	100	100	95	
Number of outcomes/effects of particular BeA	Multiple	55	60	55	55	50	0.939
	None	45	40	45	45	50	
Moderation or mediation effect of particular BeA on the outcomes of other BeA	Yes	11.3	20	5	5	15	
	Unclear	5	0	10	5	5	0.523
	No	83.7	80	85	90	80	

No statistically significant differences were observed between fields. Nevertheless, some
descriptive variations can be noted. Education showed a slightly higher proportion of
interventions with 3 or 4 BeA (45%), while clinical and social fields had somewhat higher rates
of single BeA (35% each). Clinical studies reported the highest proportion of single-level
interventions (95%), while health interventions were slightly more likely to include two
recipient levels (20%). At individual level, education showed slightly more diversity, with 10%
of studies not reporting this level, while at group/team level, education (20%) and social (15%)
included it slightly more often than clinical (5%). At system level, only education (5%) reported
it. Finally, clinical research led in testing moderation and/or mediation effects (20%), followed
by education (15%), with health and social both at 5%.

Quality reporting of implementation characteristics of the whole interventions

Table 4 and Figure 5 present the quality reporting of implementation of the intervention studies
analysed, categorized by reporting status: reported and/or replicable (RR), reported but not
replicable due to missing information (RNR), and not reported (NR).

Across fields, the reporting of implementation characteristics at the level of the whole
intervention showed a mixed picture, with notable weaknesses in several areas. Global
prerequisites of the intervention were seldom reported in sufficient detail: only 15% of studies
provided replicable information, while most either offered incomplete descriptions (77.5%) or
omitted them altogether (7.5%). This pattern was consistent across the different groups of
prerequisites. Although recipients' (57.5%) and context-related prerequisites (70%) were more
frequently addressed, information on providers (35%) and evaluators (35%) was less often
available in a replicable form. When prerequisites of the whole intervention and recipients'
behaviors were considered jointly, reporting was very limited, with only 13.8% adequately
covered. By contrast, the implementation sequence was more consistently described: nearly
three-quarters of studies (73.8%) included enough detail to allow replication. Nevertheless,
structured reporting formats that could enhance clarity and replicability were underutilized.
Only 16.3% of studies presented a diagram, and fewer than half (38.8%) provided tables or lists
outlining the intervention's structure. Costs were almost entirely neglected (3.8%).

No statistically significant differences were observed between fields in the percentage of
characteristics well reported (see Fig. 6), except for the reporting of tables or lists of
intervention structure ($p = 0.024$), which was highest in clinical (65%) and education (40%)
compared to health (20%). Recipients' prerequisites were most often reported in a replicable
manner in clinical (75%) and health (60%) compared to education (45%, and providers'
prerequisites were adequately reported more often in health (50%) than in education (30%).

Reporting of tables or lists describing the structure of the intervention program was more
frequent in clinical (65%) compared to other fields (20–40%).

Finally, it is worth noting that, regarding the section of the publications in which the previously
described general characteristics of the interventions were reported, these were not included in
the Method section of the studies in between 14.6% (administrator prerequisites) and 37%
(evaluator prerequisites) of the cases. Moreover, in 75.8% of the studies, the reporting of
contextual prerequisites (setting) was also located in sections other than the Method section.

Figure 5.

Quality reporting of implementation characteristics of the whole interventions.

Figure 6.

Percentage of implementation characteristics of the whole interventions reported and/or

replicable by field.

Table 4.

Quality reporting of implementation characteristics by field.

	All fields			Clinical field			Health field			Social field			Education field			Between fields
	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	X^2 sig.
Implementation characteristics of the whole intervention																
Global prerequisites of the whole intervention	15	77.5	7.5	10	85	5	15	75	10	15	75	10	20	75	5	0.958
Providers' prerequisites	35	11.3	53.7	40	5	55	50	0	50	20	30	50	30	10	60	0.055
Evaluators' prerequisites	35	12.5	52.5	20	15	65	45	0	55	35	25	40	40	10	50	0.208
Recipients' prerequisites	57.5	12.5	30	75	5	20	60	10	30	50	25	25	45	10	45	0.256
Context prerequisites	70	7.5	22.5	65	0	35	65	15	20	75	10	15	75	5	20	0.467
Global prerequisites of the whole intervention or for BeA	13.8	78.7	7.5	10	85	5	10	80	10	15	75	10	20	75	5	0.934
Implementation sequence	73.8	17.5	8.7	80	15	5	75	15	10	60	25	15	80	15	5	0.789
Report includes a diagram of the intervention program	16.2	0	83.8	20	0	80	25	0	75	10	0	90	10	0	90	0.479
Report includes tables/lists of the structure of the intervention program	38.8	0	61.2	65	0	35	20	0	80	30	0	70	40	0	60	0.024*
Costs	3.7	0	96.3	0	0	100	5	0	95	10	0	90	0	0	100	0.283
Implementation characteristics of recipients' behaviors or actions (BeA)																
Adaptation/Customization or not	35	15	50	40	5	55	40	20	40	15	20	65	45	15	40	0.326
Evaluation of the fidelity/adherence of the implementation	27.5	7.5	65	20	0	80	40	10	50	25	5	70	25	15	60	0.533
Global prerequisites for particular BeA	32.5	61.3	6.2	30	65	5	35	60	5	35	55	10	30	65	5	0.978
Providers' prerequisites	41.3	3.7	55	45	5	50	55	5	40	35	0	65	30	5	65	0.619
Evaluators' prerequisites	43.8	0	56.2	30	0	70	55	0	45	45	0	55	45	0	55	0.459
Recipients' prerequisites	61.2	2.5	36.3	75	5	20	60	5	35	60	0	40	50	0	50	0.466
Context prerequisites	82.5	5	12.5	85	0	15	85	5	10	75	15	10	85	0	15	0.361
Global characteristics of the implementation	13.8	86.2	0	15	85	0	15	85	0	0	100	0	25	75	0	0.146
Type of administration (e.g., remote, face-to-face)	73.7	21.3	5	80	15	5	70	30	0	60	25	15	85	15	0	0.198
Form of administration (e.g., individual, group)	80	16.2	3.8	90	5	5	80	15	5	65	30	5	85	15	0	0.446
Supervision/support (e.g., directed, autonomous, mixed)	72.5	22.5	5	80	15	5	75	25	0	65	25	10	70	25	5	0.807
Materials and support tools (e.g., multimedia, text, app)	47.5	36.3	16.2	60	20	20	50	45	5	40	35	25	40	45	15	0.405
Content and development of the sessions	36.3	55	8.7	65	25	10	25	70	5	10	80	10	45	45	10	0.011*
Periodicity (time unit; e.g., once, daily, weekly, monthly)	75	11.3	13.7	90	0	10	60	15	25	75	15	10	75	15	10	0.354
Frequency per time unit (e.g., once, a number of times)	70	13.8	16.2	75	10	15	60	20	20	70	15	15	75	10	15	0.948
Total frequency (e.g., specific number, tailored)	76.2	12.5	11.3	90	0	10	70	15	15	75	15	10	70	20	10	0.591
Duration of sessions, dose or intensity (specific, tailored)	56.3	20	23.7	55	10	35	55	25	20	50	20	30	65	25	10	0.540
Geographical, spatial, physical and/or digital place	88.7	10	1.3	90	10	0	95	5	0	70	25	5	100	0	0	0.083
Moment in time (e.g., specific, tailored, irrelevant)	95	0	5	95	0	5	100	0	0	90	0	10	95	0	5	0.551
Social, cultural, political and/or economic setting	91.2	5	3.8	90	10	0	95	5	0	85	5	10	95	0	5	0.444

Notes: RR: reported and/or replicable; RNR: reported but not replicable (missing information); NR: not reported; *: statistically significant (p<0.05)

Quality reporting of implementation characteristics of the intervention's recipients' behaviors or actions (BeA)

At the level of BeA, reporting quality of the implementation characteristics of the interventions also varied (Table 4 and Fig. 7). The reporting of BeA adaptation or customization—entailing whether the intervention was adaptable or customizable, and, if so, what adaptations or customizations were permitted—was done in a replicable manner in just over a third of studies (35%), with half failing to address this aspect at all. Similarly, fidelity and adherence were reported by 27.5% of studies with adequate detail and nearly two-thirds (65%) omitting this implementation feature.

Global prerequisites for recipients' behaviours were also weakly documented: while 32.5% reported them with replicable detail, most studies (61.3%) provided incomplete information. When broken down by type of prerequisite, reporting was uneven. Recipients' (61.3%) and context-related prerequisites (82.5%) were more frequently specified. By contrast, providers (41.3%) and evaluators (43.8%) were less consistently addressed.

Figure 7.
Quality reporting of implementation characteristics of the intervention's recipients' behaviors or actions (BeA).

In contrast, certain global characteristics of implementation were relatively comprehensively addressed. Most studies showed adequate reporting on key structural aspects, such as type (73.8%) and form of administration (80%), supervision or support (72.5%), and periodicity (75%). Frequency (70–76.3%), location (88.8%), and timing (95%) were also commonly well reported. Moreover, the social, cultural, political, and economic setting was frequently reported and/or replicable (91.3%). However, two key elements remained comparatively underreported: the content and development of sessions (36.3%) and the materials or support tools used (47.5%). The reporting of the dose or intensity of the sessions was replicable in just over half (56.3%) of the studies. Overall, the low percentage of "not reported" responses observed for most of the features analysed should be interpreted with caution, as while they indicated that

these implementation features were universally addressed, this was rarely done in a fully replicable manner in light of the observed percentage of "reported but not replicable (missing information)" results.

No statistically significant differences were found across fields (Fig. 8) except for content and development of sessions ($p = 0.011$), where the replicability of the reports was greater in the clinical field (65%) and lowest in health (25%) and social (10%) fields. Adaptation/customization was reported more frequently and accurately in clinical and health fields (40% each) than in social (15%). Reporting on implementation fidelity was most adequate in health (40%) and lowest in clinical (20%) field.

Figure 8.

Percentage of implementation characteristics of intervention's recipients' behaviors or actions (BeA) reported and/or replicable by field.

Finally, it should be noted that, concerning the section of the publications in which the characteristics previously described of the BeA were reported, the four groups of implementation prerequisites were not included in the Method section in between 70% (administrator prerequisites) and 86.4% (evaluator prerequisites) of the studies. The remaining implementation characteristics of the BeA were reported outside the Method section in less than 15% of the studies, except for the materials and support tools required for BeA implementation, their potential for adaptation/customization, and the procedures used to assess implementation fidelity, which were reported outside the Method section in 37.3%, 25%, and 17.9% of the studies, respectively.

Quality reporting of implementation characteristics of the interventions by number of intervention's recipients' behaviors or actions (BeA)

Table 5 and Figure 9 examine the quality of reporting for implementation characteristics in intervention studies, stratified by the number of BeA involved (1 BeA, 2 BeA, 3 BeA, 4 BeA,

and 5-8 BeA). The reporting of implementation characteristics for the whole intervention varied
substantially depending on the number of BeA considered. When only one BeA was addressed,
replicable reporting of global prerequisites was almost absent (4.2%), and only a minority of
studies included providers' (16.7%) or evaluators' (12.5%) prerequisites, and recipients'
prerequisites were reported in a replicable manner somewhat more frequently (25%), and
context prerequisites were well documented at a relatively high level (79.2%).

Figure 9.
Percentage of implementation characteristics of the whole interventions reported and/or
replicable by number of recipients' behaviors or actions (BeA).

As the number of BeA increased, the completeness of reporting improved. For example, in
interventions with 2 BeA, reporting of global prerequisites increased to 11.8%, providers' to
35.3%, and evaluators' to 35.3%. By the time interventions addressed 3–4 BeA, global
prerequisites were well described in 22.2% of studies, with evaluators' (44.4%) and especially
recipients' prerequisites (83.3%) reported more often. This trend culminated in interventions
with 4 BeA, where reporting of prerequisites was strongest: 25% for global prerequisites, 58.3%
for providers, 50% for evaluators, and 75% for recipients. Context prerequisites remained
relatively stable across groups (58–88%).

At the level of BeA (Fig. 10), the same pattern of improvement with increasing complexity was
observed. For interventions with only one BeA, global prerequisites were reported in a
replicable manner in 4.2% of cases, while providers', evaluators', and recipients' prerequisites
were accurately reported in 20.8%, 12.5%, and 25%, respectively. Again, context prerequisites
were relatively well covered even in the simplest interventions (75%).

Figure 10.
Percentage of implementation characteristics of the intervention's recipients' behaviors or
actions (BeA) reported and/or replicable by number of BeA.

As the number of BeA increased, reporting became progressively more complete. With 2 BeA,
global prerequisites well reported reached 29.4%, providers' 35.3%, evaluators' 41.2%, and
recipients' 47.1%, while context prerequisites climbed to 88.2%. For 3–4 BeA, adequate
reporting was higher: 38.9% for global prerequisites, 50% for evaluators, and 83.3% for
recipients. In interventions with 4 BeA, the reporting of prerequisites reached its peak, with
58.3% replicable for global prerequisites, 75% for providers and evaluators, 91.7% for
recipients, and 75% for context. For the most complex interventions (5–8 BeA), reporting levels
were also high, with recipients' prerequisites accurately reported universally (100%) and
evaluators' prerequisites in 77.8%. Global implementation characteristics followed a slightly
different pattern, generally showing low replicability (11.1–17.6%)

Table 5.

477 *Quality reporting of implementation characteristics by number of BeA.*

	1 BeA			2 BeA			3 BeA			4 BeA			5-8 BeA			Between fields
	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	RR %	RNR %	NR %	X² sig.
Implementation characteristics of the whole intervention																
Global prerequisites of the whole intervention	4.2	83.3	12.5	11.8	76.4	11.8	22.2	77.8	0	25	75	0	22.2	66.7	11.1	0.473
Providers' prerequisites	16.7	20.8	62.5	35.3	0	64.7	27.8	11.1	61.1	58.3	16.7	25	66.7	0	33.3	0.047*
Evaluators' prerequisites	12.5	8.3	79.2	35.3	17.6	47.1	44.4	5.6	50	50	33.3	16.7	55.6	0	44.4	0.016*
Recipients' prerequisites	25	20.8	54.2	47.1	11.8	41.1	83.3	5.6	11.1	75	16.7	8.3	88.9	0	11.1	0.004*
Context's prerequisites	79.2	0	20.8	70.6	5.9	23.5	77.8	11.1	11.1	58.3	16.7	25	44.4	11.1	44.4	0.412
Global prerequisites of the whole intervention or for BeA	4.2	83.3	12.5	11.8	76.4	11.8	22.2	77.8	0	25	75	0	11.1	77.8	11.1	0.488
Implementation characteristics of recipients' behaviors or actions (BeA)																
Global prerequisites for particular BeA	4.2	83.3	12.5	29.4	58.8	11.8	38.9	61.1	0	58.3	41.7	0	66.7	33.3	0	0.009
Providers' prerequisites	20.8	4.2	75	35.3	5.9	58.8	38.9	0	61.1	75	0	25	66.7	11.1	22.2	0.049
Evaluators' prerequisites	12.5	0	87.5	41.2	0	58.8	50	0	50	75	0	25	77.8	0	22.2	<.001*
Recipients' prerequisites	25	4.2	70.8	47.1	0	52.9	83.3	5.6	11.1	91.7	0	8.3	100	0	0	<.001*
Context prerequisites	75	4.2	20.8	88.2	0	11.8	88.9	5.6	5.5	75	8.3	16.7	88.9	11.1	0	0.691
Global characteristics of the implementation	12.5	87.5	0	17.6	82.4	0	11.1	88.9	0	16.7	83.3	0	11.1	88.9	0	0.974
Type of administration (e.g., remote, face-to-face)	79.2	8.3	12.5	88.2	11.8	0	55.6	38.8	5.6	66.7	33.3	0	77.8	22.2	0	0.144
Form of administration (e.g., individual, group)	83.3	12.5	4.2	88.2	5.9	5.9	72.2	22.2	5.6	75	25	0	77.8	22.2	0	0.856
Supervision/support (e.g., directed, autonomous, mixed)	79.1	16.7	4.2	88.2	5.9	5.9	55.6	44.4	0	75	25	0	55.6	22.2	22.2	0.058
Materials and support tools (e.g., multimedia, text, app)	50	29.2	20.8	47.1	23.5	29.4	38.9	50	11.1	66.7	25	8.3	33.3	66.7	0	0.219
Content and development of the sessions	45.8	37.5	16.7	41.1	47.1	11.8	27.8	72.2	0	33.3	58.4	8.3	22.2	77.8	0	0.323
Periodicity (time unit; e.g., once, daily, weekly, monthly)	66.7	4.1	29.2	70.6	17.6	11.8	83.3	5.6	11.1	91.7	8.3	0	66.7	33.3	0	0.073
Frequency per time unit (e.g., once, a number of times)	70.8	4.2	25	64.7	17.6	17.7	72.2	16.7	11.1	83.3	8.3	8.4	55.6	33.3	11.1	0.497
Total frequency (e.g., specific number, tailored)	70.8	4.2	25	70.6	17.6	11.8	88.9	11.1	0	83.3	8.3	8.4	66.7	33.3	0	0.115
Duration of sessions, dose or intensity (specific, tailored)	62.5	4.2	33.3	58.8	35.3	5.9	50	27.8	22.2	58.4	8.3	33.3	44.5	33.3	22.2	0.180
Geographical, spatial, physical and/or digital place	91.7	4.1	4.2	82.4	17.6	0	83.3	16.7	0	91.7	8.3	0	100	0	0	0.627
Moment in time (e.g., specific, tailored, irrelevant)	91.7	0	8.3	100	0	0	100	0	0	83.3	0	16.7	100	0	0	0.177
Social, cultural, political and/or economic setting	91.7	4.1	4.2	94.1	5.9	0	94.4	5.6	0	75	8.3	16.7	100	0	0	0.410

Discussion

This scoping review mapped how psychological intervention reports describe implementation at both the whole-intervention level and at the level of recipients' BeA across clinical, public health, social/organizational and educational fields. Only a minority of studies reported development-only objectives, while most combined development with evaluations of effectiveness and, to a lesser extent, feasibility. Few studies used a formal reporting quality tool or reported facilitators/barriers. Interventions typically targeted an individual-level single-category, with group and system level targeting being uncommon, and outcomes were generally well specified. Reporting of the implementation sequence was generally adequate, yet structured representations such as tables, lists or diagrams were underused, and costs were rarely reported. At the BeA level, generic structural features (such as format, supervision, periodicity, setting and timing) were usually described in a replicable way, whereas content and materials, adaptation or customization, and especially fidelity or adherence were often incompletely reported. Relevant aspects of complexity such as multi-level recipients, moderation or mediation effects were seldom analysed, although reporting quality tended to improve as the number of BeA increased.

Our findings show that, despite decades of general reporting guidelines (CONSORT, STROBE, PRISMA, SPIRIT, TREND) and intervention-specific reporting tools (TIDieR, CReDECI2, GUIDED, iCAT_SR, RIPIf), their adoption remains limited and key implementation details continue to be omitted. This is consistent with previous works showing that endorsement of reporting tools and guidelines alone does not guarantee adherence and complete reporting (Turner et al., 2012; Bastuji-Garin et al., 2013; Clayson et al., 2019). Similarly, key recommendations from the MRC guidance on developing and evaluating (complex) interventions have been largely overlooked, as most studies report both intervention development and testing within a single paper, which tends to compromise the replicability of the intervention.

Our results also resonate strongly with TIDieR's emphasis on specifying *what* was delivered (materials/content) and *how* (dose, mode, procedures). The comparatively weak reporting for content/materials (and moderate reporting for dose/intensity) indicates that, even when structural aspects are reported, some core elements of interventions are often left implicit.

Contemporary frameworks and tools such as MRC and iCAT_SR ask authors to articulate how components are expected to work (mechanisms), for whom, in what contexts, and with what interdependencies. The rarity of multi-level targeting and explicit tests of mediation/moderation among BeA suggests that many reports still treat interventions largely as packages rather than as systems of interacting parts. Explicitly modelling and reporting inter-component relations, and using logic models to link BeA to outcomes, would align practice with these frameworks and support richer cumulative science (Moore et al., 2015).

Two additional gaps stand out. First, implementation fidelity was rarely reported in a replicable way. The conceptual framework of Carroll et al. (2007) describes fidelity as a multidimensional construct (adherence, dose, quality of delivery, participant responsiveness, differentiation) that should be planned, measured, and reported systematically; the pattern we observed suggests these dimensions remain underspecified in psychological intervention reports. Moreover, meta-research consistently shows that implementation quality is tied to effect sizes, reinforcing the

importance of robust fidelity reporting (Durlak & DuPre, 2008). Second, costs were almost never reported, yet budget impact and cost-effectiveness information are essential to scaleup and policy translation. The CHEERS statement (Husereau et al., 2022) provides explicit, modernized guidance on transparent reporting of economic evaluations and is broadly applicable across intervention types; the near absence of cost reporting in our sample signals an actionable opportunity for improvement in future reports. From a practice standpoint, insufficient reporting of fidelity, adaptation policies, and costs undermines implementation planning and weakens the claim that a delivered service is truly evidence based. The taxonomy of Proctor et al. (2011) clarifies that feasibility, fidelity, penetration, sustainability, and cost are implementation outcomes in their own right; adequate reporting of these domains is therefore necessary both for replication and for implementation science to cumulate knowledge.

Reporting quality is also greatly enhanced when the structure and contents of a scientific report follow widely accepted standards (e.g., APA). In this regard, some inconsistencies have also been found in this review. Critical prerequisites and contextual details are frequently reported outside the Methods section, impeding discoverability and machine readability for evidence synthesis. Consolidating these items in Methods (with stable supplementary files hosting manuals, checklists, and materials) would align with SPIRIT/CONSORT transparency goals and facilitate automated screening by meta-research tools.

However, two encouraging findings emerge from our study. First, the implementation sequence was often reported with sufficient detail, and in more complex interventions, the reporting of prerequisites (especially for recipients and evaluators) improved. Second, generic structural elements (e.g., type/form of administration, supervision, settings) and timing-related characteristics (e.g., periodicity, frequency, duration) were frequently replicable. These strengths provide a solid foundation to build upon.

Limitations and strengths

This scoping review aimed to map reporting practices rather than quantify causal relationships or effectiveness. The review timeframe encompasses all major recent reporting frameworks, guidelines, and tools. Although selection was randomized within fields, the sample size per field limits statistical power to detect small differences and may not fully capture sub-disciplinary variations. Judgments of "replicable" versus "reported but not replicable" required coder interpretation; we mitigated this through piloting, independent double extraction, and consensus procedures, achieving a substantial interrater agreement at full-text screening. Finally, the theoretically guided and empirically grounded data extraction instrument developed for this study represents a key output. It provides powerful insights into the conceptualization of interventions' active ingredients (BeA), and a comprehensive list of appraisable properties of both these BeA and the whole intervention.

Conclusions and future directions

Reporting of psychological interventions shows encouraging coverage of structural "when/where/how" elements and implementation sequences, but persistent weaknesses in what (content/materials), how well (fidelity), whether and how it changes (adaptation/customization), and how much it costs (economic data). These deficiencies blunt replication, weaken cumulative syntheses, and impede implementation at scale, precisely the problems the reporting guideline ecosystem was designed to solve. Increasing the promotion of

intervention reporting tools and guidelines, such as TIDieR, CONSORT-SPI, RIPI-f, GUIDED,
and CReDECI-2, in psychological intervention reports, alongside intervention development and
evaluation frameworks and tools (e.g., MRC and iCAT_SR), and economic reporting standards
(e.g., CHEERS 2022), would help address the most consequential gaps identified in our scoping
review. Additionally, using content taxonomies, such as the Behavior Change Technique
Taxonomy (BCTTv1; (Michie et al., 2013)), can sharpen the granularity and comparability of
content reporting via a more precise specification of active intervention components and
thereby strengthen replication and synthesis.

Our results suggest that future research should focus primarily on:

a) Editorial interventions: evaluate editorial interventions, such as mandatory checklists,
structured tables or diagrams, and automated report audits, which have demonstrated
improvements in research integrity in other domains and adapting them to the level of
granularity required for BeA.

b) Level of BeA: jointly develop a minimum set of BeA characteristics, linking each BeA to
prerequisites, materials or content, procedures, dosage, fidelity measures, and outcomes,
including hypothesized mediation and moderation effects between BeA.

c) Complexity: increase the routine use of logic models and process evaluations to assess the
performance of components across different contexts, including multilevel recipients and
intercomponent pathways.

d) Fidelity measurement and reporting: prospectively adopt multicomponent frameworks and
evaluate whether improved fidelity reporting enhances effect estimates and reduces
heterogeneity in meta-analyses.

e) Economic transparency: integrate CHEERS 2022 elements, such as resource use, costs, and
analytical perspective, into intervention reports, even when a full economic evaluation is not
conducted, to better inform scaling-up decisions.

Beyond the scope of our findings, future research should explore the potential role of artificial
intelligence (AI) in assessing reporting quality in intervention studies. Domain-specific
language models such as BioBERT and PubMedBERT as well as recent reporting-focused work
(e.g., Chen et al., 2025) suggest promising avenues for an AI-assisted appraisal of reporting
quality.

Data availability statement

596 The data supporting the findings of this study are openly available in the Open Science
597 Framework repository at https://doi.org/10.34810/data2792.

Funding

599 This work was supported by Grant PID2022-141403NB-I00 funded by the Spanish
600 Government (MCIN/ AEI/10.13039/501100011033/FEDER, UE), and 2021SGR-00806,
601 funded by the Government of Catalonia (Spain). The authors declare no conflict of interest, and
602 the funders had no role in the study design, data collection, analysis, decision to publish, or
603 preparation of the manuscript. No additional external funding was received for this study.

Competing interests

The authors have declared that no competing interests exist.

Acknowledgements

An AI language model (Perplexity AI) was used to assist in correcting typographical and grammatical errors and to improve the clarity and fluency of English writing. The authors reviewed and approved all content.

References

Adnan NBB, Dafny HA, Baldwin C, Beccaria G, Chamberlain D. 2023. Is this the solution to wellbeing and burnout management for the critical care workforce? A parallel, interventional, feasibility and realist informed pilot randomized control trial protocol. *PLOS ONE* 18:e0285038. DOI: 10.1371/journal.pone.0285038.

Aharonovich E, Hasin DS, Nunes EV, Stohl M, Cannizzaro D, Sarvet A, Bolla K, Carroll KM, Genece KG. 2018. Modified cognitive behavioral therapy (M-CBT) for cocaine dependence: Development of treatment for cognitively impaired users and results from a Stage 1 trial. *Psychology of Addictive Behaviors* 32:800–811. DOI: 10.1037/adb0000398.

Ahlqvist-Björkroth S, Boukydis Z, Axelin AM, Lehtonen L. 2017. Close Collaboration with Parents^{TM} intervention to improve parents' psychological well-being and child development: Description of the intervention and study protocol. *Behavioural Brain Research* 325:303–310. DOI: 10.1016/j.bbr.2016.10.020.

Akkermans J, Brenninkmeijer V, Schaufeli WB, Blonk RWB. 2015. It's All About CareerSKILLS: Effectiveness of a Career Development Intervention for Young Employees. *Human Resource Management* 54:533–551. DOI: 10.1002/hrm.21633.

Almaqrami BS, Hua F, Liu Y, He H. 2020. Research waste-related editorial policies of leading dental journals: Situation 2018. *Oral Diseases* 26:696–706. DOI: 10.1111/odi.13257.

Andruschko J, Okely AD, Pearson P. 2018. A School-Based Physical Activity and Motor Development Program for Low-Fit Adolescent Females: The Sport4Fun Pilot Randomized Controlled Trial. *Journal of Motor Learning and Development* 6:345–356. DOI: 10.1123/jmld.2017-0013.

Appiah R, Wilson Fadiji A, Wissing MP, Schutte L. 2022. The *Inspired Life Program*: Development of a multicomponent positive psychology intervention for rural adults in Ghana. *Journal of Community Psychology* 50:302–328. DOI: 10.1002/jcop.22566.

Bastuji-Garin S, Sbidian E, Gaudy-Marqueste C, Ferrat E, Roujeau J-C, Richard M-A, Canoui-Poitrine F, on behalf of the European Dermatology Network (EDEN). 2013. Impact of STROBE Statement Publication on Quality of Observational Study Reporting: Interrupted Time Series versus Before-After Analysis. *PLoS ONE* 8:e64733. DOI: 10.1371/journal.pone.0064733.

Begg C. 1996. Improving the Quality of Reporting of Randomized Controlled Trials: The CONSORT Statement. *JAMA* 276:637–639. DOI: 10.1001/jama.1996.03540080059030.

Berman MI, Morton SN, Hegel MT. 2016. Health at Every Size and Acceptance and Commitment Therapy for Obese, Depressed Women: Treatment Development and Clinical Application. *Clinical Social Work Journal* 44:265–278. DOI: 10.1007/s10615-015-0565-y.

Blanco D, Schroter S, Aldcroft A, Moher D, Boutron I, Kirkham JJ, Cobo E. 2020. Effect of an
editorial intervention to improve the completeness of reporting of randomised trials: a
randomised controlled trial. *BMJ Open* 10:e036799. DOI: 10.1136/bmjopen-2020-036799.

Bork P, Bennett S. 2020. Video self-modeling, fading, and reinforcement: An effective
treatment package for children with selective mutism. *Clinical Child Psychology and*
Psychiatry 25:446–455. DOI: 10.1177/1359104519855110.

Boudreaux ED, Haskins B, Harralson T, Bernstein E. 2015. The remote brief intervention and
referral to treatment model: Development, functionality, acceptability, and feasibility. *Drug*
and Alcohol Dependence 155:236–242. DOI: 10.1016/j.drugalcdep.2015.07.014.

Caporale-Berkowitz NA, Boyer BP, Lyddy CJ, Good DJ, Rochlen AB, Parent MC. 2021. Search
inside yourself: investigating the effects of a widely adopted mindfulness-at-work
development program. *International Journal of Workplace Health Management* 14:593–
604. DOI: 10.1108/IJWHM-08-2020-0139.

Capps RE, Evans SW, Owens JS, Allan DM. 2024. A Peer-Supported School Engagement
Intervention for Youth with Attention Problems: Development and Implementation. *School*
Mental Health 16:649–666. DOI: 10.1007/s12310-024-09697-z.

Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. 2007. A conceptual framework for
implementation fidelity. *Implementation Science* 2:40. DOI: 10.1186/1748-5908-2-40.

Chan A-W, Boutron I, Hopewell S, Moher D, Schulz KF, Collins GS, Tunn R, Aggarwal R,
Berkwits M, Berlin JA, Bhandari N, Butcher NJ, Campbell MK, Chidebe RCW, Elbourne
DR, Farmer AJ, Fergusson DA, Golub RM, Goodman SN, Hoffmann TC, Ioannidis JPA,
Kahan BC, Knowles RL, Lamb SE, Lewis S, Loder E, Offringa M, Ravaud P, Richards DP,
Rockhold FW, Schriger DL, Siegfried NL, Staniszewska S, Taylor RS, Thabane L, Torgerson
DJ, Vohra S, White IR, Hróbjartsson A. 2025. SPIRIT 2025 statement: updated guideline for
protocols of randomised trials. *BMJ* 389:e081477. DOI: 10.1136/bmj-2024-081477.

Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, Hróbjartsson
A, Mann H, Dickersin K, Berlin JA, Doré CJ, Parulekar WR, Summerskill WSM, Groves T,
Schulz KF, Sox HC, Rockhold FW, Rennie D, Moher D. 2013. SPIRIT 2013 Statement:
Defining Standard Protocol Items for Clinical Trials. *Annals of Internal Medicine* 158:200–
207. DOI: 10.7326/0003-4819-158-3-201302050-00583.

Chen D, Li P, Khoshkish E, Lee S, Ning T, Tahir U, Wong HC, Lee MS, Raman S. 2025.
AutoReporter: Development of an artificial intelligence tool for automated assessment of
research reporting guideline adherence. DOI: 10.1101/2025.04.18.25326076.

Christensen DR, Landes RD, Jackson L, Marsch LA, Mancino MJ, Chopra MP, Bickel WK.
2014. Adding an Internet-delivered treatment to an efficacious treatment package for opioid
dependence. *Journal of Consulting and Clinical Psychology* 82:964–972. DOI:
10.1037/a0037496.

Chu BC, Rizvi SL, Zendegui EA, Bonavitacola L. 2015. Dialectical Behavior Therapy for
School Refusal: Treatment Development and Incorporation of Web-Based Coaching.
Cognitive and Behavioral Practice 22:317–330. DOI: 10.1016/j.cbpra.2014.08.002.

Clayson PE, Carbine KA, Baldwin SA, Larson MJ. 2019. Methodological reporting behavior,
sample sizes, and statistical power in studies of event-related potentials: Barriers to
reproducibility and replicability. *Psychophysiology* 56:1–17. DOI: 10.1111/psyp.13437.

Cohrs C, Bormann KC, Diebig M, Millhoff C, Pachocki K, Rowold J. 2019. Transformational
leadership and communication: Evaluation of a two-day leadership development program.

Leadership & Organization Development Journal 41:101–117. DOI: 10.1108/LODJ-02-2019-0097.

Coker AL, Bush HM, Brancato CJ, Clear ER, Recktenwald EA. 2019. Bystander Program Effectiveness to Reduce Violence Acceptance: RCT in High Schools. *Journal of Family Violence* 34:153–164. DOI: 10.1007/s10896-018-9961-8.

Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. 2008. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 337:a1655. DOI: 10.1136/bmj.a1655.

Des Jarlais DC, Lyles C, Crepaz N. 2004. Improving the Reporting Quality of Nonrandomized Evaluations of Behavioral and Public Health Interventions: The TREND Statement. *American Journal of Public Health* 94:361–366. DOI: 10.2105/AJPH.94.3.361.

Drake PJ, Potter SC, Clark LF, Desai M, LaSalle-Castro F, Masters T, Ayyaluru S. 2024. Evaluating a Future-Oriented Positive Youth Development Intervention to Reduce Sexual Risk Among Highly Mobile Youth: Results and Challenges. *Journal of Adolescent Health* 75:173–179. DOI: 10.1016/j.jadohealth.2024.03.003.

Duncan E, O'Cathain A, Rousseau N, Croot L, Sworn K, Turner KM, Yardley L, Hoddinott P. 2020. Guidance for reporting intervention development studies in health research (GUIDED): an evidence-based consensus study. *BMJ Open* 10:e033516. DOI: 10.1136/bmjopen-2019-033516.

Durlak JA, DuPre EP. 2008. Implementation Matters: A Review of Research on the Influence of Implementation on Program Outcomes and the Factors Affecting Implementation. *American Journal of Community Psychology* 41:327–350. DOI: 10.1007/s10464-008-9165-0.

Eklund K, Kilgus SP, Izumi J, DeMarchena SL, McCollom EM. 2021. The Resilience Education Program: Examining the efficacy of a Tier 2 internalizing intervention. *Psychology in the Schools* 58:2114–2129. DOI: 10.1002/pits.22580.

Elderson-Van Duin C, Renkema PJ, Tabbers HK. 2023. Promoting self-directed learning skills in a first-year problem-based university program with a self-development intervention. *Reflective Practice* 24:749–765. DOI: 10.1080/14623943.2023.2264195.

Emerson L-M, Rowse G, Sills J. 2017. Developing a Mindfulness-Based Program for Infant Schools: Feasibility, Acceptability, and Initial Effects. *Journal of Research in Childhood Education* 31:465–477. DOI: 10.1080/02568543.2017.1343211.

Erturk B. 2023. Program development for leaders in the post-truth era: Arts-based creative leadership communication program. *Evaluation and Program Planning* 98:102295. DOI: 10.1016/j.evalprogplan.2023.102295.

Facchini S, Martin V, Downing G. 2016. Pediatricians, Well-Baby Visits, and Video Intervention Therapy: Feasibility of a Video-Feedback Infant Mental Health Support Intervention in a Pediatric Primary Health Care Setting. *Frontiers in Psychology* 7. DOI: 10.3389/fpsyg.2016.00179.

Faretta E, Borsato T. 2016. EMDR Therapy Protocol for Oncological Patients. *Journal of EMDR Practice and Research* 10:162–175. DOI: 10.1891/1933-3196.10.3.162.

Farrand P, Woodford J, Small F, Mullan E. 2017. Behavioural activation self-help to improve depression in people living with dementia: The PROMOTE treatment protocol. *New Zealand Journal of Psychology* 46:51–62.

Ferolino AJP, Abrea SA. 2024. Pilot Study of a Positive Psychology Intervention to Reduce
Work-Related Stress Among Service Providers in a Drug Abuse Treatment Clinic.
International Perspectives in Psychology 13:202–211. DOI: 10.1027/2157-3891/a000107.

Fletcher S, Mitchell C, Moukhtarian T, Kershaw C, Russell S, Daly G, Toro CT, Meyer C. 2024.
REST 2.0: Protocol for a randomised controlled trial of a preventative psychological skills
training intervention for employee wellbeing and productivity. *Mental Health & Prevention*
34:200333. DOI: 10.1016/j.mhp.2024.200333.

Fung A. 2018. Reducing reactive aggression in schoolchildren through child, parent, and
conjoint parent-child group interventions: An efficacy study of longitudinal outcomes.
FAMILY PROCESS 57:594–612. DOI: 10.1111/famp.12323.

Georgiou K, Nikolaou I, Turban DB. 2021. The Impact of a Training Intervention Developing
Psychological Capital on Job Search Success. *Journal of Career Development* 48:369–384.
DOI: 10.1177/0894845319852425.

Glasziou P, Altman DG, Bossuyt P, Boutron I, Clarke M, Julious S, Michie S, Moher D, Wager
E. 2014. Reducing waste from incomplete or unusable reports of biomedical research. *The
Lancet* 383:267–276. DOI: 10.1016/S0140-6736(13)62228-X.

Glasziou P, Chalmers I. 2018. Research waste is still a scandal—an essay by Paul Glasziou and
Iain Chalmers. *BMJ*:k4645. DOI: 10.1136/bmj.k4645.

Goldenthal HJ, Gouze K, Raviv T, Tragoudas G, Holley C, Cicchetti C. 2024. The Ready to
Learn Through Relationships (RLR) Program: Development, Feasibility, and Acceptability.
Early Childhood Education Journal 52:587–601. DOI: 10.1007/s10643-023-01472-0.

Guise J-M, Butler ME, Chang C, Viswanathan M, Pigott T, Tugwell P. 2017a. AHRQ series on
complex intervention systematic reviews—paper 6: PRISMA-CI extension statement and
checklist. *Journal of Clinical Epidemiology* 90:43–50. DOI: 10.1016/j.jclinepi.2017.06.016.

Guise J-M, Chang C, Butler M, Viswanathan M, Tugwell P. 2017b. AHRQ series on complex
intervention systematic reviews—paper 1: an introduction to a series of articles that provide
guidance and tools for reviews of complex interventions. *Journal of Clinical Epidemiology*
90:6–10. DOI: 10.1016/j.jclinepi.2017.06.011.

Hawe P, Shiell A, Riley T. 2009. Theorising Interventions as Events in Systems. *American
Journal of Community Psychology* 43:267–276. DOI: 10.1007/s10464-009-9229-9.

Hettiarachchi S. 2022. The effectiveness of the adapted Box Clever language intervention
programme in the development of vocabulary and narrative skills of deaf and hard of hearing
children. *Deafness & Education International* 24:65–91. DOI:
10.1080/14643154.2020.1721158.

Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, Altman DG, Barbour V,
Macdonald H, Johnston M, Lamb SE, Dixon-Woods M, McCulloch P, Wyatt JC, Chan A-W,
Michie S. 2014. Better reporting of interventions: template for intervention description and
replication (TIDieR) checklist and guide. *BMJ* 348:1–12. DOI: 10.1136/bmj.g1687.

Hopewell S, Boutron I, Chan A-W, Collins GS, de Beyer JA, Hróbjartsson A, Nejstgaard CH,
Østengaard L, Schulz KF, Tunn R, Moher D. 2022. An update to SPIRIT and CONSORT
reporting guidelines to enhance transparency in randomized trials. *Nature Medicine*
28:1740–1743. DOI: 10.1038/s41591-022-01989-8.

Hopewell S, Chan A-W, Collins GS, Hróbjartsson A, Moher D, Schulz KF, Tunn R, Aggarwal
R, Berkwits M, Berlin JA, Bhandari N, Butcher NJ, Campbell MK, Chidebe RCW, Elbourne
D, Farmer A, Fergusson DA, Golub RM, Goodman SN, Hoffmann TC, Ioannidis JPA, Kahan

BC, Knowles RL, Lamb SE, Lewis S, Loder E, Offringa M, Ravaud P, Richards DP,
Rockhold FW, Schriger DL, Siegfried NL, Staniszewska S, Taylor RS, Thabane L, Torgerson
D, Vohra S, White IR, Boutron I. 2025. CONSORT 2025 Statement: Updated Guideline for
Reporting Randomized Trials. *JAMA* 333:1998. DOI: 10.1001/jama.2025.4347.

Huang C-HS, Crowther M, Allen RS, DeCoster J, Kim G, Azuero C, Ang X, Kvale E. 2016. A
Pilot Feasibility Intervention to Increase Advance Care Planning among African Americans
in the Deep South. *Journal of Palliative Medicine* 19:164–173. DOI:
10.1089/jpm.2015.0334.

Husereau D, Drummond M, Augustovski F, De Bekker-Grob E, Briggs AH, Carswell C,
Caulley L, Chaiyakunapruk N, Greenberg D, Loder E, Mauskopf J, Mullins CD, Petrou S,
Pwu R-F, Staniszewska S. 2022. Consolidated Health Economic Evaluation Reporting
Standards 2022 (CHEERS 2022) statement: updated reporting guidance for health economic
evaluations. *BMJ* 376:e067975. DOI: 10.1136/bmj-2021-067975.

Hussong AM, Samuelson PL, Richards AN, Mullane MJ. 2024. The good gifts program:
evaluation of a positive youth development intervention for opportunity youth. *The Journal*
of Positive Psychology 19:817–833. DOI: 10.1080/17439760.2023.2209537.

Ioannidis JPA, Greenland S, Hlatky MA, Khoury MJ, Macleod MR, Moher D, Schulz KF,
Tibshirani R. 2014. Increasing value and reducing waste in research design, conduct, and
analysis. *The Lancet* 383:166–175. DOI: 10.1016/S0140-6736(13)62227-8.

James S, Seidel FA, Kilian J, Trostmann J. 2020. Labor Market Integration of Young Adult
Refugees in Germany: Triangulating Perspectives Toward Program Development. *Research*
on Social Work Practice 30:553–563. DOI: 10.1177/1049731519897301.

Janka C, Stamm T, Heinze G, Dorner TE. 2022. A Training Programme for Developing Social
and Personal Resources and Its Effects on the Perceived Stress Level in Adults in Daily
Life—Study Protocol for a Prospective Cohort Study. *International Journal of*
Environmental Research and Public Health 20:523. DOI: 10.3390/ijerph20010523.

Jarero I, Artigas L, Uribe S, García LE. 2016. The EMDR Integrative Group Treatment Protocol
for Patients With Cancer. *Journal of EMDR Practice and Research* 10:199–207. DOI:
10.1891/1933-3196.10.3.199.

Jiwani Z. 2024. Assessing a Community-Based Maternal Self-Efficacy Intervention in Rural
India using a Quasi-Experimental Design. *The Counseling Psychologist* 52:166–192. DOI:
10.1177/00110000231215842.

Johnsen DB, Lomholt JJ, Heyne D, Jensen MB, Jeppesen P, Silverman WK, Thastum M. 2024.
The Effectiveness of Modular Transdiagnostic Cognitive Behavioral Therapy Versus
Treatment as Usual for Youths Displaying School Attendance Problems: A Randomized
Controlled Trial. *Research on Child and Adolescent Psychopathology* 52:1397–1412. DOI:
10.1007/s10802-024-01196-8.

Kahler CW, Spillane NS, Day A, Clerkin EM, Parks A, Leventhal AM, Brown RA. 2014.
Positive psychotherapy for smoking cessation: Treatment development, feasibility, and
preliminary results. *The Journal of Positive Psychology* 9:19–29. DOI:
10.1080/17439760.2013.826716.

Kelly EL, Siebach KF, DeHorn G, Lovejoy M. 2024. The Fulfillment Center Intervention
Study: Protocol for a group-randomized control trial of a participatory workplace
intervention. *PLOS ONE* 19:e0305334. DOI: 10.1371/journal.pone.0305334.

Kemer D, İşler Dalgıç A. 2022. Effectiveness of Sexual Abuse Prevention Training Program
Developed by Creative Drama for Preschoolers: An Experimental Study. *Journal of Child*
Sexual Abuse 31:9–32. DOI: 10.1080/10538712.2021.1994504.

Kjellström S, Areskoug Josefsson K, Fabisch A, Forsberg C, Schneider T, Avby G. 2022.
Fostering exploration and exploitation behavior in management teams to enhance
organizational performance: the LearnOvation leadership development program. *Leadership*
& Organization Development Journal 43:482–500. DOI: 10.1108/LODJ-04-2021-0162.

Knaevelsrud C, Böttche M, Pietrzak RH, Freyberger HJ, Kuwert P. 2017. Efficacy and
Feasibility of a Therapist-Guided Internet-Based Intervention for Older Persons with
Childhood Traumatization: A Randomized Controlled Trial. *The American Journal of*
Geriatric Psychiatry 25:878–888. DOI: 10.1016/j.jagp.2017.02.024.

Kokubo A, Yoshikawa K, Wu C-H. 2023. Facilitating transition from maternity leave to work
for working mothers: A self-efficacy intervention study. *Cambridge Prisms: Global Mental*
Health 10:e18. DOI: 10.1017/gmh.2023.6.

Komischke-Konnerup KB, O'Connor M, Hoijtink H, Boelen PA. 2023. Cognitive-Behavioral
Therapy for Complicated Grief Reactions: Treatment Protocol and Preliminary Findings
From a Naturalistic Setting. *Cognitive and Behavioral Practice*:S1077722923000949. DOI:
10.1016/j.cbpra.2023.11.001.

Kopelman-Rubin D, Omer H, Dar R. 2019. Brief therapy for excessive worry: Treatment
model, feasibility, and acceptability of a new treatment. *Journal of Psychotherapy*
Integration 29:291–306. DOI: 10.1037/int0000100.

Koponen T, Aro T, Peura P, Leskinen M, Viholainen H, Aro M. 2021. Benefits of Integrating an
Explicit Self-Efficacy Intervention With Calculation Strategy Training for Low-Performing
Elementary Students. *Frontiers in Psychology* 12:714379. DOI:
10.3389/fpsyg.2021.714379.

Kraiss JT, Ten Klooster PM, Chrispijn M, Trompetter HR, Stevens AWMM, Neutel E, Kupka
RW, Bohlmeijer ET. 2018. B-positive: a randomized controlled trial of a multicomponent
positive psychology intervention for euthymic patients with bipolar disorder - study protocol
and intervention development. *BMC Psychiatry* 18:335. DOI: 10.1186/s12888-018-1916-3.

Kulkarni A, Pring T, Ebbels S. 2014. Evaluating the effectiveness of therapy based around
Shape Coding to develop the use of regular past tense morphemes in two children with
language impairments. *Child Language Teaching and Therapy* 30:245–254. DOI:
10.1177/0265659013514982.

Kuschner ES, Morton HE, Maddox BB, De Marchena A, Anthony LG, Reaven J. 2017. The
BUFFET Program: Development of a Cognitive Behavioral Treatment for Selective Eating
in Youth with Autism Spectrum Disorder. *Clinical Child and Family Psychology Review*
20:403–421. DOI: 10.1007/s10567-017-0236-3.

Kwan RYC, Ng F, Lai M, Wong D, Chan S. 2023. The effects of Digital Buddy programme on
older adults' mental well-being: study protocol for a multi-centre, cluster randomized
controlled trial. *Trials* 24:95. DOI: 10.1186/s13063-023-07130-5.

La Greca AM, Ehrenreich-May J, Mufson L, Chan S. 2016. Preventing Adolescent Social
Anxiety and Depression and Reducing Peer Victimization: Intervention Development and
Open Trial. *Child & Youth Care Forum* 45:905–926. DOI: 10.1007/s10566-016-9363-0.

Lewin S, Hendry M, Chandler J, Oxman AD, Michie S, Shepperd S, Reeves BC, Tugwell P,
Hannes K, Rehfuess EA, Welch V, Mckenzie JE, Burford B, Petkovic J, Anderson LM,

Harris J, Noyes J. 2017. Assessing the complexity of interventions within systematic
reviews: development, content and use of a new tool (iCAT_SR). *BMC Medical Research*
Methodology 17. DOI: 10.1186/s12874-017-0349-x.

Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JPA, Clarke M,
Devereaux PJ, Kleijnen J, Moher D. 2009. The PRISMA statement for reporting systematic
reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and
elaboration. *BMJ* 151:W65–W94. DOI: 10.1136/bmj.b2700.

Lopez-Alcalde J, Yakoub N, Wolf M, Munder T, Von Elm E, Flückiger C, Steinert C, Liebherz
S, Rosendahl J, Witt CM, Barth J. 2022. The RIPI-f (Reporting Integrity of Psychological
Interventions delivered face-to-face) checklist was developed to guide reporting of treatment
integrity in face-to-face psychological interventions. *Journal of Clinical Epidemiology*
151:65–74. DOI: 10.1016/j.jclinepi.2022.07.013.

Losilla J-M, Oliveras I, Marin-Garcia JA, Vives J. 2018. Three risk of bias tools lead to opposite
conclusions in observational research synthesis. *Journal of Clinical Epidemiology* 101:61–
72. DOI: 10.1016/j.jclinepi.2018.05.021.

Lovett MW, Frijters JC, Steinbach KA, Sevcik RA, Morris RD. 2021. Effective intervention
for adolescents with reading disabilities: Combining reading and motivational remediation
to improve outcomes. *Journal of Educational Psychology* 113:656–689. DOI:
10.1037/edu0000639.

Maj A, Matynia M, Michalak N, Bis A, Andersson G. 2024. New in Town—An internet-based
self-efficacy intervention for internal migrants: A randomized controlled trial. *PLOS ONE*
19:e0299638. DOI: 10.1371/journal.pone.0299638.

Mannocci A, Saulle R, Colamesta V, D'Aguanno S, Giraldi G, Maffongelli E, Meggiolaro A,
Semyonov L, Unim B, La Torre G. 2014. What is the impact of reporting guidelines on
Public Health journals in Europe? The case of STROBE, CONSORT and PRISMA. *Journal*
of Public Health 37:737–740. DOI: 10.1093/pubmed/fdu108.

Matsuno E, Israel T. 2021. The Parent Support Program: Development and Acceptability of an
Online Intervention Aimed at Increasing Supportive Behaviors Among Parents of Trans
Youth. *Journal of GLBT Family Studies* 17:413–431. DOI:
10.1080/1550428X.2020.1868369.

Mesurado B, Distefano MJ, Robiolo G, Richaud MC. 2019. The Hero program: Development
and initial validation of an intervention program to promote prosocial behavior in
adolescents. *Journal of Social and Personal Relationships* 36:2566–2584. DOI:
10.1177/0265407518793224.

Michelon RC, Lucchese-Lobato F. 2024. A remote parent-led early intervention protocol to
promote motor development in infants with congenital heart disease: a feasibility pilot study
of a randomized clinical trial. *Developmental Neurorehabilitation* 27:134–144. DOI:
10.1080/17518423.2024.2365796.

Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W, Eccles MP, Cane J,
Wood CE. 2013. The Behavior Change Technique Taxonomy (v1) of 93 Hierarchically
Clustered Techniques: Building an International Consensus for the Reporting of Behavior
Change Interventions. *Annals of Behavioral Medicine* 46:81–95. DOI: 10.1007/s12160-013-
9486-6.

Moher D, Schulz KF, Altman DG. 2001. The CONSORT statement: revised recommendations
for improving the quality of reports of parallel-group randomised trials. *The Lancet*
357:1191–1194. DOI: 10.1016/S0140-6736(00)04337-3.

Möhler R, Köpke S, Meyer G. 2015. Criteria for Reporting the Development and Evaluation of
Complex Interventions in healthcare: revised guideline (CReDECI 2). *Trials* 16:204. DOI:
10.1186/s13063-015-0709-y.

Molek-Winiarska D, Kawka T. 2024. Reducing Work-Related Stress Through Soft-Skills
Training Intervention in the Mining Industry. *Human Factors: The Journal of the Human*
Factors and Ergonomics Society 66:1633–1649. DOI: 10.1177/00187208221139020.

Montgomery P, Grant S, Mayo-Wilson E, Macdonald G, Michie S, Hopewell S, Moher D. 2018.
Reporting randomised trials of social and psychological interventions: the CONSORT-SPI
2018 Extension. *Trials* 19:407. DOI: 10.1186/s13063-018-2733-1.

Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, Moore L, O'Cathain A, Tinati
T, Wight D, Baird J. 2015. Process evaluation of complex interventions: Medical Research
Council guidance. *BMJ* 350:h1258. DOI: 10.1136/bmj.h1258.

Moore GF, Evans R, Hawkins J, Littlecott H, Turley R. 2017. All interventions are complex,
but some are more complex than others: using iCAT_SR to assess complexity. *COCHRANE*
DATABASE OF SYSTEMATIC REVIEWS. DOI: 10.1002/14651858.ED000122.

Morgan JF, Lazarova S, Schelhase M, Saeidi S. 2014. Ten Session Body Image Therapy:
Efficacy of a Manualised Body Image Therapy. *European Eating Disorders Review* 22:66–
71. DOI: 10.1002/erv.2249.

Motlova V, Honsova P. 2021. The effects of a 13-week career development programme on
career-adapting thoughts and behaviours. *International Journal for Educational and*
Vocational Guidance 21:571–588. DOI: 10.1007/s10775-020-09454-z.

Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. 2018. Systematic review
or scoping review? Guidance for authors when choosing between a systematic or scoping
review approach. *BMC Medical Research Methodology* 18. DOI: 10.1186/s12874-018-
0611-x.

Newins AR, White SW. 2021. A Brief Computer-Based Sexual Assault Risk Reduction
Program: A Feasibility, Acceptability, and Preliminary Efficacy Pilot Study. *Journal of*
Interpersonal Violence 36:4854–4875. DOI: 10.1177/0886260518792974.

Nozawa K, Ishii A, Asaoka H, Iwanaga M, Kumakura Y, Oyabu Y, Shinozaki T, Imamura K,
Kawakami N, Miyamoto Y. 2022. Effectiveness of an Online Peer Gatekeeper Training
Program for Postsecondary Students on Suicide Prevention in Japan: Protocol for a
Randomized Controlled Trial. *JMIR Research Protocols* 11:e34832. DOI: 10.2196/34832.

O'Garo K-GN, Koenig HG. 2023. Spiritually Integrated Cognitive Processing Therapy for
Moral Injury in the Setting of PTSD: Initial Evidence of Therapeutic Efficacy. *Journal of*
Nervous & Mental Disease 211:656–663. DOI: 10.1097/NMD.0000000000001686.

Oldershaw A, Basra RS, Lavender T, Startup H. 2024. Specialist psychotherapy with emotion
for anorexia in Kent and Sussex: An intervention development and non-randomised single
arm feasibility trial. *European Eating Disorders Review* 32:215–229. DOI:
10.1002/erv.3034.

Olsson TM, Bergström M, Välikoski M, Karlsson M, Skoog T. 2024. Advancing Our
Understanding of Complex Intervention Development in Social Work: Co-Producing

Support for Youth Transitioning from Out-of-Home Care. *Child and Adolescent Social Work Journal*. DOI: 10.1007/s10560-024-00986-9.

Orduña-Malea E, Alonso-Arroyo A, Ontalba-Ruipérez J-A, Catalá-López F. 2023. Evaluating the online impact of reporting guidelines for randomised trial reports and protocols: a cross-sectional web-based data analysis of CONSORT and SPIRIT initiatives. *Scientometrics* 128:407–440. DOI: 10.1007/s11192-022-04542-z.

Otsuka T, Tanemura R, Noda K, Nagao T, Sakai H, Luo Z-W. 2015. Development of Computer-AidedCognitive Training Program for Elderly and Its Effectivenessthrougha 6 Months Group Intervention Study. *Current Alzheimer Research* 12:553–562. DOI: 10.2174/1567205012666150530202335.

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 372. DOI: 10.1136/bmj.n71.

Painter JM, Mote J, Peckham AD, Lee EH, Campellone TR, Pearlstein JG, Morgan S, Kring AM, Johnson SL, Moskowitz JT. 2019. A positive emotion regulation intervention for bipolar I disorder: Treatment development and initial outcomes. *General Hospital Psychiatry* 61:96–103. DOI: 10.1016/j.genhosppsych.2019.07.013.

Pakarinen A, Flemmich M, Parisod H, Selänne L, Hamari L, Aromaa M, Leppänen V, Löyttyniemi E, Smed J, Salanterä S. 2018. Protocol for digital intervention for effective health promotion of small children—A cluster randomized trial. *Journal of Advanced Nursing* 74:1685–1699. DOI: 10.1111/jan.13561.

Passoni S, Curinga T, Toraldo A, Berlingeri M, Fernandez I, Bottini G. 2018. Eye Movement Desensitization and Reprocessing Integrative Group Treatment Protocol (EMDR-IGTP) Applied to Caregivers of Patients With Dementia. *Frontiers in Psychology* 9:967. DOI: 10.3389/fpsyg.2018.00967.

Petticrew M. 2011. When are complex interventions "complex"? When are simple interventions "simple"? *The European Journal of Public Health* 21:397–398. DOI: 10.1093/eurpub/ckr084.

Petticrew M, Anderson L, Elder R, Grimshaw J, Hopkins D, Hahn R, Krause L, Kristjansson E, Mercer S, Sipe T, Tugwell P, Ueffing E, Waters E, Welch V. 2015. Complex interventions and their implications for systematic reviews: A pragmatic approach. *International Journal of Nursing Studies* 52:1211–1216. DOI: 10.1016/j.ijnurstu.2015.01.004.

Pinnock H, Barwick M, Carpenter CR, Eldridge S, Grandes G, Griffiths CJ, Rycroft-Malone J, Meissner P, Murray E, Patel A, Sheikh A, Taylor SJC. 2017. Standards for Reporting Implementation Studies (StaRI) Statement. *BMJ*:i6795. DOI: 10.1136/bmj.i6795.

Pinnock H, Epiphaniou E, Sheikh A, Griffiths C, Eldridge S, Craig P, Taylor SJ. 2015. Developing standards for reporting implementation studies of complex interventions (StaRI): a systematic review and e-Delphi. *Implementation Science* 10. DOI: 10.1186/s13012-015-0235-z.

Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R, Hensley M. 2011. Outcomes for Implementation Research: Conceptual Distinctions, Measurement Challenges, and Research Agenda. *Administration and Policy in Mental Health and Mental Health Services Research* 38:65–76. DOI: 10.1007/s10488-010-0319-7.

Ramirez MW, Woodworth CA, Evans WR, Grace GA, Schobitz RP, Villarreal SA, Howells CJ,
Gissendanner RD, Katko NJ, Jones KN, McCabe AE, Terrell DJ. 2021. A trauma-focused
intensive outpatient program integrating elements of exposure therapy with acceptance and
commitment therapy: Program development and initial outcomes. *Journal of Contextual*
Behavioral Science 21:66–72. DOI: 10.1016/j.jcbs.2021.06.002.

Rodrigo D, Foguet-Boreu Q, Juvanteny EP, Izquierdo Munuera E. 2022. Effectiveness of a
psychoeducational group intervention developed by primary care nurses on symptom control
of pediatric patients with ADHD. ADHD parent study. *Health Psychology and Behavioral*
Medicine 10:1176–1189. DOI: 10.1080/21642850.2022.2148672.

Rosengaard LO, Andersen MZ, Rosenberg J, Fonnes S. 2024. Five aspects of research waste in
biomedicine: A scoping review. *Journal of Evidence-Based Medicine* 17:351–359. DOI:
10.1111/jebm.12616.

Rutter H, Savona N, Glonti K, Bibby J, Cummins S, Finegood DT, Greaves F, Harper L, Hawe
P, Moore L, Petticrew M, Rehfuess E, Shiell A, Thomas J, White M. 2017. The need for a
complex systems model of evidence for public health. *The Lancet* 390:2602–2604. DOI:
10.1016/S0140-6736(17)31267-9.

Saffaran P, Oh P, Reitav J. 2022. Evaluating the Feasibility and Efficacy of A Novel CBTi/SMT
Treatment Protocol for Cardiac Rehab Patients: A Non-Randomized Pilot Trial. *Behavioral*
Sleep Medicine 20:716–731. DOI: 10.1080/15402002.2021.1993227.

Saulnier KG, Koscinski B, Flynt S, Accorso C, Allan NP. 2024. Brief observable anxiety
sensitivity treatment: intervention development and a pilot randomized-controlled
acceptability and feasibility trial to evaluate a brief intervention for anxiety sensitivity social
concerns. *Cognitive Behaviour Therapy* 53:190–206. DOI:
10.1080/16506073.2023.2288551.

Schneider J, Akhtar A, Boycott N, Guo B, Latimer E, Cao Z, McMurran M. 2016. Individual
placement and support versus individual placement and support enhanced with work-
focused cognitive behaviour therapy: Feasibility study for a randomised controlled trial.
British Journal of Occupational Therapy 79:257–269. DOI: 10.1177/0308022615619184.

Schulz KF, Altman DG, Moher D, for the CONSORT Group. 2010. CONSORT 2010
Statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 340. DOI:
10.1136/bmj.c332.

Sherwood A, Paynter J, Emerson L-M. 2023. A brief online mindful parenting program:
Feasibility and initial effects pilot in a community sample. *Journal of Child and Family*
Studies 32:1532–1545. DOI: 10.1007/s10826-023-02571-7.

Skivington K, Matthews L, Simpson SA, Craig P, Baird J, Blazeby JM, Boyd KA, Craig N,
French DP, McIntosh E, Petticrew M, Rycroft-Malone J, White M, Moore L. 2021. A new
framework for developing and evaluating complex interventions: update of Medical
Research Council guidance. *BMJ* 374. DOI: 10.1136/bmj.n2061.

Tanner-Smith EE, Grant S. 2018. Meta-analysis of complex interventions. *Annual Review of*
Public Health 39:135–151. DOI: 10.1146/annurev-publhealth-040617-014112.

Tran T, Nguyen HT, Shochet I, Wurfl A, Orr J, Nguyen N, La N, Nguyen H, Stocker R, Nguyen
T, Le M, Fisher J. 2020. School-based, two-arm, parallel, controlled trial of a culturally
adapted resilience intervention to improve adolescent mental health in Vietnam: study
protocol. *BMJ Open* 10:e039343. DOI: 10.1136/bmjopen-2020-039343.

Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MDJ,
Horsley T, Weeks L, Hempel S, Akl EA, Chang C, McGowan J, Stewart L, Hartling L,
Aldcroft A, Wilson MG, Garritty C, Lewin S, Godfrey CM, Macdonald MT, Langlois EV,
Soares-Weiser K, Moriarty J, Clifford T, Tunçalp Ö, Straus SE. 2018. PRISMA Extension
for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Annals of Internal*
Medicine 169:467–473. DOI: 10.7326/M18-0850.

Turner L, Shamseer L, Altman DG, Schulz KF, Moher D. 2012. Does use of the CONSORT
Statement impact the completeness of reporting of randomised controlled trials published in
medical journals? A Cochrane reviewa. *Systematic Reviews* 1:60. DOI: 10.1186/2046-4053-
1-60.

Vandenbroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, Poole C,
Schlesselman JJ, Egger M. 2007. Strengthening the Reporting of Observational Studies in
Epidemiology (STROBE). *Epidemiology* 18:805–835. DOI:
10.1097/EDE.0b013e3181577511.

Vives J, Lorente S, Casellas A, López-González E, Marín-García JA, Molla C, Rodrigo MF,
Viguer P, Losilla J-M. 2025a. Reporting quality in psychological interventions. Implications
for cumulative science and evidence-based practice: A scoping review. DOI:
10.64898/2025.11.26.25341096.

Vives J, Lorente S, Casellas-Grau A, López-González E, Marin-Garcia JA, Molla C, Rodrigo
MF, Viguer P, Losilla J-M. 2025b. Reporting quality in psychological interventions.
Implications for cumulative science and evidence-based practice. A scoping review protocol.
INPLASY - International Platform of Registered Systematic Review and Meta-analysis
Protocols. DOI: 10.37766/inplasy2025.7.0098.

Walker MR, Kallingal GJS, Musser JE, Folen R, Stetz MC, Clark JY. 2014. Treatment Efficacy
of Virtual Reality Distraction in the Reduction of Pain and Anxiety During Cystoscopy.
Military Medicine 179:891–896. DOI: 10.7205/MILMED-D-13-00343.

Ward KP, Wood DS, Young TM. 2020. Retreat Intervention Effectiveness for Female Survivors
of Child Sexual Abuse. *Research on Social Work Practice* 30:760–769. DOI:
10.1177/1049731520921936.

Wasik BA, Hindman AH. 2020. Increasing preschoolers' vocabulary development through a
streamlined teacher professional development intervention. *Early Childhood Research*
Quarterly 50:101–113. DOI: 10.1016/j.ecresq.2018.11.001.

Weigel A, Gumz A, Uhlenbusch N, Wegscheider K, Romer G, Löwe B. 2015. Preventing eating
disorders with an interactive gender-adapted intervention program in schools: Study protocol
of a randomized controlled trial. *BMC Psychiatry* 15:21. DOI: 10.1186/s12888-015-0405-1.

Zajonz P, Lautenbach F, Laborde S. 2024. Emotional intelligence training for sports coaches –
evaluation of a time-effective online intervention and its effects on coaching efficacy.
Journal of Applied Sport Psychology 36:722–739. DOI: 10.1080/10413200.2023.2296911.

Zhang M, Chan SW, You L, Wen Y, Peng L, Liu W, Zheng M. 2014. The effectiveness of a self-
efficacy-enhancing intervention for Chinese patients with colorectal cancer: A randomized
controlled trial with 6-month follow up. *International Journal of Nursing Studies* 51:1083–
1092. DOI: 10.1016/j.ijnurstu.2013.12.005.

1087

APPENDIX A. Search strategy.

Public health

PsycInfo (2609)

(((((title(("intervention*" OR "treatment*" OR "therap*" OR "program*") NEAR/5 ("develop*" OR "protocol" OR "feasibility" OR "effica*" OR "effectiv*")) NOT title("review*" OR "meta-anal*")) AND PEER(yes))) AND pd(20140101-20241231)) AND ccl.exact("Health & Mental Health Treatment & Prevention" OR "Promotion & Maintenance of Health & Wellness" OR "Health & Mental Health Services" OR "Health Psychology & Medicine" OR "Home Care & Hospice" OR "Sports & Exercise" OR "Childrearing & Child Care" OR "Nursing Homes & Residential Care" OR "Health & Mental Health Personnel Issues" OR "Aging & Older Adult Development")

Web of Science (677)

(((((TI=(("intervention*" OR treatment* OR therap* OR program*) NEAR/5 (develop* OR protocol OR feasibility OR effica* OR effectiv*))) NOT TI=(review* OR meta-anal*)) AND DOP=(2014-01-01/2024-12-31)) AND DT=(Article)) AND (TASCA==("PSYCHIATRY" OR "PSYCHOLOGY") AND TASCA==("SUBSTANCE ABUSE" OR "GERIATRICS GERONTOLOGY" OR "NURSING" OR "PEDIATRICS" OR "GERONTOLOGY" OR "HEALTH CARE SCIENCES SERVICES" OR "PUBLIC ENVIRONMENTAL OCCUPATIONAL HEALTH" OR "HEALTH POLICY SERVICES" OR "NUTRITION DIETETICS" OR "FAMILY STUDIES" OR "ONCOLOGY" OR "REHABILITATION" OR "HOSPITALITY LEISURE SPORT TOURISM" OR "SPORT SCIENCES" OR "SURGERY" OR "IMMUNOLOGY" OR "OBSTETRICS GYNECOLOGY"))

Clinical

PsycInfo (3242)

(((((title(("intervention*" OR "treatment*" OR "therap*" OR "program*") NEAR/5 ("develop*" OR "protocol" OR "feasibility" OR "effica*" OR "effectiv*")) NOT title("review*" OR "meta-anal*")) AND PEER(yes))) AND pd(20140101-20241231)) AND ccl.exact("Cognitive Therapy" OR "Psychotherapy & Psychotherapeutic Counseling" OR "Specialized Interventions" OR "Behavior Therapy & Behavior Modification" OR "Affective Disorders" OR "Neurodevelopmental & Autism Spectrum Disorders" OR "Anxiety Disorders" OR "Behavior Disorders & Antisocial Behavior" OR "Behavioral & Psychological Treatment of Physical Illness" OR "Psychological & Physical Disorders" OR "Schizophrenia & Psychotic States" OR "Psychological Disorders" OR "Eating Disorders" OR "Speech & Language Therapy" OR "Interpersonal & Client Centered & Humanistic Therapy" OR "Personality Disorders" OR "Psychoanalytic Therapy")

Web of Science (1788)

(((((TI=(("intervention*" OR treatment* OR therap* OR program*) NEAR/5 (develop* OR protocol OR feasibility OR effica* OR effectiv*))) NOT TI=(review* OR meta-anal*)) AND DOP=(2014-01-01/2024-12-31)) AND DT=(Article)) AND

1127 (TASCA=="PSYCHOLOGY CLINICAL") OR (TASCA=="PSYCHIATRY" OR
1128 "PSYCHOLOGY") AND TASCA=="CLINICAL NEUROLOGY" OR "PHARMACOLOGY
1129 PHARMACY" OR "PSYCHOLOGY PSYCHOANALYSIS" OR "GENETICS
1130 HEREDITY")))

Social and organizational

PsycInfo (865)

(((((title(("intervention*" OR "treatment*" OR "therap*" OR "program*") NEAR/5 ("develop*"
OR "protocol" OR "feasibility" OR "effica*" OR "effectiv*")) NOT title("review*" OR "meta-
anal*")) AND PEER(yes))) AND pd(20140101-20241231)) AND ccl.exact("Community &
Social Services" OR "Professional Personnel Attitudes & Characteristics" OR "Behavior
Disorders & Antisocial Behavior" OR "Occupational & Vocational Rehabilitation" OR "Social
Processes & Social Issues" OR "Organizational Psychology & Human Resources" OR
"Criminal Behavior & Juvenile Delinquency" OR "Personnel Management & Selection &
Training" OR "Working Conditions & Industrial Safety" OR "Management & Management
Training" OR "Organizational Behavior" OR "Personnel Attitudes & Job Satisfaction" OR
"Social Psychology" OR "Mass Media Communications" OR "Marriage & Family" OR
"Occupational Interests & Guidance" OR "Police & Legal Personnel" OR "Consumer Attitudes
& Behavior")

Web of Science (1098)

(((((TI=(("intervention*" OR treatment* OR therap* OR program*) NEAR/5 (develop* OR
protocol OR feasibility OR effica* OR effectiv*))) NOT TI=(review* OR meta-anal*)) AND
DOP=(2014-01-01/2024-12-31)) AND DT=(Article)) AND (TASCA=="SOCIAL SCIENCES
INTERDISCIPLINARY" OR "SOCIAL WORK" OR "SOCIAL SCIENCES BIOMEDICAL"
OR "SOCIAL ISSUES" OR "SOCIAL SCIENCES MATHEMATICAL METHODS" OR
"MANAGEMENT" OR "CRIMINOLOGY PENOLOGY"))

Education

PsycINFO (1489)

(((((title(("intervention*" OR "treatment*" OR "therap*" OR "program*") NEAR/5 ("develop*"
OR "protocol" OR "feasibility" OR "effica*" OR "effectiv*")) NOT title("review*" OR "meta-
anal*")) AND PEER(yes))) AND pd(20140101-20241231)) AND ccl.exact("Curriculum &
Programs & Teaching Methods" OR "Professional Education & Training" OR "Educational
Administration & Personnel" OR "Educational/Vocational Counseling & Student Services" OR
"Educational & School Psychology" OR "Special & Compensatory Education" OR "Classroom
Dynamics & Student Adjustment & Attitudes" OR "Developmental Psychology")

Web of Science (589)

(((((TI=(("intervention*" OR treatment* OR therap* OR program*) NEAR/5 (develop* OR
protocol OR feasibility OR effica* OR effectiv*))) NOT TI=(review* OR meta-anal*)) AND
DOP=(2014-01-01/2024-12-31)) AND DT=(Article)) AND

(TASCA=="PSYCHOLOGY DEVELOPMENTAL") OR (TASCA=="PSYCHIATRY" OR
"PSYCHOLOGY") AND TASCA=="EDUCATION SPECIAL" OR "EDUCATION
EDUCATIONAL RESEARCH")))